

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

"Off-axis" Mn-Mn bond in Mn₂(CO)₁₀ at high pressurePiero Macchi,^{a*} Nicola Casati,^{b,c} Shaun R. Evans,^{a,b} Fabia Gozzo,^{b,d} Petra Simoncic,^{a,b} and Davide Tiana^{a,e}

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

5 At variance from what previously reported, Mn₂(CO)₁₀ does not transform its conformation from *staggered* into *eclipsed* in the high pressure crystal form. X-ray powder diffraction, DFT calculations and Raman spectroscopy show that the *staggered* conformation is retained. Instead, a rotation and a translation of the Mn(CO)₅ pyramidal units produce an "off-axis" and rather shorter Mn-Mn bond.


Investigating the high pressure forms of *molecular* crystals is a relatively new field, as for many years the interest of solid state chemists and physicists was focused only on harder inorganic materials. Few studies are known before 1980,¹ whereas during the past decade we could witness an increasing curiosity for soft materials, in the range 0-10GPa, with the intent to produce new polymorphs, to modify the material properties or the intra- and intermolecular bonding. Some of these studies focused on transition metal carbonyl compounds, such as M(CO)_n complexes² or poly-metal species M_m(CO)_n.³

The current knowledge on transition metal dimers is that high pressure should favour the smaller molecular volume (hence the lower enthalpy) of an *eclipsed* conformation, in spite of the lower electronic energy of the *staggered* one, which takes advantage of smaller repulsion between the equatorial carbonyls. Adams *et al.*⁴ investigated the high pressure forms of transition metal carbonyl dimers M₂(CO)₁₀ (M = Mn, Re) with Raman and IR spectroscopy. In Mn₂(CO)₁₀ (**1**), they found a transition from *staggered* to *eclipsed* conformation (Figure 1a, 1b) at 0.8 GPa, mainly justified by the reduction of the Raman C-O stretching frequencies due to the higher symmetry. Accurate crystal structures or at least lattice parameters could not be obtained from single crystal X-ray diffraction (XRSC), because of the very fragile nature of the samples. XRSC studies and periodic density functional theory (P-DFT) calculations on other dimers, like Co₂(CO)₆(XPh₃)₂, (Ph = phenyl; X = P, As, **2a** and **2b**),⁵ revealed a change from *staggered* to almost *eclipsed* conformation within a similar pressure range, following a second order solid state phase transition. The interpretation of Raman spectra of M₂(CO)₁₀,⁴ instead, suggested a sudden transformation from D_{4d} to D_{4h} molecular symmetry. Therefore, one would anticipate a first order phase transition, implying discontinuous lattices.

At ambient pressure, **1** crystallizes in the *monoclinic* space group type C2/c, (reported by many authors in the non-standard I2/a). The molecular symmetry is just C₂, apparently close to the gas phase D_{4d} with a root mean square deviation of 0.19Å. We tentatively carried out X-ray diffraction on single crystals in a

diamond anvil cell (DAC), but we could only confirm the fragility of the samples. Nevertheless, two alternative investigations could be carried out: a) P-DFT simulations using the B3PW⁶ functional, corrected for dispersion effects,⁷ a 6-21G(d) basis set for C and O, and a small core effective core potential basis⁸ for Mn; b) x-ray powder diffraction (XRPD) in a gas membrane DAC, with synchrotron source at the powder diffraction station of the Paul Scherer Institute.

P-DFT calculations were carried out using CRYSTAL09,⁹ optimizing the geometry up to 15 GPa, by enthalpy minimization. Temperature effects were not included. In the range 0-3 GPa, we started from larger unit cells and lower symmetry but the ambient condition *space group type* was always calculated as the thermodynamically stable form, of course with a smaller volume produced by a rather anisotropic compression (Table 1 and Supporting Information). Thus, a phase change is not predicted by theory. Moreover, the *staggered* conformation is retained and no stationary point could be located for the *eclipsed* geometry, which rules out a conformational change without phase transition.



70 **Figure 1.** a) the stable gas phase *staggered* conformation of Mn₂(CO)₁₀; b) the gas phase *eclipsed* conformation (proposed for the high pressure solid state form); c) *off-axis* conformation (Mn-Mn and Mn-C_{axial} bonds are not aligned) from P-DFT predictions and XRPD at high pressure.

Two interesting geometrical changes are however calculated. One is a large contraction of the Mn-Mn bond. This is not surprising given the small force constant of this bond, but it would not occur in an *eclipsed* conformation because the repulsion between the equatorial carbonyls would force the two metals further apart, as it occurs in **2**.⁵ The other feature is a small rotation of the Mn(CO)₅ moieties about an axis *perpendicular* to Mn-Mn (and parallel to the crystallographic twofold). This rotation is different from that expected for a *staggered* → *eclipsed* transformation. Thus, the Mn-Mn bond becomes "off-axis", *i.e.* no longer aligned

with the main inertial axis of the molecule. This is not only due to the external pressure, because the feature is already evident at ambient pressure. In fact, while in the gas phase D_{4d} structure, Mn-Mn, Mn- C_{ax} and C_{ax} -O $_{ax}$ bonds are co-axial (Figure 1a), in the crystal the molecule undergoes some necessary distortions, given that D_{4d} is not a crystallographic symmetry. One of these has not been commented so far, despite many studies on **1**, including electron density determinations:¹⁰ the C_{ax} -Mn-Mn bond angle is significantly different from 180° (see Table 2). Albeit small, this feature is important and even enhanced at high pressure: at 3 GPa, P-DFT predicts C_{ax} -Mn-Mn = 170° and the distortion is quite evident (Figure 1c). Above 5 GPa, the angle returns larger (173°), but Mn-Mn still decreases. In $M_2(CO)_n$, M-M and M- C_{ax} bonds are rarely constrained on the same axis by crystallographic symmetries, few exceptions are the K^+ salt of $[Cr_2(CO)_{10}]^{2-}$, where O, C and Cr atoms lie on a twofold axis,¹¹ or the species **2** where they lie on a threefold axis. $[FeCr(CO)_9]^{2-}$, in its PPN⁺ salt,¹² is characterized by the most severe distortion (Fe-Co-C $\sim 169^\circ$). In neutral molecules, the rotation is smaller (M-M-C $> 175^\circ$) as for example in **1** (Table 2). Noteworthy, it is quite temperature dependent: in fact, C_{ax} -Mn-Mn constantly decreases from $177.0(3)^{10b}$ at 298 K to $175.3(1)$ at 74 K.^{10e}

To prove the correctness of the theoretical predictions, we carried out high pressure powder diffraction on **1**. The species undergoes a relatively rapid decomposition under the high flux of X-ray photons, necessary to obtain significant diffraction from the very few particles loaded in the DAC. At 17 KeV a sample could not survive more than 20 minutes of X-ray dose, sufficient to collect a few pressure points. At higher energy, the radiation damage was smaller. The data acquisition was quite rapid thanks to the Mythen detector.¹³ Three sets of XRPD patterns were collected: 1) 17 KeV, 0-3.5 GPa, NaCl as pressure calibrant and 4:1 methanol/ethanol as pressure transmission medium (PTM); 2) 20 KeV, 0-7.0 GPa, quartz as calibrant, 4:1 methanol/ethanol as PTM; 3) 28 KeV, 0-3.5 GPa quartz as calibrant and Daphne oil¹⁴ as PTM. After the experiments, we observed sample darkening, as previously reported,⁴ likely due to partial decomposition. The unit cells predicted by P-DFT were used to initially index the peaks, although theoretical cells are systematically smaller because carried out at 0 K. In fact, at ambient pressure a rather large volume contraction (ca. 9%) is observed when cooling from 298 K to 77 K (Table 1), thus the predicted unit cell is closer to the experimental cell at low T. The agreement between theory and experiment is better at high pressure, where the temperature has smaller influence (see S. I.). Anyway, no phase transition was observed in all data collected, proved by the I2/a phase well explaining the positions of the observed peaks. Rietveld refinements¹⁵ were carried out using TOPAS¹⁶. The $Mn(CO)_5$ unit was treated as a rigid body, allowing the rotation about the three inertial axes and translation. The results confirmed the predictions: the molecule does not modify its conformation about the Mn-Mn bond, but a substantial Mn-Mn- C_{ax} bending occurs, as the refined angle at 3.25 GPa drops to ca. 170° and the basal planes of the two pyramidal $Mn(CO)_5$ moieties are quite inclined (ϵ values in Table 2) whereas they are parallel by symmetry in the gas phase D_{4d} structure. Tentative refinements starting from the *eclipsed* conformer always converged to the *staggered* one. Although the accuracy of the refinements in DAC is small, the

trend appears clear and in agreement with the calculations.

Because of the discrepancy with respect to previous results, we collected new Raman spectra on a powder sample, up to 3.0 GPa. The ambient pressure spectrum is quite similar to that previously reported by Adams *et al.*, but above 0.7 GPa there is a huge difference in the carbonyl region (1950 - 2050 cm^{-1}). In particular, no reduction of the number of bands is observed. This is in agreement with an unmodified, low molecular symmetry, in keeping with the P-DFT predicted spectra, after re-scaling the frequencies to account for anharmonic terms (see Supporting Information). The discrepancy with Adams *et al.*⁴ could be justified by the scarce resolution of their spectra or a non-hydrostatic pressure inside the DAC that caused rather broad peaks in the region 1975 - 2030 cm^{-1} . In fact, at 2.6-2.7 GPa the FWHM is above 10 cm^{-1} in ref. 4c, but much smaller in our spectra ($< 5\text{ cm}^{-1}$). On the other hand, there is a perfect agreement for the highest energy vibration, which is quite pressure dependent and isolated (therefore easily identified): this A_{1g} mode occurs at 2115 cm^{-1} at ambient condition and at 2125 cm^{-1} at 2.6 GPa, confirming a gradient $\Delta\nu/\Delta P \sim 4\text{ cm}^{-1}/\text{GPa}$ (or $3\text{ cm}^{-1}/\text{GPa}$ from P-DFT calculations).




Figure 2 Raman spectra at 0.0001 (top), 1.6 (centre) and 2.6 GPa (bottom). In Fig. 2 of ref. 4c, only three broad peaks are visible above 1 GPa in the region 1950 - 2050 cm^{-1} . Notably, the number of predicted bands in the region 1950 - 2050 is even larger (see S.I.).

Table 1 Unit cell parameters of **1** at variable P,T conditions. More data points in S.I. For experimental data, *esd*'s are smaller than the last digit.

T (K)	P(GPa)	a(Å)	b(Å)	c(Å)	β (°)	V(Å ³)	Ref.
298	10^{-4}	14.129	7.102	14.625	105.19	1417	XRPD ^a
298	10^{-4}	14.121	7.100	14.619	105.18	1414	XRPD ^b
298	1.55	13.816	6.647	13.735	105.00	1218	XRPD ^b
298	3.25	13.528	6.452	13.293	104.84	1121	XRPD ^b
298	6.60	13.140	6.269	12.889	104.50	1028	XRPD ^b
0	10^{-4}	14.063	6.654	13.908	108.78	1232	P-DFT
0	1.0	13.864	6.623	13.344	109.04	1158	P-DFT
0	3.0	13.338	6.493	13.016	106.97	1078	P-DFT
0	5.0	13.037	6.375	12.802	105.98	1023	P-DFT
0	10.0	12.382	6.219	12.496	103.54	935	P-DFT
298	10^{-4}	14.16	7.11	14.67	105.0	1426	XRSC10 ^a
298	10^{-4}	14.135	7.010	14.628	105.2	1416	XRSC10 ^b
120°	10^{-4}	14.110	6.898	14.326	104.99	1347	XRSC10 ^c
100	10^{-4}	14.126	6.880	14.312	105.08	1343	XRSC10 ^d
74	10^{-4}	14.088	6.850	14.242	105.08	1327	XRSC10 ^e

a) glass capillary 17KeV radiation; b) gas membrane DAC 20 keV, methanol/ethanol as PTM; c) transformed in I2/a from the original data.

These results prove that pressure produces two alternative effects in $M_2(CO)_n$ species: 1) a *staggered* \rightarrow *eclipsed* transformation:

the smaller molecular volume compensates the repulsion between carbonyls and the consequently longer M-M distance (it occurs in **2**, with 6 equatorial carbonyls); 2) an approach and a slide of the $M(CO)_n$ pyramids along a OC-M-CO basal bisection together with a rotation of $M(CO)_n$ units about an axis perpendicular to M-M (it occurs in **1**, where 8 equatorial carbonyls hamper the eclipsing mechanism).

Table 2 Relevant geometrical parameters of $Mn_2(CO)_{10}$ at different P, T after refinements or PDFPT optimizations. Distances are in Å; angles and torsions in °. *esd*'s are smaller than the last digit.

T(K)	P(GPa)	Mn-Mn	Mn-Mn-C _{ax}	C-Mn-Mn-C ^a	ε ^c	Ref.
298	10 ⁻⁴	2.92	178	38	0.8	XRPD ^b
298	1.55	2.72	168	31	1.4	XRPD ^b
298	3.25	2.79	170	37	7.8	XRPD ^b
298	6.60	2.71	166	36	12.6	XRPD ^b
0	10 ⁻⁴	2.856	173.8	38.1	7.3	P-DFT
0	1.0	2.824	170.4	36.7	8.8	P-DFT
0	3.0	2.776	170.3	36.3	8.2	P-DFT
0	5.0	2.755	170.4	36.3	8.2	P-DFT
0	10.0	2.702	173.0	36.6	7.3	P-DFT
298	10 ⁻⁴	2.92	177	38.9	n.a.	XRSC ^{10a}
298	10 ⁻⁴	2.904	177.0	38.9	3.3	XRSC ^{10b}
120	10 ⁻⁴	2.904	175.9	37.6	4.6	XRSC ^{10c}
100	10 ⁻⁴	2.903	175.6	37.5	4.9	XRSC ^{10d}
74	10 ⁻⁴	2.895	175.3	37.3	5.0	XRSC ^{10e}

a) smallest torsion angle; b) DAC, 20 keV radiation, methanol/ethanol transmission medium; c) angle between the basal planes of the two $Mn(CO)_5$ pyramidal moieties (r.m.s. planes of the equatorial Oxygens).

To complete our study, we investigated the chemical bonding in **1** as a function of the pressure, using the Interacting Quantum Atom^{17a} analysis. The M-M interaction is associated with a small destabilization,^{17b} a balance between a stabilizing exchange interaction (the covalent bond) and a destabilizing Coulomb interaction (the repulsion between metals that, although formally zerovalent, are positively charged due to π -back donation). Upon compression, the exchange is stronger, the electron delocalization between the metals raises and the metal charges are smaller. This explains the significant Mn-Mn shortening. As known,¹⁸ even for unsupported M-M bonds the equatorial carbonyls are necessary to bind the two $M(CO)_n$ moieties through 1,3-M---CO interactions. In **1**, the equatorial carbonyls are more asymmetric at high pressure and one of the 1,3-M---CO interactions is particularly stabilizing. An *eclipsed* conformation would not take advantage of these two favourable effects (stronger 1,3-M---CO stabilization and smaller M-M destabilization) while having more unfavourable CO---CO contacts compared to $M_2(CO)_8$ species. We have reported on a comprehensive structural study of the high pressure form of $Mn_2(CO)_{10}$, known for more than 50 years, but still attracting interest for its intriguing bonding features. Contrary to previously published,⁴ an eclipsing of the equatorial carbonyls was not observed. Instead, a translation of the $Mn(CO)_5$ squared pyramids occurs coupled with a rotation about a direction perpendicular to Mn-Mn bond, producing a shorter and "off-axis" Mn-Mn bond. The bonding analysis confirms that this mechanism is convenient because of a better stabilization of the Mn-Mn bond and a smaller repulsion of the equatorial carbonyls compared to the eclipsed conformation. The results stimulate a re-investigation¹⁹ of $Re_2(CO)_{10}$, for which

staggered \rightarrow *eclipsed* and *eclipsed* \rightarrow *staggered* transformations were anticipated from vibrational spectroscopy at high pressure.^{4c} We thank the Swiss National Science Foundation (project 144534) and PSI for financial support. Ms. A. Lanza and Dr. M. Fisch are thanked for their assistance during experiments.

Notes and references

- ^a Department of Chemistry and Biochemistry, University of Bern, Freiestrasse 3, 3012 Bern, Switzerland. E-mail: piero.macchi@dcb.unibe.ch
^b Paul Scherrer Institute, Swiss Light Source, 5232 Villigen PSI, Switzerland
^c Diamond Light Source Ltd., Harwell Science and Innovation Campus, Didcot, Oxfordshire, OX11 0DE, UK
^d Excelsus Structural Solutions SPRL, B-1150 Brussels, Belgium
^e Department of Chemistry University of Bath, United Kingdom
- [†] Electronic Supplementary Information (ESI) available: details of the data collections and the calculations; calculated fractional coordinates. See DOI: 10.1039/b000000x/.
- G. J. Piermarini, A. D. Mighell, C. E. Weir and S. Block. *Science*, 1969, **165**, 1250.
 - S. V. Garimella, V. Drozd and A. Durygin *Chem. Phys. Lett.*, 2008, **454**, 242.
 - C. Slebodnick, J. Zhao, R. Angel, B. E. Hanson, Y. Song, Z. Liu and R. J. Hemley *Inorg. Chem.*, 2004, **43**, 5245.
 - a) D. M. Adams, P. D. Hatton, A. C. Shaw and T. K. J. Tan. *J. Chem. Soc. Chem. Comm.* 1981, 226; b) D. M. Adams and I. O. C. Ekejiuba *J. Chem. Phys.*, 1983, **78**, 5408; c) D. M. Adams, P. D. Hatton and A. C. Shaw *J. Phys.: Cond. Matter* 1991, **3**, 6145.
 - a) N. Casati, P. Macchi and A. Sironi *Chemistry, Eur. J.*, 2009, **15**, 4446; b) N. Casati, P. Macchi and A. Sironi *Angew. Chem.* 2005, **44**, 7736.
 - a) A. D. Becke *J. Chem. Phys.* 1993, **98**, 5648; b) J. P. Perdew *Electronic structure of solids*, Akademie Verlag, Berlin, 1991.
 - S. Grimme *J. Comput. Chem* 2006, **27**, 1787.
 - P. J. Hay and W. R. Wadt *J. Chem. Phys.* 1985, **82**, 270.
 - R. Dovesi, V.R. Saunders, C. Roetti, R. Orlando, C. M. Zicovich-Wilson, F. Pascale, B. Civaleri, K. Doll, N.M. Harrison, I.J. Bush, Ph. D'Arco, and M. Llunell CRYSTAL09 User's Manual, University of Torino, Torino, 2009.
 - a) L. F. Dahl and R. E. Rundle *Acta Cryst.*, 1963, **16**, 419; b) M. R. Churchill, K. N. Amoh and H. J. Wasserman *Inorg. Chem.*, 1981, **20**, 1609; c) R. Bianchi, G. Gervasio and D. Marabello *Chem. Comm.*, 1998, 1535; d) L. J. Farrugia, P. R. Mallinson and B. Stewart *Acta Cryst.*, 2003, **B59**, 234; e) M. Martin, B. Rees, and A. Mitschler *Acta Cryst.*, 1982, **B38**, 6; f) R. Bianchi, G. Gervasio and D. Marabello *Inorg. Chem.* 2000, 39, 2360.
 - E. Hey-Hawkins and H. G. Schneringer *Chem. Ber.*, 1991, **124**, 1167.
 - B. Balbach, S. Baral, H. Biersack, W. A. Herrmann, J. A. Labinger, W. R. Scheidt, F. J. Timmers and M. L. Ziegler *Organometallics*, 1988, **7**, 325.
 - A. Bergamaschi, A. Cervellino, R. Dinapoli, F. Gozzo, B. Henrich, I. Johnson, P. Philipp, A. Mozzanica, B. Schmitt and X.T. Xintian *J. Sync. Rad.* 2010, **17**, 653.
 - K. Yokogawa, K. Murata, H. Yoshino and S. Aoyama *J. Phys. Soc. Japan* 2007, **46** 3636.
 - H. M. Rietveld *J. Appl. Cryst.* 1968, **2**, 65.
 - A. Coelho. TOPAS-Academic, Coelho Software, Brisbane.
 - a) M. A. Blanco, A. Martín Pendás, and E. Francisco *J. Chem. Theory Comput.* 2005, **1**, 1096; b) D. Tiana, E. Francisco, P. Macchi, A. Sironi and A. Martín Pendás *to be submitted*.
 - a) P. Macchi, D. M. Proserpio and A. Sironi, *J. Am. Chem. Soc.* 1998, **120**, 13429; b) P. Macchi, L. Garlaschelli, and A. Sironi. *J. Am. Chem. Soc.* 2002, **124**, 141173; c) P. Macchi and A. Sironi. *Coord. Chem. Rev.* 2003, **238-239**, 383; d) P. Macchi, D. Donghi, and A. Sironi, *A. J. Am. Chem. Soc.*, 2005, **127**, 16494.
 - S. R. Evans, F. Flagiello, N. Casati, L. Quaroni, and P. Macchi in progress.