

Chronicle of a death foretold: *Plasmodium* liver stage parasites decide on the fate of the host cell

Stefanie Graewe¹, Rebecca R. Stanway², Annika Rennenberg³ & Volker T. Heussler²

¹Bernhard Nocht Institute for Tropical Medicine, Hamburg, Germany; ²Institute of Cell Biology, University of Bern, Bern, Switzerland; and ³Astra GmbH, Hamburg, Germany

Correspondence: Volker T. Heussler, Institute of Cell Biology, University of Bern, Baltzerstrasse 4, 3012 Bern, Switzerland. Tel.: +41 31 631 4650; fax: +41 31 631 4615; e-mail: heussler@izb.unibe.ch

Received 27 February 2011; accepted 22 June 2011. Final version published online 9 August 2011.

DOI: 10.1111/j.1574-6976.2011.00297.x

Editor: Gerhard Braus

Keywords

malaria; *Plasmodium* liver stage; sporozoite; parasite–host interaction; exo-erythrocytic development; merosome.

Abstract

Protozoan parasites of the genus *Plasmodium* are the causative agents of malaria. Despite more than 100 years of research, the complex life cycle of the parasite still bears many surprises and it is safe to say that understanding the biology of the pathogen will keep scientists busy for many years to come. Malaria research has mainly concentrated on the pathological blood stage of *Plasmodium* parasites, leaving us with many questions concerning parasite development within the mosquito and during the exo-erythrocytic stage in the vertebrate host. After the discovery of the *Plasmodium* liver stage in the middle of the last century, it remained understudied for many years but the realization that it represents a promising target for vaccination approaches has brought it back into focus. The last decade saw many new and exciting discoveries concerning the exo-erythrocytic stage and in this review we will discuss the highlights of the latest developments in the field.

Malaria: infection of mice and men

Malaria remains one of the most devastating infectious diseases worldwide, infecting hundreds of millions of people every year. The disease is caused by protozoan parasites of the genus *Plasmodium*, which alternate between a mosquito vector and a vertebrate host. Most of the more than 200 known *Plasmodium* species infect reptiles and birds and only a relatively small number infect mammals, with five species being considered human pathogens (White, 2008). Some *Plasmodium* species that infect rodents have become invaluable tools to study general aspects of the biology of the mammalian *Plasmodium* species as their life cycles are very similar. This is particularly true of the sporozoite and liver stages of the parasite, where species infecting rodents have widely been used. Studying the biology of such stages for human *Plasmodium* species is difficult because it requires a safety level 3 facility for the maintenance of infected *Anopheles* mosquitoes, whereas mosquitoes infected with rodent *Plasmodia* can be kept in insectaries with lower safety levels. The complete liver stage development of human *Plasmodium* species can only be studied *in vitro* in primary human

hepatocytes (Mazier *et al.*, 1985) and *in vivo* in immunocompromised chimpanzees (Daubersies *et al.*, 2000; Perlaza *et al.*, 2003). Together, these factors explain why most of our recent knowledge about the *Plasmodium* exo-erythrocytic stage is based on studies using rodent models. Although the focus of this review is the exo-erythrocytic form of the parasite, for a better understanding a brief and simplified description of the entire life cycle is provided in the following section.

The ins and outs of the *Plasmodium* life cycle

Once injected into the mammalian host by female *Anopheles* mosquitoes, the *Plasmodium* parasite must pass through a series of developmental stages to ultimately produce forms that can again infect mosquitoes. For a long time, it was postulated that mosquito-derived sporozoites directly infect red blood cells (RBCs) and replicate asexually. However, in the late 1940s, it was shown that sporozoites of mammal-infecting *Plasmodium* species initially invade hepatocytes, where they replicate asexually to form thousands of merozoites (Fonseca *et al.*, 1946;

Bastianelli, 1948; Shortt & Garnham, 1948). We now know that these merozoites are packaged into vesicles (merosomes) for safe transport into the bloodstream, where they can infect RBCs (Sturm *et al.*, 2006). After several rounds of asexual reproduction in erythrocytes, which is probably necessary to generate a critical mass of infected cells for transmission, some parasites differentiate into sexual forms (gametocytes), which are infectious to mosquitoes. During a blood meal, a mosquito ingests several microliters of blood, containing as many as several thousand gametocytes together with millions of asexual forms that, unlike gametocytes, are simply digested in the midgut of the insect. Gametocytes, in response to changes in the environment from the warm-blooded mammalian host to the midgut of the mosquito, develop into gametes. Motile male gametes (microgametes) are liberated and fuse with female gametes (macrogametes), forming zygotes that advance to ookinetes. These motile forms penetrate the midgut of the insect and encapsulate under the basal lamina to form oocysts. Through extensive asexual replication thousands of sporozoites are formed, which are liberated into the hemolymph of the insect to be passively distributed throughout the entire insect body cavity. Eventually, they reach the salivary glands and actively penetrate them. After further maturation, they migrate to the ducts of the gland and can be transmitted when the mosquito takes a blood meal. The number of sporozoites inoculated by the mosquito during each bite is relatively low (50–100) (Rosenberg *et al.*, 1990; Frischknecht *et al.*, 2004; Medica & Sinnis, 2005) and thus the infection of hepatocytes was thought to be very efficient. Surprisingly, it was recently reported that only a third of the transmitted sporozoites penetrate a blood vessel and potentially reach the liver (Amino *et al.*, 2006). This might be one of the reasons why even in hyperendemic regions not every bite of an infectious mosquito results in manifestation of the disease.

This review concentrates on the biology of the sporozoite forms injected by the mosquito and their subsequent development during the liver stage. For the blood and insect stages, many excellent reviews have been published and readers interested in this stage are referred to them. Here, three main topics will be discussed: How do sporozoites leave the place of injection and reach the liver to infect hepatocytes and what happens to those that do not make it? How can sporozoites develop into thousands of merozoites in a very short time (2 days for rodent-infectious species, 4–5 days for human-infectious species)? How do merozoites leave the liver tissue and gain access to the bloodstream where they infect RBCs? It should be kept in mind that the vast majority of work related to these questions has been performed using rodent malaria models but considering the similarities in mammalian-

infecting *Plasmodium* life cycles, it is highly likely that the human-infectious *Plasmodium* species behave similarly.

The strange journeys of *Plasmodium* sporozoites in the mammalian host

When *Plasmodium* parasites are transmitted by *Anopheles* mosquitoes into their mammalian host, they are confronted with extreme environmental changes as they move from the salivary gland of a cold-blooded insect host to the skin tissue of a warm-blooded mammalian host. Once injected into the skin, the motile sporozoites transmigrate several cells before eventually crossing endothelial cells to reach a blood vessel (Frischknecht *et al.*, 2004; Vanderberg & Frevert, 2004; Amino *et al.*, 2006). The phenomenon of transmigration is discussed in detail below. Surprisingly, only a portion of the injected sporozoites (*c.* 35%) enters a blood vessel and is carried by the bloodstream to the next destination, the liver (Fig. 1). A considerable number (*c.* 15%) ends up not in blood but in lymph vessels, which are a dead end for the parasite. An even bigger portion of sporozoites (*c.* 50%) does not leave the skin tissue at all. Interestingly, it has been shown that the parasites that do not manage to leave the skin or end up in the draining lymph node induce a strong cell-mediated immune response. There is evidence that this immune response may be the basis of protection against subsequent challenges (Sinnis & Zavala, 2008).

One-way road or the highway? Parasite development in the skin

So far it has been assumed that *in vivo*, sporozoites need to invade hepatocytes to complete exo-erythrocytic development, but recent studies suggest that there is an alternative infection route. In an experimental setup, rodent-infectious *Plasmodium berghei* sporozoites entered cells in the skin and completed development into merozoites (Gueirard *et al.*, 2010). Whether this can also take place in natural infections or for other *Plasmodium* species remains to be shown. Considering that sporozoites of avian-infecting *Plasmodium* species invade and complete their development in a variety of cells including macrophages and endothelial cells (Frevert *et al.*, 2008), infection of cell types other than hepatocytes might represent an evolutionary conserved mechanism. The capability of infecting different cell types raises the question of how sporozoites recognize their host cells. There appear to be considerable differences among *Plasmodium* species in the receptors they require for infection. It has been shown that for successful invasion by *Plasmodium falciparum* and *Plasmodium yoelii* but not *P. berghei*, expression of




Fig. 1. Fate of *Plasmodium* sporozoites injected into the skin by female *Anopheles* mosquitoes: In the skin, sporozoites become motile and either enter blood vessels to be passively transported to their final destination, the liver, or enter lymph vessels to end up in the draining lymph node where they are eliminated. The vast majority of injected sporozoites, however, remain in the skin and are removed by dendritic cells (yellow) or enter skin cells and develop into mature exo-erythrocytic forms.

CD81 on the host cell surface is required (Silvie *et al.*, 2006, 2007). It is therefore not surprising that *in vitro*, unlike *P. falciparum* and *P. yoelii*, *P. berghei* parasites can infect a wider variety of cells and it is possible that this is also the basis for their ability to infect skin cells *in vivo*. In this case, it is questionable whether the human pathogen *P. falciparum* and other species that obviously need a well-defined set of cell surface markers for recognition of their host cell can infect cells other than hepatocytes.

Parasite development in hepatocytes: first contact

Even if *Plasmodium* sporozoites can infect a wider range of cells than originally thought, it is generally agreed that the main cell type in which they complete development are hepatocytes. To access them, the motile parasites need to cross the endothelium a second time after entering the bloodstream in the skin. After being passively transported by the bloodstream through the body, they eventually reach the liver but how does the parasite know where to leave the blood vessel? In the liver sinusoids, the blood flow is very slow and sporozoites are able to adhere to the endothelium. There they bind highly sulfated heparansulfate proteoglycans (HSPGs) (Coppi *et al.*, 2007), which are presented by hepatocytes through fenestrae, small channels in endothelial cells (Fig. 2). HSPGs are presented by many cell types but the sulfation level differs and is particularly high in the liver tissue. The contact of sporozoites with HSPGs starts a signaling cascade in the parasite, involving calcium-

dependent protein kinase 6 and other kinases, which finally results in the switch to an invasion mode (Coppi *et al.*, 2007).

For many years, it was thought that the circumsporozoite surface protein (CSP) serves as the receptor on the migrating parasite that targets it to the HSPGs in the liver sinusoids (Menard, 2000; Sinnis & Nardin, 2002). However, a recent study provides clear evidence that full-length CSP does not interact specifically with HSPGs but is processed to expose the C-terminal cell-adhesive thrombospondin repeat (TSR) domain once the sporozoite recognizes HSPGs by other means (Coppi *et al.*, 2011). Exposure of the TSR then allows the sporozoite to attach to the endothelium. Thus CSP is not responsible for hepatocyte detection, but rather for an unspecific adherence to cells once processed to expose the TSR domain. This also explains very nicely how the parasite can rapidly switch to the invasion mode. Importantly, transgenic sporozoites expressing only the cell-adhesive TSR domain of CSP are constitutively in the adherence and invasion mode. They do not leave the site of injection but enter skin cells and develop into infectious merozoites, confirming a recent report suggesting that sporozoites can enter and fully develop in skin cells (Gueirard *et al.*, 2010).

However, sensing the correct environment in liver sinusoids and switching to the invasion mode is not sufficient: the parasite is still on the wrong side of the endothelium and has to cross this barrier to reach its final destination, the hepatocytes. It is therefore likely that the invasion mode is not immediately triggered but rather is




Fig. 2. Sporozoite entering the liver: in the liver sinusoids, the blood flow is slow and the sporozoites can attach to the endothelium by interacting with HSPGs presented by hepatocytes through small channels in endothelial cells. Upon crossing the endothelium, sporozoites transmute through several hepatocytes before settling in one and residing inside a parasitophorous vacuole. There it develops to a multinucleated schizont and finally, by membrane invagination, forms thousands of merozoites that are liberated from the PV into the host cell cytoplasm. PVM disruption induces host cell death and the formation of vesicles that are continuously filled with merozoites and reach into the blood vessel. Finally these vesicles (merosomes) bud off and are carried away by the bloodstream to reach the lungs, where they release merozoites to infect RBCs. The pictures on the right show representative immunofluorescence assays of exo-erythrocytic parasites. From the top: sporozoite just after invasion (green) expressing PblCP (red); schizont (PVM in green and parasite membrane in red); cytomere and merozoites inside a detached cell (staining as before); merosome (merosome membrane stained in purple); all nuclei are stained with DAPI (blue).

a progressive event. It has been suggested that sporozoites glide along the endothelium until they reach one of the numerous Kupffer cells (resident macrophages in the liver) and then transmute them to reach the other side of the endothelium (Meis *et al.*, 1983; Frevert *et al.*, 2006). Indeed, it has convincingly been shown that sporozoites can transmute Kupffer cells and macrophages *in vitro* (Pradel & Frevert, 2001). On the other hand, Meis *et al.* observed digested sporozoites in Kupffer cells suggesting that these cells actively phagocytose and destroy parasites (Meis *et al.*, 1985a). Kupffer cells are not necessarily an integral part of the endothelium but

rather sit on top of endothelial cells, meaning that crossing these cells is of no obvious advantage to the parasite. Perhaps the ability to leave Kupffer cells is used instead as an immune escape strategy by the parasite to avoid destruction by phagocytosis, whereas the endothelium is crossed via a different route. Considering that sporozoites are able to transmute endothelial cells in the skin, it would not be surprising if they could do the same thing in the liver. It has been suggested that sporozoites can migrate through cells in at least two different ways (Mota *et al.*, 2001; Pradel & Frevert, 2001). One has been described in the literature as an aggressive wounding of

cells at the point of entrance and exit, where the parasite punches holes in the membrane (Mota *et al.*, 2001). This means that some intracellular material will be released from the wounded cell, potentially attracting phagocytes to the site of transmigration. The second, alternative method of sporozoite transmigration involves an invagination of the host cell plasma membrane at the point of entry (Pradel & Frevert, 2001) and likely a membrane fusion event at the exit site of the transmigrated cell. So far this mode of transmigration has only been found in Kupffer cells but it may well be that it is used for passage through endothelial cells as well. A similar mode of transcellular migration through cells is well known for immune cells in order to rapidly cross endothelia (Muller, 2010). The transmigrating immune cell induces channel formation in endothelial cells, a process involving membrane fusion events, but in doing so does not injure the transmigrated cell. Whether sporozoites *in vivo* transmigrate the endothelium by cell wounding or by the less aggressive membrane invagination and fusion method remains to be shown, but the latter could be advantageous to the parasite because it is an immunologically quiet event. Another strong argument for silent transmigration via a membrane-surrounded channel is the *in vivo* observation that the sporozoite squeezes through a small constriction in the membrane of the transmigrated cell (Amino *et al.*, 2008). This definitely fits better with the membrane invagination and membrane fusion model than with the wounding model, especially as the wounding model would suggest two parasite constriction events, one upon entry and one upon exit, whereas *in vivo* only one is seen.

Theoretically, there are alternative explanations for the parasite passing through a constriction (Frevert *et al.*, 2006). In liver sinusoids, endothelial cells are known to have so-called *fenestrae* (De Leeuw *et al.*, 1990). These are small channels in the cell, connecting the blood vessel on one side of the endothelium and the Space of Disse on the other. The size of these channels (0.1 μm) is only a tenth of the diameter of a sporozoite (1 μm) but membranes are flexible, as are sporozoites, and it has been suggested that this route is used to directly access hepatocytes (Shin *et al.*, 1982). Transmigration of sporozoites through a tight constriction might therefore reflect the passage through fenestrae in endothelial cells. Advanced intravital imaging techniques will hopefully help us fully understand this and other aspects of sporozoite transmigration in the near future.

The molecular events underlying transmigration are not yet understood. Several proteins have been identified as being specifically expressed in transmigrating parasites and have indeed found to be essential for transmigration (Ejigiri & Sinnis, 2009). A knockout of the corresponding

genes results in parasites that cannot transmigrate but are not impaired in invasion. However, the exact functions of these proteins remain to be determined.

Not falling for the first one: hepatocyte transmigration and invasion

Following migration through the endothelium, sporozoites need to cross the Space of Disse between endothelial cells and hepatocytes before they can eventually invade their final host cell. They do not, however, infect the first hepatocyte they encounter but transmigrate a number of cells before eventually invading one (Mota *et al.*, 2001) (Supporting Information, Movies S1 and S2). Again, it is not clear which mode of transmigration the parasite uses, but it has been suggested that transmigration of sporozoites through hepatocytes by cell wounding causes the release of hepatocyte growth factor (HGF) by the wounded cells (Carrolo *et al.*, 2003; Leiriao *et al.*, 2005). HGF, in turn, could be beneficial for the invading parasite in that it promotes the survival of hepatocytes, which constitutively express the HGF receptor cMET. However, as transgenic parasite lines that are not able to transmigrate cells can still enter hepatocytes and develop successfully, HGF signaling does not appear to be essential (Ishino *et al.*, 2004). In addition, as explained above, it still remains to be demonstrated which mode of transmigration sporozoites use to cross hepatocytes. Perhaps there is no cell wounding at all and the parasite again uses the silent mode to transmigrate, avoiding unwanted immune responses at the site of infection. Although other theories exist as to why transmigration by wounding might be important for the parasite maturation (Mota *et al.*, 2002; Leiriao *et al.*, 2005), it might also be that it has no specific function. The switch from the transmigration mode to the invasion mode might instead be a progressive event. Thus sporozoites continue to transmigrate cells until the complete switch to invasion mode has taken place. How the parasite decides to invade is still a mystery. For recognition and invasion, one would expect the sporozoite to require access to the host cell surface to allow receptor-ligand interaction to occur. While migrating from cell to cell in a tightly packed space, it is difficult to conceive how the parasite would perform these necessary interactions and therefore sense the host cell that it will ultimately invade rather than transmigrate. If, however, transmigration occurs in the silent mode also in hepatocytes, there is a very simple explanation: The parasite always induces invagination of the host cell membrane and leaves the cell by membrane fusion. Once the parasite stops migrating, it is already within a vacuole inside a hepatocyte and just needs to avoid membrane fusion to stay in its final host cell.

Ejigiri *et al.*, have recently suggested how sporozoites can successfully infect different cell types (Ejigiri & Sinnis, 2009). Their hypothesis is that the parasite injects its own ligands into the host cell membrane, which then bind to receptors on the sporozoite surface. This theory is supported by the fact that despite many years of research, thus far only two host cell-derived factors (CD81, SRBI) have been identified (Silvie *et al.*, 2003; Rodrigues *et al.*, 2008; Yalaoui *et al.*, 2008), which have not been proven to bind directly to sporozoite molecules. It is possible that they facilitate parasite invasion in a different way, i.e. by allowing parasite ligands to be incorporated into the host cell membrane. Although this provides an elegant explanation for how the sporozoite can infect cells other than hepatocytes, it does not explain how the parasite incorporates the ligand through the membrane of the transmigrated cells into the membrane of the neighboring hepatocyte.

Having travelled all the way from the skin to the liver, migrated through several cells, squeezed through narrow gaps, after many signaling events and recognition of being at the correct location, the sporozoite is finally ready to invade. Although it is not clear on which basis the parasite chooses its final host cell, it is well established that during sporozoite invasion the host cell membrane invaginates to form a parasitophorous vacuole (PV) around the parasite (Movie S3). In analogy to the invasion of RBCs by merozoites, it can be assumed that the majority of host cell proteins are proteolytically removed from the parasitophorous vacuolar membrane (PVM) during the invasion process (Dowse *et al.*, 2008). The sporozoite glides into the cell by the use of a moving tight junction (Movie S4), which blocks substances from entering the forming vacuole from the outside.

For a long time, it was assumed that sporozoite entry into host cells relies only on the actin-myosin motor of the parasite, but this view has been recently been challenged. It was shown that parasite invasion induces the recruitment of host cell actin to the entry site and that this accumulation is required for successful invasion (Gonzalez *et al.*, 2009). These observations and a recent analysis of the transcriptome of infected hepatocytes (Albuquerque *et al.*, 2009) provide clear evidence that the host cell reacts strongly to the invasion event. Our own observations support these findings (Movie S5). If a cell senses infection, it could induce stress signaling and its own death to eliminate the pathogen. To avoid host cell apoptosis, *Plasmodium* parasites have developed successful strategies (Leiriao *et al.*, 2005; van de Sand *et al.*, 2005). Recently, it has been shown that *P. berghei* sporozoites secrete a potent cysteine protease inhibitor (PbICP), which is able to block cell death (Rennenberg *et al.*, 2010). This method of directly inhibiting proteases involved in cell death is probably supported by additional

anti-apoptotic signaling in the host cell, but how the parasite induces this signaling is still not known. Contrary to the related apicomplexan parasite *Theileria* (Heussler *et al.*, 2006), which induces NF- κ B-dependent survival mechanisms, it has been suggested that *Plasmodium* parasites interfere with NF- κ B activation (Singh *et al.*, 2007), similarly to another related parasite, *Toxoplasma gondii* (Butcher *et al.*, 2001; Shapira *et al.*, 2002). In case of *T. gondii*, NF- κ B-independent mechanisms to avoid apoptosis have been described (Hippe *et al.*, 2009) and it will now be interesting to investigate these pathways in *Plasmodium*-infected hepatocytes. It should be mentioned that there are also reports that some *T. gondii* strains induce a NF κ B-dependent inhibition of apoptosis (Molestina *et al.*, 2003). For *Plasmodium*-infected hepatocytes, there is evidence that CSP, which is secreted by the sporozoite into the host cell cytoplasm, out-competes nuclear import of NF- κ B and thus interferes with inflammatory responses of the infected cell. However, clearly more research is needed to clarify the role of NF- κ B in host cell survival and to determine if CSP is involved in anti-apoptotic signaling or whether it merely binds in an unspecific manner to proteins and other macromolecules by exposing the adhesive C-terminal TSR domain.

Moving in: the early phase of hepatocyte infection

Parasite proteins that are thought to interact with the signaling machinery of the host cell must cross two membranes: the parasite membrane (PM) and the PVM. For *P. falciparum* blood stage parasites, it has been demonstrated that several motifs exist that mediate secretion across these membranes (Marti *et al.*, 2004; Spielmann & Gilberger, 2010). These PEXEL (*Plasmodium* export element) and PNEP (PEXEL negative export protein) motifs have been identified in many *P. falciparum* proteins known to modify the host erythrocyte. However, in the genomes of rodent-infectious *Plasmodium* species, very few genes encoding proteins with these particular export motifs have been identified. Notably, the N-terminus of CSP contains two PEXEL motifs (Singh *et al.*, 2007) and in the N-terminus of PbICP, a PNEP-like export motif has been suggested (A. Rennenberg, unpublished observations). Although the PEXEL motifs of *P. berghei* CSP have been experimentally proven to be functional, the experiments were performed in *P. falciparum*. In fact, expression of GFP-tagged proteins with PEXEL motifs in *P. berghei* did not result in secretion from blood stage or liver stage parasites (S. Horstmann, unpublished observations). Considering this and the low number of proteins predicted to contain PEXEL motifs in rodent-infectious *Plasmodium* species, it should be considered that

rodent-infectious parasites may have developed alternative motifs to allow export of proteins.

Once inside the host cell, the parasite often localizes close to the host cell nucleus (Movie S6). It is difficult to imagine that the parasite can sense its environment through the PVM and that it can control its movement while enclosed in a membrane. Therefore, it is more likely that the vacuole attaches to the cytoskeleton of the host cell and is passively transported to the nucleus. There, it is often in close proximity to the ER (Bano *et al.*, 2007) and the Golgi apparatus (A. Rennenberg, unpublished observations) and it is an attractive hypothesis that the parasite might position itself there to benefit from the host cell secretory machinery by directing host cell vesicles to fuse with the PVM. Interestingly, some intracellular bacterial pathogens also position themselves close to the host cell Golgi apparatus and it has been suggested that in this way they benefit from the transport from and to the secretory pathway (Bakowski *et al.*, 2008).

After invasion of the host hepatocyte and formation of a PV, the parasite undergoes an initial period of comparatively subtle changes (Movie S6), at least on a morphological level. During the first 24 h after infection, the parasite remodels its PVM and transforms from its elongated form to a small, round trophozoite. Several proteins are known to be important for this early stage as their knockout led to an impairment in development. Among them are *uis3*, *uis4* and *Pb36p* (Mueller *et al.*, 2005a, b; van Dijk *et al.*, 2005).

At approximately 20 h after invasion, the parasite nucleus starts dividing repeatedly (Fig. 2) and displays one of the fastest replication rates known for eukaryotes: during the next 35 h, up to 30 000 nuclei are generated. At the same time, the PV expands in size to accommodate the growing parasite (Movie S7). During blood stage replication, the parasite is known to take up hemoglobin from its host cell through a kind of cell mouth, a so-called cytostome (Elliott *et al.*, 2008) and to digest it in a food vacuole (Rosenthal & Meshnick, 1996). Interestingly, exoerythrocytic merozoites of avian-infecting *Plasmodium* species also possess a cytostome (Aikawa *et al.*, 1966) and we have found a morphologically similar structure in the growing schizont (Fig. 3). Whether these structures have a function, however, remains to be determined. It is therefore still unknown how the parasite manages to obtain the resources necessary for its immense reproduction effort within the hepatocyte. Apart from the host cell ER, which gathers around the parasite and a loose association with the Golgi apparatus, there appears to be no constant association between the PV and host cell organelles (Bano *et al.*, 2007). For intracellular *Toxoplasma* parasites, it has been shown that host cell mitochondria are firmly associated with the PVM and




Fig. 3. Liver stage parasites possess structures resembling a cytostome. The PVM marker protein Exp1 is stained in green and the cysteine protease inhibitor PbICP is stained in red. Parasite and host cell nuclei are visualized by DAPI staining (blue). Round structures (putative cytostomes) labeled by the PVM marker and PbICP are indicated with arrows.

supply the parasite with the enzyme co-factor liponic acid (LA) (Crawford *et al.*, 2006). A similarly close association of the mitochondrion was not found in *P. berghei*-infected hepatocytes, but very careful microscopic examination revealed distinct areas at the PVM that appear to indeed interact with host cell mitochondria (C. Deschermeier, unpublished observations). We could also demonstrate that *P. berghei* needs to import LA from the host cell, most likely from host cell mitochondria. It is very surprising that liver stage parasites largely rely on LA import as they produce this co-factor themselves in the apicoplast, an essential plastid-like organelle that plays an important role in liver stage development (Stanway *et al.*, 2009a). It seems that LA cannot be easily transferred from the apicoplast to the mitochondria, which is surprising considering that other metabolites are known to be exchanged easily between organelles within the parasite, e.g. during heme detoxification (Sato *et al.*, 2004; van Dooren *et al.*, 2006; Padmanaban *et al.*, 2007). However, as hepatocytes contain many mitochondria producing LA, perhaps there has been no evolutionary pressure for the parasite to independently transport this co-factor from the apicoplast to the mitochondrion.

It has been speculated that for nutrient uptake, the parasite inserts transport channels into the PVM allowing molecules up to 800 Da to freely cross this membrane. (Bano *et al.*, 2007; Sturm *et al.*, 2009). However, larger molecules like lipids and peptides need to be actively imported and again the parasites modifies the PVM to allow this to occur. It exports a number of proteins into the PVM, which most likely interact with host cell proteins to direct nutrients toward the parasite.

However, so far only few host proteins have been identified that might support liver stage growth. One of them is the protein ApoA1 that was found to localize to the PVM. It is thought to interact with uis4 and is speculated to play a role in the synthesis of additional membrane during the enlargement of the vacuole (Prudencio *et al.*, 2006).

During its extensive growth, use of host cell resources is likely to deplete nutrients from the infected hepatocyte. In response to the resulting starvation conditions, the host cell is expected to induce autophagy and indeed we find increased autophagy in infected cells (N. Eickel, unpublished observation). However, autophagy is also a very potent mechanism to eliminate pathogens and again we have evidence that *in vitro* parasites can be destroyed in autophagosomes. Host cell autophagy therefore appears to be a double-edged sword for the parasite: on the one hand, it could provide nutrients for its extensive growth and on the other hand, it could result in parasite elimination. Further research on this highly interesting topic is needed to fully understand the function of host cell autophagy in regard to parasite survival and elimination. What has been underestimated so far is that the host cell has the capacity to eliminate the parasite. It was an accepted view that the parasite can manipulate the survival of the host cell but now it turns out that the host cell is not helpless and can successfully fight the infection. This topic will be very important to study as it may help to develop new antimalarial strategies.

For the parasite, the supply of nutrients is essential, but it must also dispose of metabolic waste products. In blood stage parasites, for example, the toxic end product of hemoglobin metabolism, hemozoin, is stored in the food vacuole (Goldie *et al.*, 1990). For liver stage parasites, so far no food vacuole has been described and the question remains how the metabolically highly active liver stage parasite deals with its waste products. It has been suggested that it employs efficient export transporters in its membrane and in the PVM (Sturm *et al.*, 2009). This might partly explain why liver stage parasites are less susceptible to drugs than blood stage parasites: they might be able to actively export these from the PV, therefore preventing them from reaching the parasite. It would be interesting to combine drug treatment with blockers of these putative transporters. Although there is evidence for the expression of such transporters in the blood stage (Valderramos & Fidock, 2006), proof of their existence and characterization in the liver stage is still missing.

When one becomes many: the challenges of replication

As liver stage parasites grow very rapidly, a classical mode of cell division and cytokinesis is most likely not possible

because it requires the expression of numerous proteins in a concerted fashion followed by the removal or inactivation of the entire machinery. It is therefore not surprising that the parasite has developed strategies to streamline proliferation and growth. The most obvious phenomenon is that the parasite avoids cytokinesis until all nuclear division is complete and thus develops into a huge syncytium, the multinucleated schizont. Less obvious is the fact that the nuclear division is not accompanied by the disappearance of the nuclear membrane. Electron microscopy studies from other parasite stages suggest that the segregating chromosomes and the spindle apparatus remain within the nuclear envelope (Bannister *et al.*, 2000) but this does not explain the highly amorphous phenotype of the nuclei during division. Could it be that the microtubule organization center is localized in a way that it can associate with the cytoskeleton in the cytoplasm of the parasite? It has indeed been shown very recently for blood stage parasites that the microtubule organization centers are embedded in the nuclear membrane (Gerald *et al.*, 2011) and it might well be possible that they are connected to the spindle apparatus in the nuclei and to the cytoskeleton outside the nucleus. Thus mitosis in *Plasmodium* parasites appears to differ in many respects from that in the mammalian host (Gerald *et al.*, 2011). This is an interesting aspect because differences in the cell biology of the parasite and its host cell might reveal new strategies for interference with parasite development. Once schizogony is completed, cytokinesis and daughter cell formation take place. As each daughter cell requires a full set of organelles, including a mitochondrion and an apicoplast, which cannot be synthesized *de novo*, the existence of a highly organized distribution machinery must be postulated. In eukaryotic cells, cell division is normally preceded by the division of organelles, which are then distributed to both daughter cells. Does the same happen in *Plasmodium* during the times of schizogony when no cytokinesis is taking place? The next section takes a closer look at the replication and distribution of several *Plasmodium* organelles and will primarily focus on the morphological and positional changes that occur for the apicoplast, mitochondrion and nuclei.

During erythrocytic development, the parasite has already been shown to form both a branched apicoplast and mitochondrion, with fission of these organelles occurring only after completion of the multiple rounds of nuclear division and with each nucleus ultimately being paired with a single apicoplast and mitochondrion (van Dooren *et al.*, 2005). During the liver stage of *Plasmodium* development, where not 16–32 but up to 30 000 daughter parasites are formed, the parasite faces an even greater challenge in terms of organelle growth and segregation into merozoites. Until recently, it was unclear

whether the liver stage parasite employs a similar mechanism of apicoplast and mitochondrial growth and segregation into daughter parasites as that in the blood stage.

The trophozoite: calm before the storm


Various transgenic *P. berghei* parasite lines have been generated that allow the visualization of the nuclei, apicoplast and mitochondrion during liver stage development (Stanway *et al.*, in press). From previous studies it was known that salivary gland sporozoites contain a single nucleus, apicoplast and mitochondrion, which do not necessarily have a clear physical connection (Stanway *et al.*, 2009a; Kudryashev *et al.*, 2010). For approximately the first 20 h after the invasion of the hepatocyte, the parasite maintains a single nucleus. During this time, the apicoplast and mitochondrion both elongate (Fig. 4). In an interesting resemblance to the blood stage, at this point the mitochondrion primarily lies at the periphery of the parasite. However, in contrast to the blood stage, where both organelles appear to maintain a continuous interaction (van Dooren *et al.*, 2005), in liver trophozoites both organelles are mainly found separated from each other. Contact between apicoplast and mitochondrion seems to be rather accidental. This observation is surprising because these two organelles are thought to share the

heme biosynthesis pathway and their physical connection is expected to be required for metabolite exchange. However, there are other apicomplexan parasites, like *Toxoplasma*, in which the connection between the apicoplast and mitochondrion is not continuous (Nishi *et al.*, 2008). Perhaps a physical connection between the organelles is not necessary to maintain a functional heme pathway or the occasional points of interaction provide enough time for metabolite exchange. Another important aspect, which has not been tackled so far, is the question how organellar genomes are distributed correctly within the huge developing network. After fission each organelle must contain the genetic material and a highly sophisticated machinery to achieve this can be predicted. Future research will hopefully shed some light on this issue.

The schizont: rapid nuclear division and extensive organelle growth

Following the trophozoite stage, the parasite begins repeated rounds of nuclear division. Based on a replication from 1 nucleus to up to 30 000 nuclei in approximately 30 h, this would imply that a round of nuclear division occurs approximately every 2 h. In parallel with this extensive nuclear division, both the apicoplast and mitochondrion become highly intertwined branched structures, but each appears to remain as a single organelle

Fig. 4. Organelle development and distribution into daughter cells: invading sporozoites contain a mitochondrion (red), apicoplast (green) and a nucleus (cyan). During schizogony, the nucleus divides but the mitochondrion and the apicoplast continue to grow and branch extensively. During the cytomere stage, nuclei become attached to the invaginating parasite membrane and the apicoplast and the mitochondrion are directed toward the forming merozoites. Each merozoite finally contains a single apicoplast, mitochondrion and nucleus. On the right are live spinning disc microscopy images of representative parasite stages, where GFP is targeted to the apicoplast and mCherry to the mitochondrion. The upper image shows a schizont with an extensive and intertwined apicoplast and mitochondrion, whereas the lower image shows a cytomere, briefly before fission of the apicoplast and mitochondrion.



(Fig. 4; Stanway *et al.*, 2011). Thus far, it is unclear how the growth of these organelles is controlled but neither branching of the mitochondrion nor the apicoplast appears to involve any clear association with nuclei. This is surprising, because in both *Toxoplasma* and *Sarcocystis* parasites, the apicoplast is connected with the nucleus (Striepen *et al.*, 2000; Vaishnav *et al.*, 2005). The observation that the mitochondrion, like the apicoplast, undergoes extensive branching during schizogony contradicts earlier EM studies that described a proliferation of the mitochondrion during schizogony based on the observation of multiple mitochondrial profiles in thin section (Meis *et al.*, 1985a, b, c). However, this observation is not contradictory to a branched network as described by us because in thin sections a highly branched structure would also result in multiple profiles. Furthermore, such branching is consistent with what has been seen in blood stage development and indeed occurs in each of the three stages of asexual replication during the *Plasmodium* life cycle (van Dooren *et al.*, 2005; Stanway *et al.*, in press).

The extensive growth of the apicoplast and mitochondrion requires a massive quantity of membrane. Coupled with the extensive growth of the parasite and presumably the replication of other cellular organelles, this may at least in part explain the reliance of the parasite on fatty acid biosynthesis pathways during exo-erythrocytic development, which stands in contrast to blood stage development and sporogony (Yu *et al.*, 2008; Pei *et al.*, 2010). Knockout of genes coding for important components of the fatty acid pathway like pyruvate dehydrogenase and FabI allow normal blood stage development but block rapid proliferation of liver stages.

The cytomere: generating order out of chaos

To manage the extensive growth of the apicoplast and mitochondrion during liver stage development is already an impressive feat but the fission of these organelles and correct segregation into forming daughter merozoites appears to be an even greater challenge. How the parasite manages this on a molecular level is not understood, but again double-fluorescent parasites have allowed us to begin to understand the morphological and positional changes in the apicoplast, mitochondrion and nucleus prior to and during merozoite formation. It appears that the processes undertaken by the liver stage parasite parallel those in the blood stage, but on a much larger scale. The large size of the liver stage parasite, however, allows a more detailed examination of these processes.

Following the completion of nuclear division, at which point the single parasite can contain many thousands of nuclei, the parasite develops to a stage known as the

cytomere. Here the plasma membrane of the parasite invaginates to form what appear to be spheres of membrane, portioning the cytoplasm into between approximately 5 to 20 units (Figs 2 and 4; Movies S7 and S8). At this stage, the nuclei of the parasite seem to be closely associated with the plasma membrane and so they are in a sphere-like arrangement, which appears like rings of nuclei when examined by two-dimensional microscopy.

The apicoplast primarily lies at the periphery of these spheres of nuclei, with a surprising resemblance to the position of the *Sarcocystis* apicoplast prior to division of the polyploid nucleus (Vaishnav *et al.*, 2005). The mitochondrion on the other hand mostly lies within these spheres of nuclei, forming clumped structures (Fig. 4). During the cytomere stage, the individual units of plasma membrane are not fully separated and mitochondrial branches connect the afore-mentioned clumps, confirming earlier EM studies on exo-erythrocytic development of the parasite (Meis *et al.*, 1981, 1985a, b, c). The cytomere stage is an intermediary parasite stage and soon after its formation, the apicoplast develops a concertinaed structure, presumably due to the constriction of the organelle at sites where fission will take place. The morphology of the apicoplast at this stage again resembles that observed in *Sarcocystis* parasites (Vaishnav *et al.*, 2005). Around this time, the morphology of the mitochondrion also changes dramatically: finger-like structures form that point to the periphery of the membrane units. The apicoplast of the parasite then divides synchronously. At the point of division or close before, an association is clearly visible between the apicoplast and mitochondrion, with each finger-like structure being associated with a divided portion of apicoplast. It remains to be seen whether these organelles are directly associated or are both connected to a third structure, for example, the same cytoskeletal element. One could hypothesize that organelles may enter the forming daughter merozoites via association with and movement along subpellicular microtubules, as has been seen in the case of *Toxoplasma* parasites (Striepen *et al.*, 2000). The existence of such microtubules, positioned within each forming daughter parasite, could explain how the mitochondrion is able to undergo such a dramatic and synchronous change in morphology. The association between the apicoplast and mitochondrion occurring primarily at the end of the liver stage would suggest that a connection between these organelles might be required to allow correct organelle segregation, as already proposed by others (Sato *et al.*, 2004; van Dooren *et al.*, 2006; Padmanaban *et al.*, 2007).

Following the change in its morphology, the mitochondrion divides. It is interesting to note that during both the asexual blood stage and the liver stage, division of the

apicoplast always precedes that of the mitochondrion. This is also true for *T. gondii* parasites (Nishi *et al.*, 2008), despite the difference in the methods used for cell division. Once mitochondrial fission is complete, the daughter merozoites are formed and released into the host cell by breakdown of the PVM.

Not just different for the sake of being different

For both mitochondrial and apicoplast fission, the underlying molecular mechanisms are far from clear. Some components of typical organelle fission machineries appear to be conserved in *Plasmodium* and *Toxoplasma* parasites, such as dynamin, which was shown to be involved in the fission of the apicoplast in both species (Charneau *et al.*, 2007; van Dooren *et al.*, 2009). However, homologs of other key players in organelle fission appear to be absent. This is particularly true of the Ftsz protein, which in other systems is responsible for the initial constriction of the organelle by allowing dynamin to bind but is reportedly absent in apicomplexan parasites (Vaishnav & Striepen, 2006). It remains to be seen whether in Apicomplexa, proteins related to division have evolved sufficiently to be elusive in homology searches or whether the parasites have developed alternative mechanisms for the division of their organelles. The latter option currently seems the most likely.

When observing the development of the apicoplast, mitochondrion and nuclei of *Plasmodium* liver stage parasites, one could be surprised by the methods used by the parasite to achieve segregation of these essential organelles. One can speculate as to why *Plasmodium* parasites at all stages of asexual reproduction develop via schizogony and not via daughter cell formation as performed during the lytic stage of the *Toxoplasma* life cycle. Indeed, it appears that none of the known apicomplexan parasite genera use an identical mechanism of asexual replication (Vaishnav & Striepen, 2006). It can only be speculated why evolution allowed the parasite to develop huge branched organelles, which undergo one fission event at the end of schizogony rather than repeated apicoplast and mitochondrial fissions along with division of the nucleus and the Golgi apparatus (Struck *et al.*, 2005; R.R. Stanway, unpublished observation). One reason might be that a majority of the proteins functioning within the apicoplast and mitochondrion are encoded in the genome of the parasite. They need to be targeted to these organelles, where import machineries allow their uptake. For a repeated fission of organelles, the parasite would need to constantly express proteins involved in fission and to import them into the relevant organelles, while at the same time maintaining a tight control of protein activity

to prevent unwanted fission events. In contrast, for a synchronous fission of the apicoplast and mitochondrion after the completion of karyokinesis, presumably only one period of protein expression is required. This would have the added advantage that control of fission could be on the level of protein expression, allowing a tight synchronicity. It is also conceivable that in the absence of repeated fission events, both the apicoplast and mitochondrion may be able to function more efficiently. For the extensive parasite growth and replication during the liver stage, the parasite must require a high level of energy and of fatty acid biosynthesis, the latter for the extensive production of membrane that accompanies this growth. Such demands on the mitochondrion and apicoplast may be incompatible with repeated divisions of these organelles.

Breaking out: egress from the parasitophorous vacuole

Once formation of merozoites is complete, they must be transported into the bloodstream where they can infect RBCs to continue the life cycle. Most text books state, probably in analogy to the events at the end of the blood stage, that the host cell ruptures and releases merozoites, which subsequently infect RBCs. However, if the host cell membrane broke down within the liver tissue, the merozoites would be on the wrong side of the endothelium (Fig. 2). In fact, it has recently been shown that the release of merozoites is a well-orchestrated, multi-step process. The first hurdle for the parasite is the PVM and it has been shown that during merozoite formation parasite proteins begin to leak into the host cell, thus demonstrating that the membrane of the PV becomes increasingly permeable before it is completely disrupted (Schmidt-Christensen *et al.*, 2008; Sturm *et al.*, 2009). As another sign of PVM breakdown, several previous *in vivo* studies had already indicated that parasite and host cell material mix freely in infected cells (Movie S8) (Meis *et al.*, 1985a,b,c; Baer *et al.*, 2007). Very recently, live imaging of a parasite strain expressing a fluorescent PVM marker protein confirmed the breakdown of the PVM toward the end of the liver stage while the host cell membrane stays intact (Graewe *et al.*, in press).

Not much is known about the molecular events resulting in the breakdown of the PVM in *Plasmodium* liver stages. It takes place within a relatively short time frame (Graewe *et al.*, in press) and therefore must be a highly efficient process. The first class of enzymes one would consider to act on membranes are lipases, but it remains to be shown whether the parasite secretes or activates lipases to destroy the PVM. However, it is known that proteases can destabilize membranes by removing

membrane-integrated proteins. Indeed, PVM breakdown can be inhibited by E64, an inhibitor of cysteine proteases, which indicates a role for this class of proteases (Sturm *et al.*, 2006). The only protein that has been identified to be involved in PVM disruption by *Plasmodium* parasites is LISP-1 in *P. berghei* (Ishino *et al.*, 2009). It localizes to the PVM and its deletion results in an inability of the parasite to escape from the PV. LISP-1 itself, however, has no recognizable functional protease domain and is therefore suspected to be either a membrane receptor for proteases or to be involved in the processing of proteases for activation (Ishino *et al.*, 2009). A protein that is proteolytically processed at this time is PbSERA3, a putative cysteine protease that is subsequently released into the host cell cytosol (Schmidt-Christensen *et al.*, 2008). The processing of PbSERA3 is E64-sensitive and it would therefore be a likely candidate for the mediation of the PVM breakdown and subsequent changes in the host cell. Despite its prediction to be a protease, so far it has not been possible to demonstrate any catalytic activity for SERA3. Therefore, like LISP-1, it might merely act as an adapter protein for the recruitment of effector proteases.

Further understanding of liver stage PVM breakdown might be reached by examining other life cycle stages of the parasite. During egress from both oocysts and erythrocytes, *Plasmodium* parasites need to break down several surrounding membranes to continue their development. In blood stage egress especially, the overall situation is similar to the late liver stage: the parasite is separated from new host cells by two membranes: the PVM and the host cell membrane. A number of molecular similarities have already been found that make a common mechanism for the disintegration of the PVM feasible. As in the liver stage (Sturm *et al.*, 2006), egress from the PV is blocked by E64 and members of the SERA family are cleaved shortly before the release of parasites (Yeoh *et al.*, 2007). The processing proteases have already been identified as PfSUB1 and DPAP3 (Yeoh *et al.*, 2007; Arastu-Kapur *et al.*, 2008) and it remains to be seen if they also act in the liver stage. An even more interesting question is whether one of the proteases in this cascade is responsible for the host cell death that is induced upon PVM breakdown. The dramatic modifications of the host cell that occur in parallel to the release of merozoites from the PV are discussed in more detail below.

It is clear that in infected hepatocytes, the host cell membrane remains intact after the disintegration of the PVM (Sturm *et al.*, 2006; Graewe *et al.*, in press), but in infected RBCs, both surrounding membranes rupture in quick succession (Blackman, 2008). This difference in parasite release makes perfect sense as hepatocyte-derived merozoites need transport to the blood vessel whereas RBC-derived merozoites can directly infect another RBC.

Still, some controversy has remained about which membrane breaks down first when parasites exit RBCs: the host cell plasma membrane or the PVM? On one hand, live imaging of parasites that express GFP in their PV has shown that the fluorescent protein spreads throughout the entire host cell toward the end of the blood stage (Wickham *et al.*, 2003). This indicates that the PVM disintegrates first. On the other hand, observations have been made of extracellular clusters of merozoites surrounded by a PVM (Soni *et al.*, 2005). Despite the uncertainty regarding merozoite release from RBCs, it is clear, however, that the disruption of the PVM and the host cell membrane are differentially inhibited. While PVM breakdown is blocked by E64, the breakdown of the host cell membrane was shown to be inhibited by the broad-spectrum cysteine and serine protease inhibitors leupeptin and chymostatin (Salmon *et al.*, 2001; Wickham *et al.*, 2003; Soni *et al.*, 2005). This indicates that the PVM and the RBC membranes are broken down by different sets of proteases (Wickham *et al.*, 2003; Blackman, 2008). As the plasma membrane of infected hepatocytes does not break down immediately upon PVM disruption, the parasite-derived protease destroying the RBC membrane might either not be synthesized in the liver stage or might be inhibited by specific parasite factors. A potential candidate protein would be the recently identified cysteine protease inhibitor PbICP, which is released into the host cell during the late liver stage (Rennenberg *et al.*, 2010). In conclusion, the data available from the study of blood stage egress indicate the involvement of specific sets of proteases that are activated in cascades (Yeoh *et al.*, 2007; Blackman, 2008). While it is likely that similar proteolytic mechanisms act in the liver stage, it is still unclear whether these activated proteases directly destabilize the PVM by cleaving integral membrane proteins, or if they act by initiating other effector molecules like lipases or pore-forming proteins.

For other intracellular parasites, like the related apicomplexan parasite *T. gondii*, it has been shown that a decrease in the intracellular potassium concentration leads to an increase in calcium concentration within the parasite, which seems to trigger egress (Moudy *et al.*, 2001; Nagamune *et al.*, 2008). The lysis of the PVM and the host cell membrane is then caused by a pore-forming perforin-like parasite protein, TgPLP1 (Kafsack *et al.*, 2009). The exit from vacuoles or host cells via use of pore-forming proteins is a common strategy for many intracellular pathogens, including organisms as diverse as *Listeria monocytogenes*, *Trypanosoma cruzi* or *Leishmania amazoniensis* (Gaillard *et al.*, 1987; Andrews, 1990; Andrews *et al.*, 1990; Noronha *et al.*, 2000). A typical feature of pore-forming proteins is the MACPF (membrane attack complex perforin) domain (Xu *et al.*, 2010). In *Plasmodium*,

several proteins containing MACPF domains have been identified and one of them has already been shown to have a role in the transmigration of Kupffer cells (Kaiser *et al.*, 2004). Further investigation will show if these proteins are expressed in the late liver stage and if they play a role in the breakdown of the PVM. It is not clear how such pore-forming activity would be limited to the PVM and would be prevented from affecting the merozoite membrane or the host cell membrane. In intracellular pathogens that escape from a phagosome, such as *T. cruzi* or *L. monocytogenes*, control is achieved by restricting pore-forming activity to occurring only at low pH (Andrews, 1990; Andrews *et al.*, 1990; Schuerch *et al.*, 2005). As soon as the parasites are liberated from the phagosome, the pH value changes and the host cell membrane is not perforated. As the PV environment of *Plasmodium* liver stage parasites is not expected to be of low pH, other means of regulation must exist. One attractive possibility is that adapter proteins that are only anchored in the PVM are necessary for the recruitment and activation of such a pore-forming protein.

Host cell death: maintaining a calm exterior

While the exact mechanism of PVM breakdown remains to be elucidated, it is obvious that escape from just the PVM does not allow the parasite to reach the blood vessel. To safely cross the endothelium, the parasite uses a trick. Once the PVM is dissolved, a parasite-dependent host cell death is initiated (Sturm *et al.*, 2006). It is characterized by cytochrome *c* release and nuclear condensation under retention of an intact cell membrane (Sturm *et al.*, 2006; Graewe *et al.*, in press). It therefore clearly differs from necrosis, which typically includes the swelling and rupture of the cell (Lemasters, 2005). This difference is not surprising as a necrotic host cell death would lead to the release of pro-inflammatory molecules and the attraction of components of the host immune system (Savill, 1998; Scaffidi *et al.*, 2002; Shi *et al.*, 2003), which could then potentially act against the parasite.

Although, at first glance, the host cell death resembles apoptosis, it differs in many respects and appears to be unique. Whether apoptosis is triggered is usually decided by the balance between pro- and anti-apoptotic stimuli. Both the integration process and the initiation of apoptosis often take place in the mitochondrion, through the formation of a mitochondrial outer membrane permeabilization pore (MAC) (Kroemer & Reed, 2000). This ultimately leads to the release of cytochrome *c* and other apoptotic mediators into the cytosol. The breakdown of the PVM could theoretically induce host cell death in

different ways. Firstly, parasite effector proteins could be released into the host cell after the PVM breaks down and this might cause death as a rather unspecific effect. As we have evidence that premature disruption of the PVM does not lead to death of the host cell, this scenario is less likely. Secondly, the host cell death might be purposefully orchestrated by specific parasite proteins secreted into the PV very late in parasite development and then released into the host cell upon PVM breakdown. Several bacteria are already known to regulate host cell death by producing factors that compromise mitochondrial integrity (Braun *et al.*, 2007; Kozjak-Pavlovic *et al.*, 2009) and *Plasmodium* might have evolved similar strategies. This hypothesis is supported by the live imaging of *Plasmodium*-infected hepatoma cells with fluorescently labeled mitochondria, which has shown these organelles to disintegrate rapidly after PVM breakdown (Graewe *et al.*, in press).

Upon closer examination, parasite-dependent host cell death lacks important hallmarks of apoptosis such as DNA fragmentation, caspase cascade activation and loss of phosphatidylserine asymmetry (Sturm *et al.*, 2006; Graewe *et al.*, in press). Interestingly, all of these features are events that take place in the mid to late stages of apoptosis. They also all require energy in the form of ATP: the fragmentation of DNA, for one, is carried out by endonucleases, which hydrolyze ATP during cleavage. Likewise, the caspase cascade is initiated through a complex composed of cytochrome *c* and Apaf-1, which undergoes an ATP-dependent conformational change that activates procaspase 9 (Zou *et al.*, 1999). While the phosphatidylserine switch itself does not appear to utilize ATP, it is linked to the activation of the caspase cascade as inhibitors of caspases were shown to prevent the phosphatidylserine asymmetry loss (Castedo *et al.*, 1996; Martin *et al.*, 1996). Based on these observations, the following model was suggested (Graewe *et al.*, in press): while the parasite undergoes schizogony, it depletes the host cell of nutrients and ATP. Upon rupture of the PVM, mitochondrial integrity is compromised, probably by activation of proteases other than caspases. This leads to the uncoupling of oxidative phosphorylation and therefore loss of the ability to produce ATP *de novo*, which aggravates the lack of accessible energy. Simultaneously, apoptotic factors that are released from the mitochondria initiate the apoptotic program, which proceeds until it reaches a point where major amounts of ATP are required. It then stalls, which leads to an aborted version of apoptosis.

The arrest of the apoptotic program might additionally be aided by inhibitors produced by the parasite. It has already been shown that *Plasmodium* is capable of suppressing host cell death pathways earlier in liver stage

development (van de Sand *et al.*, 2005) and this may also be true after merozoites are liberated into the host cell cytosol. A possible candidate for this inhibition is the previously mentioned cysteine protease inhibitor PbICP, which is not only present in the host cell cytosol early after invasion but also floods the host cell upon PVM disruption (Rennenberg *et al.*, 2010). It has been shown to inhibit apoptosis when expressed in CHO (Chinese hamster ovary) cells. As it is effective against cathepsin L-, but not B-type proteases, PbICP could in theory block host cell effector proteases while allowing parasite proteases of the cathepsin B-type to remain functional (Rennenberg *et al.*, 2010).

In addition, findings from *T. gondii* research indicate that inhibitors might be necessary to prevent rapid host cell lysis after the mass release of proteins from the PV. In infections with an avirulent *T. gondii* strain, host IRG (interferon-inducible immunity-related GTPases) proteins have been shown to disrupt the PVM prematurely (Zhao *et al.*, 2009). Although this kills the parasite, the concomitant discharge of the PV contents into the cytosol resembles *Plasmodium* PVM egress. As in *Plasmodium* infections, a caspase-independent host cell death is triggered but in contrast to the *Plasmodium* liver stage, it results in a rapid host cell death, including membrane permeabilization and the release of inflammatory proteins (Zhao *et al.*, 2009). A rapid breakdown of the host plasma membrane makes perfect sense considering the biology of *Toxoplasma* parasites. After rupture of their host cell, they immediately infect another cell and thus there is no need for the ordered cell death observed for *P. berghei*-infected hepatocytes. During liver stage development, *Plasmodium* merozoites remain within their host cell for a comparatively long time period after PVM rupture and therefore might have evolved inhibitors that slow down cell death and create a shift toward an immunologically silent outcome.

Despite the similarities between *Plasmodium* egress from hepatocytes and RBCs, there are also clear differences, most likely because of the different nature of the respective host cells and the different needs of the parasite. Upon PVM breakdown, the exo-erythrocytic merozoites stay for an extended period of time in hepatocytes until they reach their final destination where they can safely infect RBCs. In contrast, egress from RBCs needs less coordination as liberated merozoites can infect another RBC within seconds. After invasion of RBCs, *P. falciparum* activates non-selective cation channels in the erythrocyte membrane, presumably to have easy access to sodium and calcium (Kasinathan *et al.*, 2007). Usually, the activation of these channels leads to a rise in intracellular calcium levels and triggers eryptosis, the programmed cell death of erythrocytes. In *Plasmodium* infections, however, this process is

delayed, possibly through the uptake of calcium by the parasite (Kasinathan *et al.*, 2007). Characteristic features of eryptosis such as the activation of calpain do not occur until the late blood stage. Even then they seem to be tailored to support parasite development as it has been shown that in the absence of host cell calpain-I, *P. falciparum* is incapable of egressing from the erythrocyte (Chandramohanadas *et al.*, 2009). The exact mechanism is still unclear but it appears to involve a remodeling of the cytoskeleton. It has been proposed that calpain-I also becomes activated in the liver stage but this remains to be shown. As the *Plasmodium* ICPs appear to inhibit calpain-I (Pandey *et al.*, 2006) and PbICP was found in the host cell upon PVM breakdown, calpain-I is not likely to play a major role in parasite egress from hepatocytes. In addition, the increased uptake of calcium by exo-erythrocytic merozoites would further argue against activation of calpain-I. It has even been demonstrated that calcium uptake by exo-erythrocytic merozoites blocks the switch of phosphatidylserine residues from the inner leaflet to the outer leaflet of the membrane allowing the parasite to interfere with host cell signaling and to avoid attack by phagocytes (Heussler *et al.*, 2010).

Taken together, toward the end of the *Plasmodium* liver stage, the host cell undergoes an unusual cell death that appears to be an aborted version of apoptosis. It is not yet fully understood how this process occurs, but both specialized parasite inhibitors and a shortage of available energy might play a role. The result is a host cell that has detached from its surroundings and contains free exo-erythrocytic merozoites in its cytosol.

Moving out and taking the blinds: merozoite formation and re-entry into the bloodstream

Until recently, it was unclear how liver stage development was ultimately concluded. For a long time, it was believed that the host cell membrane ruptured along with the PVM to release the infectious merozoites. However, it was not known how merozoites would pass through the endothelium to reach the bloodstream and infect RBCs. Careful electron microscopical analysis revealed already that groups of merozoites are released into the bloodstream (Meis *et al.*, 1985a,b,c), but from this work it was not clear that the parasites are still surrounded by a host cell membrane. *In vitro* live imaging revealed that the host cell membrane remains intact for an extended period of time after PVM breakdown and that the entire cell detaches from its surroundings toward the end of liver stage development (Graewe *et al.*, in press). *In vivo* live imaging showed that subsequently, vesicles (merosomes) bud off, which contain up to several thousand motile merozoites

(Movie S9) (Sturm *et al.*, 2006; Baer *et al.*, 2007). These merosomes are released directly into adjacent blood vessels. The process of merosome formation has not yet been fully understood. It can be blocked by protease inhibitors and is therefore assumed to involve a protease-mediated destabilization of the surrounding membrane (Sturm *et al.*, 2006). Interestingly, there are several other intracellular pathogens such as *Chlamydia* and *L. monocytogenes* that remain wrapped within their host cell membrane during egress. While no definite underlying molecular mechanism has been identified for these either, a remodeling of the actin cytoskeleton and the active movement of the pathogen seem to play a role (Hybiske & Stephens, 2007). As *Plasmodium* merozoites have been observed to move rapidly within the detached host cell (Stanway *et al.*, 2009b; Graewe *et al.*, in press), it is in theory possible that they could push themselves into cellular extensions to form merosomes.

Apart from the specifics of the formation process, it is also puzzling that although the merosome membrane is made up of host cell membrane, it does not exhibit typical host cell markers such as ASGR1 (Baer *et al.*, 2007). Recently, however, it was shown that even a fluorescent transmembrane reporter protein is rapidly lost from the host cell membrane upon PVM breakdown (Graewe *et al.*, in press). It is proposed that this is due to the arrest of protein biosynthesis within the host cell, likely as a consequence of mitochondrial damage and a subsequent lack of energy. It appears that at this point, only a husk of the host cell remains, which is nevertheless invaluable to *Plasmodium* as the exit from the liver via merosomes is a very elegant immune evasion strategy. On its way into the blood vessel, the parasite needs to pass the resident macrophages of the liver, the Kupffer cells, which line the liver sinusoids (Arii & Imamura, 2000). Merozoites themselves would be recognized as foreign by these immune cells and it is known that phagocytes can efficiently engulf merozoites (Terzakis *et al.*, 1979). By traveling in a vesicle composed of host cell membrane, *Plasmodium* avoids recognition and gains safe passage to the bloodstream. Once the merosomes have entered the circulatory system, they are reduced by shear forces to a size of 12–18 μm (Baer *et al.*, 2007). They pass through the right side of the heart before accumulating in the pulmonary capillaries of the lung. Electron and *ex vivo* microscopy have shown that the merosomes then rupture and release their cargo of infectious merozoites (Baer *et al.*, 2007). As the local environment is rich in erythrocytes, reinvasion can take place quickly and *Plasmodium* can again take advantage of the protection of an intracellular environment. Like the liver stage, the blood stage of the parasite ends with the formation of merozoites and it is therefore not surprising that both stages

share many features and similar transcriptomes (Tarun *et al.*, 2008).

Lessons from the liver stage

Because the liver stage itself does not cause disease, at first glance the basic research discussed above may seem somewhat academic, but some important aspects should be considered. First of all, the pre-erythrocytic stage of *Plasmodium* is the main target for vaccine development because it is possible to achieve sterile protection by immunization with attenuated sporozoites which are still able to infect cells (Matuschewski, 2006). A better understanding of the biology of the liver stage may help to improve currently tested vaccination strategies using subunit vaccines or genetically attenuated parasites. Further study of liver stage development is also interesting in so far as most of the live attenuated parasites currently used for vaccination are blocked early in development. They invade cells and then degenerate in the host cell soon afterward (Mueller *et al.*, 2005a,b; van Dijk *et al.*, 2005). These parasites strains will express only early parasite antigens. Parasites attenuated late in liver stage development by, for example, interfering with the lipid biosynthesis (Yu *et al.*, 2008; Tarun *et al.*, 2009; Pei *et al.*, 2010) have the advantage of expressing a fuller set of proteins normally produced during the liver stage. Recently published work (Butler *et al.*, 2011) and our own preliminary data confirm that late attenuated parasites indeed induce a potent protective immune response, potentially allowing administration of much lower numbers of parasites during immunizations.

It was not expected that genetic manipulation of the parasite to interfere with lipid metabolism would not affect the blood stage at all but would be deleterious for the liver stage. How can it be that both stages express the same set of genes but expression of these genes is essential for only one stage? The most obvious difference between these parasite stages is the number of merozoites produced, up to 32 for the blood stage and up to 30 000 for the liver stage in a relatively short time. It is therefore plausible that the liver stage simply needs to produce more metabolites and other resources than the blood stage. In fact, during the fast formation of the enormous number of merozoites at the end of the liver stage, as well as in the rapid expansion of membranous organelles that precedes this, there is an extraordinary demand for membrane synthesis. If this is disturbed by interference with lipid biosynthesis, the parasite may not have enough resources to overcome this block, whereas blood stage parasites may be able to compensate for the lack by uptake of lipids from the host cell. Thus lipid biosynthesis-deficient knockout parasites are fully virulent in the blood stage

but are attenuated during the liver stage. This illustrates that the pathologically silent liver stage can be a good model to study certain pathways and processes in the parasite and can even help to understand features of development in the blood stage.

An even clearer example that the liver stage may serve as a model for certain aspects of the pathogenic blood stage is the aforementioned confusion about the order of PVM and host cell membrane breakdown during merozoite egress from RBCs (Salmon *et al.*, 2001; Wickham *et al.*, 2003). Barring that the parasite has invented two different methods to leave its host cell, it can be assumed that breakdown of the PVM precedes that of the host plasma membrane (Sturm *et al.*, 2009). In the blood stage, the rapid succession of these events greatly complicates their study. In the liver stage, however, the rupture of these two membranes is temporally clearly separated, allowing independent analysis of the molecular cascades preceding the two events. By studying these processes in the liver stage, we thus have the potential to uncover the underlying mechanisms, which are likely conserved between the two parasite stages but very difficult to study in the blood stage. The knowledge we will gain from this will ultimately support the continuing efforts to create new antimalarial drugs and vaccination strategies.

Acknowledgements

We would like to thank Nicole Struck and Kathleen Rankin for the movie demonstrating sporozoite transmigration and invasion of cells. We are also grateful to Jacqueline Schmuckli-Maurer for critical reading of the manuscript. We apologize to authors whose work we could not cite because of space limitations but who have regardless contributed greatly to our understanding of liver stage development. This work was supported by a DFG grant to V.T.H. (HE 4497/1-2) and the MALSIG and EviMalaR EU consortia.

Authors' contribution

S.G. and R.R.S. contributed equally to this work.

References

- Aikawa M, Hepler PK, Huff CG & Sprinz H (1966) The feeding mechanism of avian malarial parasites. *J Cell Biol* **28**: 355–373.
- Albuquerque SS, Carret C, Grosso AR *et al.* (2009) Host cell transcriptional profiling during malaria liver stage infection reveals a coordinated and sequential set of biological events. *BMC Genomics* **10**: 270.
- Amino R, Thiberge S, Martin B, Celli S, Shorte S, Frischknecht F & Menard R (2006) Quantitative imaging of *Plasmodium* transmission from mosquito to mammal. *Nat Med* **12**: 220–224.
- Amino R, Giovannini D, Thiberge S *et al.* (2008) Host cell traversal is important for progression of the malaria parasite through the dermis to the liver. *Cell Host Microbe* **3**: 88–96.
- Andrews NW (1990) The acid-active hemolysin of *Trypanosoma cruzi*. *Exp Parasitol* **71**: 241–244.
- Andrews NW, Abrams CK, Slatin SL & Griffiths G (1990) A *T. cruzi*-secreted protein immunologically related to the complement component C9: evidence for membrane pore-forming activity at low pH. *Cell* **61**: 1277–1287.
- Arastu-Kapur S, Ponder EL, Fonovic UP *et al.* (2008) Identification of proteases that regulate erythrocyte rupture by the malaria parasite *Plasmodium falciparum*. *Nat Chem Biol* **4**: 203–213.
- Arii S & Imamura M (2000) Physiological role of sinusoidal endothelial cells and Kupffer cells and their implication in the pathogenesis of liver injury. *J Hepatobiliary Pancreat Surg* **7**: 40–48.
- Baer K, Klotz C, Kappe SH, Schnieder T & Frevert U (2007) Release of hepatic *Plasmodium yoelii* merozoites into the pulmonary microvasculature. *PLoS Pathog* **3**: e171.
- Bakowski MA, Braun V & Brumell JH (2008) Salmonella-containing vacuoles: directing traffic and nesting to grow. *Traffic* **9**: 2022–2031.
- Bannister LH, Hopkins JM, Fowler RE, Krishna S & Mitchell GH (2000) A brief illustrated guide to the ultrastructure of *Plasmodium falciparum* asexual blood stages. *Parasitol Today* **16**: 427–433.
- Bano N, Romano JD, Jayabalasingham B & Coppens I (2007) Cellular interactions of *Plasmodium* liver stage with its host mammalian cell. *Int J Parasitol* **37**: 1329–1341.
- Bastianelli G (1948) Exo-erythrocytic forms of malaria parasite. *Br Med J* **1**: 520.
- Blackman MJ (2008) Malarial proteases and host cell egress: an 'emerging' cascade. *Cell Microbiol* **10**: 1925–1934.
- Braun JS, Hoffmann O, Schickhaus M *et al.* (2007) Pneumolysin causes neuronal cell death through mitochondrial damage. *Infect Immun* **75**: 4245–4254.
- Butcher BA, Kim L, Johnson PF & Denkers EY (2001) *Toxoplasma gondii* tachyzoites inhibit proinflammatory cytokine induction in infected macrophages by preventing nuclear translocation of the transcription factor NF-kappa B. *J Immunol* **167**: 2193–2201.
- Butler NS, Schmidt NW, Vaughan AM, Aly AS, Kappe SH & Harty JT (2011) Superior antimalarial immunity after vaccination with late liver stage-arresting genetically attenuated parasites. *Cell Host Microbe* **9**: 451–462.
- Carrolo M, Giordano S, Cabrita-Santos L *et al.* (2003) Hepatocyte growth factor and its receptor are required for malaria infection. *Nat Med* **9**: 1363–1369.
- Castedo M, Hirsch T, Susin SA, Zamzami N, Marchetti P, Macho A & Kroemer G (1996) Sequential acquisition of

- mitochondrial and plasma membrane alterations during early lymphocyte apoptosis. *J Immunol* **157**: 512–521.
- Chandramohanadas R, Davis PH, Beiting DP *et al.* (2009) Apicomplexan parasites co-opt host calpains to facilitate their escape from infected cells. *Science* **324**: 794–797.
- Charneau S, Bastos IM, Mouray E, Ribeiro BM, Santana JM, Grellier P & Florent I (2007) Characterization of PfDYN2, a dynamin-like protein of *Plasmodium falciparum* expressed in schizonts. *Microbes Infect* **9**: 797–805.
- Coppi A, Tewari R, Bishop JR *et al.* (2007) Heparan sulfate proteoglycans provide a signal to *Plasmodium* sporozoites to stop migrating and productively invade host cells. *Cell Host Microbe* **2**: 316–327.
- Coppi A, Natarajan R, Pradel G *et al.* (2011) The malaria circumsporozoite protein has two functional domains, each with distinct roles as sporozoites journey from mosquito to mammalian host. *J Exp Med* **208**: 341–356.
- Crawford MJ, Thomsen-Zieger N, Ray M, Schachtner J, Roos DS & Seeber F (2006) *Toxoplasma gondii* scavenges host-derived lipoic acid despite its de novo synthesis in the apicoplast. *EMBO J* **25**: 3214–3222.
- Daubersies P, Thomas AW, Millet P *et al.* (2000) Protection against *Plasmodium falciparum* malaria in chimpanzees by immunization with the conserved pre-erythrocytic liver-stage antigen 3. *Nat Med* **6**: 1258–1263.
- De Leeuw AM, Brouwer A & Knook DL (1990) Sinusoidal endothelial cells of the liver: fine structure and function in relation to age. *J Electron Microscop Tech* **14**: 218–236.
- Dowse TJ, Koussis K, Blackman MJ & Soldati-Favre D (2008) Roles of proteases during invasion and egress by *Plasmodium* and *Toxoplasma*. *Subcell Biochem* **47**: 121–139.
- Ejigiri I & Sinnis P (2009) *Plasmodium* sporozoite-host interactions from the dermis to the hepatocyte. *Curr Opin Microbiol* **12**: 401–407.
- Elliott DA, McIntosh MT, Hosgood HD III, Chen S, Zhang G, Baevova P & Joiner KA (2008) Four distinct pathways of hemoglobin uptake in the malaria parasite *Plasmodium falciparum*. *P Natl Acad Sci USA* **105**: 2463–2468.
- Fonseca F, Cambournac JC, Pinto MR, Pereira MR & Cunha A (1946) Studies on the exo-erythrocytic cycle of malaria. *Parasitology* **37**: 113–117.
- Frevert U, Usynin I, Baer K & Klotz C (2006) Nomadic or sessile: can Kupffer cells function as portals for malaria sporozoites to the liver? *Cell Microbiol* **8**: 1537–1546.
- Frevert U, Spath GF & Yee H (2008) Exoerythrocytic development of *Plasmodium gallinaceum* in the White Leghorn chicken. *Int J Parasitol* **38**: 655–672.
- Frischknecht F, Baldacci P, Martin B *et al.* (2004) Imaging movement of malaria parasites during transmission by *Anopheles* mosquitoes. *Cell Microbiol* **6**: 687–694.
- Gaillard JL, Berche P, Mounier J, Richard S & Sansonetti P (1987) In vitro model of penetration and intracellular growth of *Listeria monocytogenes* in the human enterocyte-like cell line Caco-2. *Infect Immun* **55**: 2822–2829.
- Gerald N, Mahajan B & Kumar S (2011) Mitosis in the human malaria parasite *Plasmodium falciparum*. *Eukaryot Cell* **10**: 474–482.
- Goldie P, Roth EF Jr, Oppenheim J & Vanderberg JP (1990) Biochemical characterization of *Plasmodium falciparum* hemozoin. *Am J Trop Med Hyg* **43**: 584–596.
- Gonzalez V, Combe A, David V *et al.* (2009) Host cell entry by apicomplexa parasites requires actin polymerization in the host cell. *Cell Host Microbe* **5**: 259–272.
- Graewe S, Rankin K, Lehmann C, Deschermeier C, Hecht L, Froehle U, Stanway RR & Heussler V (in press) Hostile takeover by *Plasmodium*: reorganization of parasite and host cell membranes during liver stage egress. *PLoS Pathog.*
- Gueirard P, Tavares J, Thiberge S *et al.* (2010) Development of the malaria parasite in the skin of the mammalian host. *P Natl Acad Sci USA* **107**: 18640–18645.
- Heussler V, Sturm A & Langsley G (2006) Regulation of host cell survival by intracellular *Plasmodium* and *Theileria* parasites. *Parasitology* **132** (Suppl): S49–S60.
- Heussler V, Rennenberg A & Stanway R (2010) Host cell death induced by the egress of intracellular *Plasmodium* parasites. *Apoptosis* **15**: 376–385.
- Hippe D, Weber A, Zhou L, Chang DC, Hacker G & Luder CG (2009) *Toxoplasma gondii* infection confers resistance against BimS-induced apoptosis by preventing the activation and mitochondrial targeting of pro-apoptotic Bax. *J Cell Sci* **122**: 3511–3521.
- Hybiske K & Stephens RS (2007) Mechanisms of host cell exit by the intracellular bacterium *Chlamydia*. *P Natl Acad Sci USA* **104**: 11430–11435.
- Ishino T, Yano K, Chinzei Y & Yuda M (2004) Cell-passage activity is required for the malarial parasite to cross the liver sinusoidal cell layer. *PLoS Biol* **2**: E4.
- Ishino T, Boisson B, Orito Y *et al.* (2009) LISP1 is important for the egress of *Plasmodium berghei* parasites from liver cells. *Cell Microbiol* **11**: 1329–1339.
- Kafsack BF, Pena JD, Coppens I, Ravindran S, Boothroyd JC & Carruthers VB (2009) Rapid membrane disruption by a perforin-like protein facilitates parasite exit from host cells. *Science* **323**: 530–533.
- Kaiser K, Camargo N, Coppens I, Morrisey JM, Vaidya AB & Kappe SH (2004) A member of a conserved *Plasmodium* protein family with membrane-attack complex/perforin (MACPF)-like domains localizes to the micronemes of sporozoites. *Mol Biochem Parasitol* **133**: 15–26.
- Kasinathan RS, Foller M, Koka S, Huber SM & Lang F (2007) Inhibition of eryptosis and intraerythrocytic growth of *Plasmodium falciparum* by flufenamic acid. *Naunyn Schmiedebergs Arch Pharmacol* **374**: 255–264.
- Kozjak-Pavlovic V, Dian-Lothrop EA, Meinecke M *et al.* (2009) Bacterial porin disrupts mitochondrial membrane potential and sensitizes host cells to apoptosis. *PLoS Pathog* **5**: e1000629.
- Kroemer G & Reed JC (2000) Mitochondrial control of cell death. *Nat Med* **6**: 513–519.

- Kudryashev M, Lepper S, Stanway R, Bohn S, Baumeister W, Cyrklaff M & Frischknecht F (2010) Positioning of large organelles by a membrane-associated cytoskeleton in *Plasmodium* sporozoites. *Cell Microbiol* **12**: 362–371.
- Leiriao P, Albuquerque SS, Corso S *et al.* (2005) HGF/MET signalling protects *Plasmodium*-infected host cells from apoptosis. *Cell Microbiol* **7**: 603–609.
- Lemasters JJ (2005) Dying a thousand deaths: redundant pathways from different organelles to apoptosis and necrosis. *Gastroenterology* **129**: 351–360.
- Marti M, Good RT, Rug M, Knuepfer E & Cowman AF (2004) Targeting malaria virulence and remodeling proteins to the host erythrocyte. *Science* **306**: 1930–1933.
- Martin SJ, Finucane DM, Amarante-Mendes GP, O'Brien GA & Green DR (1996) Phosphatidylserine externalization during CD95-induced apoptosis of cells and cytoplasts requires ICE/CED-3 protease activity. *J Biol Chem* **271**: 28753–28756.
- Matuschewski K (2006) Vaccine development against malaria. *Curr Opin Immunol* **18**: 449–457.
- Mazier D, Beaudoin RL, Mellouk S *et al.* (1985) Complete development of hepatic stages of *Plasmodium falciparum* *in vitro*. *Science* **227**: 440–442.
- Medica DL & Sinnis P (2005) Quantitative dynamics of *Plasmodium yoelii* sporozoite transmission by infected anopheline mosquitoes. *Infect Immun* **73**: 4363–4369.
- Meis JF, Verhave JP, Jap PH, Hess F & Meuwissen JH (1981) An ultrastructural study of developing stages of exoerythrocytic *Plasmodium berghei* in rat hepatocytes. *Parasitology* **82**: 195–204.
- Meis JF, Verhave JP, Jap PH & Meuwissen JH (1983) An ultrastructural study on the role of Kupffer cells in the process of infection by *Plasmodium berghei* sporozoites in rats. *Parasitology* **86** (Pt 2): 231–242.
- Meis JF, Verhave JP, Brouwer A & Meuwissen JH (1985a) Electron microscopic studies on the interaction of rat Kupffer cells and *Plasmodium berghei* sporozoites. *Z Parasitenkd* **71**: 473–483.
- Meis JF, Verhave JP, Jap PH & Meuwissen JH (1985b) Fine structure of exoerythrocytic merozoite formation of *Plasmodium berghei* in rat liver. *J Protozool* **32**: 694–699.
- Meis JF, Rijntjes PJ, Verhave JP, Ponnudurai T, Hollingdale MR & Yap SH (1985c) Infection of cryopreserved adult human hepatocytes with *Plasmodium falciparum* sporozoites. *Cell Biol Int Rep* **9**: 976.
- Menard R (2000) The journey of the malaria sporozoite through its hosts: two parasite proteins lead the way. *Microbes Infect* **2**: 633–642.
- Molestina RE, Payne TM, Coppens I & Sinai AP (2003) Activation of NF- κ B by *Toxoplasma gondii* correlates with increased expression of antiapoptotic genes and localization of phosphorylated I κ B to the parasitophorous vacuole membrane. *J Cell Sci* **116**: 4359–4371.
- Mota MM, Pradel G, Vanderberg JP *et al.* (2001) Migration of *Plasmodium* sporozoites through cells before infection. *Science* **291**: 141–144.
- Mota MM, Hafalla JC & Rodriguez A (2002) Migration through host cells activates *Plasmodium* sporozoites for infection. *Nat Med* **8**: 1318–1322.
- Moudy R, Manning TJ & Beckers CJ (2001) The loss of cytoplasmic potassium upon host cell breakdown triggers egress of *Toxoplasma gondii*. *J Biol Chem* **276**: 41492–41501.
- Mueller AK, Labaied M, Kappe SH & Matuschewski K (2005a) Genetically modified *Plasmodium* parasites as a protective experimental malaria vaccine. *Nature* **433**: 164–167.
- Mueller AK, Camargo N, Kaiser K, Andorfer C, Frevert U, Matuschewski K & Kappe SH (2005b) *Plasmodium* liver stage developmental arrest by depletion of a protein at the parasite-host interface. *P Natl Acad Sci USA* **102**: 3022–3027.
- Muller WA (2010) Mechanisms of leukocyte transendothelial migration. *Annu Rev Pathol* **6**: 323–344.
- Nagamune K, Moreno SN, Chini EN & Sibley LD (2008) Calcium regulation and signaling in apicomplexan parasites. *Subcell Biochem* **47**: 70–81.
- Nishi M, Hu K, Murray JM & Roos DS (2008) Organellar dynamics during the cell cycle of *Toxoplasma gondii*. *J Cell Sci* **121**: 1559–1568.
- Noronha FS, Cruz JS, Beirao PS & Horta MF (2000) Macrophage damage by *Leishmania amazonensis* cytolysin: evidence of pore formation on cell membrane. *Infect Immun* **68**: 4578–4584.
- Padmanaban G, Nagaraj VA & Rangarajan PN (2007) An alternative model for heme biosynthesis in the malarial parasite. *Trends Biochem Sci* **32**: 443–449.
- Pandey KC, Singh N, Arastu-Kapur S, Bogoy M & Rosenthal PJ (2006) Falciparum, a cysteine protease inhibitor of *Plasmodium falciparum*, facilitates erythrocyte invasion. *PLoS Pathog* **2**: e117.
- Pei Y, Tarun AS, Vaughan AM, Herman RW, Soliman JM, Erickson-Wayman A & Kappe SH (2010) *Plasmodium* pyruvate dehydrogenase activity is only essential for the parasite's progression from liver infection to blood infection. *Mol Microbiol* **75**: 957–971.
- Perla BL, Zapata C, Valencia AZ *et al.* (2003) Immunogenicity and protective efficacy of *Plasmodium falciparum* liver-stage Ag-3 in Aotus lemurinus griseimembra monkeys. *Eur J Immunol* **33**: 1321–1327.
- Pradel G & Frevert U (2001) Malaria sporozoites actively enter and pass through rat Kupffer cells prior to hepatocyte invasion. *Hepatology* **33**: 1154–1165.
- Prudencio M, Rodriguez A & Mota MM (2006) The silent path to thousands of merozoites: the *Plasmodium* liver stage. *Nat Rev Microbiol* **4**: 849–856.
- Rennenberg A, Lehmann C, Heitmann A *et al.* (2010) Exoerythrocytic *Plasmodium* parasites secrete a cysteine protease inhibitor involved in sporozoite invasion and capable of blocking cell death of host hepatocytes. *PLoS Pathog* **6**: e1000825.
- Rodrigues CD, Hannus M, Prudencio M *et al.* (2008) Host scavenger receptor SR-BI plays a dual role in the establishment of malaria parasite liver infection. *Cell Host Microbe* **4**: 271–282.

- Rosenberg R, Wirtz RA, Schneider I & Burge R (1990) An estimation of the number of malaria sporozoites ejected by a feeding mosquito. *Trans R Soc Trop Med Hyg* **84**: 209–212.
- Rosenthal PJ & Meshnick SR (1996) Hemoglobin catabolism and iron utilization by malaria parasites. *Mol Biochem Parasitol* **83**: 131–139.
- Salmon BL, Oksman A & Goldberg DE (2001) Malaria parasite exit from the host erythrocyte: a two-step process requiring extraerythrocytic proteolysis. *P Natl Acad Sci USA* **98**: 271–276.
- Sato S, Clough B, Coates L & Wilson RJ (2004) Enzymes for heme biosynthesis are found in both the mitochondrion and plastid of the malaria parasite *Plasmodium falciparum*. *Protist* **155**: 117–125.
- Savill J (1998) Apoptosis. Phagocytic docking without shocking. *Nature* **392**: 442–443.
- Scaffidi P, Misteli T & Bianchi ME (2002) Release of chromatin protein HMGB1 by necrotic cells triggers inflammation. *Nature* **418**: 191–195.
- Schmidt-Christensen A, Sturm A, Horstmann S & Heussler VT (2008) Expression and processing of *Plasmodium berghei* SERA3 during liver stages. *Cell Microbiol* **10**: 1723–1734.
- Schuerch DW, Wilson-Kubalek EM & Tweten RK (2005) Molecular basis of listeriolysin O pH dependence. *P Natl Acad Sci USA* **102**: 12537–12542.
- Shapira S, Speirs K, Gerstein A, Caamano J & Hunter CA (2002) Suppression of NF-kappaB activation by infection with *Toxoplasma gondii*. *J Infect Dis* **185** (Suppl 1): S66–S72.
- Shi Y, Evans JE & Rock KL (2003) Molecular identification of a danger signal that alerts the immune system to dying cells. *Nature* **425**: 516–521.
- Shin SC, Vanderberg JP & Terzakis JA (1982) Direct infection of hepatocytes by sporozoites of *Plasmodium berghei*. *J Protozool* **29**: 448–454.
- Shortt HE & Garnham PC (1948) Pre-erythrocytic stage in mammalian malaria parasites. *Nature* **161**: 126.
- Silvie O, Rubinstein E, Franetich JF *et al.* (2003) Hepatocyte CD81 is required for *Plasmodium falciparum* and *Plasmodium yoelii* sporozoite infectivity. *Nat Med* **9**: 93–96.
- Silvie O, Greco C, Franetich JF *et al.* (2006) Expression of human CD81 differently affects host cell susceptibility to malaria sporozoites depending on the *Plasmodium* species. *Cell Microbiol* **8**: 1134–1146.
- Silvie O, Franetich JF, Boucheix C, Rubinstein E & Mazier D (2007) Alternative invasion pathways for *Plasmodium berghei* sporozoites. *Int J Parasitol* **37**: 173–182.
- Singh AP, Buscaglia CA, Wang Q *et al.* (2007) *Plasmodium* circumsporozoite protein promotes the development of the liver stages of the parasite. *Cell* **131**: 492–504.
- Sinnis P & Nardin E (2002) Sporozoite antigens: biology and immunology of the circumsporozoite protein and thrombospondin-related anonymous protein. *Chem Immunol* **80**: 70–96.
- Sinnis P & Zavala F (2008) The skin stage of malaria infection: biology and relevance to the malaria vaccine effort. *Future Microbiol* **3**: 275–278.
- Soni S, Dhawan S, Rosen KM, Chafel M, Chishti AH & Hanspal M (2005) Characterization of events preceding the release of malaria parasite from the host red blood cell. *Blood Cells Mol Dis* **35**: 201–211.
- Spielmann T & Gilberger TW (2010) Protein export in malaria parasites: do multiple export motifs add up to multiple export pathways? *Trends Parasitol* **26**: 6–10.
- Stanway RR, Witt T, Zobiak B, Aepfelbacher M & Heussler VT (2009a) GFP-targeting allows visualization of the apicoplast throughout the life cycle of live malaria parasites. *Biol Cell* **101**: 415–430.
- Stanway RR, Graewe S, Rennenberg A, Helm S & Heussler VT (2009b) Highly efficient subcloning of rodent malaria parasites by injection of single merozoites or detached cells. *Nat Protoc* **4**: 1433–1439.
- Stanway RR, Mueller N, Zobiak B, Graewe S, Froehlke U, Zessin P, Aepfelbacher M & Heussler VT (in press) Organelle segregation into *Plasmodium* liver stage merozoites. *Cell Microbiol*.
- Striepen B, Crawford MJ, Shaw MK, Tilney LG, Seeber F & Roos DS (2000) The plastid of *Toxoplasma gondii* is divided by association with the centrosomes. *J Cell Biol* **151**: 1423–1434.
- Struck NS, de Souza Dias S, Langer C, Marti M, Pearce JA, Cowman AF & Gilberger TW (2005) Re-defining the Golgi complex in *Plasmodium falciparum* using the novel Golgi marker PfGRASP. *J Cell Sci* **118**: 5603–5613.
- Sturm A, Amino R, van de Sand C *et al.* (2006) Manipulation of host hepatocytes by the malaria parasite for delivery into liver sinusoids. *Science* **313**: 1287–1290.
- Sturm A, Graewe S, Franke-Fayard B *et al.* (2009) Alteration of the parasite plasma membrane and the parasitophorous vacuole membrane during exo-erythrocytic development of malaria parasites. *Protist* **160**: 51–63.
- Tarun AS, Peng X, Dumpit RF *et al.* (2008) A combined transcriptome and proteome survey of malaria parasite liver stages. *P Natl Acad Sci USA* **105**: 305–310.
- Tarun AS, Vaughan AM & Kappe SH (2009) Redefining the role of de novo fatty acid synthesis in *Plasmodium* parasites. *Trends Parasitol* **25**: 545–550.
- Terzakis JA, Vanderberg JP, Foley D & Shustak S (1979) Exoerythrocytic merozoites of *Plasmodium berghei* in rat hepatic Kupffer cells. *J Protozool* **26**: 385–389.
- Vaishnav S & Striepen B (2006) The cell biology of secondary endosymbiosis – how parasites build, divide and segregate the apicoplast. *Mol Microbiol* **61**: 1380–1387.
- Vaishnav S, Morrison DP, Gaji RY, Murray JM, Entzeroth R, Howe DK & Striepen B (2005) Plastid segregation and cell division in the apicomplexan parasite *Sarcocystis neurona*. *J Cell Sci* **118**: 3397–3407.
- Valderramos SG & Fidock DA (2006) Transporters involved in resistance to antimalarial drugs. *Trends Pharmacol Sci* **27**: 594–601.
- van de Sand C, Horstmann S, Schmidt A *et al.* (2005) The liver stage of *Plasmodium berghei* inhibits host cell apoptosis. *Mol Microbiol* **58**: 731–742.

- van Dijk MR, Douradinha B, Franke-Fayard B *et al.* (2005) Genetically attenuated, P36p-deficient malarial sporozoites induce protective immunity and apoptosis of infected liver cells. *P Natl Acad Sci USA* **102**: 12194–12199.
- van Dooren GG, Marti M, Tonkin CJ, Stimmler LM, Cowman AF & McFadden GI (2005) Development of the endoplasmic reticulum, mitochondrion and apicoplast during the asexual life cycle of *Plasmodium falciparum*. *Mol Microbiol* **57**: 405–419.
- van Dooren GG, Stimmler LM & McFadden GI (2006) Metabolic maps and functions of the *Plasmodium* mitochondrion. *FEMS Microbiol Rev* **30**: 596–630.
- van Dooren GG, Reiff SB, Tomova C, Meissner M, Hummel BM & Striepen B (2009) A novel dynamin-related protein has been recruited for apicoplast fission in *Toxoplasma gondii*. *Curr Biol* **19**: 267–276.
- Vanderberg JP & Frevert U (2004) Intravital microscopy demonstrating antibody-mediated immobilisation of *Plasmodium berghei* sporozoites injected into skin by mosquitoes. *Int J Parasitol* **34**: 991–996.
- White NJ (2008) *Plasmodium knowlesi*: the fifth human malaria parasite. *Clin Infect Dis* **46**: 172–173.
- Wickham ME, Culvenor JG & Cowman AF (2003) Selective inhibition of a two-step egress of malaria parasites from the host erythrocyte. *J Biol Chem* **278**: 37658–37663.
- Xu Q, Abdubek P, Astakhova T *et al.* (2010) Structure of a membrane-attack complex/perforin (MACPF) family protein from the human gut symbiont *Bacteroides thetaiotaomicron*. *Acta Crystallogr Sect F Struct Biol Cryst Commun* **66**: 1297–1305.
- Yalaoui S, Huby T, Franetich JF *et al.* (2008) Scavenger receptor BI boosts hepatocyte permissiveness to *Plasmodium* infection. *Cell Host Microbe* **4**: 283–292.
- Yeoh S, O'Donnell RA, Koussis K *et al.* (2007) Subcellular discharge of a serine protease mediates release of invasive malaria parasites from host erythrocytes. *Cell* **131**: 1072–1083.
- Yu M, Kumar TR, Nkrumah LJ *et al.* (2008) The fatty acid biosynthesis enzyme FabI plays a key role in the development of liver-stage malarial parasites. *Cell Host Microbe* **4**: 567–578.
- Zhao YO, Khaminets A, Hunn JP & Howard JC (2009) Disruption of the *Toxoplasma gondii* parasitophorous vacuole by IFN γ -inducible immunity-related GTPases (IRG proteins) triggers necrotic cell death. *PLoS Pathog* **5**: e1000288.
- Zou H, Li Y, Liu X & Wang X (1999) An APAF-1/cytochrome c multimeric complex is a functional apoptosome that activates procaspase-9. *J Biol Chem* **274**: 11549–11556.

Supporting information

Additional Supporting information may be found in the online version of this article:

Movie S1. Transmigrating sporozoite. Red fluorescent *Plasmodium berghei* sporozoites transmigrate through a HepG2 cell expressing GFP targeted to the plasma membrane.

Movie S2. Transmigrating sporozoite. Same setup as in Movie S1. Note that the parasite transmigrates from one cell to another.

Movie S3. Invading sporozoite. Same setup as in Movie S1. Note that the host cell plasma membrane tightly surrounds the invading parasite.

Movie S4. Invading sporozoite. Red fluorescent *Plasmodium berghei* sporozoites transmigrate through a HepG2 cell expressing GFP in the cytoplasm. Note the constriction of the sporozoite when entering the green host cell.

Movie S5. Invading sporozoite. Same setup as in Movie S1. Note the considerable modification of the host cell surface upon invasion.

Movie S6. Upon invasion the sporozoite localizes close to the nucleus and develops into an trophozoite. Same setup as in Movie S4. Note the position of the parasite in respect to the host cell nucleus.

Movie S7. *Plasmodium* exo-erythrocytic schizont develops into the cytomere stage and finally into thousands of merozoites. Same setup as in Movie S4.

Movie S8. *Plasmodium* cytomere development and PVM breakdown. Same setup as in Movie S4. Upon merozoite formation, the PVM breaks down and merozoites are liberated into the host cell cytoplasm. Note that the merozoites become motile upon PVM breakdown.

Movie S9. *Plasmodium* merozoite. Single phase contrast images of the same merozoite were recorded every second and combined to a movie. Note the motile merozoites inside the vesicle.

Please note: Wiley-Blackwell is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.