J. J. Hanley · J. E. Mungall · T. Pettke
E. T. C. Spooner · C. J. Bray

Ore metal redistribution by hydrocarbon–brine and hydrocarbon–halide melt phases, North Range footwall of the Sudbury Igneous Complex, Ontario, Canada

Received: 5 October 2004 / Accepted: 10 June 2005 / Published online: 9 August 2005
© Springer-Verlag 2005

Abstract We report methane-dominant hydrocarbon (fluid) inclusions ($\text{CH}_4 \pm \text{C}_2\text{H}_6\text{--C}_2\text{H}_2$, C_3H_8) coexisting with primary brine inclusions and secondary halide-melt (solid NaCl) inclusions in Au–Pt-rich quartz-sulfide-epidote alteration veins associated with the footwall-style Cu–PGE (platinum-group element)–Au deposits at the Fraser Mine (North Range of the Sudbury Igneous Complex). Evidence for coentrapment of immiscible hydrocarbon–brine, and hydrocarbon–halide melt mixtures is demonstrated. A primary CH_4 –brine assemblage was trapped during quartz growth at relatively low T (min. $T_{\text{trapping}} \sim 145\text{--}315^\circ\text{C}$) and P (max. $P_{\text{trapping}} \sim 500$ bar), prior to the crystallization of sulfide minerals in the veins. Secondary inclusions contain solid halite and a mixture of CH_4 , $\text{C}_2\text{H}_6\text{--C}_2\text{H}_2$ and C_3H_8 and were trapped at a minimum T of $\sim 710^\circ\text{C}$. The halite inclusions may represent halide-melt that exsolved from crystallizing sulfide ores that texturally postdate (by replacement) early alteration quartz hosting the primary, lower T brine– CH_4 assemblage. Laser ablation ICP-MS analyses show that the brine, hydrocarbon and halide-melt inclusions contain significant concentrations of Cu (0.1–1 wt% range), Au, Bi, Ag and Pt (all 0.1–10 ppm range). Cu:Pt and Cu:Au ratios in the inclusions are significantly (up to 4 log units) lower than in the host alter-

ation veins and adjacent massive sulfide ore veins, suggesting either (1) early Cu loss from the volatiles by chalcopyrite precipitation or (2) enhanced Au and Pt solubilities relative to Cu at the temperatures of entrapment. Concentration ratios between coexisting brine and CH_4 inclusions $[C_{\text{brine}}^{\text{elem}}/C_{\text{CH}_4}^{\text{elem}}]$ are lower for Cu, Au, Bi and Ag than for other elements (Na, Ca, Fe, Mn, Zn, Pb) indicating that during interaction with the brine, the hydrocarbon phase was enriched in ore metals. The high concentrations of ore metals in hydrocarbon, brine and halide-melt phases confirm that both aqueous and non-aqueous volatiles were carriers of precious metals in the Sudbury environment over a wide range of temperatures. Volatile evolution and magmatic sulfide differentiation were clearly part of a single, continuous process in the Sudbury footwall. The exsolution of H_2O -poor volatiles from fractionated sulfide liquid may have been a principal mechanism controlling the final distribution of PGE and Au in the footwall ore systems. The study reports the first measurements of precious metal concentrations in fluid inclusions from a magmatic Ni–Cu–PGE environment (the Sudbury district).

Keywords Hydrocarbons · Magmatic sulfide · Platinum-group elements · Gold · Brines

Electronic Supplementary Material Supplementary material is available for this article at <http://dx.doi.org/10.1007/s00126-005-0004-z>

Editorial handling: P. Lightfoot

J. J. Hanley (✉) · J. E. Mungall · E. T. C. Spooner · C. J. Bray
Department of Geology, University of Toronto,
22 Russell Street, Toronto, ON,
Canada M5S 3B1
E-mail: hanley@geology.utoronto.ca
Tel.: +1-416-9260857
Fax: +1-416-9783938

T. Pettke
Isotope Geology and Mineral Resources, ETH Zürich,
ETH Zentrum NO, 8092 Zürich, Switzerland

Introduction

The segregation of ore metals (e.g., Cu, Au) into an aqueous-carbonic vapor phase that separates from boiling NaCl-rich magmatic fluid been documented in many porphyry Cu–Au systems (Heinrich et al. 1992; Mavrogenes et al. 2002; Ulrich et al. 2001). Selective partitioning of Au and Cu into the lower density fluid phase may be facilitated by the volatilization of sulfide-bearing [e.g., $\text{Au}(\text{HS})_2^-$] rather than chloride-bearing complexes (Renders and Seward 1989; Hemley et al. 1992; Heinrich et al. 1999; Mountain and Seward 1999). Fluid inclusion evidence from unconformity-related U

deposits (Ypma and Fuzikawa 1980), REE-bearing peralkaline granites (Salvi and Williams-Jones 1992), mafic-ultramafic layered intrusions (Ballhaus and Stumpfl 1986) and high-grade calcareous metasediments (Sisson et al. 1981) confirm that aqueous-carbonic fluid immiscibility can occur over a wide range of geological P - T conditions. Additionally, aqueous-carbonic fluid immiscibility has been recognized as an important mechanism for Au deposition in Archean to Cenozoic vein Au systems (Naden and Shepherd 1989; Pettke et al. 2000). However, direct evidence for ore metal partitioning preferentially into the low-density fluid phase has not been documented in systems other than Cu-Au porphyry deposits.

In this study, we (1) describe previously unreported hydrocarbon fluid inclusions coexisting immiscibly with primary brine inclusions and secondary halide-melt inclusions in quartz from Au-PGE-rich quartz-sulfide-epidote alteration veins associated with footwall Cu-Ni-platinum-group element (PGE) ore deposits at the Fra-

ser Mine near Sudbury, Ontario and (2) demonstrate that hydrocarbon, brine and halide-melt phases segregated significant amounts of Cu and precious metals. Consequently, these volatile phases must be considered important agents for metal redistribution in the Sudbury footwall. This study presents the first in-situ measurements of precious metal concentrations in volatile-bearing inclusions from a magmatic Ni-Cu-PGE environment.

Geological setting

Sulfide deposits at the Fraser Mine (Falconbridge Ltd.) occur along the northern margin of the Sudbury Igneous Complex (SIC) (Fig. 1a, b). The SIC is a large, differentiated, hypabyssal igneous body composed of noritic to gabbroic cumulates overlain by residual granophyre. The SIC was produced by meteorite impact-induced melting of carbonaceous shale and lower crustal rocks at

Fig. 1 Geological setting of the footwall-style Cu-PGE ores and associated alteration veins at the Fraser Mine

(Falconbridge Ltd.). **a** Generalized regional geology and location of the Fraser Mine on the North Range of the Sudbury Igneous Complex (SIC), Ontario, Canada. **b** Plan view of contact-style (Ni) ore bodies (hosted in footwall breccia at the base of the SIC) and footwall-style Cu-PGE ore bodies (hosted in Sudbury Breccia within the gneissic footwall; cross-cuts at the Fraser Mine projected to the 762 m (25) level. The study area comprises the deeper footwall ore zones 37–39. **c** Enlarged area from Fig. 1b showing a drift layout (plan-view) of a footwall-style sulfide vein network hosted in Sudbury Breccia. **d** Enlargement of the tip of one of the sulfide veins showing the location of quartz-sulfide-epidote veins in their typical setting. The quartz-sulfide-epidote veins occur as splays or bifurcations off the main sulfide vein

1,850 ± 1 Ma (Dietz 1964; Krogh et al. 1984; Grieve 1994; Mungall et al. 2004). At the Fraser Mine, the SIC overlies partially molten and plastically deformed target rocks (“footwall breccia”; Greenman 1970). The footwall breccia overlies impact-brecciated, contact-metamorphosed Archean rocks of the Superior province (the Levack Gneiss complex; Coats and Snajdr 1984; Dressler 1984; Farrow 1994).

There are three major (widely distributed) magmatic sulfide deposit types associated with the SIC (cf. Farrow and Lightfoot 2002): (1) contact-style deposits (high Ni, low PGE) occurring at the base of the SIC within footwall breccia or a basal igneous unit (“sublayer”) of the SIC; (2) offset-type deposits hosted in radial or concentric quartz diorite “offset” dykes that are physically connected to the main igneous mass of the SIC (Fig. 1a); and (3) footwall-style deposits (high Cu, high PGE), hosted entirely within zones of Sudbury Breccia in the footwall rocks. At the Fraser mine, contact-style and footwall-style deposits occur within a structurally complex depression or “embayment” in the footwall (Fig. 1a, b; Morrison et al. 1994). Mineralized embayments are common along the northern and southern margins of the SIC. Contact-style mineralization was introduced into the embayments along the lower contact of the SIC during sulfide saturation and differentiation of the SIC magma. The contact-style deposits consist of lenses or blocks of sulfide-rich footwall breccia. Sulfides comprise the matrix of the footwall breccia and consist of disseminated to massive pyrrhotite–pentlandite–chalcocopyrite with minor magnetite and pyrite. The Cu/Ni ratio of contact-style ores are < 1; Pt, Pd and Au contents are generally low but may increase locally up to several g/t, and towards the footwall contact (Naldrett and Pessaran 1992). The contact-style deposits have a accepted magmatic origin.

The footwall-style deposits are widely distributed and occur up to 2 km from the SIC-footwall contact in zones of Sudbury Breccia within the Levack Gneiss Complex (Fig. 1b, c). They comprise veins (up to 1.5 m thick) of massive chalcocopyrite–cubanite (Cu content up to 27 wt%) with minor pentlandite, millerite, bornite and magnetite. Footwall-style deposits are small in comparison to contact deposits but have considerable value owing to their high Cu, Pt, Pd and Au contents. Unlike the contact-style ores, the origin of the footwall-style deposits is controversial. In the magmatic hypothesis, magmatic differentiation of sulfide liquid at the lower contact of the SIC resulted in the formation of a residual sulfide liquid phase enriched in Cu, Pt, Pd and Au which settled into zones of brecciated country rock to form stockworks of massive sulfide veins containing Pt, Pd and Au at concentrations in the 1–10 ppm range (Naldrett and Kullerud 1967; Cowan 1968; Li 1993; Li and Naldrett 1994; Morrison et al. 1994; Ebel and Naldrett 1997). Geochemical modeling of the ore compositions indicate a base and precious metal zonation within the footwall-style deposits that is

generally consistent with sulfide fractionation (Li 1993). Physical connections between the footwall-style ores and contact-style ores are observed, for example, at the McCreedy West mine on the northern margin of the SIC; this observation supports the magmatic model for footwall-style deposit formation (Farrow and Lightfoot 2002). However, in several footwall-style deposits, Au (and to a lesser extent, Pt and Pd) distribution within the footwall-style ore zones is difficult to predict rigorously using traditional models of sulfide liquid fractionation (Naldrett and Pessaran 1992; Li and Naldrett 1994; Li 1993). For example, at the Fraser Mine, Cu:Au ratios in footwall-style sulfide samples can vary by three orders of magnitude within single veins; this variation cannot be explained by magmatic fractionation processes within the vein during emplacement (Li 1993). Unlike Pt and Pd, Au concentrations tend to increase with decreasing S content in the ores and are highest in disseminated sulfide haloes surrounding massive sulfide veins, and notably, quartz-sulfide-epidote alteration veins adjacent to the ores (Li 1993; Farrow 1994).

In the hydrothermal hypothesis, Farrow (1994) originally proposed that high salinity hydrothermal fluids (driven by the heat of the cooling SIC) leached Cu, Pt, Pd and Au from contact-style mineralization and re-deposited the metals in the footwall. This model has considerable merit and is supported by fluid inclusion, structural and mineralogical evidence that implicates a complex hydrothermal history in the footwall involving primary and secondary Cl-rich brines. Major points of evidence supporting a hydrothermal model for the ores include: (1) the occurrence of high salinity fluid inclusions in both sulfides and silicate alteration minerals that contain ore metals (e.g., Ag, Cu; Farrow 1994); (2) the occurrence of metal halide minerals within the PGE–Au-rich sulfide vein assemblages (e.g., an unnamed Pd–Bi–Cl phase; Li 1993); (3) the occurrence of primarily low temperature Pt and Pd minerals (e.g., < 600°C; tellurides, bismuthide, stannides) hosted in chlorine-rich alteration assemblages along the edges of massive sulfide veins rather than in the massive sulfide veins themselves (Li 1993; Farrow 1994); and (4) the presence of oxidized and intensely altered (?Cu–Au–Pt–Pd-leached) zones of high Ni-content ores along the SIC-footwall contact (“epidote-zone” mineralization; Farrow 1994). The most comprehensive fluid inclusion study in support of the hydrothermal model was conducted by Molnar et al. (2001). A summary of the major contributions to the hydrothermal model can be found in Table 1; these will be referred to in light of our new observations below.

Although at present most researchers have generally accepted that the emplacement of fractionated sulfide liquids initially concentrate the PGE and Au in the footwall setting, a late hydrothermal component was clearly responsible for significant compositional modification of magmatic sulfides, redistribution of the base

Table 1 Principle hydrothermal studies in the footwall deposits along the northern margin of the SIC

Principle references	Key observations	Reference's major conclusions
Li (1993) Li and Naldrett (1994)	<ul style="list-style-type: none"> -Zoned metal tenor in footwall ore zones -Ca-Na-Cl-rich fluid inclusions in alteration veins associated with sulfide ores with highly variable homogenization temperatures -Cl-rich alteration selvages on sulfide vein margins -<i>Low-T</i> PGE-Au-Ag-Sn-Te-Bi ore assemblages in textural equilibrium with Pd-Bi-chlorides and hosted within Cl-rich alteration selvages 	<ul style="list-style-type: none"> -Primary magmatic control on ore formation -Sulfide liquid fractionation in footwall to produce zonation -Exsolution of poorly equilibrated (with host rocks) Cl-rich brine from sulfides as they approached their solidus -<i>Locally</i>, migration of Au and PGE into wall rocks and alteration veins (no more than metre-scale mobility) -Cl-rich fluid exsolution controlled final deposition of PGE and Au; mobilization of Au more extensive than PGE
Farrow and Watkinson (1992) Farrow (1994)	<ul style="list-style-type: none"> -Ca-Na-Fe-K-Ba-Pb-Mn-Cl-rich fluid inclusions in sulfides and associated alteration assemblages; fluids are Ca-dominant -Cl-rich alteration selvages on sulfide vein margins -<i>Low-T</i> PGE-Sn-Te-Bi assemblages 	<ul style="list-style-type: none"> -Primary hydrothermal control on ore formation -Convection of Canadian Shield saline groundwaters under the Sudbury Complex within Sudbury Breccia zones -Local rock compositions control fluid chemistry, or fluids were not homogeneous (poorly mixed); multiple pulses of ore fluid entered footwall setting -Brine transport of Cu, Au and PGE from contact ores to footwall setting; reprecipitation of ore metals and formation of Cl-rich alteration assemblages
Marshall et al. (1999)	<ul style="list-style-type: none"> -Ore metals <i>in</i> fluid inclusions (daughter minerals) -oxidized and intensely altered, Cu-PGE-Au-leached contact ore zones ("epidote" zones) -Metal chloride minerals in sulfide (e.g., FeCl₂) 	<ul style="list-style-type: none"> -Mixing of exsolved fluid from the SIC with groundwaters
Molnar et al. (2001)	<ul style="list-style-type: none"> -Alteration minerals from sulfide vein selvages have stable isotope ratios between the "magmatic water box" and Canadian Shield groundwater compositional fields -Three major fluid inclusion assemblages: <ol style="list-style-type: none"> (1) primary, high salinity brines (Ca-Na-Fe-K-Ba-Pb-Mn-Cl); hosted in footwall sulfide and alteration veins and early, high temperature magmatic residua; long evolutionary history (2) secondary aqueous-CO₂ inclusions (3) late secondary brines (Ca-Na-Cl); associated with secondary sulfides in late alteration veins -<i>Low-T</i> PGE mineral assemblages (400–575°C) 	<ul style="list-style-type: none"> -Principle fluid inclusion study -Primary fluid (~50 wt% NaCl eq.) exsolved from magmatic residua and interacted with primary magmatic sulfides, redepositing PGE and Cu in the footwall setting as massive sulfide veins -Secondary boiling due to uplift related to the Penokean Orogeny (1,850–1,700 Ma); produced immiscible aqueous-CO₂ assemblage -Late brine circulation remobilized metals (locally)

and precious metals and the formation of a halogen-enriched alteration halo around the ores.

Materials and methods

Fluid inclusion analysis was conducted on samples obtained from underground exposures of typical footwall-style Cu–PGE ore in Zone 38 of the Fraser Mine (Fig. 1; Falconbridge Ltd). The samples consisted of fragments of quartz-sulfide-epidote “alteration” veins which splay or bifurcate from massive chalcopyrite–cubanite veins (Fig. 1c, d). Although volumetrically insignificant when compared to the massive sulfide veins, the alteration veins sampled for this study are widespread in footwall-style deposits and have been documented at all footwall-style deposits along the northern margin of the SIC (Li 1993; Farrow 1994; Molnar et al. 2001). The alteration veins are rich in PGE and Au; they may contain up to 15 ppm Au, 7 ppm Pd and 5 ppm Pt on a bulk rock basis (Hanley 2002). The mineralogical and fluid inclusion characteristics of the quartz-bearing veins sampled are described below.

All fluid inclusion analysis was performed on quartz-hosted inclusions. Microthermometry was carried out on microscope-mounted Linkam THMS600 heating-cooling and Leitz 1350 high temperature stages at the Department of Geology, University of Toronto. Uncertainties associated with microthermometric measurements are $\pm 5^\circ\text{C}$ for $T_s > 560^\circ\text{C}$, $\pm 0.1^\circ\text{C}$ for T_s between -95 and 0°C and $\pm 1^\circ\text{C}$ for T_s between 0 and 560°C .

Hydrocarbon species in fluid inclusions were identified using gas chromatography (GC) at the University of Toronto. A bulk analysis of volatile compounds (N_2 , H_2O , CO_2 , $\text{C}_1\text{--C}_5$ hydrocarbons, COS , SO_2) in the inclusions was obtained by heated (110°C) crushing of quartz followed by on-line GC at the Department of Geology, University of Toronto using an HP 5890 gas chromatograph equipped with PID and TCD detectors; details of the analytical procedure and instrumentation can be found in Bray and Spooner (1992).

Element concentrations in fluid and halide-melt inclusions were determined by laser ablation ICPMS (ETH Zurich, Switzerland); the analytical approach for fluid inclusions is described in Heinrich et al. (2003). Ablation was done with a GEOLAS 193 nm ArF Excimer laser operated at 85–90 mJ output energy, and employing a pulsed beam with an energy-homogenized beam profile (Gunther et al. 1997). Ablated aerosols were transported into an ELAN 6100 quadrupole ICPMS operated in dual detector mode at conditions similar to those reported in Pettke et al. (2004). The following element masses were analyzed and detected in the inclusions: ^7Li , ^{11}B , ^{23}Na , ^{25}Mg , ^{27}Al , ^{39}K , ^{42}Ca , ^{55}Mn , ^{57}Fe , ^{65}Cu , ^{66}Zn , ^{85}Rb , ^{88}Sr , ^{107}Ag , ^{118}Sn , ^{133}Cs , ^{137}Ba , ^{195}Pt , ^{197}Au , ^{208}Pb and ^{209}Bi . Spectrometer dwell time was set to 10 ms for all elements but was increased for certain elements during short menu analytical

sessions in order to improve detection limits [^{65}Cu (20 ms), ^{107}Ag (20 ms), ^{209}Bi (30 ms), ^{197}Au (50 ms), and ^{195}Pt (50 ms)]. Data reduction followed procedures outlined by Longerich et al. (1996) and Heinrich et al. (2003). NaCl wt% eq. concentrations (from microthermometry) were converted to absolute wt% concentrations of Na using a correction for the presence of other major Cl-complexing cations (e.g., Ca^{2+} , Fe^{2+} , Ba^{2+}) in the inclusions (Heinrich et al. 2003). The resulting Na concentrations were used as the internal standard values to calculate element concentrations in the fluid inclusions.

Results

Petrography

Hydrocarbon inclusions containing mainly CH_4 (confirmed by microthermometry, and bulk GC analysis; see below) were observed in hydrothermal quartz in crack-seal type alteration veins splaying from massive sulfide veins within brecciated Levack Gneiss (Sudbury Breccia; Fig. 2a). The alteration veins extend outward to a maximum of ~ 5 m from the parent massive sulfide veins and are most abundant near the terminations of massive sulfide veins (Fig. 1d). The quartz is intergrown with minor (less than 5 vol%) large (up to cm-length) acicular epidote crystals. Occasionally, minor calcite is also observed. Texturally, these minerals appear to have crystallized at the same time. Silicate minerals were then replaced (fracture-controlled) by massive chalcopyrite with minor (< 5 vol%) millerite and bornite (Fig. 2b, c). Electrum (up to 60 elemental wt% Ag; Farrow 1994) occurs primarily (90% of occurrences) as euhedral to subhedral crystals along silicate–silicate grain margins or as fracture-infillings in the silicate phases (Fig. 2d). Sulfide content in the alteration veins decreases with increasing distance from the massive sulfide parent vein.

Hydrocarbon fluid inclusions occur with high salinity aqueous inclusions (brine) as a primary assemblage in the quartz (designated type P_{CH_4} and P_{brine} inclusions; Fig. 3a) [Note that brine inclusions of primary origin (*without* coexisting hydrocarbon inclusions) were described in detail by Li (1993), Farrow (1994) and Molnar et al. (2001)]. Commonly, inclusions in the assemblage contain both a hydrocarbon phase and brine within the same inclusion and show highly variable phase ratios (Fig. 3b), indicating heterogeneous entrapment of coexisting fluid phases. P_{CH_4} and P_{brine} inclusions occur in random, three-dimensional arrays rather than planar orientations and commonly show elongation and tapering in the inferred direction of quartz growth (Fig. 3c); these observations are consistent with established criteria for a primary origin (Goldstein 2003). In addition to halite, the P_{brine} inclusions contain at least one and as many as five additional daughter crystals. This characteristic is consistent with the classification of

Fig. 2 Textural and petrographic characteristics of Au-rich quartz-epidote-sulfide-carbonate alteration veins in the Fraser Cu Zone. **a** An underground exposure (drift wall) of a network of sub-parallel crack-seal alteration veins (qtz-sul-ep) in Sudbury Breccia, connected to a massive sulfide vein (msul). **b** Magnified view of an alteration vein showing replacement of quartz by chalcopyrite. **c** Back-scattered electron image (SEM) of fracture-controlled, pervasive replacement of earlier quartz (qtz)-epidote (ep) intergrowth by chalcopyrite. **d** Back-scattered electron image of electrum (el) along a grain boundary between a zoned epidote grain (ep) and quartz (qtz), and as a fracture filling in the epidote

primary brine inclusions in the study area by other researchers and differentiates the P_{brine} inclusions from secondary aqueous inclusions which contain only halite (e.g., Molnar et al. 2001).

Hydrocarbon inclusions also occur along curved to linear, healed fracture planes coexisting with inclusions of solid halite (designated S_{CH_4} and S_{halite} inclusions; Fig. 3d–i). The inclusion trails are not truncated by growth zones in the quartz, but cut across grain boundaries of adjoining quartz grains and are probably secondary in origin. The inclusions containing the halite are not cubic in shape; rather, they are elongated, tube-like or highly irregular (“amoeboid”) in appearance (Fig. 3d). Commonly, inclusions containing both a hydrocarbon phase and halite were observed in the trails (Fig. 3f), indicating the coexistence of both phases at the time of entrapment. Small amounts of aqueous fluid occur with the solid halite (Fig. 3g, h), visible as a thin film of ice crystals only during freezing measurements. However, no aqueous-dominant (brine) inclusions were

observed in the S_{CH_4} and S_{halite} trails, nor do any inclusions in the trails contain a significant amount (i.e., more than ~5% by vol.) aqueous fluid. Except where they contain a visible amount of brine or halite (by

Fig. 3 Photomicrographs (plane-polarized transmitted light) of hydrocarbon-bearing inclusion assemblages in quartz. Abbreviations: v vapor; b brine; h halite daughter (from brine); s additional daughter phase (from brine). **a** Primary inclusions showing coexisting brine and hydrocarbon phases (P_{CH_4} , P_{brine}). **b** A primary inclusion which has trapped both hydrocarbon and brine phases (by heterogeneous entrapment). **c** Primary inclusions (P_{brine} , P_{CH_4}) showing alignment, tapering and elongation in the inferred direction of quartz growth as a characteristic of primary origin. Inset image was slightly to the left of the main field of view. Quartz growth direction indicated with dashed arrows. Small arrows indicate tapered or “pinched” ends of the inclusions. In the main field of view, a trail of secondary brine inclusions (S_{brine}) is also visible. **d** Trail of large, halite inclusions (S_{halite}) near grain boundary between a quartz (qtz) and chalcopyrite (cpy) grain. Note that the central inclusion is highly irregular in shape (i.e., not cubic) and contains both halite and hydrocarbons ($S_{\text{halite}} + S_{\text{CH}_4}$). **e** Secondary inclusions containing solid halite (melt) and hydrocarbon phases (S_{CH_4} , S_{halite}). The halite-bearing inclusion shows a small, flattened vapor bubble. **f** A secondary inclusion which has trapped both the hydrocarbon phase and halite. **g** A S_{halite} inclusion imaged at 20°C; note the lack of contrast between the halite and host quartz due to their similar refractive indices. **h** The same S_{halite} inclusion shown in (g) but photographed at –193°C; freezing of a thin film of aqueous fluid in the wall interstices of the halite inclusion has occurred and ice crystals are visible on the surface of the halite. **i** A trail of S_{CH_4} and S_{halite} inclusions; note the hydrocarbon-rich and halite-rich segments of the trail. Image obtained at 20°C

heterogeneous entrapment; ~10% of inclusions examined), S_{CH_4} and P_{CH_4} inclusions show a single phase (either hydrocarbon L, V, or supercritical fluid, depending on density) at room temperature. Additionally, a very thin film of aqueous fluid wetting the wall of the S_{CH_4} and P_{CH_4} inclusions was identified during freezing measurements.

Secondary brine inclusions [designated S_{brine} inclusions; described by Molnar et al. (2001)] also occur in the quartz. They occur in trails which crosscut or offset,

and therefore, post-date planes of S_{CH_4} and S_{halite} inclusions. Secondary CO_2 -dominant inclusions reported previously by Molnar et al. (2001) were not observed in any of the samples examined.

The overall observed paragenetic inclusion history is (in order of decreasing age of entrapment): $P_{CH_4} + P_{brine} \rightarrow S_{CH_4} + S_{halite} \rightarrow S_{brine}$. A representative schematic map (based on a photomicrograph) showing the textural relationships among different inclusion assemblages in the quartz samples is shown in Fig. 4.

Fig. 4 Fluid inclusion map (based on a photomicrograph) showing the textural relationships between fluid inclusion assemblages observed in the alteration vein quartz. The earliest fluid assemblage consists of primary inclusions containing brine (P_{brine}) or hydrocarbon (P_{CH_4}) phases. P_{brine} inclusions contain brine, vapor, halite and at least one additional daughter mineral phase. Primary inclusions that trapped both brine and hydrocarbon phases (heterogeneous entrapment) were not observed in this area of the section. An early secondary assemblage of halite (S_{halite}), CH_4 (S_{CH_4}), and heterogeneously trapped halite + CH_4 ($S_{\text{halite}} + S_{\text{CH}_4}$) inclusions occurs as a wide trail across the mapped area, crosscutting several grain boundaries. A narrow trail of late secondary brine inclusions (S_{brine}) cuts across the early secondary halite + CH_4 inclusion trail at the center left of the map and were the last inclusions to form

Microthermometry

A detailed microthermometric study of P_{brine} inclusions (occurring *without* coexisting P_{CH_4} inclusions) in quartz-sulfide-epidote alteration veins at the Fraser Mine was conducted by Molnar et al. (2001). Some P_{brine} inclusions were heated in order to confirm that their high temperature behavior is comparable to that reported by Molnar et al. (2001). Molnar et al. (2001) reported that total homogenization (T_h) of P_{brine} inclusions in quartz-epidote sulfide veins occurred by halite dissolution between ~ 150 and 290°C ; vapor bubble disappearance

Fig. 5 Histogram showing the temperature range of final homogenization by halite dissolution (T_h) of P_{brine} inclusions occurring with hydrocarbon inclusions (this study) and without hydrocarbon inclusions (Molnar et al. 2001). The range in final homogenization temperatures of the data sets overlap with one another

Fig. 6 Histogram showing the temperature range of vapor bubble disappearance (T_v) in P_{brine} inclusions occurring with hydrocarbon inclusions (this study)

occurred slightly below the T_h , and ranged from 96 to 235°C . For comparison, P_{brine} inclusions coexisting with P_{CH_4} inclusions in this study always homogenized by halite dissolution between ~ 145 and 315°C (mean = $230 \pm 43^\circ\text{C}$ [$n=53$, 1σ]), corresponding to salinities between 29.5 and 39.3 wt% eq. NaCl (mean = 33.5 ± 2.6 wt% eq. NaCl [$n=53$, 1σ]) (Fig. 5). Vapor bubble disappearance occurred between 117 and 325°C (mean = $211 \pm 39^\circ\text{C}$ [$n=53$, 1σ]) (Fig. 6).

When cooled to -93°C , P_{CH_4} and S_{CH_4} inclusions show an inner bubble of low refractive index vapor (~ 50 – 60 vol% of the inclusion) surrounded by a more refractive liquid (~ 40 – 50 vol%). The inability to freeze the P_{CH_4} and S_{CH_4} inclusion contents using liquid N_2 indicates that additional volatile components with T_{triple} below that for pure CH_4 (-82.5°C) are probably present (see below), although significant undercooling is often required to induce freezing. Upon heating, the hydrocarbon phase in P_{CH_4} inclusions homogenizes by near-critical behavior to either liquid or vapor, or by critical behavior at T 's between -87 and -73°C (mean = $-1.4 \pm 2.1^\circ\text{C}$ [$n=40$, 1σ]), suggesting that CH_4 is the principal species ($T_{\text{critical}} = -2.6^\circ\text{C}$) (Fig. 7). The hydrocarbon phase in S_{CH_4} inclusions always homogenized to vapor at T 's between -71 and -49°C (mean = $-4.9 \pm 7.0^\circ\text{C}$ [$n=28$, 1σ]), indicating that sig-

Fig. 7 Histogram showing the range in T_h of the hydrocarbon phase in P_{CH_4} and S_{CH_4} inclusions. P_{CH_4} inclusions homogenize close to the critical T for pure CH_4 (-2.6°C), whereas S_{CH_4} inclusions homogenize at higher temperatures than the critical T and therefore contain additional volatiles

nificant quantities of volatile compounds in addition to CH_4 must be present (Fig. 7). With continued heating, the formation of a thin layer of dendritic ice crystals between -9 and -41°C was visible along the exposed (upper) wall of the S_{CH_4} and P_{CH_4} inclusions, indicating that a small amount of brine is present in the inclusions. Final melting of the ice occurred between -31 and -21°C . No further phase changes were observed above -21°C ; if final homogenization of the trace amount of aqueous brine and hydrocarbon phase occurred, it could not be observed. P_{CH_4} or S_{CH_4} inclusions that contained significant amounts of brine always decrepitated upon heating at $T > 400^\circ\text{C}$ and did not show any evidence that

Fig. 8 Photomicrographs (plane-polarized, transmitted light) of high temperature phase changes during heating of two S_{halite} inclusions. At 600°C , the inclusion walls are poorly visible. Melting begins at 772°C and is characterized by a sudden change in the relief of the inclusions due to the formation of a thin layer of melt around the solid halite. At 774°C , the melt:solid ratio begins to increase rapidly in the inclusion on the left side of the image. At 780°C , the melt:solid ratio of the inclusion on the left exceeds 50 vol% and the first melt is visible in the inclusion on the right. The inclusion on the left side of the image has melted completely by 785°C ; less than 50 vol% of the halite remains in the inclusion on the right. At 787°C , both inclusions show a single phase, high relief halide melt

homogenization was about to occur before decrepitation (i.e., no changes in phase ratios), consistent with heterogeneous entrapment.

Heating of the S_{halite} inclusions to very high T ($> 700^\circ\text{C}$) resulted in the melting of the halite (Fig. 8). Prior to melting, the disappearance of a tiny vapor bubble occurred, followed by the partial dissolution of the outer surface of the halite crystal into the small amount of aqueous brine surrounding the halite resulting in the appearance of a distinct boundary between the halite and wall of the inclusion. This occurred at $\sim 600^\circ\text{C}$ for most inclusions (Fig. 8). The melting of halite (T_m) occurred between 710 and 805°C ; melting occurred rapidly over a $\sim 10^\circ\text{C}$ interval leading up to the final disappearance of the halite; large groups of S_{halite} inclusions melted *simultaneously* between 770 and 805°C (Figs. 8, 9). Melting temperatures below the solidus T of pure halite ($\sim 815^\circ\text{C}$ at 0.5 kbar) indicate that other volatiles or cations are present in the inclusions. The presence of additional cations in the S_{halite} inclusions was confirmed by X-ray mapping and LA-ICP-MS analysis (see below).

Fig. 9 Histogram showing the range in melting temperatures in S_{halite} inclusions occurring with CH_4 . Melting T 's below that for pure halite ($\sim 815^\circ\text{C}$ at 0.5 kbar) are due to the presence of aqueous fluid containing other components (e.g., H_2O) lowering the solidus.

(Note that it is possible that during the earlier study of Molnar et al. (2001), trails of hydrocarbon inclusions were misidentified as CO₂ inclusions. Additionally, we found that halite inclusions are extremely difficult to identify without freezing or heating because the refractive indices of halite and quartz are identical).

Bulk analysis of volatiles by GC

In order to confirm the presence of CH₄ as the primary hydrocarbon species (as suggested by low-*T* microthermometric behavior), quartz containing abundant P_{brine}, P_{CH4}, S_{halite}, and S_{CH4} inclusions and relatively few S_{brine} inclusions was crushed and the released volatiles were analyzed by GC. It is important to recognize that due to the presence of two major inclusion assemblages, the results obtained represent a mixed signal with aqueous and non-aqueous volatile species from four different fluid inclusion types. Therefore, the results listed below cannot be taken as true concentrations of volatiles in any single inclusion type or assemblage, but nevertheless are valuable for the identification of compounds present.

The quartz sample yielded the following analysis (each value in mole %): 97.2% H₂O; 1.79% CH₄; 0.79% C₂H₆-C₂H₂; 0.11% N₂; 0.07% C₃H₈. Other analyzed species were present at relative concentrations of less than 0.01 mole %, or were undetected. Notably, CO₂ accounted for only 0.003 mole % of the total volatile content. This analysis does not represent the composition of only the hydrocarbon phases due to the diluting effect of H₂O from the brine inclusions on the bulk analysis. Therefore, recasting the analysis by excluding H₂O and assuming that there is relatively little of the gas species dissolved in the brine inclusions (Poling et al. 2001) yields an approximate composition of the hydrocarbon fluid from *mixed* P_{CH4} and S_{CH4} inclusions of: ~64% CH₄, 29% C₂H₆-C₂H₂, 4% N₂, and 3% C₃H₈.

The analysis is consistent with the unusual microthermometric behavior observed in the P_{CH4} and notably, the S_{CH4} inclusions. The *T*_{triple} of N₂ and C₃H₈ are -10 and -88°C, respectively; the presence of N₂ and C₃H₈ will therefore lower the bulk freezing *T* of the hydrocarbon fluid, possibly explaining our inability to freeze the hydrocarbon phase in P_{CH4} and S_{CH4} inclusions using liquid N₂. Additionally, because of the relatively high *T*_{critical} of C₂H₆-C₂H₂ (32.2°C-

Table 2 Element concentrations (laser ablation ICP-MS) in P_{brine}, S_{halite} and P_{CH4} inclusions

Element	^a P _{brine} inclusions				Element	^c S _{halite} inclusions				Element	^d P _{CH4} inclusions			
	^b Mean (<i>n</i>)	1σ	Max	Min		^b Mean (<i>n</i>)	1σ	Max	Min		^b Mean (<i>n</i>)	1σ	Max	Min
Ca (wt%)	11.08 (50)	1.93	16.42	7.27	Ca (wt%)	0.52 (4)	0.13	0.69	0.33	Ca (ppm)	3,780 (14)	1,796	6,683	375
Na	6.16 (50)	1.19	8.42	3.70	Na	37.00 (19)	0.41	37.37	35.58	Na	1,935 (18)	1,196	4,574	399
K	2.55 (38)	0.48	3.79	1.44	K (ppm)	562 (14)	603	2,002	67	K	994 (17)	618	2,373	169
Fe	1.17 (38)	0.49	2.31	0.27	Fe	9,627 (4)	15,592	36,627	198	Fe	383 (11)	326	1,047	15
Pb	1.03 (38)	0.24	1.82	0.32	Pb	53 (11)	88	322	2.2	Pb	150 (18)	154	594	24
Mn	0.64 (38)	0.16	1.27	0.24	Mn	66 (4)	71	188	22.4	Mn	75 (17)	70	262	4.0
Zn	0.44 (38)	0.15	1.10	0.16	Zn	232 (8)	329	1,073	15	Zn	188 (16)	164	572	30
Sr	0.44 (18)	0.07	0.56	0.32	Sr	24 (17)	40	178	4.1	Sr	NA	NA	NA	NA
Ba	0.43 (38)	0.08	0.69	0.31	Ba	20 (15)	19	73	4.0	Ba	NA	NA	NA	NA
Cu (ppm)	156 (37)	232	993	4.9	Cu	4,486 (14)	3,381	11,125	17	Cu	1,286 (17)	2,395	9,737	18
Al	136 (18)	116	460	12	Al	785 (5)	239	838	48	Al	NA	NA	NA	NA
Mg	115 (30)	110	472	18	Mg	515 (4)	609	1,544	12	Mg	NA	NA	NA	NA
Rb	109 (18)	19	147	78	Rb	5.6 (4)	1.6	6.8	1.9	Rb	NA	NA	NA	NA
B	84 (17)	24	153	49	B	117 (6)	110	354	33	B	NA	NA	NA	NA
Sn	85 (14)	91	296	8.1	Sn	24 (5)	16	54	11	Sn	NA	NA	NA	NA
Li	74 (14)	65	264	14	Li	955 (15)	978	2,850	4.0	Li	NA	NA	NA	NA
Ag	15 (44)	7.6	41	3.6	Ag	138 (18)	253	775	1.0	Ag	20 (15)	37	157	0.5
Cs	12 (18)	2.0	16	7.7	Cs	8.9 (5)	16	40	0.4	Cs	NA	NA	NA	NA
Pt	4.3 (11)	3.8	12	0.2	Pt	3.5 (5)	1.8	5.4	0.8	Pt	ND	ND	ND	ND
Bi	4.1 (18)	4.5	19	0.4	Bi	12 (3)	8.7	24	2.5	Bi	6.6 (9)	4.2	14	1.1
Au	3.4 (10)	3.5	11	0.2	Au	6.1 (5)	7.5	21	0.4	Au	1.4 (12)	1.2	4.6	0.1
Ca/Na	1.9 (50)	0.7	4.2	1.0	Ca/Na	0.03 (7)	0.03	0.1	0.01	Ca/Na	3.9 (14)	3.1	10	0.1
Sr/Rb	40 (18)	5.1	58	34	Sr/Rb	19 (7)	19	53	1.0	Sr/Rb	NC	NC	NC	NC
Pb/Zn	2.4 (38)	0.4	3.9	1.6	Pb/Zn	1.8 (9)	1.5	5.0	0.04	Pb/Zn	1.2 (16)	1.2	4.6	0.1
Ag/Bi	9.4 (17)	9.3	40	0.7	Ag/Bi	1.7 (4)	1.9	4.9	0.4	Ag/Bi	5.3 (9)	6.8	20	0.5
Pt/Au	0.9 (4)	0.4	1.3	0.2	Pt/Au	0.3 (1)	NC	NC	NC	Pt/Au	NC	NC	NC	NC
Cu/Pt	97 (11)	79	257	2.0	Cu/Pt	83 (7)	144	426	0.7	Cu/Pt	NC	NC	NC	NC
Cu/Au	254 (11)	504	1,821	5.8	Cu/Au	1,315 (6)	1,055	3,028	3.5	Cu/Au	1,681 (12)	2,930	9,760	15

NA not analyzed, ND not detected in any inclusions (3σ limit of detection), NC not calculated^aCalculated using measured inclusion salinity (microthermometric determination of final dissolution of halite)^bAverage concentration in “*n*” analyzed inclusions that gave element concentrations above a 3σ limit of detection (Individual detection limits for those inclusions not reported in the average

values above are listed in Electronic Supplementary Material)^cCalculated using measured inclusion salinity (microthermometric determination of final melting of halite)^dCalculated using an estimated bulk salinity of 1.15 wt% eq. NaCl (see text for explanation)

35.1°C) and C_3H_8 (96.7°C), their presence can raise the homogenization T of the P_{CH_4} and S_{CH_4} inclusions above that for pure CH_4 (−2.6°C). Thus, S_{CH_4} inclusions probably contain the highest abundances of these additional compounds, as their behavior at low T is least consistent with pure CH_4 .

LA-ICP-MS analyses of single inclusions

Table 2 summarizes average element concentrations in 50 P_{brine} inclusions and 18 P_{CH_4} inclusions (from three groups of coeval $P_{brine} + P_{CH_4}$ inclusions occurring in three different quartz crystals) and 19 S_{halite} inclusions from two different inclusion trails within a single quartz crystal. Also listed are the maximum and minimum concentrations for each element in each inclusion type, and a 1σ standard deviation on the average concentration. Tables of all individual inclusion analyses (with microthermometry measurements) can be found in Electronic Supplementary Material associated with this article. Note that for many elements, the number of analyses represented in the average concentration is less than the total number of inclusions analyzed because not all elements were detected in the inclusions at a high confidence level (3σ). Also note that the detection limit for a given element is unique to each inclusion and related to inclusion size and ablation efficiency (i.e., the mass of inclusion contents analyzed over the ablation interval). For the P_{brine} and S_{halite} inclusions, the final halite dissolution or halite melting temperature was accurately determined for each inclusion individually prior to laser ablation, to obtain the wt% eq. NaCl value required for quantification. However, to quantify element concentrations of P_{CH_4} inclusions, the wt% eq. NaCl values had to be estimated because no reliable salinity measurements could be obtained by microthermometry. Microthermometric measurements indicate that the minimum salinity of the aqueous fluid fraction was ~ 23 wt% eq. NaCl, based on the final melting of small ice dendrites on the walls of the P_{CH_4} inclusions. Using a minimum salinity of 23 wt% eq. NaCl and an estimated volume fraction of aqueous fluid of 1–5 vol %, the bulk salinity of the P_{CH_4} inclusions was estimated to be between 0.23 and 1.15 wt% eq. NaCl. This bulk salinity range was used to calculate the concentrations of all other elements in the P_{CH_4} inclusions analyzed. In Table 2, the average composition of P_{CH_4} inclusions was calculated using the higher bulk salinity of 1.15 wt% eq. NaCl; however, the full data set (in Electronic Supplementary Material) lists all analyses calculated using both the 0.23 and 1.15 wt% eq. NaCl values for comparison. Considering that ice crystals were clearly visible in the inclusions during freezing measurements, and that such inclusions can easily contain up to ~ 10 vol% aqueous fluid along their inner walls without the fluid being visible, the bulk salinity estimate of 1.15 wt% is tentatively considered most representative of the true bulk salinity of the aqueous fluid fraction in the P_{CH_4} inclusions.

Analyses of P_{brine} inclusions show that Ca is the dominant cation, with Na and K also present in wt% concentrations. Elements detected at concentrations in the thousands of ppm to wt% range are Pb, Fe, Mn, Ba, Zn and Sr. The presence of significant concentrations of these elements is consistent with the identification of the Ca–Na–K–Fe–Mn–Pb–Ba chloride daughter minerals in P_{brine} inclusions by other researchers (Farrow 1994; Molnar et al. 2001). Cu, Al, Mg, Rb, B, Sn and Li are present in the brine in the 10–1,000 ppm range. Ag, Cs, Pt, Au and Bi are present at concentrations in the 1–10 ppm range. Notably, Pt and/or Au were detected in 11 of the brine inclusions (at a 3σ confidence level), at concentrations ranging from 0.2 to 12 ppm. Ratios of Pt/Au in the P_{brine} inclusions are around ~ 1 . Most element ratios (e.g., Pb/Zn; Table 2) are very consistent from inclusion to inclusion; however, ratios involving the elements Au, Ag, Pt, and Bi show up to an order of magnitude variation within the P_{brine} assemblage; for example, Pt/Au ratios range from 0.2 to 1.3. Most variable are metal ratios involving Cu; Cu:Au ratios which range from ~ 6 to $\sim 1,800$ within the inclusion populations analyzed.

Element concentrations in P_{CH_4} inclusions (Table 2) are unremarkable. However, Cu, Ag, Au and Bi are present at concentrations comparable to or possibly higher than the coexisting P_{brine} inclusions, despite the much lower bulk salinity of the CH_4 inclusions. Figure 10a shows the transient signal obtained during the ablation of a Cu, Au, Bi, and Ag-rich P_{CH_4} inclusion (inclusion ju24a06; see full data set in Electronic Supplementary Material). Using the transient signal in Fig. 10a as an example, metal signals are clearly related to the P_{CH_4} inclusion contents and are the result of host surface contamination. Other elements (Na, K, Ca, Mn, Fe, Pb and Zn) show no relative enrichment in the P_{CH_4} inclusions. This observation is illustrated graphically in Fig. 11a in which the average element concentrations (solid circles) and ranges in concentration (black bar) in all P_{CH_4} inclusions analyzed are plotted against the respective element concentrations (average and range) of P_{brine} inclusions [Mean element concentrations in the P_{CH_4} inclusions are those calculated assuming a bulk salinity of 1.15 wt% eq. NaCl (5 vol% brine at ~ 23 wt% eq. NaCl)]. Inset Fig. 11b shows the shift in average element concentration for Cu and Zn in P_{CH_4} inclusions if a bulk salinity of 0.23 wt% eq. NaCl is used (1 vol% brine at ~ 23 wt% eq. NaCl). The solid diagonal line in Fig. 11a, b corresponds to equal element concentrations in the P_{brine} and P_{CH_4} inclusions ($C_{brine} = C_{CH_4}$); degrees of element fractionation are shown as dotted lines corresponding to lines of constant C_{brine}/C_{CH_4} for each element. In Fig. 11a, Ca, Na, K, Fe, Mn, Zn and Pb behave similarly and are always strongly enriched in the brine phase relative to the hydrocarbon fluid (mean $C_{brine}/C_{CH_4} = \sim 30$ to 100; range = ~ 10 to 1,000). Au, Bi and Ag behave differently from the strongly brine-partitioning elements; they may be variably enriched in

Fig. 10 Transient LA-ICP-MS intensity plots of inclusion ablations. **a** Ablation of a P_{CH_4} inclusion (inclusion # ju24a06; see Electronic Supplementary Material). Background is collected with the laser off for at least 50 s. The laser is switched on, monitored by the increase in the Si^{29} (grey line) count rate. At $t \sim 60$ s the inclusion is opened (increase in Na^{23} signal; white diamonds) and the pit diameter increased sequentially (corresponding to the two short drops in count rate). A mixed signal (inclusion + host) is observed for Si^{29} , Na^{23} , Cu^{65} (grey squares), Ag^{107} (dark grey triangles), Bi^{209} (crosses), and Au^{197} (solid black circles) over a ~ 20 s interval. Note the typically irregular signal for Au^{197} showing several isolated counting events which roughly coincide with the Cu^{65} signal. **B**. Ablation of a S_{halite} inclusion (inclusion # de16a07; see Electronic Supplementary Material). The inclusion is opened at ~ 120 s. Note the increase in Pt^{195} (solid black circles) counting events at the beginning and end of the inclusion signal. Also note the coincidence between the count rates for Cu^{65} (grey squares), Li^7 (grey triangles) and Na^{23} (white diamonds)

either the brine or CH_4 phase based on the range of concentrations measured and calculated (mean $C_{brine}/C_{CH_4} = \sim 0.8$ to 2; range = ~ 0.07 to 40). Cu is commonly fractionated into the CH_4 phase ($C_{brine}/C_{CH_4} = \sim 0.1$; range = ~ 0.006 –10). It is important to note that changes in the assumed bulk salinity of the P_{CH_4} inclusions will change the calculated individual element concentration ratios between brine and CH_4 (Table 2, Fig. 11a, b) will change. Therefore, in the absence of an accurate bulk salinity determination for the P_{CH_4} inclusions, it is not possible to demonstrate for certain that specific ore metals (e.g., Cu) occur in greater absolute concentration in the CH_4 phase than in the brine. However, changing the bulk salinity estimate of the P_{CH_4} inclusions does not change the relative partitioning behavior of the elements analyzed; element concentration ratios between CH_4 and brine for Cu, Au, Bi and Ag are greater than those for Na, Ca, Fe, Mn, Pb and Zn, regardless of the uncertainty in absolute element concentration. To eliminate

uncertainties in the accidental brine content and bulk salinity of the P_{CH_4} inclusions, metal enrichment in the CH_4 relative to the brine may also be expressed in a ratio normalized to Na (Heinrich et al. 1999). For example, in (coeval) group 1 of P_{CH_4} and P_{brine} inclusions (Appendix 4), the average metal enrichment ratios $[(metal_{CH_4}/Na_{CH_4})/(metal_{brine}/Na_{brine})]$ are as follows: Ca = 1.8, K = 1.4, Mn = 0.7, Fe = 0.7, Pb = 0.7, Zn = 3.0, Cu = 460.9, Ag = 54.8, Au = 11.5, Bi = 36.5. The relative tendency for elements to fractionate preferentially into CH_4 phase appears to be: Cu (strongly fractionated into the CH_4) > Ag, Bi > Au >> Ca, Na, K, Fe, Pb, Zn, Mn (strongly fractionated into the brine).

S_{halite} inclusions (Table 2) are composed primarily of NaCl, with minor amounts of Ca (up to ~ 3 wt%) and Fe (up to ~ 4 wt%). Figure 10b shows the transient signal obtained during the ablation of a S_{halite} inclusion (inclusion de16a07; see full data set in Electronic Supplementary Material). X-ray mapping (by SEM-EDS) of halite inclusions confirmed that the Ca is present in the

Fig. 11 Element concentrations in coexisting P_{brine} and P_{CH_4} inclusions. *Diagonal lines* represent constant ratios of $C_{\text{brine}}^{\text{elem}}/C_{\text{CH}_4}^{\text{elem}}$ as labeled. The *diagonal solid line* represents equal distribution of an element between the two inclusion types (no fractionation; $C_{\text{brine}}^{\text{elem}}/C_{\text{CH}_4}^{\text{elem}} = 1$). **a** *Black solid circles* are average element concentrations. *Error bars* indicate the range of concentrations for each element from inclusions containing either CH_4 or brine. Mean concentrations are based on the approximation of a bulk salinity of 1.15 wt% eq. NaCl (5 vol% aqueous film at ~23 wt% NaCl) in the P_{CH_4} inclusions. **b** Demonstration of the effect of lowering the assumed bulk salinity from 1.15 (black solid circles) to 0.23 wt% eq. NaCl (grey solid circles; 1 vol% aqueous film) on the mean concentrations of Cu and Zn in P_{CH_4} inclusions. Cu, Ag, Bi and Au are enriched in the P_{CH_4} inclusions (show lower $C_{\text{brine}}^{\text{elem}}/C_{\text{CH}_4}^{\text{elem}}$ ratios) relative to strongly brine-partitioning elements showing very similar behavior (Na, Ca, Fe, Mn, Zn, Pb). Using the 1.15 wt% eq. NaCl bulk salinity approximation, many P_{CH_4} inclusions have absolute concentrations of Cu, Ag, Bi and Au higher than the coexisting P_{brine} population

subsidiary amounts of aqueous fluid surrounding the halite, rather than as a component in the halite (i.e., in solid solution; Fig. 12). The Cu content of the S_{halite} inclusions is commonly very high (average = 4,486 ppm Cu; range 17 ppm to over 1 wt% Cu; $n=14$). Lithium contents are also very high (average = 978 ppm Li; range 4–2,850 ppm) and correlate with Cu concentration. The Cu and Li appear to be dissolved in the halite itself rather than in the subsidiary amount of aqueous fluid surrounding the halite (Fig. 10b). Other elements (Mn, K, Zn, Sr, Ba) were detected in the 10–100 ppm range and are probably present in the aqueous fluid surrounding the halite rather than in the halite itself. Several element ratios (e.g., Pb:Zn) in the P_{brine} and S_{halite} inclusions are comparable (Table 2), despite the overall much lower concentrations of these elements in the S_{halite} inclusions. Significant (ppm) concentrations of Pt and Au were detected in five of the halite-melt inclusions analyzed.

Discussion

Origin of the brine– CH_4 assemblage

The occurrence of inclusions containing CH_4 , brine and CH_4 –brine mixtures within a single, coeval assemblage suggests the entrapment of immiscible fluid phases. At ~35 wt% NaCl (comparable to the brine salinity in this study) and at a P_{trapping} of 1.1–1.5 kbar (for P_{brine} inclusions occurring without coexisting P_{CH_4} inclusions; Molnar et al. 2001), the two-phase field in the CO_2 –NaCl– H_2O system extends to ~680°C and probably higher (> 700°C) in the CH_4 –NaCl– H_2O system (Krader and Franck 1987). Graphically (Fig. 13), we estimate by the intersection of brine and CH_4 isochores that the pressure of entrapment for the $P_{\text{brine}} + P_{\text{CH}_4}$ assemblage could not have exceeded ~500 bars, a pressure considerably lower than the estimate of Molnar et al. (2001). At such a low pressure of entrapment, two immiscible fluid phases would have certainly coexisted (Bowers and Helgeson 1983; Krader and Franck 1987). On the basis of our trapping pressure estimate, the CH_4 may have been a minor dissolved component in the original one-phase brine phase (Molnar et al. 2001) that later separated from the brine during local decompression during vein formation. Alternately, simple cooling of a CH_4 -bearing brine as it came into contact with groundwater or wall rocks, or a combination of both decompression and cooling could have been responsible for CH_4 –brine unmixing.

Compositional data (Table 2) allow further constraints on the origin of the brine. The mean Ca:Na ratio in the P_{brine} inclusions is 1.9 ± 0.7 (1σ , $n=50$); this value is comparable to the range in Ca:Na ratios obtained from saline groundwaters sampled at a compa-

Fig. 12 Secondary electron and X-ray intensity images of a halide melt inclusion. **a** Secondary electron image showing opened solid inclusions of halide-melt (*center and upper left*) on a fractured surface in quartz (qtz); opened CH_4 inclusions (*arrows*) are also visible. **b** Na $K\alpha$, **c** Cl $K\alpha$, **d** Ca $K\alpha$ showing a halo of elevated X-ray intensity along the margin of the melt inclusion, corresponding to dried aqueous fluid that surrounded the solid melt in the unopened inclusion at room temperature

rable sample depth ($\sim 1,100$ m) at the Fraser Mine (Ca:Na ratio = 1.7–1.9; Frape and Fritz 1982), strengthening previous suggestions that the brine is compositionally related to deep groundwaters (i.e., formation waters with compositions that have remained relatively unchanged since their formation) have Sr:Rb ratios in the brine [mean = 40 ± 5 (1σ , $n=50$)] strongly reflect interaction with the gneissic host rocks in the study area which have an average bulk rock Sr:Rb ratio of ~ 39 (Hanley 2002). However, the brine composition has been modified to contain other cations (e.g., Fe, Mn, Pb, Ba) in wt% concentrations and significant concentrations of ore metals (Cu, Ag, Bi, Pt, Au). For comparison, Fe, Mn, Pb and Ba are present only in trace (< 10 ppm) concentrations in the deep groundwaters of the Sudbury area (Frape and Fritz 1982) and the ore metals detected in the P_{brine} inclusions could clearly not have been derived from the groundwaters. Therefore, models describing the interaction of heated groundwater with sulfide, or mixing of groundwater with a second fluid of magmatic origin exsolved from the SIC or from a sulfide liquid are

supported by our new observations (Li 1993; Farrow 1994; Marshall et al. 1999; Hanley et al. 2004).

Origin of the halite- CH_4 - C_2H_6 - C_2H_2 - C_3H_8 assemblage

Single inclusions containing the proposed halide-melt and hydrocarbon phases exhibit highly variable halite:hydrocarbon ratios, suggesting that the hydrocarbon phase and halide-melt coexisted (immiscibly) at the time of entrapment. No previous studies have reported the occurrence of hydrocarbon or solid halite inclusions associated with the Sudbury Igneous Complex or its ore deposits. Although carbonic fluid-halide melt assemblages have been reported in a few other mafic-intermediate igneous systems (e.g., Laramie Anorthosite Complex; Frost et al. 1989), none have been reported which contain pure hydrocarbon fluids (i.e., with no CO_2 present). On the basis of their non-cubic and commonly irregular morphology, narrow range of melting temperatures and a lack of any significant amount of aqueous fluid, the S_{halite} inclusions do not appear to represent accidentally trapped inclusions of halite precipitated from a halite-saturated aqueous fluid (e.g., Erwood et al. 1979). Furthermore, it is unlikely that halite precipitated from an aqueous fluid could accommodate such high concentrations of Cu dissolved in its structure (Fig. 10b), owing to the significant difference in size between the Cu^{1+} and Na^{1+} ions in sixfold coordination (Shannon 1979). The

Fig. 13 Estimation of the pressure of entrapment of the primary $P_{\text{brine}} + P_{\text{CH}_4}$ inclusion assemblage using intersecting isochores on a pressure-temperature section for the NaCl-H₂O system (position of relevant fields and phase boundaries taken from Sourirajan and Kennedy 1962; Bodnar 1994; Atkinson 2002). Note that the determination of pressure of entrapment for the $P_{\text{brine}} + P_{\text{CH}_4}$ inclusion assemblage is an approximation only because we have applied isochore calculations for the pure NaCl-H₂O system to a complex mixed salt system. Isochores for the coexisting P_{brine} inclusions cannot be determined for the majority of inclusions because the inclusions homogenize by halite dissolution and vapor disappearance temperatures are considerably lower than T_h ; for those inclusions, isochores in the H+L field of the NaCl-H₂O system are not constrained. However, $T_v \approx T_h$ in several inclusions (inclusion #s ju22a07, ju22a08, ju22a10, ss02a03, ss03a04 and ss03b07; see Electronic Supplementary Material). Isochores for the listed inclusions originate near the intersection of the halite liquid and bubble curves (for inclusions ranging from 31.1 wt% and 34.63 wt% eq. NaCl) and lie primarily in the L-only field. For the listed inclusions, isochores were calculated using Eq. 12 from Zhang and Franz (1987), the inclusion salinities, and values of T_h ; the isochores define the grey field labeled “brine isochores”. Most P_{CH_4} inclusions homogenized by critical behavior near the critical T for pure CH₄ (−2.6°C); the density of the CH₄ phase is close to critical density (0.162 g/mL) and the isochore for the CH₄ phase is relatively shallow (labeled “methane critical isochore”). The maximum pressure of entrapment (P_{trapping}) for those inclusions is constrained by the intersection of the critical CH₄ isochore with the brine inclusion isochores and is ~500 bar. For comparison, the range of P_{trapping} (1.1–1.5 kbar) for the primary assemblage where the assemblage contains only the brine phase (Molnar et al. 2001). This would suggest that a minimum fluid decompression of ~600 bar resulted in brine-CH₄ immiscibility

petrographic, microthermometric and compositional characteristics of S_{halite} inclusions all suggest that the S_{halite} inclusions represent trapped amounts of a halide melt (molten salt) phase which contained a small amount of dissolved H₂O and other trace cations at the time of entrapment.

While the occurrence of a halide-melt phase is plausible, the coexistence of a CO₂-free hydrocarbon fluid with a halide-melt is problematic: A CH₄-dominant, C–O–H gas phase (> 95 mol% CH₄) should exist stably at halide-melt entrapment temperatures (~700–800°C) *only* at unrealistically reducing conditions (QFM-4; iron-wustite buffer; Mathez et al. 1989). The presence of magnetite in massive sulfides from the footwall ore zones suggests that the sulfide liquids crystallized along a magnetite-saturated liquidus; therefore, the ambient fO_2 conditions in the vicinity of the crystallizing sulfides could not have been as reducing as QFM-4. At such temperature and moderate fO_2 , the dominant C–O–H species in the inclusions should be CO₂ (Salvi and Williams-Jones 1997; Potter and Konnerup-Madsen 2003). Additionally, the higher hydrocarbons in the S_{CH_4} inclusions would be expected to convert completely to graphite above 500°C in contact with a halide-melt phase (Shock 1990). In response to these problems, it is possible that the carbonic fluid coexisting with the halide-melt phase was originally CO₂-dominant at magmatic temperatures but was later converted to a CH₄ + C₂H₆ + C₂H₂ + C₃H₈ fluid prior to entrapment by some disequilibrium process (i.e., Fischer–Tropsch synthesis; Salvi and Williams-Jones 1997) or that the hydrocarbons survived in a metastable state long enough to be preserved in fluid inclusions. The presence of higher order hydrocarbons preserved in kimberlite pipes (Shepeleva et al. 1990) and mantle xenoliths (Mathez 1987; Sugisaki and Mimura 1994) would suggest that such compounds may survive at magmatic conditions for sufficient periods of time to be trapped within inclusions.

The sulfide ores are a favorable source for a halide-melt phase considering that: (1) S_{halite} inclusions occur only in close proximity (within a few meters) to massive sulfide ores and do not occur in the “background” fluid inclusion record in the Levack Gneiss (Molnar et al. 2001), (2) the $S_{\text{halite}} + S_{\text{CH}_4}$ assemblage and sulfide minerals texturally postdate the lower temperature $P_{\text{brine}} + P_{\text{CH}_4}$ assemblage hosted in early alteration quartz, and (3) S_{halite} inclusions contain abundant Cu and precious metals. The exsolution of halide-melt and hydrocarbon phases from crystallizing sulfides would imply that Cl, alkali metals (mainly Na), and C were dissolved (compatible) elements in the original sulfide liquid. Although the solubilities of Cl, Na and C in Cu-rich sulfide liquids are not known, trace amounts of these elements should eventually reach saturation and be ultimately rejected as the sulfide liquid reaches its solidus temperature. (Note that Li (1993) originally suggested that the sulfide-exsolved phase was the primary brine. However, we conclude from textural observations that final crystallization of the sulfides *post-dated* the entrapment of the primary brine). Unfortunately, inclusions of solid halite were not observed in any sulfide minerals in this study. However, there are several other indications that the fractionated Cu–Fe-sulfide liquids may have contain significant dissolved Cl. At the Fraser

Fig. 14 Tentative paragenetic sequence for hydrothermal/volatile events identified in this study. The formation of quartz-epidote veins, deposition of Au in the veins in association with the silicates and entrapment of primary fluid inclusions containing brine (P_{brine}) and CH_4 (P_{CH_4}) coincided with or preceded footwall sulfide vein emplacement (timing depends partly on the interpreted source of the brine; see text for description). During the final stages of sulfide fractionation and crystallization, PGE–Au-rich sulfides migrated into the earlier-formed quartz-epidote veins and replaced the silicates; metal-rich halide-melt (S_{halite}) and hydrocarbons (S_{CH_4}) exsolved from the sulfides as the sulfides approached their solidus temperature. Late-stage brines circulated through the study area after the ores had solidified and cooled.

mine, lawrencite (FeCl_2), bismoclite (BiOCl), ferropyrosomalite [$(\text{Fe}^{2+}, \text{Mn})_8\text{Si}_6\text{O}_{15}(\text{OH}, \text{Cl})_{10}$], and unnamed Pd–Bi-chlorides occur as solid inclusions (formerly, phenocrysts) within massive sulfides in the footwall (Li 1993; Farrow 1994; Hanley 2002). Recently, Mungall and Brenan (2003) confirmed field observations of the apparently chalcophile nature of the halogens by dissolving wt% concentrations of Cl (as FeCl_3) in sulfide liquid using a piston cylinder apparatus; they observed that the sulfides exsolved Si-rich, Fe-halide salts during quench. In other mafic systems, Cu-bearing halide-melt phases coexisted with stable sulfide liquid. For example, Renno et al. (2004) reported coexisting Cu-sulfide, Cu-chloride, and silicate glass within magnesiohastingsite cumulates at the TUBAF seamount, Papua New Guinea and argued that the unmixing of a complex Cu-bearing chloride-melt from mafic silicate liquid occurred at magmatic temperatures ($> 800^\circ\text{C}$).

It is possible that our interpretation of the origin of S_{halite} inclusions as a trapped halide-melt phase is incorrect and that the inclusions do represent accidentally trapped halite crystals precipitated from a much lower- T , halite-saturated fluid. The halite-saturated fluid was not aqueous in composition for the reasons discussed above. Rather, halite may have been precipitated along the walls of fractures in the quartz by a NaCl-saturated hydrocarbon fluid at lower T ($< 500^\circ\text{C}$) and more geologically realistic $f\text{O}_2$ which would favor the stability of CH_4 and other hydrocarbons as the

dominant equilibrium species. If, on cooling, a hydrocarbon fluid phase precipitated NaCl, the result would be a mixed assemblage of inclusions containing accidentally trapped halite, hydrocarbon fluid, or variable amounts of halite + hydrocarbon fluid (consistent with the observed assemblage). Although the solubility of NaCl in a non-polar solvent such as CH_4 or CO_2 has not been investigated at any P – T conditions (for all practical purposes, CH_4 and NaCl may be considered to be mutually immiscible phases), a non-polar fluid can probably carry trace quantities of NaCl at lower temperatures (~ 300 – 500°C ; Bowers and Helgeson 1983).

Relative timing of ore and volatile formation

Volatile evolution and magmatic sulfide differentiation were clearly part of a single, continuous process in the Sudbury footwall. On the basis of our new observations, metal transport and redistribution by volatiles occurred twice during the evolution of the footwall ore systems. Figure 14 and the summary below present an interpretation of the relative timing of volatile events in the footwall environment in relation to ore deposition, based on our new observations: Since the $P_{\text{brine}} + P_{\text{CH}_4}$ assemblage predates (texturally) the last stages of sulfide crystallization in the alteration veins, it is clear that hydrosaline fluids were present in the footwall immediately prior to or during massive sulfide vein emplacement (pre-solidus; Fig. 14). After the formation of the metal-rich brine (i.e., by mixing of groundwater with a magmatic ore fluid), the brine entered the footwall environment via structural pathways (Sudbury Breccia zones) and exsolved an immiscible CH_4 phase during local decompression during vein formation or by cooling of the brine as it came into contact with the host vein wall rocks. Phase separation may have been responsible for some Au precipitation in the alteration veins (Naden and Shepherd 1989). In addition to precipitating Au, phase separation caused Cu, Au, Ag and Bi to partition into the immiscible CH_4 ; these ore metals may have been

dispersed into the host rocks by the low density CH_4 fluid, explaining the presence of anomalous concentrations of Cu and Au at great distances (up to 150 m) from footwall ores (Hanley 2002). Entrapment of the $\text{P}_{\text{brine}} + \text{P}_{\text{CH}_4}$ assemblage could have predated or occurred synchronously with the emplacement of the footwall sulfide veins. As the emplacement of fractionated sulfide liquid into the footwall setting proceeded according to the quantitative models proposed by Naldrett and Pessaran (1992) and Li (1993), sulfides appear to have migrated into the quartz-epidote veins, replacing the earlier-formed silicates. As the sulfides approached their solidus temperature, molten NaCl and a hydrocarbon fluid may have exsolved from the sulfides, being trapped along secondary fractures in the quartz-epidote veins ($\text{S}_{\text{halite}} + \text{S}_{\text{CH}_4}$ assemblage). At this stage, Cu, PGE, Au and other metals were extracted from the sulfides by the exsolving volatiles and *possibly* redeposited as Pt–Pd–Te–Bi–Sn minerals, electrum, and disseminated chalcopyrite in the Cl-rich mineral assemblage that now occur along massive sulfide vein margins and in the adjacent host rocks as observed by previous authors (Table 1). The final distribution of the PGE and Au in the footwall ores may have been controlled by the halide melt-hydrocarbon exsolution, but the molten NaCl could have only persisted to temperatures as low as $\sim 700^\circ\text{C}$ (Fig. 9). Therefore, successive cooling and reequilibration of the ore mineral assemblage along the vein margins would be required to recrystallize the PGE minerals to the lower temperature assemblages observed today ($< 600^\circ\text{C}$; Table 1).

Fig. 15 Log-log plot of Cu:Pt versus Cu:Au ratios for massive sulfide ore (Li and Naldrett 1994), quartz-sulfide-epidote alteration veins, primary brine and hydrocarbon inclusions, and halide-melt inclusions. Boxes show range in Cu:Au and Cu:Pt observed in the volatile-bearing inclusions (Table 1 and Electronic Supplementary Material). Cu:Au ratios in massive sulfide samples (open circles) vary over three orders of magnitude and cannot be readily explained using traditional models for sulfide liquid fractionation within the ore zones (Li 1993). Alteration veins have lower Cu:Au ratios similar to Au-enriched massive sulfide ores whereas Cu:Pt ratios in alteration veins and massive sulfides are comparable. Inclusion compositions (brine-hydrocarbon and halide melt phases) are highly enriched in Au and Pt over Cu when compared to the massive sulfide ores and inclusion-hosting quartz-sulfide-epidote veins

Transport of ore metals by volatiles in the footwall environment

The high concentrations of Cu, Pt, Au, Ag and Bi observed in the P_{brine} , P_{CH_4} and S_{halite} inclusions confirm previous suggestions (e.g., Farrow 1994) that volatiles were important carriers of ore metals in the Sudbury environment over a wide range of temperatures. It is impossible to evaluate (quantitatively) the importance of the volatiles in modifying the original composition of the sulfide liquids or sulfide ores within the footwall environment; this would require thermochemical data for fluid-sulfide and halide melt-sulfide partitioning behavior and knowledge of the fluid:rock ratios associated with the footwall ore-forming system, which at present are non-existent. However, the data presented here offer some new insight into the behavior of metals in the Sudbury magmatic-hydrothermal environment.

For example, it is interesting to note that Cu:Pt and Cu:Au ratios in the volatile-bearing inclusions are significantly lower than in the associated massive sulfides and inclusion-hosting alteration veins (Fig. 15). In the halide-melt phase, Cu:Au and Cu:Pt ratios are ~ 1 – 2 orders of magnitude lower than the suggested parent sulfides; this may reflect the original partitioning behavior of the ore metals between coexisting sulfide and halide melt. The apparent halide melt:sulfide liquid partition coefficient is low for Cu (e.g., around ~ 0.02 based on $\sim 4,500$ ppm Cu in the halide-melt versus $\sim 270,000$ ppm Cu in the massive chalcopyrite–cubanite ore). However, Pt and Au concentrations in the halide-melt are comparable to the concentrations within the ores themselves, tentatively suggesting that partition coefficients for Pt and Au between sulfide and halide-melt are approximately unity. Consequently, models for magmatic Ni–Cu–PGE ore formation must be revised to consider that volatiles and coexisting sulfide liquids may compete for precious metals during their interaction and may fractionate the precious metals from Cu at the magmatic–hydrothermal transition.

Similarly, Au and Pt concentrations in the brine inclusions are comparable to the ores, but Cu:Au and Cu:Pt ratios are up to four orders of magnitude lower than the associated sulfides; this observation may reflect the relative solubility behavior of Cu and Au (or Pt). In a relatively low $f\text{O}_2$ system such as the footwall ore environment, Au and Pt solubility in the brine phase may have been enhanced relative to Cu if the predominant ligand was bisulfide for Au and Pt and chloride for Cu. In relatively reduced ore fluids, hydrosaline volatile phases can transport only modest quantities of Cu (10–100 ppm-range) and the solubility of Cu as a chloride complex has been shown to *decrease* with decreasing temperature, chloride concentration and $f\text{O}_2$ (Hemley et al. 1992); this observation is consistent with the higher Cu content of the higher temperature, higher salinity halide melt phase. However, Au solubility by bisulfide or chloride complexation may be significant regardless of the oxidation state of the system and has been shown to

increase upon cooling due to the transition from chloride complexation to the more soluble bisulphide complex (Gammons and Williams-Jones 1997; Mountain and Seward 1999). On the basis of the composition of alteration assemblages in the footwall environment at Sudbury, Wood (2002) showed that significant Pt transport by chloride was unlikely and that the conditions were more favorable for the transport of precious metals by bisulfide. The possibility of selective metal transport by hydrocarbon-based ligands in the lower temperature volatiles must also be considered and may explain the elevated concentrations of ore metals in the CH₄ vapor phase. Recent experimental evidence suggests that organic ligands such as acetate (Wood 2002) and porphyrin compounds (naturally occurring metabolic pigments; Williams-Jones, personal communication) may transport significant concentrations of PGE at hydrothermal conditions. Cooling of the brine, in which Cu and Au or Pt were transported by different ligands with the contrasting solubility behavior described, could lower the Cu/Au and Cu/Pt ratios in the brine considerably. (Note that Cu precipitation from a fluid may be indicated by the presence of disseminated chalcopyrite in the adjacent host rocks to the massive sulfide veins). Simple cooling has been shown to lower the Cu:Au ratio of ore-forming fluids in other magmatic-hydrothermal deposit types and may be attributed to a similar mechanism (e.g., Cu–Au porphyry systems; Ulrich et al. 2001).

Implications for exploration

During hydrocarbon, brine and halide-melt evolution in the footwall, Au and Cu transport and dispersion into the brecciated host rocks may have been substantial. Late-stage hydroxysilicate alteration throughout ore-bearing breccia zones in the footwall deposits confirms that the host rocks to the deposits were relatively fluid-permeable within ~200 m of ore (Jago et al. 1994; Morrison et al. 1994; Hanley and Mungall 2003; Hanley et al. 2004). Consequently, the occurrence of large hydrocarbon, halogen, or metal enrichment halos around massive sulfide deposits may serve as pathfinders for footwall-style ore deposit at Sudbury and fractionated ore styles in other magmatic Ni–Cu–PGE systems. Notably, CH₄ does not occur in significant concentrations in the deep groundwaters along the northern margin of the SIC (<0.2 vol%; Frape and Fritz 1982) or in brecciated host rocks from unmineralized areas along the northern margin of the SIC (Hanley 2002). If massive sulfide ore bodies are considered to be the sources of hydrocarbon fluids, then bulk analysis of host rocks for their CH₄ content may identify important hydrocarbon signatures (e.g., the total abundance of CH₄) in proximity to mineralization. Gas chromatography of volatiles released during crushing of rock samples offers an inexpensive method for detecting fluid inclusion-hosted hydrocar-

bons at extremely low concentrations (Bray and Spooner 1992).

Conclusions

The results of this study show that CH₄ and other hydrocarbon species were important volatile components during the evolution of the footwall-style Cu–PGE deposits at Sudbury. Of major significance to the origin of the footwall ore systems is the observation that the inclusion assemblages characterized in this study are very rich in Cu (up to ~1 wt%) and contain Pt and Au in the ppm-range. Together with high salinity brines and halide melts, hydrocarbons remobilized and dispersed ore metals (Cu, Pt, Au, Ag and Bi) at least twice during the evolution of the deposits: first, by coexisting (immiscible) metal-rich CH₄ and brine during or prior to sulfide emplacement into the footwall setting, and second, by a molten salt phase and hydrocarbon vapor that may have exsolved from the sulfides during their final stage of crystallization. The presence of hydrocarbons coexisting with a high-*T* halide-melt phase is, as discussed, problematic due to the equilibrium instability of pure hydrocarbon species at such high *T*. Therefore, we admit that the halite inclusions may have been formed by some other process (e.g., precipitated from a halite-saturated hydrocarbon fluid phase at lower *T*). The origin of the hydrocarbons coexisting with the S_{halite} inclusions is unclear at present and awaits future study.

The interesting partitioning of Cu, Au, Bi and Ag during early phase separation and mingling of immiscible CH₄ and brine is consistent with observations from fluid inclusion boiling assemblages in Cu–Au porphyry deposits (e.g., Heinrich et al. 1999; Ulrich et al. 2001) and identifies similar processes operating in two very different magmatic-hydrothermal ore deposit systems. The high concentrations of Cu, Pt, Au, Ag and Bi observed in the P_{brine}, P_{CH₄} and S_{halite} inclusions add considerable support to hydrothermal models (e.g., Farrow 1994) for the formation of footwall ore zones at Sudbury and confirm that volatiles were important carriers of ore metals over a wide range of temperatures. Overall, the early unmixing of CH₄ and brine, and later exsolution of H₂O-poor volatiles (hydrocarbons, halide melt) from the sulfide ores may have been two principal mechanisms controlling the final distribution of PGE and Au in the footwall ore systems at Sudbury and may explain the occurrence of distal metal anomalies in ore-hosting breccia zones, the decoupling of Au from S and the PGE, and the occurrence of PGE minerals with Cl-rich alteration minerals along sulfide vein margins and in adjacent host rocks.

Acknowledgements The authors wish to thank Dave King and Kevin Chisholm (Falconbridge Ltd.) for authorizing access to underground workings at the Fraser Mine and assisting with sample collection. Funding to J. M. from NSERC and PREA

(Premiere's Research Excellency Awards) was greatly appreciated. Support for J.H. came from an NSERC postgraduate scholarship. The Swiss National Science Foundation is acknowledged for their continued support of the LA-ICP-MS facility at ETH Zurich (Isotope Geology and Mineral Resources). The manuscript was improved through critical reviews by D. Marshall, D. Ebel and L. Meinert and benefited from discussions with B. Sherwood-Lollar and I. Samson.

References

- Atkinson B (2002) A model for the PTZ properties of H_2O – NaCl . PhD Thesis, Virginia Polytechnic Institute, pp 133
- Ballhaus CG, Stumpfl EF (1986) Sulphide and platinum mineralization in the Merensky Reef: evidence from hydrous silicates and fluid inclusions. *Contrib Mineral Petrol* 94:193–204
- Bodnar RJ (1994) Synthetic fluid inclusions; XII, the system H_2O – NaCl ; experimental determination of the halite liquidus and isochores for a 40 wt% NaCl solution. *Geochim Cosmochim Acta* 53:725–733
- Bowers TS, Helgeson HC (1983) Calculation of the thermodynamic and geochemical consequences of nonideal mixing in the system H_2O – CO_2 – NaCl on phase relations in geological systems; equation of state for H_2O – CO_2 – NaCl fluids at high pressures and temperatures. *Geochim Cosmochim Acta* 47:1247–1275
- Bray CJ, Spooner ETC (1992) Fluid inclusion volatile analysis by gas chromatography with photoionization/micro-thermal conductivity detectors; applications to magmatic MoS_2 and other H_2O – CO_2 and H_2O – CH_4 fluids. *Geochim Cosmochim Acta* 56:261–272
- Coats CJA, Snajdr P (1984) Ore deposits of the North Range, Onaping-Levack Area. In: Pye EG, Naldrett AJ, Giblin PE (eds) The geology and ore deposits of the Sudbury structure. Ontario Geological Survey special volume 1:327–346
- Cowan JC (1968) Geology of the Strathcona ore deposit. *Can Inst Mining Metallurg Bull* 61:38–45
- Dietz RS (1964) Sudbury structure an Astrobleme. *J Geol* 72:412–434
- Dressler BO (1984) The effects of the Sudbury event and the intrusion of the Sudbury Igneous Complex on the Footwall Rocks of the Sudbury structure. In: Pye EG, Naldrett AJ, Giblin PE (eds) The geology and ore deposits of the Sudbury structure. Ontario Geological Survey spec vol 1:97–136
- Ebel DS, Naldrett AJ (1997) Crystallization of sulfide liquids and the interpretation of ore composition. *Can J Earth Sci* 34:352–365
- Erwood RJ, Kesler SE, Cloke PL (1979) Compositionally-distinct, saline hydrothermal solutions, Naica Mine, Chihuahua, Mexico. *Econ Geol* 74:95–108
- Farrow CEG (1994) Geology, alteration, and the role of fluids in Cu–Ni–PGE mineralization of the footwall rocks to the Sudbury Igneous Complex, Levack and Morgan Townships, Sudbury District, Ontario. PhD Thesis, Carleton University
- Farrow CEG, Lightfoot PC (2002) Sudbury PGE revisited: towards and integrated model. In: Cabri LJ (ed) The geology, geochemistry, mineralogy and mineral beneficiation of platinum-group elements. Canadian Institute of Mining, Metallurgy and Petroleum, Montreal, pp 273–297
- Farrow CEG, Watkinson DH (1992) Alteration and the role of fluids in Ni, Cu and platinum-group element deposition, Sudbury igneous complex contact, Onaping–Levack Area, Ontario. *Mineral Petrol* 46:67–83
- Frape SK, Fritz P (1982) The chemistry and isotopic composition of saline groundwaters from the Sudbury basin, Ontario. *Can J Earth Sci* 19:645–661
- Frost BR, Touret JLR (1989) Magmatic CO_2 and saline melts from the Sybille Monzosyenite, Laramie anorthosite complex, Wyoming. *Contrib Mineral Petrol* 103:178–186
- Gammons CH, Williams-Jones AE (1997) Chemical mobility of gold in the porphyry-epithermal environment. *Econ Geol* 92:45–59
- Goldstein RH (2003) Petrographic analysis of fluid inclusions. In: Samson I, Marshall D, Anderson A (eds) Fluid inclusions: analysis and interpretation. Mineralogical Association of Canada, Short Course Series Vol 32:9–54
- Greenman L (1970) The petrology of the footwall breccias in the vicinity of the Strathcona Mine, Levack, Ontario. PhD Thesis, University of Toronto
- Grieve RAF (1994) An impact model of the Sudbury structure: proceedings of the Sudbury. In: Lightfoot PC, Naldrett AJ (eds) Proceedings of the Sudbury–Noril'sk symposium. Ontario Geol Surv Spec Vol 5:119–132
- Gunther D, Frischknecht R, Heinrich CA (1997) Capabilities of a 193 nm ArF excimer laser for LA-ICP-MS microanalysis of geological materials. *J Anal At Spectrom* 12:939–944
- Hanley JJ (2002) The distribution of the halogens in Sudbury Breccia matrix as as pathfinder elements for footwall Cu–PGE mineralization at the Fraser Cu Zone, Barnet Main Copper Zone, and surrounding margin of the Sudbury Igneous Complex, Onaping-Levack Area, Ontario, Canada. MSc Thesis, University of Toronto
- Hanley JJ, Mungall JE (2003) Chlorine enrichment and hydrous alteration of the Sudbury Breccia hosting footwall Cu–Ni–PGE mineralization at the Fraser mine, Sudbury, Ontario, Canada. *Can Mineral* 41:857–881
- Hanley JJ, Mungall JE, Bray CJ, Gorton MP (2004) The origin of bulk and water-soluble Cl and Br enrichments in ore-hosting Sudbury Breccia in the Fraser Copper Zone, Strathcona Embayment, Sudbury, Ontario, Canada. *Can Mineral* 42:1777–1798
- Heinrich CA, Ryan CG, Mernagh TP, Eadington PJ (1992) Segregation of ore metals between magmatic brine and vapor; a fluid inclusion study using PIXE microanalysis. *Econ Geol* 87:1566–1583
- Heinrich CA, Gunther D, Audetat A, Ulrich T, Frischknecht R (1999) Metal fractionation between magmatic brine and vapor, determined by microanalysis of fluid inclusions. *Geology* 27:755–758
- Heinrich CA, Pettke T, and Halter WE (2003) Quantitative multi-element analysis of minerals, fluid and melt inclusions by laser-ablation inductively-coupled-plasma mass-spectrometry. *Geochim Cosmochim Acta* 67:3473–3497
- Hemley JJ, Cygan GL, Fein JB, Robinson GR, D'Angelo WM (1992) Hydrothermal ore-forming processes in the light of studies in rock-buffered systems: I. Fe–Cu Zn–Pb sulphide solubility relationships. *Econ Geol* 87:1–22
- Jago BC, Morrison GG, Little TL (1994) Metal zonation patterns and microtextural and micromineralogical evidence for alkali- and halogen-rich fluids in the genesis of the Victor Deep and McCreedy East Footwall Copper Ore bodies, Sudbury Igneous Complex. In: Lightfoot PC, Naldrett AJ (eds) Proceedings of the Sudbury–Noril'sk Symposium. Ontario Geol Surv Spec Vol 5:65–75
- Krader T, Franck EU (1987) The ternary systems H_2O – CH_4 – NaCl and H_2O – CH_4 – CaCl_2 to 800 K and 250 MPa. *Ber Bunsenges Phys Chem* 9:627–634
- Krogh TE, Davis DW, Deutch A (1984) Precise U–Pb zircon and baddeleyite ages for the Sudbury Area. In: Pye EG, Naldrett AJ, Giblin PE (eds) The geology and ore deposits of the Sudbury structure. Ontario Geological Survey Special Volume 1:431–446
- Li C (1993) A quantitative model for the formation of sulfide ores at Sudbury and a study on the distributions of platinum-group elements in the Strathcona copper-rich zones, Sudbury, Ontario. PhD Thesis, University of Toronto
- Li C, Naldrett AJ (1994) A numerical model for the compositional variations of Sudbury sulfide ores and its application of exploration. *Econ Geol* 89:1599–1607
- Longerich HP, Jackson SE, Gunther D (1996) Laser ablation inductively coupled plasma mass spectrometry transient signal data acquisition and analyte concentration calculation. *J Anal Atom Spectrom* 11:899–904

- Marshall D, Watkinson D, Farrow CEG, Molnar F, Fouillac AM (1999) Multiple fluid generations in the Sudbury igneous complex: fluid inclusion, Ar, O, H, Rb and Sr evidence. *Chem Geol* 154:1–19
- Mathez EA (1987) Carbonaceous matter in mantle xenoliths; composition and relevance to the isotopes. *Geochim Cosmochim Acta* 51:2339–2347
- Mathez EA, Deitrich VJ, Holloway JR, Boudreau AE (1989) Carbon distribution in the Stillwater Complex and evolution of vapor during crystallization of Stillwater and Bushveld magmas. *J Petrol* 30:153–173
- Mavrogenes JA, Berry AJ, Newville M, Sutton SR (2002) Copper speciation in vapor-phase fluid inclusions from the Mole Granite, Australia. *Am Mineral* 87:1360–1364
- Molnar F, Watkinson DH, Jones PC (2001) Multiple hydrothermal processes in footwall units of the North Range, Sudbury igneous complex, Canada, and implications for the genesis of vein-type Cu–Ni–PGE deposits. *Econ Geol* 96:1645–1670
- Morrison GG, Jago BC, White TL (1994) Footwall mineralization of the Sudbury Igneous Complex. In: Lightfoot PC, Naldrett AJ (eds) *Proceedings of the Sudbury–Noril'sk symposium*. Ontario Geol Surv Spec Vol 5:57–64
- Mountain BW, Seward TM (1999) The hydrosulphide/sulphide complexes of copper (I); experimental determination of stoichiometry and stability at 22°C and reassessment of high temperature data. *Geochim Cosmochim Acta* 63:11–29
- Mungall JE, Brennan JM (2003) Experimental evidence for the chalcophile behaviour of the halogens. *Can Mineral* 41:207–220
- Mungall JM, Ames DE, Hanley JH (2004) Geochemical evidence from the Sudbury structure for crustal redistribution by large bolide impacts. *Nature* 429:546–548
- Naden J, Shepherd T (1989) Role of methane and carbon dioxide in gold deposition. *Nature* 342:793–795
- Naldrett AJ, Kullerud G (1967) A study of the Strathcona Mine and its bearing on the origin of the nickel–copper ores of the Sudbury District, Ontario. *J Petrol* 8:453–531
- Naldrett AJ, Pessaran A (1992) Compositional variation in the Sudbury ores and prediction of the proximity of footwall copper–PGE orebodies. In: Milne V (ed) *Ontario Geol Survey Misc. Paper* 159:47–62
- Pettke T, Heinrich CA, Ciocan AC, Gunther D (2004) Quadrupole mass spectrometry and optical emission spectroscopy: detection capabilities and representative sampling of short transient signals from laser-ablation. *J Anal Atom Spectrom* 9:1149–1155
- Pettke T, Diamond LW, Kramers JD (2000) Mesothermal gold lodes in the north-western Alps; a review of genetic constraints from radiogenic isotopes. *Eur J Mineral* 12:213–230
- Poling BE, Prausnitz JM, O'Connell JP (2001) *The properties of gases and liquids*. McGraw-Hill, New York, pp 752
- Potter J, Konnerup-Madsen J (2003) A review of the occurrence and origin of abiogenic hydrocarbons in igneous rocks. In: Petford N, McCaffrey KJW (eds) *Hydrocarbons in crystalline rocks*. Geological Society of London, Spec Pub 214:151–173
- Renders PJ, Seward TM (1989) The stability of hydrosulphido- and sulphido-complexes of Au (I) and Ag (I) at 25°C. *Geochim Cosmochim Acta* 53:245–253
- Renno AD, Leander F, Witzke T, Herzig PM (2004) The coexistence of melts of hydrous copper chloride, sulfide and silicate compositions in a magnesiohastingsite cumulate, TUBAF seamount, Papua New Guinea. *Can Mineral* 42:1–16
- Salvi S, Williams-Jones AE (1992) Reduced orthomagmatic C–O–H–N–NaCl fluids in the Strange Lake rare-metal granitic complex, Quebec/Labrador, Canada. *Eur J Mineral* 4:1155–1174
- Salvi S, Williams-Jones AE (1997) Fisher-Tropsch synthesis of hydrocarbons during sub-solidus alteration of the Strange Lake peralkaline granite, Quebec/Labrador, Canada. *Geochim Cosmochim Acta* 61:83–99
- Shepeleva NN, Ogloblina AJ, Pikovskiy YI (1990) Polycyclic aromatic hydrocarbons in Carbonaceous matter of Daldyn-Alakit area of the Siberian Platform. *Geokhimiya* 5:731–741
- Shock E (1990) Geochemical constraints on the origin of organic compounds in hydrothermal systems. *Orig Life Evol Biosph* 20:331–367
- Sisson VB, Crawford ML, Thompson PH (1981) CO₂-brine immiscibility at high temperatures, evidence from calcareous metasedimentary rocks. *Contrib Mineral Petrol* 78:371–378
- Sourirajan S, Kennedy GC (1962) The system NaCl–H₂O at elevated temperatures and Pressures. *Am J Sci* 260:115–141
- Sugisaki R, Mimura K (1994) Mantle hydrocarbons; abiotite or biotic? *Geochim Cosmochim Acta* 58:2527–2542
- Ulrich T, Gunther D, Heinrich CA (2001) The evolution of a porphyry Cu–Au deposit, based on LA-ICP-MS analysis of fluid inclusions; Bajo de la Alumbrera, Argentina. *Econ Geol* 96:1743–1774
- Wood SA (2002) The aqueous geochemistry of the platinum-group elements with applications to ore deposits. In: Cabri LJ (eds) *The geology, geochemistry, mineralogy and mineral beneficiation of platinum-group elements*. Canadian Institute of Mining Metallurgy and Petroleum, Montreal, pp 211–249
- Ypma PJM, Fuzikawa K (1980) Fluid inclusion and oxygen isotope studies of the Nabarlek and Jabiluka uranium deposits, Northern Territory, Australia. In: Ferguson J, Goleby A (eds) *Proceedings of the International Uranium symposium on the Pine Creek Geosyncline*. Proc Series Intern Atomic Energy Agency 555:375–395
- Zhang Y, Franz JD (1987) Determination of the homogenization temperatures and densities of supercritical fluids in the system NaCl–KCl–CaCl₂–H₂O using synthetic fluid inclusions. *Chem Geol* 64:335–350