



Monopole quivers and new 3D $\mathcal{N} = 2$ dualities

Antonio Amariti, Domenico Orlando, Susanne Reffert*

Albert Einstein Center for Fundamental Physics, Institute for Theoretical Physics, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland

Received 24 July 2017; accepted 10 September 2017

Available online 15 September 2017

Editor: Stephan Stieberger

Abstract

We present a new family of dualities for three-dimensional gauge theories, motivated by the brane realization of the reduction of four-dimensional dualities on a circle. This family can be understood as a generalization of Aharony duality to quiver gauge theories whose nodes interact via monopole terms in the superpotential. We refer to this family of theories as monopole quivers. We corroborate the new dualities by checking the equivalence of the three-sphere partition functions, obtained from the standard circle reduction of the four-dimensional superconformal index. As a special case, we recover some dualities recently discussed in the literature.

© 2017 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The description of the low-energy dynamics of ultra-violet (UV) free quantum field theories (QFTs) often requires the use of non-perturbative techniques. A different paradigm consists in describing the theory in terms of dual degrees of freedom, hopefully weakly coupled at low energy scales. Large classes of such dualities have been worked out in the case of supersymmetric QFTs, mostly thanks to the power of holomorphy. When considering four-dimensional theories with the minimal amount of supersymmetry, a rich duality web has been obtained, based on extensions and deformations of Seiberg duality [1]. It has been quickly realized that similar dualities exist for three-dimensional $\mathcal{N} = 2$ theories [2–7]. The similarity has been explained

* Corresponding author.

E-mail address: sreffert@gmail.com (S. Reffert).

in [8], where three-dimensional dualities have been derived from a circle compactification of four-dimensional Seiberg dualities.

The web of three-dimensional dualities turns out to be richer than the one in four dimensions, mostly due to the presence of a Coulomb branch for three-dimensional theories with four supercharges. This Coulomb branch is described in terms of monopole operators, associated to chiral fields consisting of a combination of a real adjoint scalar in the vector multiplet and of the dual photon. These operators have been used to enlarge the spectrum of the three-dimensional dualities.

In general, new dualities emerge if the compactification limit is taken while performing real mass flows and Higgsing the gauge group in non-trivial vacua. A complete classification is still lacking to date and a full understanding of the behavior of the monopole operators in a circle compactification is a necessary step in this direction.

The reduction of four-dimensional dualities to three dimensions simplifies when the gauge theories are engineered in a setup of intersecting branes [9]. The circle reduction corresponds to a T-duality from the type IIA description of the four-dimensional theory to the type IIB description of the three-dimensional theory. The power of this picture stems from the fact that the monopole operators have a straightforward realization in terms of Euclidean D1-branes. Dual configurations in presence of non-trivial vacuum structures can be directly obtained via a Hanany–Witten [10] (HW) transition.

In this paper, we show that a very rich spectrum of new three-dimensional $\mathcal{N} = 2$ dualities emerges from reducing four-dimensional dualities in the brane setup and considering non-trivial vacuum configurations. In field theory, they correspond to products of three-dimensional supersymmetric quantum chromodynamics (SQCD)-like sectors interacting through Affleck–Harvey–Witten [11] (AHW) superpotentials. They can be represented as generalized quiver gauge theories, where the gauge nodes interact through monopole operators. These monopole operators, responsible for the coupling of the various gauge nodes, reconstruct the original Kaluza–Klein (KK) monopole of the circle compactification.

In terms of the brane set-up, we will be using the same technique of the circle compactification as in [9] and consider vacuum configurations obtained by spreading several stacks of D3 branes on the circle, attaching them to D5 branes. The resulting theory is thus akin to a quiver gauge theory in which each of the stacks of D3s corresponds to a node in the quiver. We will however take the circle radius to be very large, in which case the bifundamental fields, corresponding to fundamental strings stretched between the stacks vanish, while the monopoles, corresponding as before to D1 branes stretched between the stacks of D3s remain. The stacks of D3s thus only interact via the monopoles, which is why we coin the term *monopole quiver* for this brane construction.

Using this set-up as a starting point, we can perform a variety of duality transformations via HW transitions just as in [9,12], resulting in new dualities.

The plan of this paper is as follows. In Section 2, we review the brane construction introduced in [9] on which the dualities are realized as HW moves. In Section 3, we derive a new duality between two monopole quivers with unitary gauge groups by reducing four-dimensional Seiberg duality on S^1 and by performing a large real mass and Higgs flow to recover the three-dimensional limit. In Section 4 we discuss how to recover the dualities described in [7] via our brane set-up. The generalization of the picture to monopole quivers with Sp gauge groups is discussed in Section 5. In Sec. 6, we end with concluding remarks and an outlook. In Appendix A, we discuss the three-dimensional limit for the monopole quiver duality with a generic amount of unitary gauge groups, showing the matching of the partition function between the dual phases af-

Table 1
 Brane configuration for $\mathcal{N} = 1$ SYM on $\mathbb{R}^3 \times S^1$. The D3-branes are suspended between the two NS5s.

	0	1	2	3	4	5	6	7	8	9
NS	×	×	×	×	×	×				
NS'	×	×	×	×					×	×
D3	×	×	×				×			




Fig. 1. D1-branes (in grey) stretched between D3-branes and NS5 branes.

ter the real mass and Higgs flow. [Appendix B](#) is a review of the reduction of the four-dimensional superconformal index to the three-dimensional partition function on a squashed three-sphere, serving also to fix notation and conventions.

2. The brane set-up on the circle

Circle compactification Let us first review the circle compactification as introduced in [9]. We start with the brane set-up for $\mathcal{N} = 1$ $U(N)$ super Yang–Mills (SYM) on $\mathbb{R}^3 \times S^1$, denoting by R_3 the radius of S^1 . It can be arrived at by a T-duality from a stack of N D4 branes suspended between an NS5 and an NS5' brane at distance ℓ_6 in the x^6 direction. Performing the T-duality along the compact x^3 direction, the D4s turn into D3s, while the NS5 branes remain unchanged. The resulting configuration is summarized in [Table 1](#).

The corresponding gauge theory has N isolated vacua, corresponding to stable supersymmetric configurations of the brane system. There is a repulsive force between the D3s and, in a stable configuration, the D3 branes are distributed along the circle direction x^3 at equal distances. All moduli are lifted as the D3s cannot move freely.

Let us discuss a moment the origin of this repulsive force. From the field theory point of view in three dimensions, instantons induce a non-perturbative superpotential. The instantons are represented in the brane picture by Euclidean D1s stretched between each pair of D3 branes along x^3 and in x^6 and the NS and NS' branes (see [Fig. 1](#)).

The D1-branes give rise to the superpotential [13]

$$W = \sum_i e^{-S_{D1_i}} = \sum_{i=1}^{N-1} \exp[\Sigma_{i+1} - \Sigma_i], \tag{2.1}$$

with $\Sigma_i = \sigma_i/e_3^2 + i\phi_i$, where the scalar σ_i parameterizes the position of the i -th D3, ϕ is the dual photon, and $e_3^2 = 2\pi\sqrt{\alpha'}/(R_3\ell_6)$ is the three-dimensional gauge coupling.¹ Similarly, a Euclidean fundamental string stretched between the NS5 and the D3 branes will give a contribution

$$e^{-S_{F1}} = \exp\left[-\frac{\sqrt{\alpha'} \Delta \Sigma}{R_3 e_3^2}\right], \tag{2.2}$$

which vanishes in the $R_3 \rightarrow 0$ limit that we take.

Monopole quivers We can avoid the breaking of the gauge group to $U(1)^N$ due to the mutual repulsion of the D3 branes by attaching them to D5 branes. We will consider only the stable case $\#D5 > \#D3$ as we are interested in the dimensional reduction of Seiberg duality. In the type IIA frame, we start with F D6-branes extended along 0123789 and sitting on the NS'-brane. The D6 branes become D5-branes after T-duality. The strings stretched between the stack of D3-branes and the D5s correspond to F massless fundamentals Q and anti-fundamentals \tilde{Q} .

When D5-branes sitting at $x_3 = 0$ intersect the worldsheet of the D1-strings, they contribute two additional zero modes to the D1-instanton and the superpotential in Eq. (2.1) is not generated. So the D5-branes have the effect of screening the repulsive force between the D3-branes [16,17].

The effective three-dimensional theories on S^1 can undergo non-trivial real mass or Higgs flows. These flows are often necessary in order to recover conventional three-dimensional theories, allowing to preserve dualities when taking the zero-radius limit. In the brane setup, this corresponds to moving stacks of n_i D3s and f_i D5s along the circle, with $f_i > n_i$. The resulting theory corresponds to a product of $U(n_i)$ gauge factors, each with f_i flavors. In the infrared (IR), these gauge sectors do not interact through matter fields. Considering each sector as decoupled, the 1-st and the n_i -th D3-brane in each stack are free to move without being subjected to any force, so two directions in the moduli space seem to remain unlifted in each sector. These directions are however lifted by an AHW superpotential between the i -th and the $i + 1$ -th gauge sectors. Such a superpotential has the form

$$W = \exp\left[i \left(\Sigma_i^{(1)} - \Sigma_{i+1}^{(n_{i+1})}\right)\right] = T_i \tilde{T}_{i+1}, \tag{2.3}$$

where $\Sigma_i^{(j)}$ refers to the j -th brane in the i -th stack. In gauge theoretical terms, $\Sigma_i^{(1)}$ and $\Sigma_i^{(n_i)}$ are the bare monopoles of the $U(n_i)$ theory with flux $(1, 0, \dots, 0)$ and $(0, 0, \dots, -1)$, respectively. We will use the notation T for the former and \tilde{T} for the latter.

The resulting theory is thus akin to a quiver gauge theory in which each of the stacks of D3s corresponds to a node in the quiver. When we take the (dual) circle radius to be very large, the bifundamental fields corresponding to fundamental strings stretched between the stacks vanish, while the monopoles corresponding to the D1 branes stretched between the stacks of D3s remain, due to their different R_3 -dependence in the action, as pointed out above. The stacks of D3s thus only interact via the monopoles, which is why we refer to the resulting quiver as *monopole quiver*. The simplest case, with two gauge groups and superpotential

$$W = T_1 \tilde{T}_2 + \tilde{T}_1 T_2, \tag{2.4}$$

¹ In this configuration (without D5-branes) there is also an extra contribution coming from the D1s between the N -th and the first D3 brane which gives rise to the η -superpotential [14,15] $W_\eta = \eta e^{\Sigma_N - \Sigma_1}$ as discussed in [9]. This will in general not play a role for our monopole quivers where η amounts to a field redefinition.




Fig. 2. A monopole quiver. The gauge groups $U(n_1)$ and $U(n_2)$ are represented as circles, the flavors as squares and the monopole interactions as darts joining the first brane in the first group to the last in the second group and the first brane in the second group to the last one in the first group.

is represented in Fig. 2. Using this set-up as a starting point, we can perform a variety of duality transformations via HW transitions just as in [9,12,18], resulting in new dualities.

Adding orientifolds Adding orientifold planes to our brane set-up allows us to construct theories with real orthogonal and symplectic gauge groups and matter fields in the symmetric and antisymmetric representation of the gauge group, see e.g. the reviews [19,18]. An orientifold is the combined action of a parity inversion σ of the coordinates transverse to the plane, a world-sheet parity Ω and $(-1)^{F_L}$, F_L being the left-moving fermion number. There are p -dimensional orientifold planes (O_p planes) with even p in type IIA and odd p in type IIB. We will study the case of $p = 4$ in type IIA. There are in total four possibilities, O_p^\pm and \widetilde{O}_p^\pm [20,21]. Specifying the \mathbb{Z}_2 charges characterizes the action of the orientifold on the gauge theory completely.

We will add an $O4$ plane on top of the stack of D4 branes realizing the four-dimensional gauge theory. It can be shown that

- for $O4^-$ we get gauge group $SO(2N)$,
- for $O4^+$ we have $Sp(2N)$,
- for $\widetilde{O}4^-$ we have $SO(2N + 1)$ and
- for $\widetilde{O}4^+$ we get again $Sp(2N)$ but with a different non-perturbative sector.

The three-dimensional system is obtained by compactifying the x_3 -direction and T-dualizing. In the presence of a compact direction, orientifold planes come in pairs located at $x_3 = 0$ and at the so-called mirror point $x_3^\circ = \frac{\alpha'}{R_3}\pi$ [22]. Here we restrict our attention to the $O4^+$ -plane, that will turn into a pair of $(O3^+, O3^+)$ -planes after T-duality.

3. A new 3D duality from 4D

In this section we derive a new duality by reducing four-dimensional Seiberg duality on S^1 and performing a large real mass flow to recover the three-dimensional limit.

The final duality can be summarized as follows:

- The electric theory is a $U(n_1) \times U(n_2)$ gauge theory with f_1 flavors Q_1 and \widetilde{Q}_1 in the first sector and f_2 flavors Q_2 and \widetilde{Q}_2 in the second sector, with $f_i > n_i$. There is also a superpotential

$$W = \exp[i \left(\Sigma_1^{(1)} - \Sigma_2^{(n_2)} \right)] + \exp[i \left(\Sigma_2^{(1)} - \Sigma_1^{(n_1)} \right)] \equiv T_1 \widetilde{T}_2 + \widetilde{T}_1 T_2, \tag{3.1}$$

Table 2

Field charges in the electric phase of the duality for the $U(n_1) \times U(n_2)$ monopole quiver.

	$U(n_1)$	$U(n_2)$	$SU(f_1)$	$SU(f_2)$	$U(1)_A$	$U(1)_J$	$U(1)_R$
Q_1	n_1	1	f_1	1	$1/f_1$	0	Δ_1
\tilde{Q}_1	\bar{n}_1	1	\bar{f}_1	1	$1/f_1$	0	Δ_1
Q_2	1	n_2	1	f_2	$-1/f_2$	0	Δ_2
\tilde{Q}_2	1	\bar{n}_2	1	\bar{f}_2	$-1/f_2$	0	Δ_2
T_1	1	1	1	1	1	1	$f_1(1 - \Delta_1) - n_1 + 1$
\tilde{T}_1	1	1	1	1	1	-1	$f_1(1 - \Delta_1) - n_1 + 1$
T_2	1	1	1	1	-1	1	$f_2(1 - \Delta_2) - n_2 + 1$
\tilde{T}_2	1	1	1	1	-1	-1	$f_2(1 - \Delta_2) - n_2 + 1$

where $\Sigma_1^{(1)}$ and $\Sigma_1^{(n_1)}$ are the bare monopoles of the $U(n_1)$ theory with flux $(1, 0, \dots, 0)$ and $(0, 0, \dots, -1)$, and $\Sigma_2^{(1)}$ and $\Sigma_2^{(n_2)}$ are the bare monopoles of the $U(n_2)$ theory with flux $(1, 0, \dots, 0)$ and $(0, 0, \dots, -1)$. The charges of the fields are given in Table 2.

- The magnetic theory is a $U(f_1 - n_1) \times U(f_2 - n_2)$ gauge theory with f_1 flavors q_1 and \tilde{q}_1 in the first sector and f_2 flavors q_2 and \tilde{q}_2 in the second sector. There are also the mesons $M_1 = Q_1 \tilde{Q}_1$ and $M_2 = Q_2 \tilde{Q}_2$. The superpotential takes the form

$$W = M_1 q_1 \tilde{q}_1 + M_2 q_2 \tilde{q}_2 + t_1 \tilde{t}_2 + \tilde{t}_1 t_2, \tag{3.2}$$

where t_1 and \tilde{t}_1 are the bare monopoles of the $U(f_1 - n_1)$ theory with flux $(1, 0, \dots, 0)$ and $(0, 0, \dots, -1)$, and t_2 and \tilde{t}_2 are the bare monopoles of the $U(f_2 - n_2)$ theory with flux $(1, 0, \dots, 0)$ and $(0, 0, \dots, -1)$.

In the following we will derive the above duality via the reduction of the four-dimensional duality on the brane system and independently via the reduction of the four-superconformal index (SCI) to the three-dimensional partition function on the squashed three-sphere.

3.1. Derivation from the brane setup

Consider the reduction of four-dimensional Seiberg duality on S^1 in the brane picture [19]. This is done by T-dualizing in the direction x_3 . The theory on the circle has one NS and one NS' brane, with one compact direction. There are N D3s extended along x_6 and F D5s. The D-branes are at the origin of the circle. In order to perform the three-dimensional limit, we select a vacuum in which we move n_1 D3s and f_1 D5s to $x_3 = \pi\alpha' f_2/r$ and n_2 D3s and f_2 D5s to $x_3 = -\pi\alpha' f_1/r$ on the circle. The final configuration corresponds to a product of two gauge factors interacting via the superpotential in Eq. (3.1), as can be understood by visualizing the Euclidean D1 branes discussed above between the two sectors.

The dual phase is obtained via a HW transition. There are $f_1 - n_1$ D3s and f_1 D5s at $x_3 = \pi\alpha' f_2/r$ and $f_2 - n_2$ D3s and f_2 D5s at $x_3 = -\pi\alpha' f_1/r$ on the circle. This theory corresponds to a product of two gauge factors interacting through the superpotential (3.2).

3.2. Derivation from the partition function

Here we show that the duality discussed above can be obtained by implementing the real mass flow on the partition function on S^3_b . The relevant formulas are given in Appendix B. When the integral identity between the four-dimensional Seiberg-dual phases is reduced on the circle, the partition functions are related by the relation

$$Z_{U(N)}(\Lambda; \mu, \nu) = \prod_{a,b=1}^F \Gamma_h(\mu_a + \nu_b) Z_{U(F-N)}(-\Lambda; \omega - \nu, \omega - \mu) \tag{3.3}$$

with the balancing condition

$$\sum_{a=1}^F \mu_a = \sum_{a=1}^F \nu_a = \omega(F - N) \tag{3.4}$$

that signals the presence of a three-dimensional effective duality with an η -superpotential in both phases. In the vacuum chosen above, the real masses split as

$$\mu \rightarrow \begin{cases} m_a^1 + f_2 s \\ m_a^2 - f_1 s \end{cases} \quad \nu \rightarrow \begin{cases} n_a^1 - f_2 s, & a = 1, \dots, f_1 \\ n_a^2 + f_1 s, & a = 1, \dots, f_2. \end{cases} \tag{3.5}$$

The gauge group is broken by the choice of the vacuum

$$\sigma \rightarrow \begin{cases} \sigma_i^1 - f_2 s, & i = 1, \dots, n_1 \\ \sigma_i^2 + f_1 s, & i = 1, \dots, n_2. \end{cases} \tag{3.6}$$

The balancing condition becomes

$$\sum_{a=1}^{f_1} m_a^1 + \sum_{a=1}^{f_2} m_a^2 = \sum_{a=1}^{f_1} n_a^1 + \sum_{a=1}^{f_2} n_a^2 = \omega(F - N). \tag{3.7}$$

The real masses can be written in terms of the global symmetries as

$$m_a^1 = M_a^1 + \frac{m_A}{f_1} + \omega \Delta_1 \quad m_a^2 = M_a^2 + \frac{m_A}{f_1} + \omega \Delta_1 \tag{3.8}$$

$$n_a^1 = N_a^1 - \frac{m_A}{f_2} + \omega \Delta_2 \quad n_a^2 = N_a^2 - \frac{m_A}{f_2} + \omega \Delta_2 \tag{3.9}$$

with the constraint on the non-Abelian gauge symmetries

$$\sum_{a=1}^{f_1} M_a^1 = \sum_{a=1}^{f_1} N_a^1 = \sum_{a=1}^{f_2} M_a^2 = \sum_{a=1}^{f_2} N_a^2 = 0. \tag{3.10}$$

The balancing condition forces the R -charges to be constrained by

$$f_1(1 - \Delta_1) + f_2(1 - \Delta_2) - n_1 - n_2 = 0. \tag{3.11}$$

The dual theory has real masses inherited from the electric theory and the gauge group is broken as

$$\tilde{\sigma} \rightarrow \begin{cases} \tilde{\sigma}_i^1 - f_2 s, & i = 1, \dots, f_1 - n_1 \\ \tilde{\sigma}_i^2 + f_1 s, & i = 1, \dots, f_2 - n_2. \end{cases} \tag{3.12}$$

At large s , there is a divergent phase on both sides of (3.3). This divergent term coincides in the electric and in the magnetic phase. One can ignore it and can compare only the finite terms, arriving to the equality

$$Z_{U(n_1) \times U(n_2)}(\Lambda_1, \Lambda_2; m^1, m^2, n^1, n^2) = \prod_{a,b}^{f_1} \Gamma_h(m_a^1 + n_a^1) \prod_{a,b}^{f_2} \Gamma_h(m_a^2 + n_a^2) \\ Z_{U(f_1-n_1) \times U(f_2-n_2)}(-\Lambda_1, -\Lambda_2; \omega - n^1, \omega - n^2, \omega - m^1, \omega - m^2), \tag{3.13}$$




Fig. 3. The duality for the $U(n_1) \times U(n_2)$ monopole quiver can be understood as a sequence of two Aharony dualities. Starting from the electric configuration (a) we can either Aharony-dualize the second node (b) and then the first one, arriving to the configuration (d), or Aharony-dualize the first node (c) and then the second one, arriving again to the magnetic configuration (d). In the intermediate phases (b) and (c), some monopoles are singlets of the theory.

where the effective Fayet–Iliopoulos (FI) terms are

$$\Lambda_1 = \Lambda + 2m_A - 2\omega(n_2 - f_2(1 - \Delta_2)), \quad \Lambda_2 = \Lambda + 2m_A + 2\omega(n_1 - f_2(1 - \Delta_1)). \tag{3.14}$$

These FI terms are compatible with the presence of the monopole superpotentials in both phases. The relation (3.13) corresponds to the new duality between the monopole quivers discussed on the field theory side.

3.3. Relation to Aharony dualities

We can also consider the initial configuration and perform a series of Aharony dualities, first on the $U(n_1)$ node and then on the $U(n_2)$ node (see Fig. 3). This gives us a consistency check of the duality discussed above, because the final configuration should coincide with the duality derived by a HW transition on the brane setup.

Let us study the first duality. The new quiver consists of a $U(f_1 - n_1) \times U(n_2)$ theory with f_1 flavors q_1 and \tilde{q}_1 in the $U(f_1 - n_1)$ sector, with a meson $M_1 = Q_1 \tilde{Q}_1$ and f_2 flavors Q_2 and \tilde{Q}_2 in the $U(n_2)$ sector. The superpotential of this theory is

$$W = M_1 q_1 \tilde{q}_1 + T_1 \tilde{t}_1 + \tilde{T}_1 t_1 + T_1 \tilde{T}_2 + \tilde{T}_1 T_2. \tag{3.15}$$

In this phase, the partition function is given by

$$\Gamma_h \left(\pm \frac{\Lambda_1}{2} - \frac{1}{2} \sum_{a=1}^{f_1} (m_a + n_a) + \omega(f_1 - n_1 + 1) \right) \prod_{a < b} \Gamma_h(m_a + n_a) \times Z_{U(f_1 - n_1) \times U(n_2)}(-\Lambda_1, \Lambda_2; \omega - n_a, \tilde{m}_a, \omega - m_a, \tilde{n}_a). \tag{3.16}$$

Now we perform the second Aharony duality on the second node. In this case the theory is the one discussed above, with superpotential

$$W = M_1 q_1 \tilde{q}_1 + M_2 q_2 \tilde{q}_2 + T_1 \tilde{t}_1 + \tilde{T}_1 t_1 + T_1 \tilde{T}_2 + \tilde{T}_1 T_2 + T_2 \tilde{t}_2 + \tilde{T}_2 t_2, \tag{3.17}$$

where the monopoles T_i and \tilde{T}_i in (3.17) should be treated as massive singlets and integrated out. This leads to the expected dual superpotential

$$W = M_1 q_1 \tilde{q}_1 + M_2 q_2 \tilde{q}_2 + t_1 \tilde{t}_2 + \tilde{t}_1 t_2. \tag{3.18}$$

We can also reproduce this result on the partition function. In this case we have

$$\Gamma_h \left(\pm \frac{\Lambda_1}{2} - \frac{1}{2} \sum_{a=1}^{f_1} (m_a^1 + n_a^1) + \omega(f_1 - n_1 + 1) \right) \prod_{a < b} \Gamma_h(m_a^1 + n_a^1) \times \Gamma_h \left(\pm \frac{\Lambda_2}{2} - \frac{1}{2} \sum_{a=1}^{f_2} (m_a^2 + n_a^2) + \omega(f_2 - n_2 + 1) \right) \prod_{a < b} \Gamma_h(m_a^1 + n_a^1) \times Z_{U(f_1 - n_1) \times U(f_2 - n_2)}(-\Lambda_1, -\Lambda_2; \omega - n_a^1, \omega - n_a^2, \omega - m_a^1, \omega - m_a^2). \tag{3.19}$$

The monopoles can be integrated out by imposing the balancing condition and using the relation $\Gamma_h(\omega \pm x) = 1$. In this way formula (3.13) is recovered.

Observe that the duality for the monopole quiver discussed here reduces to the ordinary Aharony duality for $f_2 = 1$ and $n_2 = 0$. In this case, the dual theory corresponds to a $U(f_1 - n_1) \times U(1)$ quiver, where the supersymmetric quantum electrodynamics (SQED) sector has one flavor and it is mirror-dual to the XYZ model. The fields Y and Z , correspond to the monopoles of the $U(1)$ sector, t_2 and \tilde{t}_2 while the field X is identified with the singlet $q_2 \tilde{q}_2$. The dual superpotential becomes

$$W = M_1 q_1 \tilde{q}_1 + M_2 X + XYZ + t_1 Y + \tilde{t}_1 Z. \tag{3.20}$$

By integrating out the massive singlets M_2 and X , we end up in the expected Aharony-dual theory. The monopoles Y and Z act as singlets in the dual theory and they have the same quantum numbers of the monopoles T_1 and \tilde{T}_1 .

The reduction of the monopole quiver duality to Aharony duality for $f_2 = 1$ and $n_2 = 0$ can be obtained also from the partition function. In this case one has to use the identity [23]

$$\int d\sigma e^{-i\pi\Lambda\sigma} \Gamma_h(\sigma + \omega - \tilde{m}) \Gamma_h(-\sigma + \omega - \tilde{n}) = \Gamma_h(2\omega - \tilde{m} - \tilde{n}) \Gamma_h \left(\frac{\pm\Lambda + \tilde{m} + \tilde{n}}{2} \right). \tag{3.21}$$

The first term on the RHS of (3.21) corresponds to the field X and it cancels the contribution of the dual meson M_1 in the partition function, as can be seen by using the identity $\Gamma_h(2\omega - x) \Gamma_h(\omega) = 1$. The second term in (3.21) corresponds to the contribution of the monopoles Y and Z , corresponding to the singlets T_1 and \tilde{T}_1 in the Aharony duality. This can be shown with the help of the balancing condition, which corresponds to substituting the relation

$$\tilde{m} + \tilde{n} = 2\omega(f_1 - n_1 + 1) - \sum_{a=1}^{f_1} (m_a^1 + n_a^1) \tag{3.22}$$

Table 3
Brane setup realizing the four-dimensional IP duality.

	0	1	2	3	4	5	6	7	8	9
D4	×	×	×	×			×			
D6	×	×	×	×				×	×	×
NS	×	×	×	×	×	×				
NS'	×	×	×	×					×	×
O4 ⁺	×	×	×	×			×			

into (3.21). This gives the monopole contribution in the Aharony-dual phase as expected.

We conclude this section by commenting on the general situation. One can indeed construct a monopole quiver with a generic amount of K $U(n_i)$ gauge factors, by separating the N D3 branes on the compact x_3 direction into K stacks. Stability requires that f_i D5 branes are attached to each sector, such that $f_i > n_i$. The sectors interact through D1 branes connecting the n_i -th and n_{i+1} -th sector, realizing a generic monopole quiver. The dual theory is obtained via a HW transition and it generalizes the construction discussed in this section. One has a set of $U(f_i - n_i)$ SQCD sectors, with dual fundamentals interacting with mesons. Again these sectors interact with each other through AHW terms, represented by D1 branes in this picture. In this general case, one can check the matching between the partition functions by engineering the flow on $Z_{S_b^3}$. We show this derivation in Appendix A.

4. SQCD dualities and orientifolds

In this section we show how to derive SQCD dualities from four-dimensional theories with real gauge groups via the brane picture, which in these cases also includes orientifold planes. First we consider the two new dualities that have been derived in [7]. In order to reproduce them via a brane set-up, we will use the non-trivial vacuum structure discussed above, this time starting from the four-dimensional brane representation of the Intriligator-Pouliot [24] (IP) duality. We show that both three-dimensional dualities can be obtained in the brane picture by shifting some D-branes on the T-dual circle while sending its radius to infinity. We conclude the section showing that also the flow leading to the conventional Aharony duality can be engineered from the very same brane picture.

Four dimensions Before starting the three-dimensional analysis, we review the basic aspects of the four-dimensional IP duality that will be necessary in the following.

The electric theory has an $Sp(2N)$ gauge group with $2F$ fundamentals Q and vanishing superpotential. The dual theory has $Sp(2(F - N - 2))$ gauge group with $2F$ fundamentals q and $F(2F - 1)$ mesonic operators $M = QQ$, interacting through the superpotential $W = Mqq$.

In a type IIA brane setup, there are $2N$ D4 branes stretched between one NS and one NS' brane. An O4⁺ plane on the stack of D4 branes realizes the projection of $SU(2N)$ to $Sp(2N)$. Flavor is introduced by the addition of $2F$ D6 branes. The brane set-up is summarized in Table 3. The dual theory is obtained via a HW transition. In this case, one has $2(F - N - 2)$ D4 branes stretched between the NS and the NS' brane.²

² The factor of 2 is necessary to preserve the linking number in presence of the O4⁺ plane.

Table 4
Brane setup realizing the reduction of IP duality to three dimensions.

	0	1	2	3	4	5	6	7	8	9
D3	×	×	×				×			
D5	×	×	×					×	×	×
NS	×	×	×	×	×	×				
NS'	×	×	×	×					×	×
O3 ⁺	×	×	×				×			

Reduction The IP duality can be reduced on a circle, leading to a duality between an $Sp(2N)$ gauge theory with $2F$ fundamentals and an $Sp(2(F - N - 2))$ gauge theory with $2F$ fundamentals interacting with $F(2F - 1)$ mesons. The presence of the circle prevents the generation of the axial symmetry by inducing an η superpotential in both phases. This is the usual circle reduction of four-dimensional dualities to three dimensions discussed in [8].

One can flow to three dimensions with a real mass and/or a Higgs flow in a conventional way, obtaining the Aharony duality with symplectic gauge groups, as discussed in [12]. By performing a different real mass and Higgs flow in the two dual phases, we reproduce the new dualities of [7]. We refer to these dualities as BBP_I and BBP_{II} duality.

4.1. BBP_I

Field theory The BBP_I duality is obtained by splitting the $2F$ real masses μ_a into two sets, m_a and \tilde{m}_a , each with F elements, and performing the shifts

$$m_a \rightarrow m_a + s, \quad n_a \rightarrow n_a - s. \tag{4.1}$$

A Higgs flow $\sigma_i \rightarrow \sigma_i + s$ is taken for both real scalars in the $Sp(2N)$ and in the dual $Sp(2(F - N - 2))$ gauge groups. At large s , the massive fields can be integrated out, leading to a three-dimensional duality between two unitary gauge theories.

- The electric theory is a three-dimensional $U(N)$ $\mathcal{N} = 2$ theory with F fundamental and antifundamental flavors, Q and \tilde{Q} . There is a superpotential interaction

$$W = T + \tilde{T}, \tag{4.2}$$

where T and \tilde{T} are the $(1, 0, \dots, 0)$ and the $(0, \dots, 0, -1)$ monopoles, respectively.

- The dual theory is a three-dimensional $\mathcal{N} = 2$ $U(F - N - 2)$ theory with F fundamental and antifundamental flavors, q and \tilde{q} , and F^2 singlets M . There is a superpotential interaction

$$W = Mq\tilde{q} + t + \tilde{t}, \tag{4.3}$$

where t and \tilde{t} are the $(1, 0, \dots, 0)$ and the $(0, \dots, 0, -1)$ monopoles, respectively.

Brane picture The duality just reviewed can be obtained by brane engineering the IP duality reduced on the circle. By compactifying x_3 and performing a T-duality, we arrive at a system consisting of an NS brane, an NS' brane, $2N$ D3 branes, $2F$ D5s and two $O3^+$ planes. The branes are extended as shown in Table 4.

The $2N$ D3s, the $2F$ D5s and one $O3^+$ are initially placed at the origin of x_3 . The second $O3^+$ is at the point $x_3^0 = \pi\alpha'/R_3$. Now we shift the branes along x^3 : the configuration of interest has $(F - 2)$ D5s and N D3 branes moved to $x_3 = \pi\alpha'/(2R_3)$, and symmetrically $(F - 2)$ D5s

Table 5

Brane content at four points in the direction x_3 for the different phases in the BBP_I duality. In all phases, there is an $O3^+$ in $x_3 = 0$ and x_3^o .

	$x_3 = 0$	$x_3 = \frac{\alpha\pi}{2R_3}$	$x_3 = x_3^o$	$x_3 = -\frac{\alpha\pi}{2R_3}$
IP on S^1	$2N$ D3 $2F$ D5	–	–	–
electric	2 D5	N D3 $(F - 2)$ D5	2 D5	N D3 $(F - 2)$ D5
after HW	2 D5	$(F - N - 2)$ D3 $(F - 2)$ D5	2 D5	$(F - N - 2)$ D3 $(F - 2)$ D5
magnetic	–	$(F - N - 2)$ D3 $(F - 2)$ D5	–	$(F - N - 2)$ D3 $(F - 2)$ D5

and N D3 branes moved to $x_3 = 3\pi\alpha'/(2R_3) = -\pi\alpha'/(2R_3)$. Two D5s are left at $x_3 = 0$ and the last two are at $x_3^o = \pi\alpha'/R_3$.

After a HW transition we obtain $F - 2$ D5s and the $F - N - 2$ D3 branes at $x_3 = \pi\alpha'/(2R_3)$, $F - 2$ D5 and the $(F - N - 2)$ D3 branes at $x_3 = -\pi\alpha'/(2R_3)$. There are again two D5s at $x_3 = 0$ and two are at $x_3 = \pi\alpha'/R_3$.

In both phases, we can finally shift the D5s so that the resulting configuration has F D5s and the $(F - N - 2)$ D3 branes at $x_3 = \pi\alpha'/(2R_3)$ and the same amounts at $x_3 = -\pi\alpha'/(2R_3)$. A crucial aspect of this construction is that the real mass flow associated to this last shift of the D5 branes only involves flavor degrees of freedom (DOF) so that the two theories, originally obtained from a HW transition, remain dual at the end of the flow. The monopole superpotential can be inferred from the D1 branes in the usual manner as explained in Sec. 2. It is $W = T + \tilde{T}$ on the electric side and $W = t + \tilde{t}$ on the magnetic side. The brane content of the different phases is given in Table 5 and Fig. 4.

In the $R_3 \rightarrow 0$ limit, this reproduces the BBP_I duality.

4.2. BBP_{II}

Field theory The second duality discussed in [7] was obtained by considering the BBP_I model with $(F + 1)$ flavors, shifting the $(F + 1)$ -th flavor by

$$m_{F+1} \rightarrow m_{F+1} + s, \quad n_{F+1} \rightarrow n_{F+1} - s, \tag{4.4}$$

and taking the large- s limit.

- The electric theory is a three-dimensional $\mathcal{N} = 2$ $U(N)$ theory with F fundamental and antifundamental flavors, Q and \tilde{Q} . There is a superpotential interaction,

$$W = T, \tag{4.5}$$

where T is the $(1, 0, \dots, 0)$ monopole.

- The dual theory is a three-dimensional $\mathcal{N} = 2$ $U(F - N - 1)$ theory with F fundamental and antifundamental flavors, q and \tilde{q} , F^2 singlets M , and a singlet S . There is a superpotential interaction

$$W = Mq\tilde{q} + t + S\tilde{t}, \tag{4.6}$$

where t and \tilde{t} are the $(1, 0, \dots, 0)$ and the $(0, \dots, 0, -1)$ monopoles.




Fig. 4. Brane content at four points in the direction x_3 for the different phases in the BBP_{II} duality. The configuration obtained from dimensional reduction is in (a), then the D3s and some D5s are moved to realize the electric phase (b); a HW transition is performed (c) and finally all the D5s are moved to the same point to realize the magnetic phase (d). The D5 branes are projected onto the line.

Brane picture Here we show that the BBP_{II} duality can be obtained from the brane setup as well. Start with the reduction of the IP duality on the circle as done above. In this case, we have $2(F + 1)$ D6 branes and, on the circle, $2N$ D3s, $2(F + 1)$ D5s plus one $O3^+$ at the origin $x_3 = 0$ with its dual $O3^+$ at $x_3^\circ = \pi\alpha'/R_3$.

The three-dimensional limit is obtained by shifting $(F - 1)$ D5s and N D3 branes to $x_3 = \pi\alpha'/(2R_3)$ and, symmetrically, $(F - 1)$ D5s and N D3 branes to $x_3 = 3\pi\alpha'/(2R_3) = -\pi\alpha'/(2R_3)$. Two D5s are left at $x_3 = 0$ and the last two are at $x_3^\circ = \pi\alpha'/R_3$. We can make a HW transition on this configuration, obtaining $(F - 1)$ D5s and $(F - N - 1)$ D3 branes at $x_3 = \pi\alpha'/(2R_3)$ and similarly at $x_3 = -\pi\alpha'/(2R_3)$. There are again two D5s at $x_3 = 0$ and two at $x_3^\circ = \pi\alpha'/R_3$.

In both phases we can now shift the D5s to $x_3 = 0$ (or equivalently the D5s at x_3°) so that the final configuration has F D5s and all the $(F - N - 1)$ D3 branes at $x_3 = \pi\alpha'/(2R_3)$ (and at $x_3 = -\pi\alpha'/(2R_3)$). There are also two D5 branes at $x_3 = \pi\alpha'/R_3$ (respectively two D5s at $x_3 = 0$) in both phases.

- In the electric phase, the three-dimensional limit gives rise directly to the superpotential (4.5), in terms of the D1 branes connecting the two stacks of D3 branes through the orientifold at $x_3 = 0$. Furthermore, the absence of massless singlets from the D5 branes placed at $x_3 = \pi\alpha'/R_3$ allows us to ignore this sector.

Table 6

Brane content at four points in the direction x_3 for the different phases in the BBP_{II} duality. In all phases, there is an O3⁺ in $x_3 = 0$ and x_3^o .

	$x_3 = 0$	$x_3 = \frac{\alpha\pi}{2R_3}$	$x_3 = x_3^o$	$x_3 = -\frac{\alpha\pi}{2R_3}$
IP on S^1	2N D3 2(F + 1) D5	–	–	–
electric	2 D5	N D3 (F – 1) D5	2 D5	N D3 (F – 1) D5
after HW	2 D5	(F – N – 1) D3 (F – 1) D5	2 D5	(F – N – 1) D3 (F – 1) D5
magnetic	–	(F – N – 1) D3 F D5	2 D5	(F – N – 1) D3 F D5

- On the magnetic side, the situation is different. We still have the D1 branes connecting the two stacks of D3 branes through the orientifold at $x_3 = 0$ which, like in the electric phase, give a contribution $W = T$ to the superpotential. This time, on the other hand, there is also a massless singlet arising from the D5 branes placed at $x_3 = \pi\alpha'/R_3$, since the D5s are parallel to the NS' brane along the directions (8, 9). This is the same as the situation discussed in [7] for the reduction of the IP duality to the Aharony duality for symplectic gauge group. In the UV, i.e. for small T-dual radius, the contribution of the D1 branes from this sector is $W \simeq t_{high}$. Flowing to the IR, a scale-matching relation identifies the operator \tilde{t}_{high} with the combination $\tilde{t}_{low}S$, where S is a singlet arising from the D5 branes which, on the field theory side, is identified with the massless component of the broken meson after the real mass flow. All in all, the singlet S interacts with the monopole of the magnetic gauge theory and it has the same quantum number as the electric monopole. The final result coincides with the superpotential (4.6) of the magnetic phase of the BBP_{II} duality.

The brane content of the different phases is given in Table 6.

4.3. Aharony duality

We conclude this section by showing that it is also possible to recover the duality of Aharony by reducing the IP duality to three dimensions. The picture is similar to the one discussed above, so we will be brief in many aspects of the derivation, referring the reader to the details discussed above when necessary.

Field theory As already observed in [7], Aharony duality can be obtained from BBP_I by a real mass deformation. One starts from $(F + 2)$ flavors and shifts the masses as

$$\begin{aligned}
 m_{F+1} &\rightarrow m_{F+1} + s, n_{F+1} \rightarrow n_{F+1} - s, \\
 m_{F+2} &\rightarrow m_{F+1} - s, n_{F+2} \rightarrow n_{F+2} + s.
 \end{aligned}
 \tag{4.7}$$

In the large- s limit, the usual Aharony duality is recovered. The field content and interactions of Aharony duality can be summarized as follows:

- The electric theory is a three-dimensional $\mathcal{N} = 2$ $U(N)$ theory with F fundamental and antifundamental flavors, Q and \tilde{Q} with vanishing superpotential.

Table 7

Brane content at four points in the direction x_3 for the different phases in the Aharony duality. In all phases, there is an $O3^+$ in $x_3 = 0$ and x_3° .

	$x_3 = 0$	$x_3 = \frac{\alpha\pi}{2R_3}$	$x_3 = x_3^\circ$	$x_3 = -\frac{\alpha\pi}{2R_3}$
IP on S^1	2N D3 2(F + 2) D5	–	–	–
electric	2 D5	N D3 F D5	2 D5	N D3 F D5
after HW	2 D5	(F – N) D3 F D5	2 D5	(F – N) D3 F D5

- The dual theory is a three-dimensional $\mathcal{N} = 2$ $U(F - N)$ theory with F fundamental and anti-fundamental flavors, q and \tilde{q} , F^2 singlets M , and singlets S and \tilde{S} . There is a superpotential interaction

$$W = Mq\tilde{q} + S\tilde{t} + \tilde{S}t, \tag{4.8}$$

where t and \tilde{t} are the $(1, 0, \dots, 0)$ and the $(0, \dots, 0, -1)$ monopoles. The singlets S and \tilde{S} are identified respectively with the $(0, \dots, 0, -1)$ and the $(1, 0, \dots, 0,)$ monopoles, \tilde{T} and T , of the electric theory.

Brane picture Aharony duality can be reproduced at the brane level as above. In this case we can consider the brane realization of the four-dimensional IP duality with $2(F + 2)$ D6 branes plus $2N$ D3s, $2(F + 2)$ D5s, one $O3^+$ at the origin $x_3 = 0$ and a second $O3^+$ at $x_3^\circ = \pi\alpha'/R_3$.

The three-dimensional limit is obtained by shifting F D5s and N D3 branes to $x_3 = \pi\alpha'/(2R_3)$, and symmetrically F D5s and N D3 branes to $x_3 = 3\pi\alpha'/(2R_3) = -\pi\alpha'/(2R_3)$, and we have again two D5s at $x_3 = 0$ and two at $x_3^\circ = \pi\alpha'/R_3$.

We can make a HW transition on this configuration obtaining F D5s and the $F - N$ D3 branes at $x_3 = \pi\alpha'/(2R_3)$, and similarly at $x_3 = -\pi\alpha'/(2R_3)$.

The final step consists in implementing the large- s limit, considering the massless modes in the spectrum that survive because of the mesons, as discussed in the derivation of the BBP_{II} duality. The brane content of the different phases is given in Table 7.

In this case there are two mesonic sectors in which extra massless matter emerges, leading to the monopole superpotential in the dual phase. This can be visualized as follows. At large s , the electric theory is $U(N)$ SQCD with F flavors.

On the magnetic side, the situation is analogous to the one described above. We have a $U(F - N)$ SQCD with F dual flavors interacting with a meson M . There are also extra singlets, arising from the original mesonic operators, because the D5s at $x_3 = 0$ and x_3° are parallel to the NS' branes along the directions $(8, 9)$. The monopole superpotential can be reconstructed as follows. One starts by placing the D1 branes in the UV description, *i.e.* when the T-dual radius is small. They give rise to a UV monopole superpotential on both the electric and the magnetic side, $W_{UV}^{(ele)} = t_{high} + \tilde{t}_{high}$ and $W_{UV}^{(mag)} = t_{high} + \tilde{t}_{high}$. The flow to the IR is done by sending the T-dual radius to be large. In the electric theory, in absence of massless singlets at $x_3 = 0$ and x_3° , we can safely remove the monopole superpotential. In the dual phase, one has to consider a scale-matching relation for the two monopoles t and \tilde{t} , in terms the massless singlets arising from the D5 branes. They correspond, on the field theory side, to some massless components of the original meson M . By looking at the charge structure, the scaling can be formulated as

$$t_{high} = t_{low}\tilde{S}, \quad \tilde{t}_{high} = \tilde{t}_{low}S. \tag{4.9}$$

The singlets appearing in these relations have the same charges as the electric monopoles T and \tilde{T} . By substituting the rescaled monopole into the superpotential $W_{UV}^{(mag)}$, one recovers the expected results for Aharony duality.

5. Monopole quiver dualities with orientifolds

The monopole quivers considered in Section 3 can be also be constructed in the presence of orientifolds. By considering the reduction of four-dimensional theories on S^1 one ends up with pairs of $O3$ planes at $x_3 = 0$ and $x_3 = x_3^o$, as discussed above. In this case, one can consider non-trivial vacuum structures with stacks of D3- and D5-branes along the compact directions (consistently with the identifications imposed by the orientifolds). When flowing to the three-dimensional limit, a monopole quiver with both real and unitary gauge groups is generated. The real groups arise if some stacks of D3 branes are placed at $x_3 = 0$ and/or $x_3 = x_3^o$, while the D3 branes at different positions in x_3 give rise to unitary gauge groups.

In this section, we discuss the simplest realization of such a configuration, a duality involving a monopole quiver with $Sp(2n_1) \times U(n_2)$ gauge group, obtained by reducing the four-dimensional IP duality on S^1 and performing a large mass flow.

Field theory Let us consider the flow on the gauge theory side. In the electric theory we assign the masses to the $2F$ fundamentals as

$$\mu \rightarrow \begin{cases} \mu_a & \text{if } a = 1, \dots, 2f_1 \\ m_a + s & \text{if } a = 1, \dots, f_2 \\ n_a - s & \text{if } a = 1, \dots, f_2. \end{cases} \tag{5.1}$$

We also Higgs the gauge group as

$$\sigma \rightarrow \begin{cases} \sigma_i & \text{if } i = 1, \dots, n_1 \\ \tilde{\sigma}_i & \text{if } i = 1, \dots, n_2. \end{cases} \tag{5.2}$$

This breaks the gauge theory to an $Sp(2n_1)$ theory with $2f_1$ fundamentals and a $U(n_2)$ theory with f_2 fundamental flavors. The global symmetry is broken to a non-Abelian part $SU(2f_1) \times SU(f_2)^2$ and an Abelian $U(1)_A \times U(1)_R$. The axial $U(1)_A$ symmetry is a combination of the two axial symmetries of the unitary and of the symplectic sector and of the topological symmetry arising from shifting the dual photon of $U(1) \subset U(n_2)$. These symmetries are broken to a single $U(1)_A$ by the AHW interaction between the monopoles Y of the $Sp(2n_1)$ sector and the T of the $U(n_2)$ sector. This interaction is

$$W = Y\tilde{T} + T. \tag{5.3}$$

The dual theory is obtained by an opportune real mass flow on the $2f_1$ dual fundamentals and by the Higgsing

$$\sigma \rightarrow \begin{cases} \sigma_i & i = 1, \dots, \tilde{n}_1 \\ \tilde{\sigma}_i & i = 1, \dots, \tilde{n}_2, \end{cases} \tag{5.4}$$

where $\tilde{n}_1 = f_1 - n_1 - 1$ and $\tilde{n}_2 = f_2 - n_2 - 1$. The dual theory becomes an $Sp(2\tilde{n}_1)$ theory with $2f_1$ fundamentals and a $U(\tilde{n}_2)$ theory with f_2 fundamental flavors. There are also $f_1(2f_1 - 1)$ mesons M_1 in the symplectic sector and f_2^2 mesons M_2 in the unitary sector. The superpotential of this dual theory is

Table 8

Brane content at four points in the direction x_3 for the different phases in the duality for the $Sp(n_1) \times U(n_2)$ quiver. In all phases, there is an $O3^+$ at $x_3 = 0$ and at x_3° .

	$x_3 = 0$	$x_3 = \frac{\alpha\pi}{2R_3}$	$x_3 = x_3^\circ$	$x_3 = -\frac{\alpha\pi}{2R_3}$
IP on S^1	2N D3 2F D5	–	–	–
electric	$2n_1$ D3 $2f_1$ D5	n_2 D3 $(f_2 - 1)$ D5	2 D5	n_2 D3 $(f_2 - 1)$ D5
after HW	$2(f_1 - n_1 - 1)$ D3 $2f_1$ D5	$(f_2 - n_2 - 1)$ D3 $(f_2 - 1)$ D5	2 D5	$(f_2 - n_2 - 1)$ D3 $(f_2 - 1)$ D5
magnetic	$2(f_1 - n_1 - 1)$ D3 $2f_1$ D5	$(f_2 - n_2 - 1)$ D3 f_2 D5	–	$(f_2 - n_2 - 1)$ D3 f_2 D5

$$W = M_1 q_1 q_1 + M_2 q_2 \tilde{q}_2 + y \tilde{t} + t, \tag{5.5}$$

where the monopoles y of the $Sp(2n_1)$ sector and t, \tilde{t} of the $U(n_2)$ sector interact through the AHW superpotential leaving the axial $U(1)_A$ symmetry as in the electric theory.

Brane picture This duality can be guessed from the brane picture by considering the reduction of the $Sp(2N)$ theory with $2F$ fundamentals on S^1 . After the duality, one can shift n_2 D3s and $(f_2 - 1)$ D5 branes to $x_3 = \pi\alpha'/(2R_3)$, and symmetrically to $x_3 = -\pi\alpha'/(2R_3)$ plus two D5 branes at $x_3 = x_3^\circ = \pi\alpha'/r$, thus leaving $2N - 2n_2 = 2n_1$ D3 branes and $2F - 2f_2 = 2f_1$ D5 branes at $x_3 = 0$, on the $O3^+$ plane. After a HW transition we have $(f_2 - n_2 - 1)$ D3 and $(f_2 - 1)$ D5 branes at $x_3 = \pm\pi\alpha'/(2R_3)$, $2(f_1 - n_1 - 1)$ D3 and $2f_1$ D5 at $x_3 = 0$ and two D5 branes at x_3° . Finally, we are free to move one D5 placed at x_3° to $x_3 = \pi\alpha'/(2R_3)$ and the other one to $x_3 = -\pi\alpha'/(2R_3)$, to recover the magnetic phase. This motion preserves the duality, being just a real mass flow on the flavor side. By placing the D1 branes between the stacks of D3 branes we can also read the monopole superpotential in both phases.

The various configurations are collected in Table 8.

Partition function We can corroborate the duality by performing the real mass flow on the partition function. In this case we use trick of [7], based on the symmetry of the integrals. The large- s behavior cancels in the two theories, because the s -dependent phase is

$$-4s \left(\omega \left(2f_2 n_1 + n_2^2 + n_2 \right) - n_1 \sum_{a=1}^{f_2} (m_a + n_a) \right). \tag{5.6}$$

After eliminating this phase we obtain the equality

$$\begin{aligned} & \int \prod_{i=1}^{n_1} d\sigma_i \prod_{i=1}^{n_2} d\tilde{\sigma}_i e^{\lambda e \tilde{\sigma}_i} \frac{\prod_{i=1}^{n_1} \prod_{a=1}^{2f_1} \Gamma_h(\pm\sigma_i + \mu_a) \prod_{i=1}^{n_2} \prod_{a=1}^{f_2} \Gamma_h(\tilde{\sigma}_i + m_a) \Gamma_h(-\tilde{\sigma}_i + n_a)}{\prod_{i < j}^{n_1} \Gamma_h(\pm\sigma_i \pm \sigma_j) \prod_{i=1}^{n_1} \Gamma_h(\pm 2\sigma_i) \prod_{i < j}^{n_2} \Gamma_h(\pm(\tilde{\sigma}_i - \tilde{\sigma}_j))} \\ &= e^{\phi_m} \prod_{a < b}^{2f_1} \Gamma_h(\mu_a + \mu_b) \prod_{a, b=1}^{f_2} \Gamma_h(m_a + n_b) \int \prod_{i=1}^{\tilde{n}_1} d\sigma_i \prod_{i=1}^{\tilde{n}_2} d\tilde{\sigma}_i e^{\lambda_m \tilde{\sigma}_i} \\ & \frac{\prod_{i=1}^{\tilde{n}_1} \prod_{a=1}^{2f_1} \Gamma_h(\pm\sigma_i + \omega - \mu_a) \prod_{i=1}^{\tilde{n}_2} \prod_{a=1}^{f_2} \Gamma_h(\tilde{\sigma}_i + \omega - n_a) \Gamma_h(-\tilde{\sigma}_i + \omega - m_a)}{\prod_{i < j}^{\tilde{n}_1} \Gamma_h(\pm\sigma_i \pm \sigma_j) \prod_{i=1}^{\tilde{n}_1} \Gamma_h(\pm 2\sigma_i) \prod_{i < j}^{\tilde{n}_2} \Gamma_h(\pm(\tilde{\sigma}_i - \tilde{\sigma}_j))} \end{aligned} \tag{5.7}$$

with the balancing condition

$$\sum_{a=1}^{2f_1} \mu_a + \sum_{a=1}^{f_2} (m_a + n_a) = 2\omega(F - N - 1). \tag{5.8}$$

The phase ϕ_m and the FI terms are given by

$$\begin{aligned} \phi_m &= \sum_{a=1}^{f_2} (m_a - n_a) \left(\sum_{a=1}^{f_2} (m_a + n_a) - 2\omega(f_2 - 3n_2 - 1) \right) - (2n_2 + 1) \sum_{a=1}^{f_2} (m_a^2 - n_a^2), \\ \lambda_e &= 2 \sum_{a=1}^{f_2} (m_a + n_a) - 4\omega(f_2 - n_2 - 1), \\ \lambda_m &= 4\omega(f_2 - n_2) - 6 \sum_{a=1}^{f_2} (m_a + n_a). \end{aligned}$$

6. Conclusions and further developments

In this paper we have discussed large classes of three-dimensional dualities for $\mathcal{N} = 2$ quivers, defined as sets of decoupled SQCD sectors interacting through AHW interactions. These dualities are obtained by reducing four-dimensional Seiberg-dual theories on a circle and triggering real mass and Higgs flows.

We have derived the dualities from the brane-engineering of the gauge theories and corroborated our results via the matching of the three-dimensional partition functions, computed from the circle reduction of the four-dimensional superconformal index. We have also shown that in presence of orientifolds, our brane construction allows to recover the dualities recently discovered in [7] from the reduction of the Intriligator–Pouliot [24] duality.

The IP duality is realized at the brane level by adding an $O4^+$ plane to the usual setup of SQCD. After T-duality it turns into a pair of $(O3^+, O3^+)$ planes on S^1 . When considering pairs of $O3$ planes on S^1 , there are in general six possibilities. Three of them correspond to the compactification of four-dimensional theories with $SP(2N)$, $SO(2N + 1)$ or $SO(2N)$ gauge groups, while the others correspond to twisted compactifications with an outer automorphism. It would be interesting to study the reduction of these theories and the flow in the non-trivial vacua of the type discussed here. However, an immediate problem arises when studying the reduction of the index to the partition function for four-dimensional dualities with orthogonal gauge groups. This procedure is well-defined on the field theory side and on the brane side but the reduction of the index on the circle produces a divergent partition function [25]. We expect that a double-scaling limit would have to be performed in order to recover the three-dimensional limit discussed here. In the orthogonal case it would be also interesting to study the reduction of theories with different global properties [26,25] and to study implications for unitary theories obtained after the three-dimensional real mass flow.

One can also study the reduction of four-dimensional theories with tensor matter. These have a known D-brane realization and the identities between the superconformal index of the dual phases have been listed in [27,28]. It should be possible, for example, to study the reduction of the duality of [29] for $Sp(2N)$ gauge theories with tensor matter and trigger a flow to the duality of [30] for three-dimensional $U(N)$ SQCD with adjoint matter. Another important aspect that

we did not study here regards the $U(N)$ dualities with higher powers in the monopole superpotential, discussed in [7]. In the brane picture this should be related to multiple stacks of D1 branes extended along the D3- and the NS-branes along the directions x_3 and x_6 . It would be interesting to check this guess and study the HW transition in presence of such an effect. It would be also interesting to test the dualities proposed here by matching other partition functions obtained through localization on different compact manifolds, as for example the three-dimensional superconformal index [31] and the topologically twisted index of [32].

In general, we have proposed new dualities without focusing on the possible presence of accidental symmetries in the IR [33–36]. They can arise from the presence of gauge singlets with scaling dimension Δ below the unitarity bound, $\Delta = \frac{1}{2}$. This possibility can be checked by maximizing the free energy obtained from the S^3 partition function in terms of the R -charges. A complete understanding would require finding a UV completion of our models in the spirit of [7], which deserves further investigation but is beyond the scope of our present analysis.

We wish to conclude our discussion with a comment on the real mass and Higgs flows from the brane construction. As already observed in [22], the displacement of the branes to generic points on the circle requires switching on a Wilson line in the gauge theory. This can be understood in the brane picture in terms of repulsive interactions between D3 branes. One realization of this phenomenon is obtained if we place our brane construction into a curved background that preserves the right symmetries and provides the (unique) Wilson line parameter for the quiver on the circle. This role is played by the fluxtrap background introduced and studied in [37,38].

Acknowledgements

A.A. is grateful to the Galileo Galilei Institute for Theoretical Physics (GGI) for hospitality and INFN for partial support during the completion of this work, within the program “New Developments in AdS₃/CFT₂ Holography”.

The work of A.A. and S.R. is supported by the Swiss National Science Foundation (SNF) under grant number PP00P2_157571/1.

Appendix A. The general duality for $U(n)$ product groups

In this appendix we study the flow leading to the generic monopole quiver duality for K $U(n_I)$ SQCD sectors each with f_I pairs of flavors Q_I and \tilde{Q}_I and a monopole superpotential coupling the gauge sectors of the form

$$W = T_1 \tilde{T}_2 + T_2 \tilde{T}_3 + \cdots + T_K \tilde{T}_1. \quad (\text{A.1})$$

The dual theory has K $U(\tilde{n}_I = f_I - n_I)$ SQCD sectors each with f_I pairs of flavors q_I and \tilde{q}_I , f_I^2 singlets M_I for each SQCD sector, and a monopole superpotential coupling the gauge sectors. In this case, the dual superpotential is

$$W = \sum_{I=1}^K M_I q_I \tilde{q}_I + t_1 \tilde{t}_2 + t_2 \tilde{t}_3 + \cdots + t_K \tilde{t}_1. \quad (\text{A.2})$$

This duality can be obtained from the reduction of four-dimensional Seiberg duality for $U(N)$ SQCD with F fundamentals on the circle. After the reduction, one has to perform a real mass flow on the F masses μ_a and ν_a and choosing a non-trivial vacuum on the scalars σ_i in the vector multiplet, Higgsing the gauge theory and reconstructing the quiver. The AHW interactions

associated to this Higgsing reconstruct the monopole superpotential. The dual theory is recovered by performing the opportune real mass flow and Higgsing in the dual phase.

In the following we perform the real mass flow and the Higgsing on the equality relating the partition functions of the dual phases, obtained from the reduction of the equality between the SCI of the four-dimensional Seiberg duality. In this way we automatically obtain the integral identity matching the three-sphere partition functions of the proposed duality.

In the electric theory the real masses μ_a and ν_a can be shifted in K different sectors

$$\mu_a \rightarrow m_a^I + s_I, \quad \nu_a \rightarrow n_a^I - s_I, \quad a = 1, \dots, f_I, \tag{A.3}$$

where each s_I is a divergent real contribution. The scalar σ_i can also be shifted in K sectors as

$$\sigma_i \rightarrow \sigma_i^I - s_I, \quad i = 1, \dots, n_I. \tag{A.4}$$

The ranks f_I and n_I and the infinite shifts s_I are constrained by the relations

$$F = \sum_{I=1}^K f_I, \quad N = \sum_{I=1}^K n_I, \quad \sum_{I=1}^K f_I s_I = 0. \tag{A.5}$$

The balancing conditions on the real masses μ_a and ν_a becomes

$$\sum_{I=1}^K \sum_{a=1}^{f_I} m_a^I = \sum_{I=1}^K \sum_{a=1}^{n_I} n_a^I = \omega(F - N). \tag{A.6}$$

In the dual side the real masses are shifted accordingly. The dual Higgsing is

$$\tilde{\sigma}_i \rightarrow \tilde{\sigma}_i^I - s_I \quad i = 1, \dots, \tilde{n}_I. \tag{A.7}$$

Imposing the real mass and Higgs flows discussed above, the electric and the magnetic partition function can be computed at large s_I by integrating out the divergent contributions. Formally, we are left with a relation between the electric and the magnetic partition functions of the form

$$\prod_{I=1}^K \lim_{s_I \rightarrow \pm\infty} e^{\Phi_e} Z_{U(n_I)} = \prod_{I=1}^K \prod_{a,b=1}^{f_I} \Gamma_h(m_a^I + n_b^I) \lim_{s_I \rightarrow \pm\infty} e^{\Phi_m} Z_{U(\tilde{n}_I)}. \tag{A.8}$$

The electric phase picks up contributions from integrating out the charged matter and the vector multiplet. The magnetic phase has contributions from the charged matter, the meson and the vector multiplet. Summarizing, we have

$$\Phi_e = \Phi_Q + \Phi_V \quad \Phi_m = \Phi_q + \Phi_{\tilde{v}} + \Phi_M, \tag{A.9}$$

where in the electric theory the relevant contributions are

$$\begin{aligned} \Phi_Q = 2 \sum_{I \neq J} \text{sign}(s_I - s_J) & \left[\left(2\omega f_J - \sum_{a=1}^{f_J} (m_a^J + n_a^J) \right) \sum_{i=1}^{n_I} \sigma_i^I + \omega n_I \sum_{a=1}^{f_J} (m_a^J - n_a^J) \right. \\ & \left. + \left(n_I \sum_{a=1}^{f_J} (m_a^J + n_a^J) - 2\omega n_I f_J \right) (s_I - s_J) - \frac{n_I}{2} \sum_{a=1}^{f_J} ((m_a^J)^2 - (n_a^J)^2) \right] \end{aligned} \tag{A.10}$$

$$\Phi_V = -2\omega \sum_{I \neq J} \text{sign}(s_I - s_J) \left[2n_J \sum_{i=1}^{n_I} \sigma_i^I - n_I n_J (s_I - s_J) \right], \tag{A.11}$$

while in the magnetic theory we have

$$\Phi_q = \sum_{I \neq J} \text{sign}(s_I - s_J) \left[\tilde{n}_I \sum_{a=1}^{f_J} \left((m_a^J)^2 - (n_a^J)^2 \right) + 2 \sum_{i=1}^{\tilde{n}_I} \tilde{\sigma}_i^I \sum_{a=1}^{f_J} (m_a^J + n_a^J) - 2\tilde{n}_I \sum_{a=1}^{f_J} (m_a^J + n_a^J)(s_I - s_J) \right] \tag{A.12}$$

$$\Phi_{\tilde{v}} = -2\omega \sum_{I \neq J} \text{sign}(s_I - s_J) \left[2\tilde{n}_J \sum_{i=1}^{\tilde{n}_I} \tilde{\sigma}_i^I - \tilde{n}_I \tilde{n}_J (s_I - s_J) \right] \tag{A.13}$$

$$\Phi_M = \sum_{I \neq J} \text{sign}(s_I - s_J) \left[2 \sum_{a=1}^{f_I} m_a^I \sum_{a=1}^{f_J} n_a^J - f_I \sum_{a=1}^{f_J} \left((m_a^J)^2 - (n_a^J)^2 \right) + 2\omega f_I \sum_{a=1}^{f_J} (m_a^J - n_a^J) - 2f_I \left(\omega f_J - \sum_{a=1}^{f_I} (m_a^J + n_a^J) \right) (s_I - s_J) \right]. \tag{A.14}$$

The phases can be reorganized into a divergent term, an FI term and a real mass contribution. The divergent term has to be the same in the electric and in the magnetic phase. In the electric phase the divergent term is

$$2 \sum_{I \neq J} n_I |s_I - s_J| \left[\sum_{a=1}^{f_J} (m_a^J + n_a^J) - \omega (2f_J - n_J) \right], \tag{A.15}$$

and it is straightforward to check that it coincides with the one obtained in the magnetic phase. The electric and the magnetic FI terms are

$$\lambda_e^I \sum_{i=1}^{n_I} \sigma_i^I \equiv 2 \sum_{i=1}^{n_I} \sigma_i^I \sum_{J(\neq I)=1}^K \text{sign}(s_I - s_J) \left[2\omega (f_J - n_J) - \sum_{a=1}^{f_J} (m_a^J + n_a^J) \right], \tag{A.16}$$

$$\lambda_m^I \sum_{i=1}^{\tilde{n}_I} \tilde{\sigma}_i^I \equiv -2 \sum_{i=1}^{\tilde{n}_I} \tilde{\sigma}_i^I \sum_{J(\neq I)=1}^K \text{sign}(s_I - s_J) \left[2\omega (f_J - n_J) - \sum_{a=1}^{f_J} (m_a^J + n_a^J) \right], \tag{A.17}$$

where in the relations above, I is fixed. As expected, $\lambda_e^I = -\lambda_m^I$ for each gauge group $U(n_I)$ in the quiver. By summing the other real mass contributions to the phase, most of the terms cancel among the electric and the magnetic theory. We are left with the contribution

$$2 \sum_{I \neq J} \text{sign}(s_I - s_J) \left[\omega (f_I - n_I) \sum_{l=1}^{f_J} (m_a^J - n_a^J) + \sum_{a=1}^{f_I} m_a^I \sum_{a=1}^{f_J} n_a^J \right]. \tag{A.18}$$

In order to show that it is vanishing as well let us parameterize the real masses as

$$m_a^I = \tilde{m}_a^I + m_A^I \frac{\alpha_I}{f_I}, \quad n_a^I = \tilde{n}_a^I + m_A^I \frac{\alpha_I}{f_I}, \tag{A.19}$$

with the constraints on the non-Abelian symmetries and on the axial symmetries

$$\sum_{a=1}^{f_I} M_a^I = \sum_{a=1}^{f_I} N_a^I = 0, \quad I = 1, \dots, K; \quad \sum_{I=1}^K m_A^I \alpha_I = \omega(F - N). \tag{A.20}$$

Observe that the last constraint shows that there are $K - 1$ axial symmetries and one is redundant. The phase (A.18) becomes

$$2 \sum_{I \neq J} \text{sign}(s_I - s_J) \alpha_I \alpha_J m_A^I m_A^J = 0. \tag{A.21}$$

Appendix B. Reduction of the superconformal index to 3D

The reduction of four-dimensional dualities as effective dualities on S_r^1 and the $r \rightarrow 0$ limit can be, in general, reproduced by localization. By considering an opportune scaling limit on the fugacities weighting the SCI I_{4d} , it has indeed been shown [39–41,8] that the three-dimensional partition function on a (possibly squashed) three-sphere $Z_{S_b^3}$ [42,23,43,44] can be recovered. The integral identity relating the SCI of a pair of four-dimensional Seiberg dual phases translates in an identity relating the partition functions of a pair of effective three-dimensional dualities on S_r^1 . The presence of an η -superpotential, constraining the three-dimensional duality on S_r^1 , translates to the reduction of a constraint among the four-dimensional fugacities into a constraint among the three-dimensional real masses. Such constraints are referred to in the literature as balancing conditions, both in four and in three dimensions. These constraints avoid the generation of symmetries in three dimensions which are anomalous in four dimensions (e.g. axial symmetries), playing essentially the role of the η -superpotential in localization.

Recovering the usual three-dimensional limit on $Z_{S_b^3}$ requires triggering a real mass and/or a Higgs flow. This corresponds to shifting some (real) parameters in the partition function by an infinite amount. In these cases some care is necessary to show that the divergent parts of the identities coincide. These real mass flows also modify the balancing condition, allowing, in general, the generation of axial-like symmetries, forbidden on S_r^1 . In our discussion we made large use of these ideas to corroborate the new three-dimensional dualities obtained from the brane picture. In this appendix we review the basic formalism, both to make the discussion self-contained and to provide some references for the main formulas used in the body of the paper.

In the following, we briefly summarize some formal aspects of the reduction of I_4 to $Z_{S_b^3}$. Let us start by introducing the notion of the index (see [45–47,27] for details). It can be defined as

$$I = \text{Tr}(-1)^F e^{-\beta H} (pq)^{\frac{\Delta}{2}} p^{j_1+j_2-R/2} q^{j_1-j_2-R/2} \prod_a u_a^{q_a}, \tag{B.1}$$

where F represents the fermion number, the Hamiltonian H is defined on $S^3 \times \mathbb{R}$, the fugacities p and q refer to the $SO(4) = SU(2)_l \times SU(2)_r$ isometry of S^3 , where j_1 and j_2 are the third spin components, and R is the $U(1)$ R-charge. There are also fugacities u_a , referring to the Cartan of the global (and gauge) symmetry group. The fugacities p and q satisfy the conditions

$$\text{Im}(pq) = 0, \quad |p/q| = 1, \quad |pq| < 1. \tag{B.2}$$

The SCI receives non-vanishing contributions only from states satisfying $H = 0$. The calculation of the SCI for a gauge theory proceeds by computing the single particle index and then taking the plethystic exponential. This corresponds in localization to calculating the one-loop determinants of the matter and the vector multiplets. They can be formulated as elliptic Gamma functions,

$$\Gamma_e(y; p, q) \equiv \Gamma_e(y) \equiv \prod_{j,k=0}^{\infty} \frac{1 - p^{j+1}q^{k+1}/y}{1 - p^j q^k y}, \tag{B.3}$$

where y refers to the fugacities of the global and gauge symmetries. The gauge-invariant quantities contributing to the index are found by integrating over the holonomy of the gauge group. Finally, one arrives at the formula

$$I_G = \frac{\kappa^{r_G}}{|W|} \oint_{T^G} \frac{dz_i}{2\pi i z_i} \prod_{\alpha \in G_+} \Gamma_e^{-1}(z^{\pm\alpha}) \prod_{\rho \in \mathcal{R}_I, \tilde{\rho} \in \tilde{\mathcal{R}}_I} \Gamma_e(z^\rho u^{\tilde{\rho}} (pq)^{\frac{R_I}{2}}), \tag{B.4}$$

where $\kappa = (p; p)(q; q)$ and $(x; p) = \prod_{k=0}^{\infty} (1 - xp^k)$. Here, the label $\alpha \in G_+$ refers to the positive roots of G , and $|W|$ is the dimension of the Weyl group. The weight ρ_I stands for the representation of the matter multiplets under the gauge group and the weight $\tilde{\rho}_I$ refers to the representation of the matter multiplets under the flavor symmetry group. The fugacity z is in the Cartan of the gauge symmetry and the fugacity u is in the Cartan of the flavor symmetry. The integral is over the maximal Abelian torus of G , denoted as T^G . The R charge is denoted by R_I .

The partition function is obtained by a KK reduction on S^1 of the states contributing to the four-dimensional index. The reduction is done on $S_b^3 \times \tilde{S}^1$, where S_b^3 represents a squashed three-sphere preserving $U(1)^2 \subset SO(4)$ and b is the squashing parameter. Defining as \tilde{r}_1 the radius of \tilde{S}^1 the fugacities above can be expressed as

$$p = e^{2\pi i \tilde{r}_1 \omega_1}, \quad q = e^{2\pi i \tilde{r}_1 \omega_2}, \quad u_a = e^{2\pi i \tilde{r}_1 \mu_a}, \quad z_i = e^{2\pi i \tilde{r}_1 \sigma_i}. \tag{B.5}$$

In the three-dimensional language, μ_a are real masses for the flavor symmetries and σ_i are the real scalars in the three-dimensional vector multiplets. The parameters $\omega_{1,2}$ are related to the squashing parameter b by $\omega_1 = ib$ and $\omega_2 = ib^{-1}$. We also define the linear combination $\omega \equiv \frac{\omega_1 + \omega_2}{2}$.

The four-dimensional superconformal index reduces to the three-dimensional partition function computed on the squashed three-sphere S_b^3 . The Bogomol'nyi–Prasad–Sommerfield (BPS) states contributing to the four-dimensional index have to be KK reduced on the circle. The massless modes in this reduction are the states contributing to the partition function. In order to perform the KK reduction, it is necessary that all the fugacities appearing in the index flow to unity. The limit $\tilde{r}_1 \rightarrow 0$ corresponds to

$$\lim_{\tilde{r}_1 \rightarrow 0} \Gamma_e(e^{2\pi i \tilde{r}_1 x}; e^{2\pi i \tilde{r}_1 \omega_1}, e^{2\pi i \tilde{r}_1 \omega_2}) = e^{\frac{i\pi^2}{6\tilde{r}_1 \omega_1 \omega_2} (x-\omega)} \Gamma_h(x; \omega_1, \omega_2). \tag{B.6}$$

This formula reduces the one-loop determinants of the four-dimensional fields to the ones of the three-dimensional fields, or more formally, the elliptic gamma functions Γ_e to the hyperbolic gamma function Γ_h defined as

$$\Gamma_h(x; \omega_1, \omega_2) \equiv \Gamma_h(x) \equiv e^{\frac{i\pi}{2\omega_1 \omega_2} ((x-\omega)^2 - \frac{\omega_1^2 + \omega_2^2}{12})} \prod_{j=0}^{\infty} \frac{1 - e^{\frac{2\pi i}{\omega_1} (\omega_2 - x)} e^{\frac{2\pi i \omega_2 j}{\omega_1}}}{1 - e^{-\frac{2\pi i}{\omega_2} x} e^{-\frac{2\pi i \omega_1 j}{\omega_2}}}. \tag{B.7}$$

Observe that the divergent prefactor in (B.6) represents the four-dimensional gravitational anomalies and it coincides in the four-dimensional Seiberg-dual phases. The general expression for the partition function of a three-dimensional gauge theory on S_b^3 is given by

$$Z_{G;k}(\lambda; \vec{\mu}) = \frac{1}{|W|} \int \prod_{i=1}^G \frac{d\sigma_i}{\sqrt{-\omega_1 \omega_2}} e^{\frac{ik\pi \sigma_i^2}{\omega_1 \omega_2} + \frac{2\pi i \lambda \sigma_i}{\omega_1 \omega_2}} \frac{\prod_I \Gamma_h(\omega \Delta_I + \rho_I(\sigma) + \tilde{\rho}_I(\mu))}{\prod_{\alpha \in G_+} \Gamma_h(\pm\alpha(\sigma))}. \tag{B.8}$$

The integral is performed over the eigenvalues of the scalar σ in the Cartan of the gauge group G . The parameter μ represents the real mass in the Cartan of the flavor symmetry. The parameter λ is an FI term while k refers to the Chern–Simons (CS) action, if present in the dynamics (observe that an analogue CS term for the flavor symmetry can be turned on and it is related to the contact terms of the global currents [48,49]). The R charge is denoted by Δ_I .

References

- [1] N. Seiberg, Electric–magnetic duality in supersymmetric non Abelian gauge theories, Nucl. Phys. B 435 (1995) 129–146, arXiv:hep-th/9411149.
- [2] O. Aharony, et al., Aspects of $N = 2$ supersymmetric gauge theories in three-dimensions, Nucl. Phys. B 499 (1997) 67–99, arXiv:hep-th/9703110.
- [3] O. Aharony, IR duality in $d = 3$ $N = 2$ supersymmetric $USp(2N(c))$ and $U(N(c))$ gauge theories, Phys. Lett. B 404 (1997) 71–76, arXiv:hep-th/9703215.
- [4] A. Karch, Seiberg duality in three-dimensions, Phys. Lett. B 405 (1997) 79–84, arXiv:hep-th/9703172.
- [5] A. Giveon, D. Kutasov, Seiberg duality in Chern–Simons theory, Nucl. Phys. B 812 (2009) 1–11, arXiv:0808.0360 [hep-th].
- [6] F. Benini, C. Closset, S. Cremonesi, Comments on 3d Seiberg-like dualities, JHEP 1110 (2011) 075, arXiv:1108.5373 [hep-th].
- [7] F. Benini, S. Benvenuti, S. Pasquetti, SUSY monopole potentials in 2+1 dimensions, arXiv:1703.08460 [hep-th], 2017.
- [8] O. Aharony, S.S. Razamat, N. Seiberg, B. Willett, 3d dualities from 4d dualities, JHEP 1307 (2013) 149, arXiv:1305.3924 [hep-th].
- [9] A. Amariti, et al., The braneology of 3D dualities, J. Phys. A 48 (26) (2015) 265401, arXiv:1501.06571 [hep-th].
- [10] A. Hanany, E. Witten, Type IIB superstrings, BPS monopoles, and three-dimensional gauge dynamics, Nucl. Phys. B 492 (1997) 152–190, arXiv:hep-th/9611230.
- [11] I. Affleck, J.A. Harvey, E. Witten, Instantons and (super)symmetry breaking in $(2 + 1)$ -dimensions, Nucl. Phys. B 206 (1982) 413.
- [12] A. Amariti, et al., 4D/3D reduction of dualities: mirrors on the circle, JHEP 10 (2015) 048, arXiv:1504.02783 [hep-th].
- [13] J. de Boer, et al., Mirror symmetry in three-dimensional theories, $SL(2, Z)$ and D-brane moduli spaces, Nucl. Phys. B 493 (1997) 148–176, arXiv:hep-th/9612131.
- [14] N.M. Davies, T.J. Hollowood, V.V. Khoze, M.P. Mattis, Gluino condensate and magnetic monopoles in supersymmetric gluodynamics, Nucl. Phys. B 559 (1999) 123–142, arXiv:hep-th/9905015.
- [15] S.H. Katz, C. Vafa, Geometric engineering of $N = 1$ quantum field theories, Nucl. Phys. B 497 (1997) 196–204, arXiv:hep-th/9611090.
- [16] J. de Boer, K. Hori, Y. Oz, Dynamics of $N = 2$ supersymmetric gauge theories in three-dimensions, Nucl. Phys. B 500 (1997) 163–191, arXiv:hep-th/9703100.
- [17] S. Elitzur, et al., Brane dynamics and $N = 1$ supersymmetric gauge theory, Nucl. Phys. B 505 (1997) 202–250, arXiv:hep-th/9704104.
- [18] A. Amariti, D. Orlando, S. Reffert, String theory and the 4D/3D reduction of Seiberg duality. A review, arXiv:1611.04883 [hep-th], 2016.
- [19] A. Giveon, D. Kutasov, Brane dynamics and gauge theory, Rev. Mod. Phys. 71 (1999) 983–1084, arXiv:hep-th/9802067.
- [20] A. Hanany, A. Zaffaroni, Issues on orientifolds: on the brane construction of gauge theories with $SO(2n)$ global symmetry, JHEP 9907 (1999) 009, arXiv:hep-th/9903242.
- [21] A. Hanany, B. Kol, On orientifolds, discrete torsion, branes and M theory, JHEP 0006 (2000) 013, arXiv:hep-th/0003025.
- [22] A. Hanany, J. Troost, Orientifold planes, affine algebras and magnetic monopoles, JHEP 0108 (2001) 021, arXiv:hep-th/0107153.
- [23] D.L. Jafferis, The exact superconformal R-symmetry extremizes Z, JHEP 1205 (2012) 159, arXiv:1012.3210 [hep-th].
- [24] K.A. Intriligator, P. Pouliot, Exact superpotentials, quantum vacua and duality in supersymmetric $SP(N(c))$ gauge theories, Phys. Lett. B 353 (1995) 471–476, arXiv:hep-th/9505006.

- [25] O. Aharony, S.S. Razamat, N. Seiberg, B. Willett, $3d$ dualities from $4d$ dualities for orthogonal groups, JHEP 1308 (2013) 099, arXiv:1307.0511.
- [26] O. Aharony, N. Seiberg, Y. Tachikawa, Reading between the lines of four-dimensional gauge theories, JHEP 1308 (2013) 115, arXiv:1305.0318.
- [27] V. Spiridonov, G. Vartanov, Elliptic hypergeometry of supersymmetric dualities, Commun. Math. Phys. 304 (2011) 797–874, arXiv:0910.5944 [hep-th].
- [28] V.P. Spiridonov, G.S. Vartanov, Elliptic hypergeometry of supersymmetric dualities II. Orthogonal groups, knots, and vortices, Commun. Math. Phys. 325 (2014) 421–486, arXiv:1107.5788 [hep-th].
- [29] K.A. Intriligator, R. Leigh, M. Strassler, New examples of duality in chiral and nonchiral supersymmetric gauge theories, Nucl. Phys. B 456 (1995) 567–621, arXiv:hep-th/9506148.
- [30] H. Kim, J. Park, Aharony dualities for $3d$ theories with adjoint matter, JHEP 06 (2013) 106, arXiv:1302.3645 [hep-th].
- [31] S. Kim, The Complete superconformal index for $N = 6$ Chern–Simons theory, Nucl. Phys. B 821 (2009) 241–284, arXiv:0903.4172 [hep-th], Erratum: Nucl. Phys. B 864 (2012) 884.
- [32] F. Benini, A. Zaffaroni, A topologically twisted index for three-dimensional supersymmetric theories, JHEP 07 (2015) 127, arXiv:1504.03698 [hep-th].
- [33] T. Morita, V. Niarchos, F-theorem, duality and SUSY breaking in one-adjoint Chern–Simons–Matter theories, Nucl. Phys. B 858 (2012) 84–116, arXiv:1108.4963 [hep-th].
- [34] P. Agarwal, A. Amariti, M. Siani, Refined checks and exact dualities in three dimensions, JHEP 10 (2012) 178, arXiv:1205.6798 [hep-th].
- [35] B.R. Safdi, I.R. Klebanov, J. Lee, A crack in the conformal window, JHEP 04 (2013) 165, arXiv:1212.4502 [hep-th].
- [36] J. Lee, M. Yamazaki, Gauging and decoupling in $3d \mathcal{N} = 2$ dualities, JHEP 06 (2016) 077, arXiv:1603.02283 [hep-th].
- [37] S. Hellerman, D. Orlando, S. Reffert, String theory of the Omega deformation, JHEP 01 (2012) 148, arXiv:1106.0279 [hep-th].
- [38] S. Hellerman, D. Orlando, S. Reffert, The Omega deformation from string and M-theory, JHEP 1207 (2012) 061, arXiv:1204.4192 [hep-th].
- [39] A. Gadde, W. Yan, Reducing the $4d$ index to the S^3 partition function, JHEP 12 (2012) 003, arXiv:1104.2592 [hep-th].
- [40] F.A.H. Dolan, V.P. Spiridonov, G.S. Vartanov, From $4d$ superconformal indices to $3d$ partition functions, Phys. Lett. B 704 (2011) 234–241, arXiv:1104.1787 [hep-th].
- [41] Y. Imamura, Relation between the $4d$ superconformal index and the S^3 partition function, JHEP 09 (2011) 133, arXiv:1104.4482 [hep-th].
- [42] A. Kapustin, B. Willett, I. Yaakov, Exact results for Wilson loops in superconformal Chern–Simons theories with Matter, JHEP 03 (2010) 089, arXiv:0909.4559 [hep-th].
- [43] N. Hama, K. Hosomichi, S. Lee, Notes on SUSY gauge theories on three-sphere, JHEP 1103 (2011) 127, arXiv:1012.3512 [hep-th].
- [44] N. Hama, K. Hosomichi, S. Lee, SUSY gauge theories on squashed three-spheres, JHEP 05 (2011) 014, arXiv:1102.4716 [hep-th].
- [45] J. Kinney, J.M. Maldacena, S. Minwalla, S. Raju, An index for 4 dimensional super conformal theories, Commun. Math. Phys. 275 (2007) 209–254, arXiv:hep-th/0510251.
- [46] C. Romelsberger, Counting chiral primaries in $N = 1$, $d = 4$ superconformal field theories, Nucl. Phys. B 747 (2006) 329–353, arXiv:hep-th/0510060.
- [47] F.A. Dolan, H. Osborn, Applications of the superconformal index for protected operators and q -hypergeometric identities to $N = 1$ dual theories, Nucl. Phys. B 818 (2009) 137–178, arXiv:0801.4947 [hep-th].
- [48] C. Closset, et al., Contact terms, unitarity, and F-maximization in three-dimensional superconformal theories, JHEP 10 (2012) 053, arXiv:1205.4142 [hep-th].
- [49] C. Closset, et al., Comments on Chern–Simons contact terms in three dimensions, JHEP 09 (2012) 091, arXiv:1206.5218 [hep-th].