

# **tRNAs and proteins use the same import channel for translocation across the mitochondrial outer membrane of trypanosomes**

**Moritz Niemann<sup>1)</sup>, Anke Harsman<sup>1)</sup>, Jan Mani<sup>1)</sup>, Christian D. Peikert<sup>3)</sup>, Silke Oeljeklaus<sup>3,4)</sup>, Bettina Warscheid<sup>3,4)</sup>, Richard Wagner<sup>2)</sup> and André Schneider<sup>1)</sup>**

<sup>1)</sup> Department of Chemistry and Biochemistry, University of Bern, Freiestrasse 3, Bern CH-3012, Switzerland

<sup>2)</sup> Department of Life Sciences and Chemistry, Jacobs University Bremen, 28719 Bremen, Germany

<sup>3)</sup> Department of Biochemistry and Functional Proteomics, Institute of Biology II, Faculty of Biology, University of Freiburg, Schänzlestr. 1, 79104 Freiburg, Germany

<sup>4)</sup> BIOSS Centre for Biological Signalling Studies, University of Freiburg, Schänzlestr. 18, 79104 Freiburg, Germany

## Abstract

Mitochondrial tRNA import is widespread, but the mechanism by which tRNAs are imported remains largely unknown. The mitochondrion of the parasitic protozoan *Trypanosoma brucei* lacks tRNA genes and thus imports all tRNAs from the cytosol. Here we show that in *T. brucei* in vivo import of tRNAs requires four subunits of the mitochondrial outer membrane protein translocase but not the two receptor subunits, one of which is essential for protein import. The latter shows that it is possible to uncouple mitochondrial tRNA import from protein import. Ablation of the intermembrane space domain of the translocase subunit, ATOM14, on the other hand while not affecting the architecture of the translocase, impedes both protein and tRNA import. A protein import intermediate arrested in the translocation channel prevents both protein and tRNA import. In the presence of tRNA, blocking events of single channel currents through the pore formed by recombinant ATOM40 were detected in electrophysiological recordings. These results indicate that both types of macromolecules use the same import channel across the outer membrane. However, while tRNA import depends on the core subunits of the protein import translocase, it does not require the protein import receptors indicating that the two processes are not mechanistically linked.

## **Significance**

**In most eukaryotes mitochondrial function requires not only import of proteins but also import of at least some tRNAs. Trypanosomes are extreme in that they lack mitochondrial tRNA genes and therefore must import all of their mitochondrial tRNAs from the cytosol. Here we show that in trypanosomes both proteins and tRNAs use the same  $\beta$ -barrel protein pore to be translocated across the mitochondrial outer membrane. Moreover, we show that tRNA import can be uncoupled from protein import. Based on these results we propose the new "alternate import model", in which tRNAs use the same outer membrane import pore than proteins but are imported as naked molecules. The new model combines features of the previously proposed "co-import" and "direct import" models.**

\body

Mitochondria derive from an endosymbiotic event between an archaeon and a bacterium approximately 2 billion years ago (1-3). While they have maintained their own genome it encodes only a small number of proteins (4, 5). Most mitochondrial proteins (>95%) are encoded in the nucleus and after their synthesis in the cytosol imported into the organelle. Mitochondrial protein import has been studied in great detail and the machineries as well as the mechanisms of the process are well understood (6-9).

The mitochondrial genomes of many eukaryotes lack a complete set of tRNA genes and it has been shown that this lack is compensated for by import of a small fraction of the corresponding cytosolic tRNAs (10-12). Moreover, it has been postulated that even in organisms such as mammals, whose mitochondrial genome encodes a complete set of mitochondrial tRNAs, some apparently redundant tRNAs may be imported into mitochondria (13, 14). Furthermore, some studies indicate that mammals import the 5S rRNA (15-17) and the RNA subunit of RNase P (18-20). The function of these imported RNAs is unclear as the imported 5S rRNA has not been detected in the structure of the mammalian mitoribosome (21, 22) and a protein-only RNase P has been characterized in mammalian mitochondria (23).

In contrast to mitochondrial protein import, mitochondrial tRNA import is poorly understood. It has experimentally mainly been studied in yeast, plants and trypanosomatids and two general models have been proposed of how the process might work. In the first one, the tRNA is coimported with a mitochondrial precursor protein along the protein import pathway. This model applies to the tRNA<sup>Lys</sup> of yeast and has been elucidated in detail (24-26). In the second more vaguely defined model, which appears to apply to plants and protozoa (27-29) as well as to some tRNAs that appear to be imported into mammalian mitochondria (13, 14), tRNAs are directly imported independent of ongoing protein import and thus do not use the same translocation channel as proteins (11, 12).

The parasitic protozoan *T. brucei* is an excellent system to study mitochondrial tRNA import. Unlike most other eukaryotes its mitochondrial genome does not encode any tRNAs, indicating that all of its organellar tRNAs, more than 30 different species, must be imported from the cytosol in quantities sufficient to support mitochondrial translation (28-30).

The protein import machinery of the *T. brucei* mitochondrion has recently been characterized and was shown to be highly diverged when compared to other eukaryotes (31). Thus, protein import across the outer membrane (OM) is mediated by the archaic translocase of the OM (ATOM), a complex consisting of 6 subunits (32, 33). (more recently TbLOK1, initially identified as a mitochondrial morphology protein, was also shown to be a component of the ATOM complex raising the number of subunits to 7 (34)). The core subunits of the ATOM complex include ATOM40 and ATOM14, which are remotely related to Tom40 and Tom22 of other eukaryotes, as well as ATOM12 and ATOM11 which are exclusively found in trypanosomatids (31). These core subunits are associated with the two receptors ATOM46 and ATOM69 which are unique to trypanosomatids. With the exception of ATOM46, which is required for growth at elevated temperature only, all ATOM subunits are essential. To elucidate whether tRNA import in *T. brucei* complies with the direct import or the co-import model we investigated, whether subunits of the OM protein translocase are required for the process and whether tRNA import is coupled to protein import. The results show that: i) some but not all essential subunits of the OM protein translocase are required for tRNA import, ii) that tRNAs and proteins use the same pore for import and iii) that tRNA import likely is not coupled to protein import.

## **Results**

### **In vivo import of tRNAs depends on ATOM core subunits**

Mitochondrial tRNA import in trypanosomatids has been studied using both in vivo (35-37) and in vitro assays (27, 38-41). However, even though it is known that in vivo import of tRNAs requires cytosolic factors (42) in vitro import has only been analyzed in the absence of cytosolic fractions. It therefore remains unclear to which extent these in vitro systems reflect the in vivo situation. In order to be sure to obtain physiologically meaningful results we were focusing on an in vivo approach which consisted in measuring tRNA steady state levels in RNAi cell lines ablated for the individual ATOM components.

Inhibition of tRNA import will result in reduced steady state levels of mitochondrial tRNAs. However, a possible confounding factor might be that in absence of protein import mitochondrial translation will eventually also be inhibited. If tRNAs are not used in translation they might be unstable which would result in a decreased steady state

level even though tRNA import is not affected. Fig. S1 shows that this is not the case as the steady state levels of mitochondrial tRNAs is not affected in a RNAi cell line that is unable to synthesize mitochondrial proteins due to the absence of the mitochondrial translation elongation factor Tu (EF-Tu) (43). This indicates that the steady state levels of mitochondrial tRNAs are indeed a good proxy to estimate the efficiency of import.

Thus to determine tRNA steady state levels, mitochondrial RNA from different uninduced and induced ATOM subunit-RNAi cell lines (32) was isolated from equal cell equivalents of digitonin-extracted pellets. To prevent pleiotropic effects the induced cell lines were analyzed at the onset of the growth phenotype (Fig. S2). As an internal control for the yield of the multistep RNA purification protocol 1 pmol of a short synthetic RNA molecule was added to each pellet prior to RNA isolation. Subsequently the RNA samples were separated on gels, visualized by ethidium bromide-staining and analyzed by Northern hybridization for the presence of the efficiently imported tRNA<sup>Ile</sup> (30). In addition, protein samples of the digitonin-extracted pellets were analyzed on immunoblots for the levels of lipoamide dehydrogenase (LDH). LDH is a soluble protein of the mitochondrial matrix. Its recovery in the pellet fractions indicates the structural integrity of the mitochondria after digitonin extraction. Moreover, LDH has a long half-life and its mitochondrial steady state levels, in contrast to many other mitochondrial proteins, are only affected at late time points after ablation of protein import. Thus, its abundance at early times of induction after RNAi is proportional to the amounts of mitochondria. The steady state levels of the tRNA<sup>Ile</sup> in uninduced and induced RNAi cell lines was quantified on Northern blots and normalized with the levels of the synthetic RNA molecule that serves as a control for the RNA extraction. All experiments were replicated four to six times to allow for statistical analysis.

In a first set of experiments we focused on the pore-forming ATOM40 and the core subunits ATOM11, ATOM12 and ATOM14. As previously shown, knockdown of these proteins leads to a growth arrest under all tested conditions including in an engineered bloodstream form cell line that can grow in the absence of mitochondrial DNA. Moreover, concomitant with the growth arrest an inhibition of mitochondrial protein import was observed both in vitro and in vivo (32) (Fig. S2A).

As evident from Figure 1A, down-regulation of any of the four tested ATOM subunits resulted in a significant reduction of the amounts of mitochondrial tRNA<sup>Ile</sup> which likely is due to an inhibition of mitochondrial tRNA import. The ethidium bromide-stained gel shows that the same reduction is seen for the entire tRNA region indicating that all tRNAs behave the same. With 75% the reduction of the tRNA<sup>Ile</sup> levels in the absence of

ATOM40 was slightly higher than the approximately 60% observed for ATOM11, ATOM12 and ATOM14. Thus, these results show that tRNA import becomes drastically reduced in the absence of any of the four tested ATOM complex core subunits suggesting they are directly or indirectly required for mitochondrial tRNA import.

In many RNAi cell lines a set of bands larger than the mitochondrial rRNAs are visible on the ethidium bromide-stained gel. In some cases these bands are enriched in the induced RNAi cell lines (Fig. 1A). The Northern analysis in Fig. S3 shows that these bands are not unprocessed mitochondrial rRNAs as they do not hybridize with a mitochondrial rRNA-specific probe. Rather they correspond to cytosolic rRNAs suggesting that in the absence of protein import cytosolic ribosomes are recruited to the surface of mitochondria.

### **In vivo import of tRNAs does not require ATOM receptor subunits**

In a second set of experiments we asked, whether the protein receptor subunits ATOM46 and ATOM69 are also required for tRNA import. The role of these two proteins in protein import has previously been studied (32). Their large cytosolic domains bind mitochondrial precursor proteins with distinct but in part overlapping specificities. ATOM69, as all ATOM core subunits, is essential for growth under all conditions and for mitochondrial protein import in vivo. ATOM46-depleted cells grow normally at the standard temperature of 27°C but show a growth arrest at the elevated temperature of 33°C (Fig. S2B) (32). In the ATOM69/ATOM46-double knockdown cell line the growth phenotype is stronger than in the ATOM69-RNA cell line and in vitro import of proteins is inhibited as well (32). Thus, regarding growth and protein import the ATOM69/ATOM46-double knock down cell line shows the same phenotypes as the ones observed after ablation of any of the ATOM core subunits discussed in the previous paragraph.

Figure 1B shows that, in contrast to the core subunits, ablation of the receptor subunit ATOM69 did not lead to reduced steady state levels of mitochondrial tRNAs. This was not only the case at the time when the growth arrest becomes evident (Fig. 1B) but also two days after appearance the growth phenotype (Fig. S4). At this later time point protein import is completely abolished and the cells may start to suffer from secondary effects caused by the lack of imported proteins.

The same results were obtained when cells ablated for ATOM46 were assayed at 33°C. Importantly, also the simultaneous down-regulation of both receptor subunits, had only

a negligible effect on mitochondrial tRNA levels (Fig. 1B). These data indicate that while tRNA import requires the ATOM core subunits, it is independent of the protein import receptors ATOM46 and ATOM69. Thus in the ATOM46/ATOM69 double knock down cell line tRNA import has been uncoupled from protein import. These results strongly suggest that the inhibition of mitochondrial tRNA import observed after ablation of the individual ATOM core subunits (Fig. 1A) is a consequence of a non functional ATOM complex

### **OM proteins reduced in abundance in ATOM subunit-lacking cells**

The analysis of the results in Fig. 1 is complicated by the fact that ablation of a specific ATOM subunit affects the steady state levels of the other subunits in a complex way (32). To better define the role of ATOM40, ATOM11 and ATOM12 in tRNA import we compared the effects their ablation has on the previously characterized OM proteome (44) using mitochondria-enriched digitonin pellets of uninduced and induced RNAi cell lines in combination with SILAC. Fig. 2 shows all proteins of the OM proteome whose levels were reduced more than two-fold upon ablation of the indicated ATOM subunits:

- Ablation of ATOM40 lead to greatly reduced levels of ATOM14, ATOM11, ATOM46 and ATOM69 (Fig. 2A). The downregulated protein import receptors ATOM46 and ATOM69 cannot be responsible for the phenotype as their individual or combined ablation does not affect tRNA import (Fig. 1B) (Fig. S4). Thus, the proteomic analysis of the ATOM40 RNAi cell line firmly links tRNA import to the ATOM complex. However, it is not clear whether the inhibition of import is due to individual or combined ablation of ATOM40, ATOM14 and ATOM11.

- Ablation of ATOM11, while affecting tRNA import, does not reduce the levels of any other ATOM subunits besides the receptors ATOM46 and ATOM69, suggesting it might have a direct role in mitochondrial tRNA import (Fig. 2B).

- The inhibition of tRNA import observed in the ATOM12-RNAi cell line is likely due to ATOM12 itself which is most efficiently downregulated. The only other ATOM subunit whose level is more than 2-fold reduced is ATOM14 (Fig. 2C).

The data shown in Fig. 2 allow a critical examination of the possibility that the observed tRNA import inhibition in the ATOM40-, ATOM11-, and ATOM12-RNAi cell lines might be an indirect effect, caused by reduced import of a hypothetical OM tRNA import factor that is not an ATOM subunit. In all RNAi cell lines shown in Fig. 1A tRNA import is reduced by at least 2.5-fold indicating that, if a hypothetical tRNA import factor would



be responsible for the phenotype, its level should be reduced to the same extent. However except for the ATOM subunits (Fig. 2, red bars) there are only very few proteins that are more than 2-fold downregulated in any of the three proteomically analyzed RNAi cell lines. Four candidates are found in the ATOM40-RNAi cell lines, namely MICU, POMP20, the NADH-ubiquinone oxidoreductase Tb927.11.9930 and POMP15 (Fig. 2A). In the ATOM11- and ATOM12-RNAi cell lines, finally, no proteins, other than ATOM11 and ATOM12 itself, are sufficiently downregulated to explain the observed reduction in tRNA import (Fig. 2BC).

More than 35 different tRNAs need to be imported into the trypanosome mitochondrion in quantities sufficient to support organellar translation (30). Thus, in analogy to mitochondrial protein import, factors that mediate tRNA import across the OM are likely abundant proteins. The coverage of the OM proteome we reached in our proteomic analyses was 76 to 87% indicating that all abundant OM proteins were detected. Thus, the results shown in Fig. 2 strongly suggest that the tRNA import phenotypes observed in the RNAi cell lines shown in Fig. 1A are a direct consequence of the ablation of ATOM complex subunits and not caused by reduced levels of a hypothetical as yet unknown tRNA import factor in the OM.

### **tRNA import requires the intermembrane space domain of ATOM14**

Depletion of individual ATOM core subunits has variable effects on the architecture of the high molecular weight ATOM complex (32). It has previously been shown that replacement of full length ATOM14, the remote Tom22 orthologue of *T. brucei*, with a truncated variant that lacks the C-terminal intermembrane space (IMS) exposed domain results in a growth arrest and inhibits mitochondrial protein import (45). However, in contrast to the cell line where the entire ATOM14 is depleted, the assembly of the high molecular weight ATOM complex is not affected in the cells expressing the truncated ATOM14 variant (45). We thus asked the question if this protein-import incompetent ATOM complex is still able to translocate tRNA. As shown in Figure 3 and Fig. S5, this was not the case. Knockdown of ATOM14 using an RNAi cell line targeting the 3'UTR of the ATOM14 mRNA reduced the tRNA steady state levels to similar values as the cell line used in Figure 1A, where the RNAi was directed against the ORF. Expression of the truncated ATOM14 variant in the background of the ATOM14 3'UTR-RNAi cell line however showed the same reduction of tRNA import, indicating that the protein-import defective ATOM complex is also defective for tRNA import. The control in which the 3'UTR cell line was complemented with full length ATOM14 behave like wild-type, as

would be expected. - In summary these results suggest a direct involvement of the IMS-domain of ATOM14 in both protein and tRNA import.

### **Obstruction of the ATOM40 pore abolishes tRNA import**

The  $\beta$ -barrel protein ATOM40, which has some similarity to Tom40 of other eukaryotes (46, 47), forms the pore across which proteins are translocated across the OM. Fig. 1A shows that ablation of ATOM40 impedes mitochondrial tRNA import, which raises the question whether import of tRNAs and proteins uses the same OM pore.

We have recently shown that by expression of a fusion protein comprised of the N-terminal 160 aa of mitochondrial LDH fused to mouse dihydrofolate reductase (DHFR) followed by a C-terminal hemagglutinin (HA)-tag we can produce an import intermediate that is stuck in the import pore (48). Under standard conditions the fusion protein (LDH-DHFR-HA) is completely translocated into the mitochondrial matrix across both mitochondrial membranes. However, in the presence of the folic acid analogue aminopterin the DHFR adopts a conformation that is structurally rigid and therefore cannot be imported. Addition of aminopterin to a cell line expressing LDH-DHFR-HA therefore results in the formation of an intermediate that blocks the import channels in both the ATOM complex and the protein translocase of the inner membrane. It has been shown that in these cell lines further import of mitochondrial proteins is abolished (48).

In the experiment shown in Fig. 4 we compared the kinetics of mitochondrial tRNA abundance between two cell cultures. Both were incubated with aminopterin but only one received tetracycline and thus expressed the LDH-DHFR-HA fusion protein. Aminopterin slows down cell growth, however for the duration of the experiment the mitochondrial integrity and the membrane potential were not compromised, as verified by Mitotracker staining (Fig. S6). The bottom panels of Figure 4 show that expression of LDH-DHFR-HA in the presence of aminopterin lead to the formation of the LDH-DHFR-HA import intermediate, which blocked further import of proteins as evidenced by the accumulation of unprocessed cytochrome oxidase subunit 4 (cox4) precursor on immunoblots. Northern analysis shows a dramatic decline of the levels of the imported tRNA<sup>Ile</sup> that is specific for the cells in which LDH-DHFR-HA is expressed and the import intermediated is formed. The decline increases the longer the channel gets blocked by the import intermediate and is seen for the entire tRNA region visualized on the

ethidium bromide-stained gels. These results strongly suggest that proteins and tRNAs use the same pore for import across the OM.

### **tRNA influences conductance of the recombinant ATOM40 pore**

From the *in vivo* data we concluded that ATOM40 might represent the channel that translocates tRNAs across the OM. Our efforts to demonstrate a direct interaction between tRNAs and ATOM40 by gel shift assays or flotation experiments using vesicles containing recombinant ATOM40 were unsuccessful. This is likely due to the fact that the interaction between the tRNA and its import channel is transient.

Thus in order to measure direct interaction between ATOM40 and tRNAs we performed electrophysiology experiments. We have previously shown that ATOM40 forms a hydrophilic pore of large conductance and high open probability, and that addition of the N-terminal presequence of a substrate protein induced transient pore closure (49). Thus, we analogously expressed and purified recombinant ATOM40 to near homogeneity (Fig. 5A) and reconstituted individual ATOM40 molecules into planar lipid bilayers for single channel current measurement. Figure 5B shows a representative experiment with ATOM40 displaying the typical conductivity of ca. 1.5 nS with few gating events at a holding potential of 100 mV (Figure 5B, upper panel). Instead of adding a presequence peptide as in the previous study we added isolated tRNAs to the sample. Fig. 5B shows that, just as for the peptide, this resulted in channel flickering, indicating that interaction of the added tRNAs with recombinant ATOM40 induced temporary closure or obstruction of the ATOM40 channel (Figure 5B, middle panel). These results are consistent with the results of the LDH-DHFR-HA/aminopterin experiment (Fig. 4) and indicate that import of tRNAs and proteins across the mitochondrial OM use the same import pore.

## Discussion

We have investigated how trypanosomal tRNAs are translocated across the mitochondrial OM under physiological conditions. Our results reveal a tight connection between mitochondrial tRNA and protein import. Ablation of any of the core subunits of the ATOM complex, the trypanosomal analogue of the TOM complex in other eukaryotes, also abolished tRNA import. Replacement of ATOM14 with a truncated variant did not disturb the overall architecture of the ATOM complex (45) but inhibited both protein and tRNA import. Finally, when the mitochondrial protein import channels are blocked by an import substrate whose C-terminal moiety cannot be unfolded (48) we see a concomitant reduction of protein and tRNA import.

At a first glimpse these results appear to support the co-import model, which posits that tRNAs associate with mitochondrial precursor proteins and that the resulting tRNA/protein complexes are imported along the protein import pathway. However, a closer look also reveals discrepancies that are not in line with the co-import model. Ablation of the protein import receptor ATOM69 or combined ablation of both receptors ATOM69 and ATOM46 inhibited protein import, caused a growth arrest (32) but did not affect mitochondrial tRNA import, showing that in *T. brucei* it is possible to uncouple protein from tRNA import. This contrasts with import of the tRNA<sup>Lys</sup> into yeast mitochondria, the best studied example of the co-import model. Here ablation of the protein import receptor Tom20, the trypanosomal analogue of which is ATOM69, abolished import of the tRNA<sup>Lys</sup> since the precursor of mitochondrial lysyl-tRNA synthetase, which serves as carrier for the process, could no longer be imported (24).

Nevertheless it is presently not possible to exclude the co-import model for the translocation of trypanosomal tRNAs across the OM with certainty. However, should it apply, our results impose strong constraints on the properties of a hypothetical tRNA-binding carrier protein. It would need to be imported into mitochondria independently of the two protein import receptors ATOM46 and ATOM69. Presently no such protein is known.

Based on in vitro import experiments it has been proposed that the direct import model applies for mitochondrial tRNA import in plants. Antibody inhibition experiments indicate that the voltage dependent anion channel (VDAC) is the pore, or at least a component thereof, that translocates tRNAs across the mitochondrial OM (50). However, interestingly components of the protein import system such as Tom20 and Tom40 appear also to be involved in the process. The direct import model implies that the tRNAs which will be imported into mitochondria can directly interact with the

translocation pore. This prediction was experimentally confirmed in potato where imported tRNAs were shown to differentially bind to recombinant versions of the two VDAC isoforms (51).

In summary, our results show that mitochondrial tRNA import across the OM in *T. brucei* has features of both the co-import model as well as the direct import model. Thus, it might be useful to introduce a third intermediate model, where tRNAs and proteins are imported by the same machinery and even use the same import pore, but where tRNA import is nevertheless not coupled to protein import (Fig. S7). This novel model implies that imported tRNAs will directly, although probably transiently, interact with the protein import channel and possibly with other components of the protein import machinery. Indeed the results of electrophysiological single channel measurements indicate that bulk tRNAs can transiently block isolated ATOM40 pores in the absence of any of the other ATOM subunits. Moreover, our results suggest that tRNAs may also interact with the IMS domain of ATOM14.

The present study focuses on the mitochondrial OM and provides no information how tRNAs are translocated across the inner membrane. In a highly controversial study in *L. tropica*, a close relative of *T. brucei*, a protein complex of the inner membrane termed RIC, one component of which is the  $\alpha$ -subunit of the F1-ATP synthase complex, has been implicated in tRNA import (52, 53). A more recent in vivo study on the other hand challenges these results and suggests that import of tRNAs across the inner membrane requires TbTim17, the core subunit of the trypanosomal inner membrane translocase (TIM complex) (48), which is consistent with the idea that tRNAs and proteins may also use the same pore for transport across the inner membrane (54).

We conclude that the trypanosomal ATOM complex has a dual function: it translocates not only proteins but also tRNAs across the OM of mitochondria. However, a single ATOM complex likely either transports proteins or tRNAs but not both simultaneously. In wildtype cells there are up to four distinct ATOM complexes of different molecular weights. The main differences between them is the extent to which they are associated with the protein import receptors ATOM46 and ATOM69 (32). It is tempting to speculate that the ATOM complex showing the lowest molecular weight, that is termed the core complex and that essentially lacks the receptor subunits, might be specialized for mitochondrial tRNA import, whereas the ATOM complexes of higher molecular weight might mainly be engaged in protein import.

In *T. brucei* mitochondrial protein import is essential under all conditions, as in all other eukaryotes. Mitochondrial translation and therefore mitochondrial tRNA import was

shown to be essential in both procyclic and bloodstream forms of *T. brucei*. The trypanosomal ATOM complex which mediates both protein and tRNA import and to a large part is composed of trypanosome-specific subunits is therefore an excellent novel drug target.

## Material and Methods

### Transgenic cell lines

All cell lines are based on procyclic *T. brucei* strain 29-13. Cells were grown at 27°C in SDM-79 containing 10% (v/v) fetal calf serum. The tetracycline-inducible RNAi cell lines for ATOM12, ATOM14, ATOM40, ATOM46, ATOM69, the double knockdown ATOM46/69 and the conditional knockout cell line ATOM11 have been describe before (32, 33). Transfection and generation of clonal cell lines was achieved as described previously (55). The ATOM14 3'UTR RNAi cell line and the complementation of this cell line with the truncated ATOM14 lacking the IMS domain is described in (45). The cell line expressing the import intermediate LDH-DHFR-HA fusion protein used to produce the import intermediate was described in (48).

### Monitoring of mitochondrial tRNA abundance

The RNAi cell lines were induced by addition of 1µg/ml tetracycline to the medium. Induction times for ATOM12, ATOM14, ATOM40 and ATOM46/A69 RNAi cell lines were three days. The ATOM69 RNAi cell line was induced for four and six days whereas the ATOM46 cell line was induced for three and a half days at 33°C. The conditional knockout cell line for ATOM11 was assayed three days after removal of tetracycline. Induction times were chosen to coincide with the onset of the growth phenotypes (45). Equal numbers ( $1-2 \times 10^8$ ) of uninduced and induced cells were harvested and a mitochondria-enriched fraction was prepared using digitonin extraction and RNase A digestion (30). Subsequently 1pmol of an RNA oligonucleotide (5'-GGAGCUCGCCCCGGCGAGGCCGUGCCAGCUCUUCGGAGCAAUACUCGGC-3', 49nt) was added to each sample. It serves as internal standard for the yield of the RNA extraction. Finally, mitochondrial RNA was isolated using the acid guanidinium method (56). The resulting RNA fractions were separated on denaturing 10% polyacrylamide gels containing 8M urea and stained with ethidium bromide. Gels were electroblotted (200mA for 14-18 hours) onto charged nylon membranes (Genescreen Plus Hybridization Transfer Membrane), UV cross-linked (250mJ/cm<sup>2</sup>) and probed for tRNA<sup>Ile</sup> and the RNA oligonucleotide (49-mer, see above), which served as internal standard. The blots were hybridized in 6x SSPE (60 mM NaH<sub>2</sub>PO<sub>4</sub>, 0.9 M NaCl, 6 mM EDTA) containing 5x Denhardt's and 0.5% (w/v) SDS at 55°C for 14-18hrs with 10 pmol of 5'-labeled oligonuceotides (tRNA<sup>Ile</sup>: 5'-TGCTCCCGCGGGTTCGAA-3'; internal

standard: 5'-GCCGAGTATTGCTCCGAAG-3') using  $\gamma^{32}\text{P}$ -ATP and the polynucleotide kinase forward reaction. The Northern blots were analyzed and quantified using a Phosphorimager. The signal corresponding to the tRNA<sup>Le</sup> was normalized to the signal of the internal standard. For each quantified signal the mean and standard errors of four to six independent replicates were calculated and plotted on the graphs. Samples for SDS-PAGE and subsequent immunoblot analysis were collected after the digitonin fractionation. Immunoblots were probed for LDH and cox4. For the stalling of the ATOM complex by the import intermediate, formed by LDH-DHFR-HA, the cells were grown in the presence of 250  $\mu\text{M}$  aminopterin, and induced with tetracycline for 24-51 hours prior to mitochondrial RNA isolation.

### Mass spectrometric analysis

LC-MS analysis of mitochondria-enriched fractions generated from tetracycline-treated and untreated SILAC-labeled ATOM40-, ATOM11- or ATOM12-RNAi cells ( $n=3$ ) was performed as previously described (57). Briefly, proteins were acetone-precipitated and separated by SDS-PAGE followed by colloidal Coomassie staining. Gel lanes were cut and proteins were in-gel digested using trypsin (37°C, overnight) after reduction and alkylation of cysteine residues. Peptide samples were dried in vacuo and resuspended in 0.1% TFA for LC-MS analysis on an Orbitrap Elite instrument (Thermo Fisher Scientific, Bremen, Germany) connected to an UltiMate 3000 RSLCnano HPLC system (Thermo Fisher Scientific, Dreieich, Germany). Full MS spectra ( $m/z$  370-1,700) were acquired in the orbitrap at a resolution of 120,000 (at  $m/z$  400). For acquisition of MS/MS spectra, low-energy collision-induced dissociation of isolated multiply charged peptides was performed in the linear ion trap. Data were analyzed using MaxQuant (58) (version 1.5.5.1) and its integrated search engine Andromeda (59). MS/MS data were searched against all entries for *T. brucei* TREU927 (TriTryp database, version 8.1; 11,067), to which the 18 mitochondrially encoded proteins (<http://dna.kdna.ucla.edu/trypanosome/seqs/index.html>) were added, using the following parameters: tryptic specificity (maximum of two missed cleavages; Trypsin/P); mass tolerances, 4.5 ppm for precursor and 0.5 Da for fragment ions; fixed modification, carbamidomethylation of cysteine; variable modifications, N-terminal acetylation and methionine oxidation; heavy labels, Arg10 and Lys8. Proteins were identified based on at least one unique peptide (minimum length, 7 aa) and a false discovery rate (FDR) of 0.01 applied to both peptide and protein level. Proteins were



quantified based on unique peptides and with a minimum ratio count of one. The options 're-quantify' and 'match between runs' were enabled.

### **Immunofluorescence**

$1 \times 10^7$  of aminopterin-treated cells that do or do not express LDH-DHFR-HA were incubated with 250 nM MitoTracker Red CM-H<sub>2</sub>-XRos (Molecular Probes) for 20min at 27°C. As negative control, uninduced cells were incubated with 100  $\mu$ M CCCP for 20min to dissipate the membrane potential. Subsequently the cells were fixed in 1x PBS/4% (w/V) paraformaldehyde for 10min and permeabilized with 1xPBS/0.2% Triton-X-100 for 10min and immunofluorescence was done as described (60). Slides were mounted using Vectashield containing 1.5  $\mu$ g/ml DAPI (Vector Laboratories). The images were obtained from a Leica DMI6000 B microscope.

### **Recombinant protein purification and reconstitution**

Recombinant (r) ATOM40 was expressed in *E. coli* strain BL21 (DE3) and purified from inclusion bodies using immobilized metal ion affinity chromatography in combination with Ni<sup>2+</sup> at denaturing conditions (8M urea) and an additional anion exchange chromatography step as described (49). Affinity purified rATOM40 was solubilized in 2% (w/v) SDS and reconstitution was carried out as described (49).

### **Single channel recordings from planar lipid bilayers**

Planar lipid bilayer measurements of reconstituted rATOM40 were essentially performed as described previously (49, 61). Briefly, single channel currents were measured in standard buffer conditions (250 mM KCl, 10 mM Mops/Tris pH 7) in the absence and presence of 150  $\mu$ M tRNA (bulk tRNA from bakers yeast, Roche) on both sides of the bilayer using a GeneClamp 500 amplifier and the CV-5-1G headstage (Axon Instruments). Filtering was conducted with the inbuilt four-pole Bessel low pass filter (5kHz). Data were recorded at a sampling rate of 50 kHz using a DigiData 1200 (Axon Instruments) and Clampex 9 software.

**Acknowledgments**

M. Niemann and A. Harsman gratefully acknowledge fellowships from the Peter und Traudl Engelhorn foundation. Research in the lab of A. Schneider was supported by grant 138355 and in part by the NCCR "RNA & Disease" both funded by the Swiss National Science Foundation. We thank Elke Horn and Kurt Lobenwein for technical assistance. Research in the Warscheid group was funded by the Deutsche Forschungsgemeinschaft, the Excellence Initiative of the German Federal & State Governments (EXC 294 BIOSS Centre for Biological Signalling Studies) and the European Research Council (ERC, Consolidator Grant Number 648235).

## References

1. Archibald JM (2015) Endosymbiosis and Eukaryotic Cell Evolution. *Curr. Biol.* 25(19):R911-921.
2. Williams TA, Foster PG, Cox CJ, & Embley TM (2013) An archaeal origin of eukaryotes supports only two primary domains of life. *Nature* 504(7479):231-236.
3. Koonin EV & Yutin N (2014) The dispersed archaeal eukaryome and the complex archaeal ancestor of eukaryotes. *Cold Spring Harb. Perspect. Biol.* 6(4):a016188.
4. Gray MW, Burger G, & Lang BF (1999) Mitochondrial Evolution. *Science* 283:1476-1481.
5. Allen JF (2003) Why chloroplasts and mitochondria contain genomes. *Comp. Funct. Genomics* 4(1):31-36.
6. Schulz C, Schendzielorz A, & Rehling P (2015) Unlocking the presequence import pathway. *Trends Cell Biol.* 25(5):265-275.
7. Chacinska A, Koehler CM, Milenkovic D, Lithgow T, & Pfanner N (2009) Importing mitochondrial proteins: machineries and mechanisms. *Cell* 138:628-644.
8. Schmidt O, Pfanner N, & Meisinger C (2010) Mitochondrial protein import: from proteomics to functional mechanisms. *Nat. Rev. Mol. Cell Biol.* 11:655-667.
9. Marom M, Azem A, & Mokranjac D (2011) Understanding the molecular mechanism of protein translocation across the mitochondrial inner membrane: still a long way to go. *Biochim. Biophys. Acta* 1808(3):990-1001.
10. Salinas T, Duchêne AM, & Maréchal-Drouard L (2008) Recent advances in tRNA mitochondrial import. *Trends Biochem. Sci.* 33:320-329.
11. Alfonzo JD & Söll D (2009) Mitochondrial tRNA import--the challenge to understand has just begun. *Biol. Chem.*:717-722.
12. Schneider A (2011) Mitochondrial tRNA Import and Its Consequences for Mitochondrial Translation. *Ann. Rev. Biochem.* 80:1033-1053.
13. Rubio MA, *et al.* (2008) Mammalian mitochondria have the innate ability to import tRNAs by a mechanism distinct from protein import. *Proc. Natl. Acad. Sci. USA* (105):9186-9191.
14. Mercer TR, *et al.* (2011) The human mitochondrial transcriptome. *Cell* 146:645-658.
15. Smirnov A, *et al.* (2010) Mitochondrial enzyme rhodanese is essential for 5 S ribosomal RNA import into human mitochondria. *J. Biol. Chem.* 285:30792-30803.
16. Smirnov A, *et al.* (2008) Two distinct structural elements of 5S rRNA are needed for its import into human mitochondria. *RNA* 14:749-759.
17. Entelis NS, Kolesnikova OA, Dogan S, Martin RP, & Tarassov IA (2001) 5 S rRNA and tRNA import into human mitochondria. *J. Biol. Chem.* 276:45642-45653.
18. Wang G, Shimada E, Koehler CM, & Teitell MA (2012) PNPASE and RNA trafficking into mitochondria. *Biochim. Biophys. Acta* 1819(9-10):998-1007.

19. von Ameln S, *et al.* (2012) A mutation in PNPT1, encoding mitochondrial-RNA-import protein PNPase, causes hereditary hearing loss. *Am. J. Hum. Genet.* 91(5):919-927.
20. Wang G, *et al.* (2010) PNPASE regulates RNA import into mitochondria. *Cell* 142:456-467.
21. Amunts A, Brown A, Toots J, Scheres SH, & Ramakrishnan V (2015) Ribosome. The structure of the human mitochondrial ribosome. *Science* 348(6230):95-98.
22. Greber BJ, *et al.* (2015) Ribosome. The complete structure of the 55S mammalian mitochondrial ribosome. *Science* 348(6232):303-308.
23. Holzmann J, *et al.* (2008) RNase P without RNA: identification and functional reconstitution of the human mitochondrial tRNA processing enzyme. *Cell* 135:462-474.
24. Tarassov I, Entelis N, & Martin RP (1995) An intact protein translocating machinery is required for mitochondrial import of a yeast cytoplasmic tRNA. *J. Mol. Biol.* 245:315-323.
25. Tarassov I, Entelis N, & Martin RP (1995) Mitochondrial import of a cytoplasmic lysine-tRNA in yeast is mediated by cooperation of cytoplasmic and mitochondrial lysyl-tRNA synthetases. *EMBO J.* 14:3461-3471.
26. Kolesnikova OA, *et al.* (2000) Suppression of mutations in mitochondrial DNA by tRNAs imported from the cytoplasm. *Science*.
27. Kapushoc ST, Alfonzo JD, Rubio MAT, & Simpson L (2000) End processing precedes mitochondrial importation and editing of tRNAs in *Leishmania tarentolae*. *J. Biol. Chem.* 275:37907-37914.
28. Simpson AM, Suyama Y, Dewes H, Campbell DA, & Simpson L (1989) Kinetoplastid mitochondria contain functional tRNAs which are encoded in nuclear DNA and also contain small minicircle and maxicircle transcripts of unknown function. *Nucleic Acids Res.* 17:5427-5445.
29. Hancock K & Hajduk SL (1990) The mitochondrial tRNAs of *Trypanosoma brucei* are nuclear encoded. *J. Biol. Chem.* 265:19208-19215.
30. Tan THP, Pach R, Crausaz A, Ivens A, & Schneider A (2002) tRNAs in *Trypanosoma brucei*: genomic organization, expression and mitochondrial import. *Mol. Cell. Biol.* 22:3707-3717.
31. Mani J, Meisinger C, & Schneider A (2016) Peeping at TOMs-Diverse Entry Gates to Mitochondria Provide Insights into the Evolution of Eukaryotes. *Mol. Biol. Evol.* 33:337-351
32. Mani J, *et al.* (2015) Novel mitochondrial protein import receptors in Kinetoplastids reveal convergent evolution over large phylogenetic distances. *Nat. Commun.* 6:6646.
33. Pusnik M, *et al.* (2011) Mitochondrial preprotein translocase of trypanosomatids has a bacterial origin. *Curr. Biol.* 21:1738-1743.
34. Desy S, Mani J, Harsman A, Käser S, & Schneider A (2016) TbLOK1/ATOM19 is a novel subunit of the non-canonical mitochondrial outer membrane protein translocase of *Trypanosoma brucei*. *Mol. Microbiol.* 10.1111/mmi.13476.
35. Crausaz-Esseiva A, Marechal-Drouard L, Cosset A, & Schneider A (2004) The T-stem determines the cytosolic or mitochondrial localization of trypanosomal methionyl-tRNAs. *Mol. Biol. Cell.* 15:2750-2757.

36. Hauser R & Schneider A (1995) tRNAs are imported into mitochondria of *Trypanosoma brucei* independent of their genomic context and of their genetic origin. *EMBO J.* 14:4212-4220.
37. Lima BD & Simpson L (1996) Sequence-dependent in vivo import of tRNAs into the mitochondrion of *Leishmania tarentolae*. *RNA* 2:429-440.
38. Sherrer RL, Yermovsky-Kammerer AE, & Hajduk SL (2003) A sequence motif within trypanosome precursor tRNAs influences abundance and mitochondrial localization. *Mol. Cell. Biol.* 23:9061-9072.
39. Rubio MA, Liu X, Yuzawa H, Alfonzo JD, & Simpson L (2000) Selective importation of RNA into isolated mitochondria from *Leishmania tarentolae* RNA. *RNA* 6:988-1003.
40. Nabholz CE, Horn EK, & Schneider A (1999) tRNAs and proteins are imported into mitochondria of *Trypanosoma brucei* by two distinct mechanisms. *Mol. Biol. Cell.* 10:2547-2557.
41. Kaneko T, *et al.* (2003) Wobble modification differences and subcellular localization of tRNAs in *Leishmania tarentolae*: implication for tRNA sorting mechanism. *EMBO J.* 22:657-667.
42. Bouzaidi-Tiali N, Aeby E, Charrière F, Pusnik M, & Schneider A (2007) Elongation factor 1a mediates the specificity of mitochondrial tRNA import in *T. brucei*. *EMBO J.* 26:4302-4312.
43. Cristodero M, *et al.* (2013) Mitochondrial translation factors of *Trypanosoma brucei*: elongation factor-Tu has a unique subdomain that is essential for its function. *Mol. Microbiol.* 90:744-755.
44. Niemann M, *et al.* (2013) Mitochondrial outer membrane proteome of *Trypanosoma brucei* reveals novel factors required to maintain mitochondrial morphology. *Mol. Cell. Proteomics* 12:515-528.
45. Mani J, Rout S, Desy S, & Schneider A (2016) Mitochondrial protein import - Functional analysis of the highly diverged Tom22 orthologue of *Trypanosoma brucei*. *Sci. Rep.* 7:40738.
46. Zarsky V, Tachezy J, & Dolezal P (2012) Tom40 is likely common to all mitochondria. *Curr. Biol.* 22:R479-R481.
47. Pusnik M, *et al.* (2012) Response to Zarsky *et al.* *Curr. Biol.* 22:R481-R482.
48. Harsman A, *et al.* (2016) The non-canonical mitochondrial inner membrane presequence translocase of trypanosomatids contains two essential rhomboid-like proteins. *Nat. Commun.* 19:13707.
49. Harsman A, *et al.* (2012) Bacterial origin of a mitochondrial outer membrane protein translocase: New perspectives from comparative single channel electrophysiology. *J. Biol. Chem.* 287:31437-31445.
50. Salinas T, *et al.* (2006) The voltage-dependent anion channel, a major component of the tRNA import machinery in plant mitochondria. *Proc. Natl. Acad. Sci. USA* 103:18362-18367.
51. Salinas T, *et al.* (2014) Molecular basis for the differential interaction of plant mitochondrial VDAC proteins with tRNAs. *Nucleic acids research* 42(15):9937-9948.
52. Goswami S, *et al.* (2006) A bifunctional tRNA import receptor from *Leishmania* mitochondria. *Proc. Natl. Acad. Sci. USA* 103:8354-8359.
53. Schekman R (2010) Editorial Expression of Concern: A bifunctional tRNA import receptor from *Leishmania* mitochondria. *Proc. Natl. Acad. Sci. USA* 107:9476.

54. Tschopp F, Charrière F, & Schneider A (2011) In vivo study in *Trypanosoma brucei* links mitochondrial transfer RNA import to mitochondrial protein import. *EMBO Rep.* 12:825-832.
55. McCulloch R, Vassella E, Burton P, Boshart M, & Barry JD (2004) Transformation of monomorphic and pleomorphic *Trypanosoma brucei*. *Methods Mol. Biol.* 262:53-86.
56. Chomczyinski P & Sacchi N (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 162:156-159.
57. Peikert CD, *et al.* (2017) Charting Organellar Importomes by Quantitative Mass Spectrometry. *Nat. Commun.* 8:15272.
58. Cox J & Mann M (2008) MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.* 26:1367–1372.
59. Cox J, *et al.* (2011) Andromeda: A Peptide Search Engine Integrated into the MaxQuant Environment. *J. Proteome Res.* 10:1794-1805.
60. Schneider A, *et al.* (1987) Subpellicular and flagellar microtubules of *Trypanosoma brucei brucei* contain the same alpha tubulin isotypes. *J. Cell Biol.* 104:431-438.
61. Bartsch P, Harsman A, & Wagner R (2013) Single channel analysis of membrane proteins in artificial bilayer membranes. *Methods Mol. Biol.* 1033:345-361.

## Figure Legends

### Figure 1. In vivo import of tRNAs depends on ATOM core but not on ATOM receptor subunits

(A) The sketch at the top shows a model of the ATOM complex and denotes, which of the ATOM core subunits were knocked down (dashed red line). The table summarizes the previously established phenotypes of the indicated cell lines regarding growth and mitochondrial protein import (32) (downward arrow, reduced) (see Fig. S2). All experiments are based on tetracycline (Tet)-inducible RNAi cell lines, except for ATOM11 in procyclic cells for which a conditional knockout cell line was used. This cell line allows depletion of ATOM11 in the absence of Tet. The ethidium bromide-stained (EthBr) gels below the table show mitochondrial RNA extracts separated on 8 M urea/10% polyacrylamide gels that were isolated from the indicated uninduced and induced cell lines. The positions of the 12S and 9S mitochondrial rRNAs and the tRNA region are indicated. The two panels below show the corresponding Northern blots probed either for tRNA<sup>Ile</sup> or the short synthetic RNA molecule (stand.) that was used as an internal standard for the yield of the RNA preparation. The panel below the Northern blots shows an immunoblot of the same mitochondrial fractions probed for LDH. It serves as a quality control for the intactness and yield of the mitochondria present in the digitonin-extracted pellet fractions. The graph at the bottom shows the quantification of the abundance of the tRNA<sup>Ile</sup> normalized to the levels of the short synthetic RNA molecule that was used to control for the yield of the RNA extraction. The mean of four to six replicate experiments (as indicated) is shown. Bar, average deviation. (B) As in (A), but RNAi cell lines for the ATOM receptor subunits ATOM46 and ATOM69 were tested. Ablation of ATOM46 did not impair growth at 27°C. However a growth arrest was observed at 33°C which is why the experiment was performed at 33°C.

### Figure 2. OM proteins reduced in abundance in ATOM subunit-lacking cells. (A)

Abundance changes in OM proteins of procyclic *T. brucei* induced by conditional ablation of ATOM40 (A), ATOM11 (B) and ATOM12 (C), respectively, were analyzed by SILAC-MS. Time of induction is indicated by the arrows in sup. Fig. S2. The graph shows all OM proteins quantified in at least two of the three replicates ( $P$  value < 0.05) that were down-regulated more than 2-fold in ablated cells (standard deviations are indicated). ID, indicates how many of the 82 mitochondrial OM proteins (44) were identified in the corresponding experiments. Grey columns, target of ablation; red columns, ATOM

subunits; white columns, other proteins. POMP, protein of the OM proteome; MICU1, mitochondrial calcium uptake 1. Data for ATOM40 are from (57).

**Figure 3. tRNA import requires the intermembrane space domain of ATOM14.** As in Fig. 1 but tRNA import was compared between an induced ATOM14 3'UTR-RNAi cell line (right panel) and the same cell line complemented with either full length ATOM14, or a ATOM14 variant lacking the IMS domain (ATOM14-ΔIMS). It has previously been shown that the IMS domain of ATOM14 is required for growth and mitochondrial protein import as shown in Fig. S5 (45).

**Figure 4. Obstruction of the ATOM40 pore abolishes tRNA import.** The sketch at the top shows how in cells treated with aminopterin (AMT) expression of the LDH-DHFR-HA fusion protein generates an import intermediate that is stuck in the ATOM40 protein import channel (48). The ethidium bromide-stained (EthBr) gels below the sketch show mitochondrial RNA extracts isolated from cells grown in the presence of aminopterin that either do (+tet) or do not express the LDH-DHFR-HA fusion protein (-tet) for the time indicated. The two panels below show the corresponding Northern blots probed either for tRNA<sup>Ile</sup> or the short synthetic RNA molecule (stand.) that was used as an internal standard. Below the Northern blots immunoblots of the same mitochondrial fractions probed for LDH are shown. The lowest panel depicts an immunoblot probed for the imported protein cytochrome oxidase subunit 4 (Cox4). The positions of unprocessed Cox4 (pre), indicative of the non-imported protein, and mature Cox4, corresponding to the imported protein are indicated. The graph at the bottom shows the mean of the abundance of the tRNA<sup>Ile</sup> normalized to the levels of the short synthetic RNA molecule in six replicates of either uninduced cells (-tet) or cells induced (+tet) for the indicated times. Bar, average deviation.

**Figure 5. tRNA influences conductance of the recombinant ATOM40 pore as measured by single channel electrophysiology experiments.** (A) Recombinant ATOM40 (rATOM40) was purified from *E. coli* to near homogeneity under denaturing conditions (WC, whole cell lysate). (B) Upper panel, rATOM40 was reconstituted into a planar lipid bilayer and displays its typical electrophysiological characteristics (1.5 nS, few gating events) (49). Lower panel, addition of tRNA to both sides of the chamber induced frequent but temporary closure of the rATOM40 channel.








Fig. 1

**Fig. 2**

**Fig. 3**





Fig. 4

**Fig. 5**