1 **First Results of the Swiss National Surgical Site Infection Surveillance**

2 **Program: Who Seeks Shall Find**

3 Nicolas Troillet, MD, MSc;^{1,2,3} Emin Aghayev, MD;⁴ Marie-Christine Eisenring, CNS, ICP;^{1,2} Andreas F.
4 Widmer, MD, MSc;^{1,5} Swissnoso

5
6 ¹Swissnoso-National Center for Infection Control, Bern, Switzerland

7 ²Service of Infectious Diseases, Central Institute, Valais Hospital, Sion, Switzerland

8 ³Services of Preventive Medicine and Infectious Diseases, Centre Hospitalier Universitaire Vaudois and
9 University of Lausanne, Switzerland;

10 ⁴Swiss RDL-Medical Registries and Data Linkage, Institute for Social and Preventive Medicine,
11 University of Bern, Switzerland

12 ⁵Division of Infectious Diseases and Hospital Epidemiology, University Hospital of Basel, Switzerland

13
14 Address correspondence: Nicolas Troillet, Central Institute, Valais Hospital, Ave. Grand-Champsec 86,
15 Sion, CH 1950, Switzerland (nicolas.troillet@hopitalvs.ch).

16
17 Financial support: No financial support was provided relevant to this article.

18
19 Potential conflicts of interest: All authors report no conflicts of interest relevant to this article.
20

21 **4 Tables and 1 Figure**

22 **Table 1:** CDC/NHSN Criteria Used for the Diagnosis of Superficial Incisional SSI in Colon Surgery

23 **Table 2:** Number of Operations and Hospitals Included in the Surveillance, Timing of Antibiotic
24 Prophylaxis, and Surgical Technique, by Type of Operation

25 **Table 3 A, B, C:** Adjusted Risk and Protective Factors for SSI After herniorraphy and C-Section, Including
26 the Effect of the Time From the Initiation of Surveillance to the Operation (Time to Operation)

27 **Table 4:** Adjusted Risk and Protective Factors for SSI After herniorraphy and C-Section, Including the
28 Effect of the Time From the Initiation of Surveillance to the Operation (Time to Operation)

29 **Figure 1:** Crude rates of surgical site infections (SSI) after herniorrhaphy and C-section according to
30 the time from the initiation of surveillance to the date of the operation, by surgical procedure.

31

ABSTRACT

Objectives: To report on the results of the Swiss national surgical site infection (SSI) surveillance program, including temporal trends, and to describe methodological characteristics that may influence SSI rates.

Design: Countrywide survey of SSI over a 4-year period. Analysis of prospectively collected data including patient and procedure characteristics as well as aggregated SSI rates stratified by risk categories, type of SSI, and time of diagnosis. Temporal trends were analyzed using stepwise multivariate logistic regression models with adjustment of the effect of the duration of participation in the surveillance program for confounding factors.

Setting: The study included 164 Swiss public and private hospitals with surgical activities.

Results: From October 2011 to September 2015, a total of 187,501 operations performed in this setting were included. Cumulative SSI rates varied from 0.9% for knee arthroplasty to 14.4% for colon surgery. Postdischarge follow-up was completed in >90% of patients at 1 month for surgeries without an implant and in >80% of patients at 12 months for surgeries with an implant. High rates of SSIs were detected postdischarge, from 20.7% in colon surgeries to 93.3% in knee arthroplasties. Overall, the impact of the duration of surveillance was significantly and independently associated with a decrease in SSI rates in herniorraphies and C-sections but not for the other procedures. Nevertheless, some hospitals observed significant decreases in their rates for various procedures.

Conclusions: Intensive post-discharge surveillance may explain high SSI rates and cause artificial differences between programs. Surveillance per se, without structured and mandatory quality improvement efforts, may not produce the expected decrease in SSI rates.

INTRODUCTION

Surgical site infection (SSI) is the most frequent nosocomial infection in surgical patients. Depending on the type of surgical procedure, SSIs develop in <1% to >20% of patients after operation, and SSIs account for 38% of all nosocomial infections in this population.¹⁻⁴ SSIs prolong hospital stay and increase costs, morbidity, and mortality.⁵⁻⁸ Secondary to the landmark study by Haley et al,⁹ surveillance is considered an essential tool for the prevention of SSI.^{5,9,10} Many national programs have been set up for this purpose during the past 40 years.

In Switzerland, a first multicenter surveillance system for SSIs was developed in the mid-1990s and progressively included hospitals from the southwestern part of the country until 2010.^{4,11,12} In 2009, the Swiss National Association for the Development of Quality in Hospitals and Clinics (ANQ) asked Swissnoso, the National Center for Infection Control, to implement a countrywide surveillance system for SSIs. In 2011, SSI surveillance became mandatory for Swiss hospitals with departments of surgery, and the 2 programs merged. This article reports on the results of this national surveillance system and describes its method, which includes continued on-site quality check of data generated by individual hospitals and high rates of follow-up.

METHODS

Description of the Surveillance Program

The Swissnoso SSI surveillance system was developed according to the principles of the US National Nosocomial Infections Surveillance (NNIS) system, currently known as the National Healthcare Safety Network (NHSN).^{13,14} It is described in a practical guide written in the 3 national languages (German, French, and Italian), which is available for participating hospitals on the Swissnoso website (www.swissnoso.ch).

Infection control nurses (ICNs), supervised by infectious diseases specialists or other physicians without a hierarchical link within the departments of surgery, orthopedics, or gynecology are in charge of the surveillance in each participating hospital. Surgeons do not take part actively in the process of

documenting SSIs. All ICNs and supervising physicians must attend a 1-day special training course before surveillance begins. SSIs are diagnosed according to the Centers for Diseases Control and Prevention (CDC) definitions and are classified as superficial incisional, deep incisional, or organ-space infections.¹⁵ Patients are followed-up during their hospital stay by ICNs through systematic reviews of patient charts, including medical and nursing notes, antibiotic use, microbiology and other laboratory or radiology results, and reports regarding operations, consultations, and discharges. The postdischarge follow-up is performed by ICNs through standardized phone interviews with each patient 1 month after the operation. A second interview takes place 12 months after surgeries with implants such as arthroplasties and cardiac surgeries. The follow-up survey comprises 6 questions regarding unplanned medical visits, rehospitalization, antibiotic prescription, and clinical symptoms of infection. At least 5 telephone attempts must be documented before a patient can be considered lost to follow-up. Any suspected or unclear case triggers further contacts with the family or hospital physician to gather any available additional information. Any suspicion of SSI or unclear situation is presented to the supervising physician for decision according to the CDC criteria. As an example, Table 1 shows the criteria used for superficial SSIs secondary to colon surgery. Approximately every other year, 1-day onsite audits by Swissnoso supervisors take place in each participating hospital to evaluate the quality of the surveillance. These audits include comprehensive reviews of 15 randomly selected operations that have been included in the surveillance.

Hospitals are free to choose at least 3 among 15 surgical procedures. However, those doing colon surgery must include this procedure. All patients undergoing the included procedures during given surveillance periods must be registered and followed-up. The variables collected include characteristics of the patients (sex, age, American Society of Anesthesiologists [ASA] score, and delay from admission to operation) and characteristics of the operations: contamination class, duration of the operation, planned or unplanned (ie, emergency) procedure, type and timing of antibiotic prophylaxis, minimally invasive or laparoscopic operation, multiple procedures during the surveyed operation, reoperation for noninfectious complications. Data are entered online in a university-

owned, webbased database (www.ispm.unibe.ch), and each hospital has direct individual access to its results. Various data validation rules automatically apply during data entry. Inventory reports are available for every hospital, signaling cases to be completed or cases with possible errors. A hotline, staffed by Swissnos collaborators, is available 5 of 7 days per week.

Detailed reports are edited yearly for every participating hospital, allowing crude and adjusted comparisons with others, using the NNIS/NHSN risk index.¹⁴ The implementation or reinforcement of preventive measures is left to each hospital's discretion. Participants are invited to an annual meeting where the results are presented and discussed. Since 2014, starting with the 2011 data, the Swiss National Association for the Development of Quality in Hospitals and Clinics (ANQ) has openly published the surveillance results by hospital (<http://www.anq.ch/messengergebnisse/ergebnisse-akutsomatik/>), including their names, NNIS/NHSN adjusted SSI rates, and quality of surveillance as rated during onsite visits.

Statistical Analyses

Analyses were performed on data collected from October 2011 to September 2015. Characteristics of the patients and operations, 4-year cumulative SSI rates (3-year for surgeries with implants), proportions of postdischarge diagnoses, and proportions of superficial incisional, deep incisional, and organ-space infections were calculated by type of surgical procedure.

Because not all participating hospitals began the surveillance simultaneously, temporal trends in SSI rates were calculated for each surgical procedure taking the duration of participation in the surveillance program into account rather than calendar years. This duration was determined for each hospital and was stratified in 1-year periods, as described in previous studies.^{4,16}

Forward stepwise logistic regression models were developed for each surgical procedure that had been included for >2 years in the surveillance program to identify independent risk or protective factors for SSI, including the duration of participation in the program. The following covariates were entered in the initial models: time from the initiation of the surveillance to the surveyed operation,

patient sex, age, ASA score, delay from admission to operation, contamination class, duration of the operation, emergency procedure, antibiotic prophylaxis <1 h before incision, laparoscope use for digestive surgery, multiple procedures during the operation, and reoperation within the follow-up period for non-infectious complications. In surgical procedures where the time from the initiation of the surveillance to the surveyed operation was not initially retained as a significant covariate in the final model, it was added to obtain notwithstanding the effect of this variable independently of changes in patient characteristics over time. In addition, individual hospitals with significantly increasing or decreasing SSI rates over these time periods were identified using the Cochran-Armitage test.¹⁷

Analyses were performed using SAS version 9.4 software (SAS Institute, Cary, NC). All tests were 1-tailed, and α was set at 0.05 throughout the study.

RESULTS

Overall, 187,501 operations in 164 hospitals had been included in the surveillance system and could be analyzed (Table 2). Patient and operation characteristics, including age, sex, NNIS/NHSN risk index, contamination of the surgical wound, ASA score, and duration of the operation appeared similar to those reported in other large populations (Online Supplemental Table 1). Hysterectomies and laminectomies were included only in 2015 in the surveillance, explaining the smaller numbers of operations. Prophylactic antibiotics were often administered >1 hour before incision or after incision, showing room for improvement. In C-sections, antibiotics were still given late in 42.6% of the patients, after umbilical cord clamping.

The aggregated rates of SSIs are shown in Table 3 by operation, globally, and stratified by NNIS/NHSN risk index category, by surgical technique (laparoscopic vs open) for digestive surgery, and by type of infection. Also shown in Table 3, 90% of the operations in digestive surgery were followed-up for 1 month and >80% of surgeries with orthopedic implants and cardiac surgeries with foreign-material implants were followed-up for 1 year. Many SSIs occurred after discharge; these rates varied from

20.7% in colon surgery to 93.3% in knee arthroplasty. Taking all operations into account, 3,292 of 6,953 SSIs (47.4%) were detected after discharge: 1,599 (48.6%) were superficial incisional, 521 (15.8%) were deep incisional, and 1,172 (35.6%) were organ-space infections. Rehospitalization occurred in 455 of 1,599 (29.4%) patients with postdischarge superficial incisional SSIs, in 412 of 521 (79.1%) patients with postdischarge deep incisional SSIs, and in 1,129 of 1,172 (96.3%) patients with postdischarge organ-space infections. Mean and median times to the diagnosis of SSI varied respectively from 9.8 to 17.6 days and 9 to 16 days for surgeries without implant, and from 29.9 to 83.2 and 17 to 34 days for surgeries with implant (Online Supplemental Table 2). A significant decrease was observed in operation-specific SSI rates of 27 hospitals or surgical departments over the observed time period, whereas a significant increase was observed in 11 hospitals (Online Supplemental Table 2). The effect of the time from the start of surveillance to the operation on SSI rates, as estimated by logistic regression models, is shown in Table 4 for herniorrhaphy and C-section, the 2 operations for which it remained an independent protective factor after adjustment for other factors. This effect is shown for all the types of operations in Online Supplemental Table 3. The trends of SSI rates according to this time are shown in Figure 1 for herniorrhaphy and C-section, and SSI trends are shown in Online Supplemental Figure 1 for each operation that had been included in the surveillance for >2 years.

DISCUSSION

Some SSI rates reported by our surveillance system may appear high in comparison with those reported in France, England, Germany, and the European Union and in the United States by the CDC's National Healthcare Safety Network (NHSN).^{18–22} These rates are quite similar to those reported in the Netherlands by the PREZIES system and those reported in the United States by the American College of Surgeons National Surgical Quality Improvement Program (ACS NSQIP).^{23,24} Indeed, the Swissnoso system shares several characteristics with these 2 programs that distinguish their approach from other SSI surveillance systems and might contribute to the finding of higher SSI rates.

First, the Swissnoso program puts particular efforts in postdischarge surveillance (PDS) and reaches complete follow-up in the majority of included operations (from a minimum 82.8% in cardiac surgeries to 97% in rectal surgeries). Indeed, PDS is an important component of SSI surveillance, particularly in the current setting where hospital stays tend to be shorter and surgical procedures are more frequently performed in an ambulatory setting.^{5,25–27} As in other systems that also perform active PDS,^{24,26} high rates of SSI were detected postdischarge in our program, varying from 20.7% in colon surgeries to >90% in arthroplasties. In addition, 51.4% of the SSIs detected through PDS were severe (deep incisional or organ-space infections) and some of them, if not rehospitalized in the same hospital where the first operation took place, would not have been recognized without active PDS.

However, no consensus exists on a method for PDS, and methods may differ between systems. We chose an indirect active method based on standardized phone interviews with the patients by trained ICNs, completed by a secondary contact with his/her general practitioner (GP) in case of a suspected SSI or an unclear situation. Written or oral questionnaires administered to patients or surgeons have low sensitivity and specificity for detecting SSIs.^{28,29} Nonetheless, Whitby et al²⁹ found that the patient's recall of prescription of an antibiotic by their GP for SSI during the postoperative period correlated very well with diagnoses made by experienced ICNs, particularly when confirmed by the GP in those patients reporting an infection. Our PDS is based on both a questionnaire administered to the patients by an ICN and a subsequent contact with their GPs, and we believe that it is a reliable and necessary tool for getting the correct figures to feed back to hospitals and surgical teams. Other systems based on electronic algorithms and/or administrative data may be less resource consuming, but they still need development to be used with heterogeneous information systems.³⁰ Automated telephony could also help decrease the workload of IC nurses.³¹

Second, since public reporting of SSI rates by hospitals became mandatory in Switzerland, important efforts were made to periodically audit hospitals to minimize risks of underreporting due to various possible biases that have been reported elsewhere and recently motivated specific recommendations from the US Healthcare Infection Control Practices Advisory Committee.³² Indeed, studies in Scotland,

the Netherlands, Australia, and New York state found 0.6% to 4.3% of SSIs among cases initially not reported as such.^{33–36} Our audits revealed differences between hospitals, particularly in the access to the necessary clinical information for ICNs to detect cases. They found 1.4% of false negatives, and 0.09% of false positives (Kuster et al., Structure, Process and Outcome Quality of Surgical Site Infection Surveillance in Switzerland. Submitted. April 2017). Thus, to create incentives to perform better, Swissnoso gives marks to hospitals for the quality of their surveillance that are openly published together with their infection rates. Moreover, the same SSI case definitions may allow variable interpretations between persons and countries and could also result in artificial differences in SSI rates.³⁷

The main goal of SSI surveillance is to decrease SSI rates by providing hospitals, surgical teams, and stakeholders with data that can be used for benchmarking, monitoring, and, if deemed necessary, implementation of better preventive measures. Reaching this goal depends on factors that are external to the surveillance system and linked to hospitals or surgical departments themselves. Indeed, some individual hospitals or departments participating in our surveillance system experienced decreasing SSI rates in some operations over a 4-year observation period and others did not. But a general, statistically significant and independent protective effect on SSI of the duration of participation in the Swissnoso system was seen only for herniorrhaphy and C-section, whereas no significant trends were found for other surgeries. Such observations have already been made in other surveillance programs where SSI rates tend to decrease over time for some operations but not for others or even do increase.^{4,16,38,39} Various requirements have been listed for a surveillance system to succeed.⁴⁰ They may be implemented at different levels of adherence among hospitals, within the same hospital, or among particular surgical teams. Overall, our results showed that the timing of antibiotic prophylaxis should be improved in most surgeries. Secondary to the initiation of surveillance, Swissnoso launched a new program aiming at introducing standardized process measures in hospitals for monitoring and improving some preventive measure for SSI such as hair removal, skin disinfection, and the dosing and timing of antibiotics.

In conclusion, a high-quality nationwide SSI surveillance program was implemented in Switzerland in 2011. Its first results show high SSI rates as compared with other programs, but these differences may be, at least in part, artificial and due to methodological differences, particularly with respect to an effective PDS system that detected up to 47% of all SSIs, but this system is resource consuming and could benefit from future developments in information technology. In addition, regular on-site audits guarantee the quality of the collected data. To date, the ultimate goal of SSI surveillance (a decrease in SSI rates) has only been reached in a subset of patients and hospitals, but temporal trends are difficult to predict over a relatively short period of time. A longer observation period and additional efforts by hospitals may thus be necessary to reach this goal through projects based on a culture of safety such as those developed in the United States by the Comprehensive Unit-based Safety Program (CUSP) of the Armstrong Institute for Patient Safety and Quality.

ACKNOWLEDGMENTS

Members of Swissnoso: Carlo Balmelli, Marie-Christine Eisenring, Stephan Harbarth, Stefan Kuster, Jonas Marschall, Virginie Masserey Spicher, Didier Pittet, Christian Ruef, Hugo Sax, Matthias Schlegel, Alexander Schweiger, Nicolas Troillet, Andreas Widmer, and Giorgio Zanetti.

REFERENCES

1. Sax H, Uçkay I, Balmelli C, et al. Overall burden of healthcare associated infections among surgical patients: results of a national study. *Ann Surg* 2011;253:365–370.
2. Smyth ET, McIlvenny G, Enstone JE, et al. Four country healthcare associated infection prevalence survey 2006: overview of the results. *J Hosp Infect* 2008;69:230–248.
3. Edwards JR, Peterson KD, Mu Y, et al. National healthcare safety network (NHSN) report: data summary for 2006 through 2008, issued December 2009. *Am J Infect Control* 2009;37:783–805.
4. Staszewicz W, Eisenring MC, Bettschart V, Harbarth S, Troillet N. Thirteen years of surgical site infection surveillance in Swiss hospitals. *J Hosp Infect* 2014;88:40–47.
5. Anderson DJ, Podgorny K, Berrios-Torres I, et al. Strategies to prevent surgical site infections in acute care hospitals: 2014 update. *Infect Control Hosp Epidemiol* 2014;35:605–627.
6. Broex EC, van Asselt AD, Bruggeman CA, van Tiel FH. Surgical site infections: how high are the costs? *J Hosp Infect* 2009;72: 193–201.
7. Weber WP, Zwahlen M, Reck S, et al. Economic burden of surgical site infections at a European university hospital. *Infect Control Hosp Epidemiol* 2008;29:623–629.

- 266 8. Astagneau P, Rioux C, Golliot F, Brückner G, INCISO Network Study Group. Morbidity and mortality
267 associated with surgical site infections: results from the 1997–1999 INCISO surveillance. *J Hosp*
268 *Infect* 2001;48:267–274.
- 269 9. Haley RW, Culver DH, White JW, et al. The efficacy of infection surveillance and control programs
270 in preventing nosocomial infections in US hospitals. *Am J Epidemiol* 1985;121:182–205.
- 271 10. Surgical site infections. 2008. clinical guideline 74. National Institute for Clinical Excellence
272 website. <http://www.nice.org.uk/CG74>. Published 2008. Accessed December 9, 2016.
- 273 11. Romy S, Eisenring MC, Bettschart V, Petignat C, Francioli P, Troillet N. Laparoscope use and
274 surgical site infections in digestive surgery. *Ann Surg* 2008;247:627–632.
- 275 12. Hubner M, Diana M, Zanetti G, Eisenring MC, Demartines N, Troillet N. Surgical site infections in
276 colon surgery: the patient, the procedure, the hospital, and the surgeon. *Arch Surg*
277 2011;146:1240–1245.
- 278 13. Emori TG, Culver DH, Horan TC, et al. National nosocomial infections surveillance system (NNIS):
279 description of surveillance methods. *Am J Infect Control* 1991;19:19–35.
- 280 14. Culver DH, Horan TC, Gaynes RP, et al. Surgical wound infection rates by wound class, operative
281 procedure, and patient risk index. National Nosocomial Infections Surveillance System. *Am J Med*
282 1991;91(Suppl B):152S–157S.
- 283 15. Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated
284 infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control*
285 2008;36:309–332.
- 286 16. Mannien J, van der Hof S, Muilwijk J, van der Broek PJ, van Benthem B, Wille JC. Trends in the
287 incidence of surgical site infection in the Netherlands. *Infect Control Hosp Epidemiol*
288 2008;29:1132–1138.
- 289 17. Armitage P. Tests for linear trends in proportions and frequencies. *Biometrics* 1955;11:375–386.
- 290 18. Edwards JR, Peterson KD, Mu Y, et al. National Healthcare Safety Network (NHSN) report: data
291 summary for 2006 through 2008, issued December 2009. *Am J Infect Control* 2009;37:783–805.
- 292 19. Surveillance des infections du site opératoire, France 2013. Institut de Veille sanitaire website.
293 [http://www.invs.sante.fr/fr/Publications-et-outils/Rapports-et-syntheses/Maladies-infectieuses/](http://www.invs.sante.fr/fr/Publications-et-outils/Rapports-et-syntheses/Maladies-infectieuses/2015/Surveillance-des-infections-du-site-operatoire-France-2013)
294 [2015/Surveillance-des-infections-du-site-operatoire-France-2013](http://www.invs.sante.fr/fr/Publications-et-outils/Rapports-et-syntheses/Maladies-infectieuses/2015/Surveillance-des-infections-du-site-operatoire-France-2013). Published 2014. Accessed
295 December 9, 2016.
- 296 20. Surveillance of surgical site infections in NHS hospitals in England, 2014–15. Public Health
297 England website.
298 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/484874/Surv](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/484874/Surveillance_of_Surgical_Site_Infections_in_NHS_Hospitals_in_England_report_2014-15.pdf)
299 [eillance_of_Surgical_Site_Infections_in_NHS_Hospitals_in_England_report_2014-15.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/484874/Surveillance_of_Surgical_Site_Infections_in_NHS_Hospitals_in_England_report_2014-15.pdf).
300 Published 2015. Accessed December 9, 2016.
- 301 21. Surveillance of surgical site infections in Europe 2010–2011. European Centre for Diseases
302 prevention and Control website. [http://ecdc.europa.eu/en/healthtopics/healthcare-](http://ecdc.europa.eu/en/healthtopics/healthcare-associated-infections/surgical-site-infections/pages/ssi.aspx)
303 [associated_infections/surgical-site-infections/pages/ssi.aspx](http://ecdc.europa.eu/en/healthtopics/healthcare-associated-infections/surgical-site-infections/pages/ssi.aspx). Published 2013. Accessed
304 December 9, 2016.
- 305 22. National and state healthcare-associated infections. Progress report. Centers for Disease Control
306 and Prevention website. [http://www.cdc.gov/HAI/pdfs/progress-report/hai-progress-](http://www.cdc.gov/HAI/pdfs/progress-report/hai-progress-report.pdf)
307 [report.pdf](http://www.cdc.gov/HAI/pdfs/progress-report/hai-progress-report.pdf). Published 2016. Accessed December 9, 2016.
- 308 23. Mannien J, van den Hof S, Brandt C, Behnke M, Wille JC, Gastmeier P. Comparison of the national
309 surgical site infection surveillance data between the Netherlands and Germany: PREZIES versus
310 KISS. *J Hosp Infect* 2007;66:224–231.

24. Ju MH, Ko CY, Hall BL, Bosk CL, Bilmoria KY, Wick EC. A comparison of 2 surgical site infection monitoring systems. *JAMA Surg* 2015;150:51–57.
25. Huotari K, Lyytikäinen O. Impact of post-discharge surveillance on the rate of surgical site infection after orthopedic surgery. *Infect Control Hosp Epidemiol* 2006;27:1324–1329.
26. Manniën J, Wille JC, Snoeren RLMM, van den Hof S. Impact of postdischarge surveillance on surgical site infection rates for several surgical procedures: results from the nosocomial surveillance network in the Netherlands. *Infect Control Hosp Epidemiol* 2006;27:809–816.
27. Reilly J, Allardice G, Bruce J, Hill R, McCoubrey J. Procedurespecific surgical site infection rates and post-discharge surveillance in Scotland. *Infect Control Hosp Epidemiol* 2006;27:1318–1323.
28. Sands K, Vineyard G, Platt R. Surgical site infections occurring after hospital discharge. *J Infect Dis* 1996;173:963–970.
29. Whitby M, McLaws ML, Collopy B, et al. Post-discharge surveillance: can patients reliably diagnose surgical wound infections? *J Hosp Infect* 2002;52:155–160.
30. Gastmeier P, Behnke M. Electronic surveillance and using administrative data to identify healthcare associated infections. *Curr Opin Infect Dis* 2016;29:394–399.
31. McNeish J, Lyle D, McCowan M, et al. Post-discharge surgical site infection surveillance by automated telephony. *J Hosp Infection* 2007;66:232–236.
32. Talbot TR, Bratzler DW, Carrico RM, et al. Public reporting of health care-associated surveillance data: recommendations from the healthcare infection control practices advisory committee. *Ann Intern Med* 2013;159:631–635.
33. Mc Coubrey J, Reilly J, Mullings A, Pollock KG, Johnston F. Validation of surgical site infection surveillance data in Scotland. *J Hosp Infect* 2005;61:194–200.
34. Manniën J, van der Zeeuw AE, Wille JC, van der Hof S. Validation of surgical site infection surveillance in the Netherlands. *Infect Control Hosp Epidemiol* 2007;28:36–41.
35. Friedman ND, Russo PL, Bull AL, Richards MJ, Kelly H. Validation of coronary artery bypass graft surgical site infection surveillance data from a statewide surveillance system in Australia. *Infect Control Hosp Epidemiol* 2007;28:812–817.
36. Haley VB, van Antwerown C, Tserenpuntsag B, et al. Use of administrative data in efficient auditing of hospital-acquired surgical site infections, New York state 2009–2010. *Infect Control Hosp Epidemiol* 2012;33:565–571.
37. Birgand G, Lepelletier D, Baron G, et al. Agreement among healthcare professionals in ten European countries in diagnosing case-vignettes of surgical site infections. *PLoS One* 2013;8: e68618.
38. Brandt C, Sohr D, Behnke M, Daschner F, Rüden H, Gastmeier P. Reduction of surgical site infection rates associated with active surveillance. *Infect Control Hosp Epidemiol* 2006;27:1347–1351.
39. Astagneau P, L’Hériveau F, Daniel F, et al. Reducing surgical site infection through a network: results from the French ISO-RAISIN surveillance system. *J Hosp Infect* 2009;72: 127–134.
40. Gastmeier P, Sohr G, Schwab F, et al. Ten years of KISS: the most important requirements for success. *J Hosp Infection* 2008;70(Suppl 1):11–16.

352 **TABLES**353 **Table 1**

354 CDC/NHSN Criteria Used for the Diagnosis of Superficial Incisional SSI in Colon Surgery

355 *Note:* CDC, Centers for Disease Control and Prevention; NHSN, National Health Safety Network; SSI, surgical site infection.356 ^aMore than one criteria can be used for the same infection.

357

Criteria ^a	Frequency, No. (%) (N = 1,124)
Purulent drainage from the superficial incision	641 (57.0)
Organisms isolated from an aseptically obtained culture of fluid or tissue from the superficial incision	423 (37.6)
At least 1 of the following signs or symptoms of infection: pain or tenderness, localized swelling, redness, or heat, and superficial incision is deliberately opened by surgeon and is culture positive or not cultured	776 (69.0)
Diagnosis of superficial incisional SSI by the surgeon or attending physician	957 (85.1)

358

359

Table 2Number of Operations and Hospitals Included in the Surveillance, Timing of Antibiotic Prophylaxis, and Surgical Technique, by Type of Operation^a*Note:* CABG, coronary arterial bypass.^aTotal percentage may not equal 100 due to missing values or rounding

Operation	No. of Participating Hospitals	No. of Operations Included	Antibiotics (%)			Laparoscopy or Minimal Invasive Technique, %
			>1 h Before Incision	≤1 h Before Incision	After Incision	
Appendectomy	94	15,439	17.6	65.8	8.8	88.7
Cholecystectomy	61	20,402	14.1	63.4	5.4	91.3
Herniorraphy	60	17,030	5.1	79.0	2.8	33.8
Gastric bypass	15	3,077	15.7	81.7	1.5	93.7
Colon surgery	116	22,889	28.3	65.0	4.5	42.1
Rectum surgery	25	1,835	49.5	47.1	2.4	51.6
C-section	63	32,814	6.0	43.4	42.6	...
Abdominal hysterectomy	9	841	7.6	80.3	3.2	65.2
Vaginal hysterectomy	10	452	7.4	80.3	3.5	...
Laminectomy w/o implant	10	2,050	11.3	85.0	2.1	4.7
Laminectomy with implant	4	430	35.3	60.5	2.8	2.1
Cardiac surgery, overall	15	11,974	24.1	71.7	2.8	1.6
CABG, no other cardiac surgery	14	5,541	25.5	71.1	2.2	2.9
Hip arthroplasty	120	32,102	11.8	86.0	1.3	7.0
Knee arthroplasty	86	20,625	14.5	82.8	1.7	...

366 **Table 3 A, B, C**

367 Adjusted Risk and Protective Factors for SSI After herniorrhaphy and C-Section, Including the Effect of the Time From the Initiation of Surveillance to the
 368 Operation (Time to Operation)

369 *Note:* SSI, surgical site infection; CI, confidence interval; NNIS, National Nosocomial Infection Surveillance; NHSN, National Healthcare

370 Security Network.

A	Appendectomy	Cholecystectomy	Herniorrhaphy	Gastric Bypass	Colon Surgery	Rectal Surgery
No. of SSIs/total	609/15,439	448/20,402	169/17,030	135/3,077	3,304/22,889	212/1,835
SSI rate, % (95% CI)	3.9 (3.6–4.2)	2.2 (2.0–2.4)	1.0 (0.9–1.2)	4.4 (3.7–5.2)	14.4 (14.0–14.9)	11.6 (10.1–13.1)
SSI rate by NNIS/NHSN index category, % (95% CI)						
NNIS/NHSN 0	0.8 (0.4–1.5)	1.2 (1.0–1.4)	0.6 (0.5–0.8)	4.5 (3.6–5.7)	8.6 (7.9–9.4)	5.7 (3.0–9.5)
NNIS/NHSN 1	2.9 (2.6–3.3)	2.1 (1.8–2.5)	1.8 (1.4–2.3)	4.0 (3.1–5.2)	12.4 (11.8–13.1)	9.4 (7.5–11.6)
NNIS/NHSN 2	5.4 (4.9–6.1)	4.5 (3.8–5.3)	6.9 (4.3–10.4)	5.0 (1.9–10.6)	19.7 (18.7–20.8)	13.7 (11.1–16.7)
NNIS/NHSN 3	10.5 (7.9–13.7)	8.4 (6.4–10.7)	13.0 (4.7–33.6)	25.0 (3.2–65.1)	27.0 (24.8–29.2)	24.0 (17.4–31.7)
SSI rate by surgical technique, % (95% CI)						
Laparoscopy	3.5 (3.2–3.9)	1.6 (1.5–1.8)	0.5 (0.3–0.7)	4.2 (3.5–5.0)	8.9 (8.4–9.5)	7.3 (5.7–9.1)
Open surgery	7.2 (6.1–8.6)	8.0 (6.8–9.3)	1.3 (1.1–1.5)	6.7 (3.6–11.2)	18.5 (17.8–19.1)	16.1 (13.8–18.7)
Type of SSI, N (%)						
Superficial incisional	200 (33.0)	236 (52.7)	119 (70.4)	67 (50.0)	1124 (34.0)	58 (27.4)
Deep incisional	59 (9.7)	49 (10.9)	38 (22.5)	11 (8.1)	460 (13.9)	31 (14.6)
Organ-space	350 (57.4)	163 (36.4)	12 (7.1)	57 (41.9)	1720 (52.1)	123 (58.0)
Completed follow-up (%)	92.1	92.4	93.5	91.9	92.1	97.1
Postdischarge SSI (%)	71.8	63.0	84.6	66.7	20.8	22.2

371

B	C-Section	Hysterectomy Abdominal	Hysterectomy Vaginal	Laminectomy Without Implant	Laminectomy With Implant
No. of SSIs/total	511/32,814	25/841	11/452	30/2,050	22/430
SSI rate, % (95% CI)	1.6 (1.4–1.7)	3.0 (1.9–4.4)	2.4 (1.2–4.3)	1.5 (1.0–2.1)	5.1 (3.0–7.6)
SSI rate by NNIS/NHSN index category, % (95% CI)					
NNIS/NHSN 0	1.2 (1.1–1.4)	1.1 (0.4–2.5)	1.9 (0.7–4.1)	0.9 (0.4–1.6)	3.4 (0.7–9.7)
NNIS/NHSN 1	2.1 (1.9–2.5)	3.8 (2.0–6.5)	3.4 (0.9–8.5)	2.2 (1.3–3.6)	4.4 (2.3–7.5)
NNIS/NHSN 2	2.5 (1.8–3.4)	11.3 (4.7–21.9)	5.9 (0.1–28.7)	2.3 (0.6–5.7)	10.5 (4.3–20.4)
NNIS/NHSN 3	4.3 (1.4–9.7)	12.5 (0.3–52.5)	0 (0–97.5)	0 (0–45.9)	0 (0–97.5)
Type of SSI, N (%)					
Superficial incisional	340 (66.5)	11 (44.0)	2 (18.2)	9 (30.0)	4 (18.2)
Deep incisional	50 (9.8)	4 (16.0)	2 (18.2)	11 (36.7)	5 (22.7)
Organ-space	121 (23.7)	10 (40.0)	7 (63.6)	10 (33.3)	13 (59.1)
Completed follow-up (%)	91.0	93.3	96.2	96.9	88.7
Postdischarge SSI (%)	86.5	72.0	72.7	90.0	68.2

C	Cardiac Surgery Overall	CABG Without Other Cardiac Surgery	Hip Arthroplasty	Knee Arthroplasty
No. of SSIs/total	566/11,974	281/5,541	436/32,102	194/20,625
SSI rate, % (95% CI)	4.7 (4.4–5.1)	5.1 (4.5–5.7)	1.4 (1.2–1.5)	0.9 (0.8–1.1)
SSI rate by NNIS/NHSN index category, % (95% CI)				
NNIS/NHSN 0	1.8 (0.8–3.5)	2.1 (0.6–5.3)	1.0 (0.9–1.2)	0.8 (0.6–0.9)
NNIS/NHSN 1	4.3 (3.9–4.8)	4.8 (4.2–5.4)	1.8 (1.6–2.1)	1.0 (0.8–1.2)
NNIS/NHSN 2	6.6 (5.6–7.8)	7.2 (5.6–9.1)	3.3 (2.4–4.3)	2.0 (1.3–2.8)
NNIS/NHSN 3	13.2 (8.0–20.1)	6.7 (0.2–32.0)	0 (0–84.2)	0 (0–97.5)
Type of SSI, N (%)				
Superficial incisional	207 (36.6)	104 (37.0)	124 (28.4)	87 (44.8)
Deep incisional	162 (28.6)	90 (32.0)	61 (14.0)	17 (8.8)
Organ-space	197 (34.8)	87 (31.0)	251 (57.6)	90 (46.4)
Completed follow-up (%)	83.0	84.2	91.4	89.8
Postdischarge SSI (%)	61.8	60.1	90.8	93.3

373

374

375 **Table 4**

376 Adjusted Risk and Protective Factors for SSI After herniorrhaphy and C-Section, Including the Effect of the Time From the Initiation of Surveillance to the
 377 Operation (Time to Operation)

378 *Note:* OR, odds ratio; CI, confidence interval; ASA, American Society of Anesthesiologists

Intervention	Variable	OR (95% CI)	P Value
Herniorrhaphy	Time to operation, per year	0.72 (0.62–0.83)	<.001
	Sex, female vs male	1.91 (1.36–2.68)	<.001
	ASA score, per category	1.73 (1.40–2.13)	<.001
	Antibiotics within 1 h, yes vs no	0.48 (0.34–0.67)	<.001
	Wound class, per class	1.57 (1.16–2.12)	.004
	Laparoscopy, yes vs no	0.40 (0.27–0.61)	<.001
	Duration of surgery, per min	1.009 (1.007–1.012)	<.001
C-section	Time to operation, per year	0.92 (0.84–1.00)	.05
	Age, per year	0.98 (0.96–1.00)	.02
	ASA score, per category	1.19 (1.00–1.42)	.05
	Unplanned operation, yes vs no	1.75 (1.41–2.16)	<.001
	Wound class, per class	1.41 (1.17–1.71)	<.001
	Multiple interventions, yes vs no	1.85 (1.26–2.73)	<.001
	Duration of surgery, per min	1.006 (1.002–1.009)	<.001

379

380

381 **FIGURE**382 **Figure 1**

383 Crude rates of surgical site infections (SSI) after herniorrhaphy and C-section according to the time
384 from the initiation of surveillance to the date of the operation, by surgical procedure.

385