

Oral microbiota in Swiss adolescents

Sigrun Eick · Małgorzata Pietkiewicz · Anton Sculean

Received: 30 September 2011 / Accepted: 13 February 2012 / Published online: 28 February 2012
© Springer-Verlag 2012

Abstract

Objectives The purpose of the study was to determine the prevalence of different oral microbes in gingival plaque samples and in samples from the dorsum of the tongue in a Swiss adolescent population.

Materials and methods Ninety-nine adolescents between 15 and 18 years were enrolled. Plaque index, bleeding on probing (BOP), the periodontal screening index, and decayed missed filled tooth (DMFT) index were recorded. Samples from subgingival plaque and swabs from the tongue were analyzed by the Checkerboard DNA-DNA hybridization method. Additionally, counts of *Streptococcus mutans* and *Aggregatibacter actinomycetemcomitans*, *Porphyromonas gingivalis*, *Tannerella forsythia*, and *Treponema denticola* were determined by real-time PCR.

Results Periodontitis was not diagnosed in any of the subjects but all of them presented signs of gingival inflammation displaying a mean BOP of 28%. Ten (10.1%) subjects were tested positive for *P. gingivalis*, each 22 (22.2%) for *A. actinomycetemcomitans* and *T. forsythia*, (47.5%) for *T. denticola*. *T. denticola* and *S. mutans* showed a high affinity to the gingival plaque, whereas *T. forsythia* was often detected from the dorsum of the tongue. DMFT was associated with *S. mutans* counts, and BOP correlated with counts of *P. gingivalis* and *T. denticola*.

Conclusions The present data indicate that: (a) gingivitis but not periodontitis is a common finding among Swiss adolescents, and (b) bacteria associated with periodontitis were frequently detected in the subgingival dental plaque and on the dorsum of the tongue in Swiss adolescents with gingivitis.

Clinical relevance Although gingivitis was a frequent finding in Swiss adolescents, periodontitis was not detected in this population. The dorsum of the tongue appears to represent an important reservoir for periodontopathic bacteria.

Keywords Adolescents · Periodontal health · Microflora · Periodontitis-associated bacteria · *Streptococcus mutans*

Introduction

Initiation and progression of periodontal disease is closely related with the colonization of the microorganisms, including *Aggregatibacter actinomycetemcomitans*, as well as the members of the so called “red complex”: *Porphyromonas gingivalis*, *Tannerella forsythia*, and *Treponema denticola* [1–3]. An inverse relationship between caries and periodontitis has been suggested, underlined by the fact that counts of *P. gingivalis* and *Streptococcus mutans* correlate negatively in saliva [4]. Bacteria associated with periodontitis are detectable already in early childhood and adolescents [5, 6]. Not only plaque seems to be a reservoir for bacteria associated with periodontitis and *S. mutans* but also tongue might be colonized by those species already in the very early age of 6 months [7, 8].

In older children and adolescents, the detection rates of the periodontopathogens vary. Studies in the 1980s reported a rare detection of *P. gingivalis* in a study population in London [9] and a presence of *A. actinomycetemcomitans* in 10% of the subjects in London and Finland [9, 10]. In contrast, detection rates of 100% of those species were

S. Eick · A. Sculean (✉)
Department of Periodontology, Dental School, University of Bern,
Freiburgstrasse 7,
CH-3010 Bern, Switzerland
e-mail: anton.sculean@zmk.unibe.ch

M. Pietkiewicz
School Dental Service,
Bern, Switzerland

described recently in a Greece population [8]. In a previous study, 33 subjects were included in a follow-up study in Switzerland during puberty [11]. In that study, culture techniques have failed to detect *P. gingivalis* through the whole observation period while *A. actinomycetemcomitans* was only rarely detected in some of the subjects developing a puberty gingivitis [11].

The purpose of the present study was to evaluate the prevalence of oral bacteria primarily associated with periodontitis in subgingival plaque and on the tongue dorsum of Swiss adolescents in relation to clinical variables by using nucleic-acid based methods.

Materials and methods

The ethical committee of the Canton of Bern, CH approved the study (KEK-BE: 235/08). Written informed ascent and consent was obtained from each subject and from their parents respectively prior to participation.

Assessment of epidemiological and clinical data

Ninety-nine adolescents aged between 15 and 18 years and involved in the oral health monitoring protocol of a dental service for children and adolescents in Bern (Switzerland) were enrolled in the present study. Subjects were asked about their smoking habits as well as for significant systemic disease (e.g., diabetes mellitus, coronary heart disease). Subjects with antibiotic therapy within the last 6 months and pregnant or lactating females were excluded. Further, an ongoing orthodontic treatment at the time-point of the investigation was an exclusion criterion.

From all participants of the study, decayed missed filled teeth (DMFT) index was recorded. Further, periodontal screening index (PSI [12]) was used to characterize periodontal health. Probing depths were measured with a periodontal probe (PCP-UNC 15, Hu Friedy, Leimen, Germany). Bleeding on probing (BOP) was calculated as the percentage of positive sites per subject based on a measurement of four sites/tooth. Oral hygiene was recorded by using the plaque index described by O'Leary et al. [13].

All clinical recordings were performed by the same calibrated examiner (MP). Examiner calibration was performed as follows: five adolescents, not enrolled in the study, were evaluated by the examiner on two separate occasions, 48 h apart. Calibration was accepted if measurements at baseline and at 48 h were similar to the millimeter at $\geq 90\%$.

Sample collection

For sampling, each of the mesiobuccal site of the first molar was selected. Without removing supragingival plaque [14],

the test site was air dried and kept dry using cotton rolls. Each one paper point (ISO 50) was inserted into the selected gingival sulcus for 20 s to obtain plaque. The samples from the four different sites were pooled in the transport vials. Furthermore, samples from the tongue were taken by swabbing of about 1 cm^2 of the center of the dorsum of the tongue with sterile cotton sticks (Applimed SA, Châtel-St-Denis, Switzerland). Samples were sent to the laboratory, where they were stored at -20°C until analyzed.

Microbiological analysis

Counts of 21 oral species were determined in each sample using a modification of the checkerboard DNA-DNA hybridization technique [15, 16]. DNA was extracted by using Chelex method [17] and then 100 μl of the extract (half of the total extract) were placed in lanes on a nylon membrane using a Minislot device (Immunetics, Cambridge, MA, USA). (Before applying Chelex to checkerboard, the suitability of the method has been tested. This method is easy to make and allows direct comparison of different methods by using the same DNA.) After fixation of the DNA to the membrane, the membrane was placed in a Miniblitter 45 (Immunetics, Cambridge, MA, USA) with the lanes of DNA at 90° to the lanes of the device. Digoxigenin-labeled whole genomic DNA probes to 21 bacterial species were hybridized in individual lanes of the Miniblitter. After hybridization, the membranes were washed at high stringency, and the DNA probes were detected using antibody to digoxigenin conjugated with alkaline phosphatase and chemiluminescence detection. The probes and their source strains were compared as described in a methodological paper by Ximenez-Fyvie et al. [18]. Signals were detected using AttoPhos substrate (Amersham Life Science, Arlington Heights, Illinois, USA) and a Storm Fluorimager (Molecular Dynamics, Sunnyvale, CA, USA). Two lanes in each run contained standards at concentrations of 10^5 and 10^6 cells of each species. The sensitivity of the assay was adjusted to permit detection of 7.5×10^4 cells of a given species by adjusting the concentration of each DNA probe. Signals were evaluated using the Storm Fluorimager and converted to absolute counts by comparison with the standards on the same membrane. Because of the methodology, cross-reactions cannot be completely excluded between closely related taxa [19]. Following, that closely related species were grouped (Table 1) by using the highest signal as result.

Additionally to the Checkerboard technique, real-time PCRs of four bacterial species associated with periodontitis (*A. actinomycetemcomitans*, *P. gingivalis*, *T. forsythia*, and *T. denticola*) and of *S. mutans* as a cariogenic species were performed as described recently [20, 21] modified by using GoTaq[®] qPCR Master mix (Promega Corporation, Madison, WI, USA). The detection level (cut-off) was set to 10^2

Table 1 Grouping of microorganisms used as probes in Checkerboard technique

Group	Species
Oral Streptococci	<i>S. mitis</i> ATCC 49456, <i>S. oralis</i> ATCC 35037, <i>S. gordonii</i> ATCC 10558
<i>Actinomyces</i> sp.	<i>A. odontolyticus</i> ATCC 17929, <i>A. viscosus</i> ATCC 43146/ <i>A. naeslundii</i> ATCC 12104
<i>Neisseria</i> sp.	<i>N. mucosa</i> ATCC 19696
<i>Veillonella</i> sp.	<i>V. parvula</i> ATCC 10790
Black pigmented (Bp) <i>Prevotella</i> sp.	<i>P. intermedia</i> ATCC 25611, <i>P. melaninogenica</i> ATCC 25845
<i>Fusobacteria</i> sp.	<i>F. nucleatum</i> ssp. <i>nucleatum</i> ATCC 25586, <i>F. periodonticum</i> ATCC 33693
<i>Capnocytophaga</i> sp.	<i>C. gingivalis</i> ATCC 33612, <i>C. ochraceae</i> ATCC 33596
<i>Campylobacter</i> sp.	<i>C. rectus</i> ATCC 33238, <i>C. gracilis</i> ATCC 33236
<i>Eikenella corrodens</i>	<i>E. corrodens</i> ATCC 23834

bacteria, although often, lower counts were also measured. In the final analysis, only results of real-time PCR for these species were used. Due to technical problems, the obtained DNA from tongue samples did not cover all PCRs from all subjects. Thus, in part, a few samples are missing in analysis; the exact number of included samples is always given in the results.

Data analysis

All data were entered into the PASW 18.0 (SPSS Inc, Chicago, IL, USA) program, and were analyzed using parameter-free methods. Tongue samples and plaque were compared by Wilcoxon test. Mann–Whitney *U* test was used for comparisons of subgroups within the study population. The level of significance was set to $p<0.05$.

Results

Clinical data

Demographic and clinical data are presented in Table 2. All subjects were in the age range between 15 and 18 years. Seventeen (17.1%) were already smokers. Thirty-three (33.3%) of the adolescents were immigrants (13 from

Table 2 Demographic and clinical data of the study population

Study group	
<i>n</i>	99
Gender (m:f)	52:47
Age (mean±SD) (years)	16.8±1.1
Smoking habits (smoker:non-smoker)	17:82
Plaque index (mean±SD)	19.34%±15.48%
BOP (mean±SD)	27.52%±12.42%
DMFT (mean±SD)	2.21±3.02
PSI (mean±SD)	1.91±0.29

former Yugoslavia, 6 from South Europe, 5 of the other Europe, 6 from Asia and 3 from Africa). The study population consisted of adolescents aged between 15 and 18 years and enrolled in a dental service program. The adolescents were students of regular schools and schools that provided vocational education. Therefore, the age range was relatively small.

None of the study subjects was diagnosed with periodontitis. All subjects were diagnosed with gingivitis; the BOP was 27.52%±12.42%. Only nine (9.1%) adolescents had PSI=1, in 90 (90.9%) additionally calculus and/or defected margins were present (PSI=2). The high mean BOP was associated with a mean plaque index of nearly 20%.

In thirty-eight (38.4%) of the participants (DMFT=0), no tooth with clinical evidence of tooth decay or respective treatment was detected. In three (3%) of the adolescents, a DMFT >10 was found, the highest score was DMFT=15. DMFT differed significantly between natives (1.35±1.56) and immigrants (3.94±4.28; $p=0.003$).

Microorganisms identified in samples from gingival sulci and from tongue samples

Using a detection level of about 7.5×10^4 microbes in Checkerboard technique, the percentages of the positive samples range between 37.4% (black pigmented *Prevotella* sp.) and 82.4% (*Eikenella corrodens*) by analyzing gingival plaque samples, and between 43.3% (*Campylobacter* sp.) and 97.9% (*Actinomyces* sp., oral streptococci, *Veillonella* sp.) by analyzing tongue swabs. Most groups of microbes were detected in higher amounts on tongue than in gingival sulcus. This might depend on the sample size. No difference was found for *Campylobacter* sp. and *Capnocytophaga* sp. suggesting an affinity of those for gingival region (Table 3).

The proportions of subjects defined as carrying the studied species at different detection levels by real-time PCR are presented (Fig. 1). Using real-time PCR as reference and considering the detection level of Checkerboard technique, comparison of both results showed a sufficient sensitivity

Table 3 Detection of different species by Checkerboard technique

Species (group)	Gingival sulcus		Tongue		Difference P	Correlation	
	Positive ($\geq 7.5 \times 10^4$) (%)	High load ($\geq 10^6$) (%)	Positive ($\geq 7.5 \times 10^4$) (%)	High load ($\geq 10^6$) (%)		R	P
<i>Actinomyces</i> sp.	42/99 (42.4)	0/99 (0)	97/99 (98.0)	27/99 (27.3)	<0.001	0.037	0.358
<i>Campylobacter</i> sp.	39/99 (39.4)	0/99 (0)	43/99 (43.4)	0/99 (0)	0.298	0.377	<0.001
<i>Capnocytophaga</i> sp.	74/99 (74.7)	0/99 (0)	75/99 (75.6)	0/99 (0)	0.998	0.162	0.054
<i>E. corrodens</i>	82/99 (82.8)	0/99 (0)	93/99 (93.9)	9/99 (9.1)	<0.001	0.187	0.032
<i>Fusobacterium</i> sp.	64/99 (64.6)	0/99 (0)	85/99 (85.9)	1/99 (1.0)	<0.001	0.207	0.020
<i>Neisseria</i> sp.	76/99 (76.8)	2/99 (2.0)	90/99 (90.9)	17/99 (17.2)	<0.001	0.249	0.006
Bp <i>Prevotella</i> sp.	37/99 (37.4)	0/99 (0)	85/99 (85.9)	27/99 (27.3)	<0.001	0.146	0.075
Oral streptococci	55/99 (55.6)	0/99 (0)	97/99 (98.0)	1/99 (1.0)	<0.001	0.152	0.066
<i>Veillonella</i> sp.	74/99 (74.7)	5/99 (5.1)	97/99 (98.0)	2/99 (2.0)	<0.001	0.039	0.350

(close to 100%) but in part an inadequate specificity for Checkerboard technique (*A. actinomycetemcomitans* 34%). Real-time PCR results showed that 78 (78.8%) of all subjects were positive when tested for *S. mutans*; in 22 subjects, the species was found only in plaque; in nine, only on the tongue; whereas in 47 adolescents, it was detectable both in plaque and tongue swabs. Among the ten subjects being positive for *P. gingivalis*, and the 22 being positive for *A. actinomycetemcomitans*, each about one third is colonized only on the tongue, in another one third, the species is found only in plaque, and in the last third, in both samples. In case of *T. forsythia*, only the tongue samples are mostly positive (14 (16.1% of all subjects)); in four subjects (4.6%), the species was detected in plaque and on tongue; whereas in only one subject, the plaque sample exclusively had *T. forsythia*. Contrary, only in two of the 47 *T. denticola* positive subjects, the species was detectable only on tongue. Except for *T. forsythia*, the counts in gingival plaque correlated with those on tongue (Table 4).

Clinical data and microflora

Clinical data were set into association with the microflora. The microflora did not depend on age and gender of the adolescents. Data analysis failed to demonstrate differences

in bacterial counts and smoking status. The BOP scores were, however, higher in smokers ($p=0.05$).

The counts of *S. mutans* correlated with DMFT index (Table 5, Fig. 2). Subjects with PSI=2 had significantly more Fusobacteria sp. ($p=0.022$) and bp *Prevotella* sp. ($p=0.020$) in their plaque compared with those having PSI=1. The BOP was found to be correlated with *P. gingivalis* and *T. denticola* (Fig. 3), whereas the PI was associated with increasing counts of *Capnocytophaga* sp., *T. denticola*, and *Veillonella* sp. (Table 5). The counts of *S. mutans* were positively correlated with the numbers of *T. forsythia* ($R=0.370$, $p<0.001$).

Discussion

The present study aimed to collect information about the colonization of bacteria associated with periodontitis in subgingival sulci and from the dorsum of the tongue in Swiss adolescents. *A. actinomycetemcomitans* was found by real-time PCR in approximately 20% of the study subjects. Using culture techniques, a similar prevalence of *A. actinomycetemcomitans* was found in patients without periodontitis in USA [22] and in Finland [10]. Lower detection rates were reported in 1–15%

Fig. 1 Detection of *S. mutans*, *A. actinomycetemcomitans*, *P. gingivalis*, *T. forsythia* and *T. denticola* in gingival sulcus samples and swab samples obtained from the dorsum of the tongue




Table 4 Detection of different species by Real-time PCR

Species	Gingival plaque		Tongue		Difference <i>P</i>	<i>R</i>	<i>P</i>
	Positive ($\geq 10^2$) (%)	Moderate–high load ($\geq 10^4$) (%)	Positive ($\geq 10^2$) (%)	Moderate–high load ($\geq 10^4$) (%)			
<i>S. mutans</i>	69/99 (69.7)	12/99 (12.1)	56/97 (57.7)	3/97 (3.1)	0.001	0.401	<0.001
<i>A. actinomycetemcomitans</i>	17/99 (17.2)	2/99 (2.0)	16/96 (16.8)	1/96 (1.0)	0.686	0.466	<0.001
<i>P. gingivalis</i>	6/99 (6.1)	1/99 (1.0)	7/97 (7.2)	1/97 (1.0)	0.877	0.380	<0.001
<i>T. forsythia</i>	8/99 (8.1)	8/99 (8.1)	18/87 (20.7)	4/87 (4.6)	0.909	-0.030	0.777
<i>T. denticola</i>	45/99 (45.5)	4/99 (4.0)	24/85 (28.2)	4/85 (4.7)	0.379	0.458	<0.001

year-old children in Japan [23] using PCR techniques. Remarkable higher rates have been determined in 13-year-old children in Thailand [24] and in less than 3-year-old children in Northern Mariana Islands [7], in a Greek population [8], and in USA [25]. Taken together, these findings appear to suggest geographic or ethnical differences in susceptibility. It should, however, be pointed out that the highest prevalence was found following the use of hybridization assays. On the other hand, our group has found a low specificity for *A. actinomycetemcomitans* when using Checkerboard technique. The method which allows determination of several species at the same time uses the whole genomic DNA of selected species for hybridization. In our laboratory, it was found that cross-reactivity occurs when genetically similar species are present in high numbers (data not shown). As recently demonstrated, the real-time PCRs used in the present study has shown a good correlation with another nucleic-based method [20]. Furthermore, the method was additionally validated by using primers against other regions in the *A. actinomycetemcomitans* genome (leukotoxin, cytolethal toxins), thus confirming the correct identification of that species. In this study, we did not determine the serotypes of *A. actinomycetemcomitans*. In general, serotype b is thought to be associated particularly with periodontal disease [26] especially with aggressive periodontitis

[27], but serotype distribution depends also on geographical regions and ethnicity of the subjects [27, 28].

The presence of *P. gingivalis* in about 10% of the subjects is consistent with other data [6, 23, 29]. It has been shown that in children, the prevalence and load of *P. gingivalis* increases with age [30]. While in children, *P. gingivalis* colonizes only transiently; in adolescents, colonization becomes stable [25]. In our study, nearly half of the subjects were colonized by *T. forsythia* and *T. denticola*, which were the most frequently identified major pathogens.

Cortelli et al. [5] analyzed *T. forsythia*, *P. gingivalis*, and *A. actinomycetemcomitans* within gingival sulci and on tongue in adolescents; prevalence of the species in gingival sulcus was similar to our results, whereas they found an affinity of those species to gingival region which is not confirmed in the present study. The anatomy of the tongue provides favorable conditions for anaerobes. Six months after full-mouth tooth extraction, periodontopathogens were still detectable on tongue [31]. The high prevalence of periodontopathic bacteria (e.g., fusobacteria, *E. corrodens*, and *T. forsythia*) on tongue might not only play a role in the development of periodontal disease. Halitosis is clearly seen in association with tongue coatings containing a high





Fig. 2 Detection of *S. mutans* in gingival sulcus samples in dependence of DMFT index (DMFT 0: $n=38$; DMFT=1–3: $n=36$; DMFT ≥ 4 : $n=23$)

Fig. 3 Detection of *P. gingivalis* and *T. denticola* in gingival sulcus samples in dependence of BOP (BOP≤20: n=33; BOP 21–30: n=30; BOP 31–40: n=21; BOP >40: n=15)



number of anaerobes [32]. The proportion of *T. forsythia* in tongue dorsum samples was higher in patients with malodor than in those without [33]. In this respect, *T. denticola* and *F. nucleatum* have been shown to be significantly associated with indicator measurements of halitosis [34]. The problem of halitosis has been reported already in adolescents [35].

The present study has failed to show any gender-related differences in terms of clinical and microbiological parameters which is in contrast to an earlier report [10]. On the other hand, our data are comparable to those from a recent study, which has found higher BOP values in adolescent smokers [36]. In the present study, however, smoking was not associated with different microflora. One reason for this discrepancy may be related to the young age of the subjects. It is obvious that due to their age, the examined subjects were not able to smoke for a long time. On the other hand, it should be kept in mind that also in adult periodontitis patients, no clear difference in the subgingival microflora between smokers and non-smokers was observed [37, 38].

Very recent epidemiological data suggest that the DMFT is 0.9 in 12-year-old children in Switzerland and among the lowest in Europe [39]. The relatively high mean DMFT measured in this study was, most probably, influenced by the included immigrants. Even in adolescents, the DMFT is higher in immigrants compared to Swiss residents confirming the higher DMFT values in 7-year-old children [40]. The counts of *S. mutans* as an important causative agent for caries correlated positively with DMFT and with the counts of *T. forsythia*. An inverse association of *S. mutans* with clinical indices as well as with other species was never found.

In the present study population, no periodontal pocket depth exceeded 3 mm, but signs of gingival inflammation were visible in each subject. A study by Asikainen et al. [10] has also failed to detect periodontitis in 12–17-year-old residents of Finland. In contrast, Schiffner et al. [41] reported a rate of 13.4% of periodontitis patients in German adolescents. Previous reports have also indicated that gingivitis is a common disease in adolescents [9, 10, 42, 43]. In a study analyzing 624 Swiss army recruits in the age of 18–24 years, at least one site with pathological periodontal

probing depth and a mean rate of 27% on bleeding at gentle sulcular probing were found in 5.1% of the study population [44]. These findings are in accordance with those from the present study, despite the fact that the study was conducted in a slightly younger Swiss population.

Onset and severity of gingivitis in children has been associated with *P. gingivalis* and *A. actinomycetemcomitans* [45]. In a study group of 14–15-year-old subjects with gingivitis, *P. gingivalis*, spirochetes, and black pigmented *Prevotella* sp. but not *A. actinomycetemcomitans* have been detected in high counts [9]. In a Swiss study analyzing 33 subjects over 6 years during puberty, high counts of spirochetes were detected in subjects developing a pubertal gingivitis [11]. In the present study, counts of *P. gingivalis* and *T. denticola* were associated with BOP. Especially, the high prevalence of *T. denticola* in gingival plaque supports the role of spirochetes in the etiology of gingivitis. Although periodontopathic bacteria are an important causative agent for periodontitis, follow-up studies supporting or neglecting a risk in development are extremely rare. This view is supported by findings from a large, long-term (i.e., up to 7 years), longitudinal epidemiological study in 167 untreated Indonesian adolescents where the presence or absence of *A. actinomycetemcomitans* in subgingival plaque was associated with disease development and progression [46, 47].

In conclusion, the present data have provided evidence that gingivitis but not periodontitis is a common clinical finding among Swiss adolescents. Bacteria associated with periodontitis are frequently present in this population on the dorsum of the tongue, representing an important reservoir for those species. Follow-up studies should focus on the prevalence of periodontitis-associated species as a possible predictive marker for development of periodontitis.

Acknowledgments We are grateful to Marianne Weibel and Regula Hirschi for their excellent assistance in performing the microbiological analysis. Professor Dr. Wolfgang Strübig is acknowledged for supporting the adolescents' examinations at the Dental Service Bern. The Department of Periodontology at the University of Bern, Bern, Switzerland funded the study.

Conflicts of interest The authors declare that they have no conflicts of interest.

References

- Holt SC, Ebersole JL (2005) *Porphyromonas gingivalis*, *Treponema denticola*, and *Tannerella forsythia*: the “red complex”, a prototype polybacterial pathogenic consortium in periodontitis. *Periodontol* 2000 38:72–122
- (1996) Consensus report. Periodontal diseases: pathogenesis and microbial factors. *Ann Periodontol* 1:926–32
- Meng S, Zhao L, Yang H, Wu Y, Ouyang Y (2009) Prevalence of *Actinobacillus actinomycetemcomitans* in Chinese chronic periodontitis patients and periodontally healthy adults. *Quintessence Int* 40:53–60
- Iwano Y, Sugano N, Matsumoto K, Nishihara R, Iizuka T, Yoshinuma N, Ito K (2010) Salivary microbial levels in relation to periodontal status and caries development. *J Periodontal Res* 45:165–169
- Cortelli JR, Aquino DR, Cortelli SC, Fernandes CB, de Carvalho-Filho J, Franco GC, Costa FO, Kawai T (2008) Etiological analysis of initial colonization of periodontal pathogens in oral cavity. *J Clin Microbiol* 46:1322–1329
- Kulekci G, Leblebicioglu B, Keskin F, Ciftci S, Badur S (2008) Salivary detection of periodontopathic bacteria in periodontally healthy children. *Anaerobe* 14:49–54
- Tanner AC, Milgrom PM, Kent R Jr, Mokeem SA, Page RC, Riedy CA, Weinstein P, Bruss J (2002) The microbiota of young children from tooth and tongue samples. *J Dent Res* 81:53–57
- Papaioannou W, Gizani S, Haffajee AD, Quirynen M, Mamai-Homata E, Papagiannoulis L (2009) The microbiota on different oral surfaces in healthy children. *Oral Microbiol Immunol* 24:183–189
- Ashley FP, Gallagher J, Wilson RF (1988) The occurrence of *Actinobacillus actinomycetemcomitans*, *Bacteroides gingivalis*, *Bacteroides intermedius* and spirochaetes in the subgingival microflora of adolescents and their relationship with the amount of supragingival plaque and gingivitis. *Oral Microbiol Immunol* 3:77–82
- Asikainen S, Alaluusua S, Kari K, Kleemola-Kujala E (1986) Subgingival microflora and periodontal conditions in healthy teenagers. *J Periodontol* 57:505–509
- Mombelli A, Rutar A, Lang NP (1995) Correlation of the periodontal status 6 years after puberty with clinical and microbiological conditions during puberty. *J Clin Periodontol* 22:300–305
- Cutress TW, Hunter PB, Beck DJ, de Souza P (1978) A comparison of WHO periodontal status index with the periodontal and oral hygiene indices. *Community Dent Oral Epidemiol* 6:245–252
- O’Leary TJ, Drake RB, Naylor JE (1972) The plaque control record. *J Periodontol* 43:38
- Beikler T, Schnitzer S, Abdeen G, Ehmke B, Eisenacher M, Flemmig TF (2006) Sampling strategy for intraoral detection of periodontal pathogens before and following periodontal therapy. *J Periodontol* 77:1323–1332
- Socransky SS, Smith C, Martin L, Paster BJ, Dewhirst FE, Levin AE (1994) “Checkerboard” DNA-DNA hybridization. *Biotechniques* 17:788–792
- Haffajee AD, Cugini MA, Dibart S, Smith C, Kent RL Jr, Socransky SS (1997) The effect of SRP on the clinical and microbiological parameters of periodontal diseases. *J Clin Periodontol* 24:324–334
- Yang JL, Wang MS, Cheng AC, Pan KC, Li CF, Deng SX (2008) A simple and rapid method for extracting bacterial DNA from intestinal microflora for ERIC-PCR detection. *World J Gastroenterol* 14:2872–2876
- Ximenez-Fyvie LA, Haffajee AD, Socransky SS (2000) Microbial composition of supra- and subgingival plaque in subjects with adult periodontitis. *J Clin Periodontol* 27:722–732
- Socransky SS, Haffajee AD, Smith C, Martin L, Haffajee JA, Uzel NG, Goodson JM (2004) Use of checkerboard DNA-DNA hybridization to study complex microbial ecosystems. *Oral Microbiol Immunol* 19:352–362
- Eick S, Straube A, Guentsch A, Pfister W, Jentsch H (2011) Comparison of real-time polymerase chain reaction and DNA-strip technology in microbiological evaluation of periodontitis treatment. *Diagn Microbiol Infect Dis* 69:12–20
- Yoshida A, Suzuki N, Nakano Y, Kawada M, Oho T, Koga T (2003) Development of a 5' nuclease-based real-time PCR assay for quantitative detection of cariogenic dental pathogens *Streptococcus mutans* and *Streptococcus sobrinus*. *J Clin Microbiol* 41:4438–4441
- Slots J, Reynolds HS, Genco RJ (1980) *Actinobacillus actinomycetemcomitans* in human periodontal disease: a cross-sectional microbiological investigation. *Infect Immun* 29:1013–1020
- Umeda M, Miwa Z, Takeuchi Y, Ishizuka M, Huang Y, Noguchi K, Tanaka M, Takagi Y, Ishikawa I (2004) The distribution of periodontopathic bacteria among Japanese children and their parents. *J Periodontal Res* 39:398–404
- Dahlen G, Konradsson K, Eriksson S, Teanpaisan R, Piwat S, Carlen A (2010) A microbiological study in relation to the presence of caries and calculus. *Acta Odontol Scand* 68:199–206
- Lamell CW, Griffen AL, McClellan DL, Leys EJ (2000) Acquisition and colonization stability of *Actinobacillus actinomycetemcomitans* and *Porphyromonas gingivalis* in children. *J Clin Microbiol* 38:1196–1199
- Socransky SS, Haffajee AD, Smith C, Dibart S (1991) Relation of counts of microbial species to clinical status at the sampled site. *J Clin Periodontol* 18:766–775
- Jentsch H, Cachovan G, Guentsch A, Eickholz P, Pfister W, Eick S (2012) Characterization of *Aggregatibacter actinomycetemcomitans* strains in periodontitis patients in Germany. *Clin Oral Investig*. 2012 Jan 14. [Epub ahead of print]
- Kim TS, Frank P, Eickholz P, Eick S, Kim CK (2009) Serotypes of *Aggregatibacter actinomycetemcomitans* in patients with different ethnic backgrounds. *J Periodontol* 80:2020–2027
- Sirinian G, Shimizu T, Sugar C, Slots J, Chen C (2002) Periodontopathic bacteria in young healthy subjects of different ethnic backgrounds in Los Angeles. *J Periodontol* 73:283–288
- Bimstein E, Sapir S, Houri-Haddad Y, Dibart S, Van Dyke TE, Shapira L (2004) The relationship between *Porphyromonas gingivalis* infection and local and systemic factors in children. *J Periodontol* 75:1371–1376
- Van Assche N, Van Essche M, Pauwels M, Teughels W, Quirynen M (2009) Do periodontopathogens disappear after full-mouth tooth extraction? *J Clin Periodontol* 36:1043–1047
- Quirynen M, Avontroodt P, Soers C, Zhao H, Pauwels M, van Steenberghe D (2004) Impact of tongue cleansers on microbial load and taste. *J Clin Periodontol* 31:506–510
- Tanaka M, Yamamoto Y, Kuboniwa M, Nonaka A, Nishida N, Maeda K, Kataoka K, Nagata H, Shizukuishi S (2004) Contribution of periodontal pathogens on tongue dorsa analyzed with real-time PCR to oral malodor. *Microbes Infect* 6:1078–1083
- Yasukawa T, Ohmori M, Sato S (2010) The relationship between physiologic halitosis and periodontopathic bacteria of the tongue and gingival sulcus. *Odontology* 98:44–51
- Kanli A, Kanbur NO, Dural S, Derman O (2008) Effects of oral health behaviors and socioeconomic factors on a group of Turkish adolescents. *Quintessence Int* 39:e26–e32
- Hermann P, Gera I, Borbely J, Fejery P, Madlena M (2009) Periodontal health of an adult population in Hungary: findings of a national survey. *J Clin Periodontol* 36:449–457

37. Apatzidou DA, Riggio MP, Kinane DF (2005) Impact of smoking on the clinical, microbiological and immunological parameters of adult patients with periodontitis. *J Clin Periodontol* 32:973–983
38. Van der Velden U, Varoufaki A, Hutter JW, Xu L, Timmerman MF, Van Winkelhoff AJ, Loos BG (2003) Effect of smoking and periodontal treatment on the subgingival microflora. *J Clin Periodontol* 30:603–610
39. Malmö, University (2011) <http://www.mah.se/CAPP/Country-Oral-Health-Profiles/EURO/#5>
40. Marthaler TM (2004) Changes in dental caries 1953–2003. *Caries Res* 38:173–181
41. Schiffner U, Hoffmann T, Kerschbaum T, Micheelis W (2009) Oral health in German children, adolescents, adults and senior citizens in 2005. *Community Dent Health* 26:18–22
42. Campus G, Cagetti MG, Senna A, Spano G, Benedicenti S, Sacco G (2009) Differences in oral health among Italian adolescents related to the type of secondary school attended. *Oral Health Prev Dent* 7:323–330
43. Ericsson JS, Abrahamsson KH, Ostberg AL, Hellstrom MK, Jonsson K, Wennstrom JL (2009) Periodontal health status in Swedish adolescents: an epidemiological, cross-sectional study. *Swed Dent J* 33:131–139
44. Rothlisberger B, Kuonen P, Salvi GE, Gerber J, Pjetursson BE, Attstrom R, Joss A, Lang NP (2007) Periodontal conditions in Swiss army recruits: a comparative study between the years 1985, 1996 and 2006. *J Clin Periodontol* 34:860–866
45. Morinushi T, Lopatin DE, Van Poperin N, Ueda Y (2000) The relationship between gingivitis and colonization by *Porphyromonas gingivalis* and *Actinobacillus actinomycetemcomitans* in children. *J Periodontol* 71:403–409
46. Timmerman MF, Van der Weijden GA, Arief EM, Armand S, Abbas F, Winkel EG, Van Winkelhoff AJ, Van der Velden U (2001) Untreated periodontal disease in Indonesian adolescents. Subgingival microbiota in relation to experienced progression of periodontitis. *J Clin Periodontol* 28:617–627
47. Timmerman MF, Van der Weijden GA, Abbas F, Arief EM, Armand S, Winkel EG, Van Winkelhoff AJ, Van der Velden U (2000) Untreated periodontal disease in Indonesian adolescents. Longitudinal clinical data and prospective clinical and microbiological risk assessment. *J Clin Periodontol* 27:932–942