Activation of proton translocation by respiratory Complex I

Nikolai P. Belevich, Christoph von Ballmoos, and Marina Verkhovskaya

Biochemistry, **Just Accepted Manuscript** • DOI: 10.1021/acs.biochem.7b00727 • Publication Date (Web): 29 Sep 2017

Downloaded from <http://pubs.acs.org> on October 1, 2017

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

ACS Publications

Biochemistry is published by the American Chemical Society, 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Activation of proton translocation by respiratory Complex I

Nikolai Belevich¹, Christoph von Ballmoos², Marina Verkhovskaya^{3*}

^{1,3} Institute of Biotechnology, PO Box 65 (Viikinkaari 1) FIN-00014 University of Helsinki,
Finland

* Author for correspondence: e-mail: Marina.Verkhovskaya@Helsinki.Fi

² Department of Chemistry and Biochemistry, University of Bern, Freiestrasse 3, CH-3012
Bern,
Switzerland

Abstract

Activation of proton pumping by reconstituted and native membrane-bound Complex I was studied using optical electric potential- and pH-sensitive probes. We find, that reconstituted Complex I has a delay in proton translocation, which is significantly longer than the delay in quinone reductase activity, indicating an initially decoupled state of Complex I. Studies of the amount of NADH required for the activation of pumping indicate the prerequisite of multiple turnovers. Proton pumping by Complex I was also activated by NADPH, excluding significant reduction of Complex I or pre-existing $\Delta\psi$ as activation factors. Co-reconstitution of Complex I and ATPase did not indicate an increased membrane permeability for protons in the uncoupled Complex I state. The delay in Complex I proton pumping activation was also observed in subbacterial vesicles. While it is negligible at room temperature, it strongly increases at a lower temperature. We conclude that Complex I undergoes a conversion from decoupled to coupled state upon the activation. The possible origins and importance of the observed phenomenon are discussed.

Introduction

NADH:ubiquinone oxidoreductase (Complex I) is the electron input enzyme in the respiratory chain of mitochondria and many bacterial species, which couples electron transfer from NADH to ubiquinone with transmembrane proton translocation. The electron transfer proceeds along the intraprotein redox chain located in the hydrophilic domain and proton translocation is performed by antiporter-like membrane subunits. No redox centers are found in the membrane fragment

where the antiporter-like subunits are disposed in a row so that the energy of the redox reaction must be transduced over a distance of 100 Å. Such high spatial separation of electron and proton transfers with no obvious coupling site indicates an intricate molecular mechanism of proton pumping. Elucidation of this mechanism could be significantly advanced by monitoring the enzyme operation upon turnover in real time, which prerequisites the enzyme in its active state. However, oxidized mitochondrial Complex I exists in a de-active state (D) which spontaneously converts to an active state (A) upon turnover¹⁻³, after which the oxidized enzyme slowly relaxes to the D-state. The D/A transition is believed to be physiologically relevant since it could serve as a protective mechanism preventing the oxidative damage of tissues (see for a review⁴). In contrast to mitochondrial complex I, the D/A transition was not observed in bacterial Complex I from *Paracoccus denitrificans*⁵, *Thermus thermophilus* and *Escherichia coli*⁶ and it was suggested that the D/A transition might be a unique property of the former^{7, 8}. Subunit composition and molecular structures of Complex I from various organisms demonstrate that bacterial enzymes are “minimal versions” of the mitochondrial ones, i.e. they exert the exact same molecular function with similar rates and stoichiometry⁷⁻¹⁰. Bacterial Complex I is thus a good model for studying the coupling mechanism between NADH oxidation and proton translocation, although it might not share all properties, such as D/A transition. Upon this transition, mitochondrial Complex I is proposed to undergo conformational changes occurring within three subunits at the junction of the hydrophilic and membrane fragments of the enzyme^{11, 12}. Although the D/A transition has been proposed to be a unique property of the mitochondrial enzyme^{7, 8}, dynamics and flexibility of the region at the hydrophilic/membrane domain¹⁰ introduces some uncertainty in modeling of

the conformational changes involved and interconversion of these states in bacterial enzymes cannot be excluded and must be studied experimentally.

We have recently shown that *E. coli* Complex I may also exist in two states, resting (R) and active (A), characterized by a delayed maximal NADH:ubiquinone oxidoreductase activity¹³. In a subsequent study using fluorescent probes, we obtained results indicating that upon the activation from state R to state A, Complex I undergoes conformational changes, which can be mapped to the junction of the hydrophilic and membrane domains in the region of the assumed acetogenin-binding site¹⁴. The observed R/A transition was quick (1-2 s)¹³, regardless whether Complex I was solubilized, reconstituted into liposomes or native membrane-bound and thus faster than reported for the mitochondrial enzyme ($\tau_{1/2} \sim 10$ s). The lifetime of A state was too short to be determined, which is in contrast to an active state of several hours in the mitochondrial enzyme at 0° C¹⁵.

In the previous report, R/A transition with solubilized or reconstituted enzyme was analyzed under conditions that neglected the proton pumping activity of Complex I¹³. Here, we study the activation profiles of proton pumping activity using reconstituted and native membrane-bound Complex I. We find that the activation properties of the two reactions catalyzed by Complex I, NADH:ubiquinone oxidoreduction and proton translocation across the membrane, are significantly different.

Materials and Methods

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Bacterial growth and purification of Complex I and ATP synthase. The *E. coli* MWC215 (Sm^R *ndh::Cm^R*) strain was grown in LB medium at 37°C in a 25 L fermenter and harvested at the late exponential growth phase. The membranes for Complex I purification were prepared by passing the cells through an APV Gaulin homogenizer. Then Complex I was purified by two consecutive chromatography steps using DEAE-Trisacryl M (Bio-Septra) anion exchanger columns and gel filtration on Superdex 200 prep grade (GE Healthcare), respectively, as described¹⁶. ATP synthase of *E. coli* was expressed and purified as described previously¹⁷

Reconstitution. Purified *Complex I* was reconstituted into liposomes as previously described¹⁸ except that azolectin was suspended in the buffer containing MOPS-BTP 100 mM, pH 7.0, n-dodecyl β-D-maltopyranoside (DDM) 0.4%, Na-cholate 0.1%. A ratio of 1:10 (w/w), of the enzyme to azolectin was used unless other specified. ATP synthase and ATP synthase/Complex I co-reconstitution were performed in the same way.

Preparation of subbacterial vesicles. The vesicles were prepared as described in¹⁹, but vesicles were loaded with 100 mM MOPS-BTP, 7.2, 50 mM KCl and 1 mM MgSO₄, frozen and stored at -80°C until use.

Kinetic measurements.

Monitoring Δψ in proteoliposomes was carried out at 24°C using a high-resolution CCD array spectrometer (HR2000+, Ocean Optics) combined with a DH-2000-BAL light source (Ocean Optics). Upon the reaction, optical spectra in the range of 250-680 nm were collected for 600 s. The difference in absorption at 630-590 nm presents Δψ-sensitive Oxonol VI response. Optical changes at 340 nm reflect simultaneous NADH consumption. The assay buffer comprised 100 mM

MOPS-BTP, 7.0, 1 mM MgSO₄, 5 mM (NH₄)₂SO₄, 2.5 μM Oxonol VI, 10 nM solubilized bo₃ oxidase, decylubiquinone (DQ) 100 μM. Prior to measurements proteoliposomes (protein concentration 7-15 μg/ml) were incubated in the assay buffer for 7 min at RT for equilibration of the DQ distribution.

Monitoring ΔpH in proteoliposomes were performed using entrapped pH-sensitive probe trisodium 8-hydroxypyrene-1,3,6-trisulfonate (pyranine) as described in¹⁸.

Monitoring ΔpH in membrane vesicles was carried out following fluorescence changes of pH-sensitive probe acridine orange (AO) with a Hitachi F-7000 fluorescence spectrophotometer, λ_{ex} = 493 nm, λ_{em} = 530 nm at 24°C unless other specified. The assay buffer comprised 50 mM MOPS-BTP, pH 7.2, 100 mM KCl, 1 mM MgCl₂, 1 μM valinomycin, 100 μM DQ, 1 mM KCN, 3.5 μM AO. The final protein concentration was 150-180 μg/ml.

Fast measurements of reconstituted Complex I NADH:ubiquinone oxidoreductase activity.

Following the initial stage of NADH oxidation was performed by rapid mixing of equal volumes of solubilized Complex I and buffer containing NADH (Unisoku Stopped-flow RSP-2000 apparatus) using a high-resolution CCD-array spectrometer (HR2000+, Ocean Optics) combined with a Xenon lamp as described in¹³.

Results

Delay of pumping activity in resting Complex I. In our previous work, activation of Complex I was followed by monitoring NADH oxidation and quinone reduction¹³. In functional Complex I this reaction is coupled to electrogenic proton transfer across the membrane that results in generation of a Δψ and a ΔpH. In order to monitor transmembrane events, purified Complex I was

reconstituted into liposomes and energized by addition of NADH in the presence of DQ, and the generation of $\Delta\psi$ or ΔpH was monitored (Figure 1). Accumulation of reduced ubiquinone was prevented by addition of a small amount of quinol oxidase bo_3 in detergent micelles, regenerating

oxidized ubiquinone without contributing to transmembrane proton transport.

Figure1. Following the $\Delta\psi$ (A) and ΔpH (B) generation in proteoliposomes by reconstituted Complex I. Following the first additions of NADH were made after full dissipation of $\Delta\psi$ or ΔpH and complete NADH consumption. 100 μM DQ and bo_3 oxidase was added to proteoliposomes prior to measurements.

Proton pumping initiated by addition of external NADH generates an electric potential (inside positive) across the proteoliposomes membrane that was followed using the $\Delta\psi$ -sensitive probe,

Oxonol VI. As shown in Figure 1A, initial addition of NADH resulted in $\Delta\psi$ generation followed by its relaxation to zero level after all NADH was consumed. Notably, subsequent NADH additions caused much faster formation of $\Delta\psi$. A similar effect was observed when acidification of the liposome lumen was followed with the pH-sensitive dye pyranine entrapped in the liposomes (Figure 1B). Similar to $\Delta\psi$ generation, the second addition of NADH caused a faster rate of acidification than the first addition. While the kinetics of the two experiments are different (fewer protons are required to establish a membrane potential than to significantly acidify the buffered liposome interior), the similar trend strongly indicates that initial proton pumping (in the resting enzyme) is much slower than in the activated enzyme.

Correlation of pumping and NADH:ubiquinone oxidoreductase activity. In the next series of experiments, activation profiles of both NADH oxidation (as reported in¹³) and $\Delta\psi$ generation were compared simultaneously (Figure 2A) by two consecutive NADH additions as described above.

Figure 2. A. Kinetics of $\Delta\psi$ generation and NADH consumption by reconstituted Complex I upon the first and the second addition of 50 μ M NADH. Upper panel, simultaneous $\Delta\psi$ and NADH consumption monitoring. NADH addition is indicated by an arrow. $\Delta\psi$ generation on 1st NADH addition, blue, on 2nd NADH addition, red. Corresponding NADH consumption tracks, black and magenta.

B. Fast measurements of

NADH:ubiquinone oxidoreduction by reconstituted Complex I. At zero time, the solution containing proteoliposomes equilibrated with 100 μM DQ was mixed 1:1 with the same buffer containing 100 μM NADH. Non-activated proteoliposomes, black; pre-activated with 30 μM NADH, magenta. Gray lines, exponential fits of the kinetics.

The data in Figure 2A show that the difference in rates between the first and second addition is much more pronounced for proton pumping than for NADH oxidation. While the NADH consumption rate in the time interval of 5-10 s at the first NADH addition was $\sim 70\%$ of the second addition, the maximal rate of $\Delta\psi$ response on the first NADH addition was only $\sim 15\%$ of the second.

The discrepancy between the rates of proton and electron transport indicates that Complex I is partially uncoupled during the first seconds of the measurements, i.e. electron transport is not tightly coupled to transmembrane proton transport. In the used setup, NADH oxidation could not be reliably followed in the time interval of 0-5 s because of a strong artificial response at 340 nm upon NADH dilution. To exclude the possibility that the strong activation of NADH consumption occurs during this time range, we performed stopped-flow measurements to resolve the initial time range (Figure 2B). At zero time, suspensions containing proteoliposomes, either untreated or activated with 30 μM NADH for several minutes, were mixed 1:1 with the same buffer containing 100 μM NADH. The results in Figure 2B show that the difference in the initial kinetics of NADH:ubiquinone oxidoreductase activities did not exceed 10%.

Dependence the proton pumping activation on the ratio Complex I:azolectin upon reconstitution.

The energization of proteoliposomes membrane is always a result of a superposition of two

oppositely directed fluxes, one being active H^+ pumping by the enzyme and the other being H^+ (or another ion) leakage through the membrane. The signal amplitude of active pumping is thus expected to increase with the number of Complex I proteins in the liposomes, while the membrane leakage contribution remains unaffected (Figure 3A).

Figure 3. Kinetics of $\Delta\psi$ generation by reconstituted Complex I upon the first (A) and the second (B) addition of NADH at 50 μM concentration indicated by arrows. The ratio Complex I/azolectin (w/w) upon reconstitution 1:50, 1:30, 1:20, 1:15, 1:10, from top to bottom. NADH additions are indicated by arrows.

Purified Complex I preparation contains some amount of annular lipids that could be important for the enzyme behavior. Reconstitution into azolectin liposomes may result in a replacement of these lipids dependently on the proportion of the lipid phase, therefore it was important to test whether proteoliposome content affects the observed activation of Complex I. We carried out monitoring

$\Delta\psi$ generation in proteoliposomes with different Complex I:lipid ratios. In Figure 3 the responses on 1st (A) and 2nd (B) NADH addition to proteoliposomes were Complex I was reconstituted in ratios of 1:50, 1:30, 1:20, 1:15, 1:10 (w/w) are presented. Although the amplitude of the response changed in accordance with protein content, as expected, the proportion between the initial rates after 1st and 2nd NADH additions stayed unaltered. When the ratio Complex I:lipid exceeds 1:10 (w/w) the signal becomes lower, most probably due to Complex I aggregation and poor reconstitution. Taken together, the results show that the activation phenomenon is not affected by changes in protein:lipid ratio.

Dependence of Complex I activation on concentration of initially added NADH. Next, the effect of NADH concentrations varying from 2 to 100 μM as the first addition was tested (Figure 4A), while the second addition was always 50 μM . The second portion was added only after the $\Delta\psi$

generated

Figure 4. Dependence of reconstituted Complex I activation on concentration of initially added NADH. **A.** Generation of $\Delta\psi$ on the 2nd NADH addition at concentration of 50 μM (indicated by an arrow). The first NADH addition was 0, 5, 10, 20, 50, 100 μM , from top to bottom. **B.** Dependence of the initial rate of $\Delta\psi$ generation by activated Complex I on the concentration of NADH added for the activation.

upon the first addition had completely dissipated. We found that the full activation of proton pumping required 10-15 μM NADH during the first addition, while further increase in NADH concentration resulted in the same activation (Figure 4B). Upon the third NADH addition, which is performed immediately after full NADH consumption of the second addition the profile of $\Delta\psi$ generation was similar to the second NADH addition.

Activation of Complex I pumping activity by NADPH. The results from the experiments above indicate that preceding enzyme turnover directly influences the extent of activation. We therefore tested, if activation is possible with NADPH, which is able to reduce Complex I, showing very slow turnover. NADPH was found to activate mitochondrial Complex I^{15, 20}. As expected, Complex I incubation with 100 μM NADPH resulted in negligible NADPH:ubiquinone oxidoreductase activity and almost non-detectable $\Delta\psi$ generation, since the turnover is so slow that H^+ pumping cannot

overcome the membrane leakage (Figure 5, black trace).

Figure 5. Activation of Complex I proton pumping activity by NADPH. Almost no generation of $\Delta\psi$ after 100 μM NADPH addition (black line) was detected, however the enzyme was activated.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Response to the primary addition of 50 μM NADH after preincubation with NADPH, red. Generation of $\Delta\psi$ without preincubation with NADPH, blue curves: 1st 50 μM NADH addition, dashed; 2nd 50 μM NADH addition, solid. Nucleotides addition is indicated by an arrow.

However, after an incubation of the proteoliposomes with NADPH for ~ 2 minutes, the addition of 50 μM NADH shows that Complex I has been significantly (but not completely) activated (Figure 5, red trace) if compared with non-activated complex I (blue dotted trace) or fully activated enzyme (blue trace). Incubation of the proteoliposomes with 100 μM NAD⁺ as a substrate analogue that cannot be turned over, did not affect the slow proton pumping on the 1st NADH addition (Figure S1) and thus not activate Complex I. These data show that the enzyme activation is rather dependent on the turnover than on the incubation with a substrate analogue.

Co-reconstitution of ATPase and Complex I. As mentioned above, a possible explanation for the observed results is that initial NADH oxidation is uncoupled from proton pumping. However, another possibility would be that in the inactive state, Complex I confers a higher proton permeability to the membrane. The observed effect would then be due to a delayed recovery of the low permeability (tightening of the membrane) upon enzyme activation. We tested this hypothesis by co-reconstitution of complex I with ATP synthase, a separate proton pump. In these experiments, ATP is added and the ATP synthase works in hydrolysis direction as a primary proton pump.

As seen in Figure 6 the amplitude and the kinetics of $\Delta\psi$ generated by ATP synthase were similar regardless of whether it was reconstituted alone (blue trace) or together with Complex I, activated (red trace) or non-activated (green trace). The data show that the presence of Complex I in the

proteoliposomes, either “as prepared” or activated, does not affect the membrane permeability of the liposomes upon ATP driven proton pumping. It should be noted that the amplitude of $\Delta\psi$ generated by the ATP synthase is higher than that of Complex I, which is explained by the lower turnover number of the complex I that is limited by the slow exchange of ubiquinone between lipid and water phase²¹.

Figure 6. Generation of $\Delta\psi$ by reconstituted ATPase in the presence or absence of Complex I. ATPase reconstituted alone, red curve; ATPase co-reconstituted with Complex I, green curve; ATPase was reconstituted with Complex I, which was activated by 50 μ M NADH, blue curve. 0.4 mM ATP was added as indicated. The dissipation of $\Delta\psi$ was achieved by an addition of 1.5 μ g/ml gramicidin D.

Taken together, these data indicate that the different activation profiles of NADH oxidation and proton pumping are indeed due to a lowered proton pumping stoichiometry after the first addition.

Activation of proton pumping by Complex I in membrane vesicles. Finally, inverted membranes vesicles of *E. coli* were prepared to test whether the different activation profiles can also be observed in native membranes. To suppress proton pumping by the *bo*₃ oxidase initiated by NADH, the medium was supplemented with 1 mM KCN, which specifically blocks *bo*₃ oxidase. Instead, 100 μ M DQ was added as an electron acceptor. In these experiments, only a limited amount of NADH could be used, since the reduced ubiquinone, which is accumulated in the

absence of *bo*₃ oxidase activity, notably inhibits Complex I, similar to the inhibiting effect of ubiquinol Q₁ with the mitochondrial Complex I²². Due to high density and activity of Complex I in the membranes, it was impossible to follow the fast development of $\Delta\psi$, thus we studied proton translocation by following the acidification of the vesicle interior by the Δ pH sensitive fluorescent probe acridine orange (AO). To convert $\Delta\psi$ to Δ pH, the assay buffer was supplemented with 1 μ M

valinomycin and 100 mM KCl. In parallel, NADH consumption was determined optically in separate samples but under identical conditions. At 24 °C, an insignificant acceleration of Δ pH generation by pre-activated Complex I was observed and no reliable difference in NADH consumption by non-activated and activated Complex I was detected as reported previously¹³. However, at lower temperatures the difference in the activity of activated and non-activated Complex I became evident. At 15 °C, the maximal acidification rate is 2-3 times higher if the enzyme was activated and the delay in pumping is much more pronounced (Figure 7).

Figure 7. Acidification of the membrane vesicles interior and NADH consumption at 24 °C (**A**) and 15 °C (**B**). Before measurements the membranes were equilibrated with AO, 100 μM DQ and 1 mM KCN for 5 min at corresponding temperatures. AO response to: the 1st 20 μM NADH addition, blue; to the 2nd 20 μM NADH addition, red. NADH consumption at: 1st 20 μM NADH addition, black; 2nd 20 μM NADH addition, magenta. NADH was added at zero time as indicated by arrow.

NADH consumption as well proceeds with slightly longer delay at lower temperatures, however, it does not correlate with the delay in acidification. The maximal rate of NADH consumption was identical in both conditions, whereas there was still a significant difference in the maximal acidification rates between untreated and activated complex I (Figure 7).

Discussion

The previously found R/A transition of Complex I from *E. coli* is a relatively fast process; it occurs in the time range of 1-2 s¹³, after which the rate of NADH:ubiquinone oxidoreduction has reached its steady-state value. Here, we found that in contrast to the full activation of NADH oxidation, coupled proton translocation in reconstituted Complex I stays significantly depressed much longer (Figure 2). Since Complex I contains solvent-accessible, low-potential FMN and therefore produces ROS upon reduction, it could be assumed that at the initial stage of the reaction, accelerated electron leakage to oxygen occurs. Production of ROS at significant rate could result in a mismatch between NADH oxidation and proton pumping at this stage. However, previous measurements of ROS production by Complex I revealed that the rate of ROS production does not exceed 0.2% of NADH oxidation rate regardless whether Complex I was solubilized or

1
2
3 reconstituted²³. Such low electron leakage cannot affect proton pumping. More likely explanations
4
5 of the different activation profiles of electron and proton translocations could be that inactive
6
7 Complex I either provides higher proton permeability to the membrane, or it undergoes a
8
9 transition from de-coupled to coupled state (D/C transition) upon turnover. In our experiments,
10
11 the former possibility was excluded by co-reconstitution of Complex I with ATP synthase, a
12
13 separate proton pump. Irrespective of the presence or the state of Complex I, proton permeability
14
15 tested by ATP driven proton pumping was unaffected. Based on our findings, we therefore
16
17 propose that bacterial Complex I undergoes a transition from decoupled to coupled state upon
18
19 turnover.
20
21
22
23
24

25
26 Properties of the system did not let us determine the activation time of proton pumping directly.
27
28 The rate of NADH consumption may stay at a constant steady state relatively long time if the
29
30 concentration of substrates is high and of the enzyme is low, allowing the observation of
31
32 NADH:ubiquinone oxidoreduction activation. In contrast, the membrane energization occurs
33
34 relatively fast and the final level of $\Delta\mu_{H^+}$ is approached before the enzyme activation (Figure 1).
35
36 Therefore, only the comparison of the initial rates of the energization reliably indicates the
37
38 enzyme state. Some information on activation time can be obtained from data on the
39
40 dependence of full activation of pumping on the amount of NADH. Complex I concentration in the
41
42 sample was approximately 15 nM and the addition of 2 μ M NADH guaranteed over 100 turnovers,
43
44 yet full activation was not achieved. The obtained data showed that the activation can be reached
45
46 by multiple turnovers during approximately 80 s required for the consumption of 10 μ M NADH
47
48 (Figure 4). The activation of pumping by NADPH also indicates a necessity of multiple turnovers
49
50 and rules out a requirement of Complex I reduction to the level that can be approached by NADH
51
52
53
54
55
56
57
58
59
60

addition. In the presence of DQ, the enzyme reduction by NADPH is insignificant due to the much lower affinity of the electron donor NADPH as compared to the electron acceptor DQ. These findings are in agreement with the data on NADPH effect on A/D transition of mitochondrial Complex I: the enzyme reduction by NADH or NADPH in the absence of electron acceptors does not initiate its activation, which can be achieved by NADPH only under conditions when electron transfer from the substrate to quinone acceptor is permitted²⁰. Pre-existing $\Delta\psi$ maybe also excluded as an essential activation factor, as incubation with NADPH did not result in $\Delta\psi$ generation due to the slow turnover, yet Complex I was activated.

It is likely that the fast acceleration of NADH:ubiquinone oxidoreduction¹³ and simultaneous conformational changes¹⁴ are the early stages of an establishment of the fully functional state of Complex I. The observed lack of direct correlation between electron and proton translocation implies a decoupled state of Complex I that was not detected previously. It is reasonable to raise the question whether this feature is an intrinsic property of Complex I or it is acquired due to enzyme treatment upon purification and reconstitution. To test this, we studied Complex I functioning in native membranes of subbacterial vesicles. Previously we found that NADH:ubiquinone oxidoreductase activity of the membrane-bound enzyme has a short (1-2 s) delay as well as purified Complex I, and no activation was detected at RT¹³. In this study, monitoring of the proton pumping activity showed a small, but highly reproducible acceleration after initial activation by NADH. However, lowering the temperature clearly revealed the phenomenon of pumping activation (Figure 7). These results imply that the D/C transition is indeed a built-in property of Complex I, but further studies are required to manifest this statement.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The proposed transition could be a precursor of active/de-active transition of mitochondrial Complex I, which evolved in evolution to ensure the enzyme’s regulation in eukaryotes. Testing proton pumping by Complex I in the natural environment, in membrane vesicles, indicated that D/C transition is an intrinsic property of the enzyme. However, the significantly prolonged decoupled state of the reconstituted enzyme could be of artificial nature, i.e. acquired during the enzyme isolation and purification and might be caused by a loss of an unidentified factor, which provides fast Complex I activation. Nevertheless, it is clear that both native and purified, reconstituted Complex I from *E. coli* may exist in resting, non-pumping state, which converts to pumping upon slow turnover. Finding of two states of *E. coli* Complex I should be taken into account when planning future research, it provides a powerful tool to determine properties of coupled and decoupled enzyme and, therefore, opens up opportunities for studying the molecular mechanism of Complex I proton translocation.

Funding

We acknowledge the financial support of the Biocentrum Helsinki 7919530, the Sigrid Juselius Foundation 4702217, and Magnus Ehrnrooth Foundation 96133, and the Swiss National Science Foundation 153351.

Supporting Information

Figure S1 showing that NAD⁺ is incapable to activate proton pumping by Complex I.

References

(1) Gavrikova, E. V., and Vinogradov, A. D. (1999) Active/de-active state transition of the mitochondrial complex I as revealed by specific sulfhydryl group labeling, *FEBS Lett.* 455, 36-40.

- (2) Gostimskaya, I. S., Cecchini, G., and Vinogradov, A. D. (2006) Topography and chemical reactivity of the active-inactive transition-sensitive SH-group in the mitochondrial NADH : ubiquinone oxidoreductase (Complex I), *Biochim. Biophys. Acta* 1757, 1155-1161.
- (3) Grivennikova, V. G., Kapustin, A. N., and Vinogradov, A. D. (2001) Catalytic activity of NADH-ubiquinone oxidoreductase (Complex I) in intact mitochondria - Evidence for the slow active/inactive transition, *J. Biol. Chem.* 276, 9038-9044.
- (4) Dröse, S., Stepanova, A., and Galkin, A. (2016) Ischemic A/D transition of mitochondrial complex I and its role in ROS generation, *Biochim. Biophys. Acta* 1857, 946-957.
- (5) Kotlyar, A. B., Albracht, S. P. J., and van Spanning, R. J. M. (1998) Comparison of energization of complex I in membrane particles from *Paracoccus denitrificans* and bovine heart mitochondria, *Biochim. Biophys. Acta* 1365, 53-59.
- (6) Maklashina, E., Kotlyar, A. B., and Cecchini, G. (2003) Active/de-active transition of respiratory complex I in bacteria, fungi, and animals, *Biochim. Biophys. Acta* 1606, 95-103.
- (7) Zickermann, V., Wirth, C., Nasiri, H., Siegmund, K., Schwalbe, H., Hunte, C., and Brandt, U. (2015) Structural biology. Mechanistic insight from the crystal structure of mitochondrial complex I, *Science* 347, 44-49.
- (8) Wirth, C., Brandt, U., Hunte, C., and Zickermann, V. (2016) Structure and function of mitochondrial complex I, *Biochim. Biophys. Acta* 1857, 902-914.
- (9) Fiedorczuk, K., Letts, J. A., Degliesposti, G., Kaszuba, K., Skehel, M., and Sazanov, L. A. (2016) Atomic structure of the entire mammalian mitochondrial complex I, *Nature* 538, 406-410.
- (10) Zhu, J. P., Vinothkumar, K. R., and Hirst, J. (2016) Structure of mammalian respiratory complex I, *Nature* 536, 354-359.
- (11) Galkin, A., Meyer, B., Wittig, I., Karas, M., Schagger, H., Vinogradov, A., and Brandt, U. (2008) Identification of the mitochondrial ND3 subunit as a structural component involved in the active/deactive enzyme transition of respiratory complex I, *J. Biol. Chem.* 283, 20907-20913.
- (12) Babot, M., Labarbuta, P., Birch, A., Kee, S., Fuszard, M., Botting, C. H., Wittig, I., Heide, H., and Galkin, A. (2014) ND3, ND1 and 39kDa subunits are more exposed in the de-active form of bovine mitochondrial complex I, *Biochim. Biophys. Acta* 1837, 929-939.
- (13) Belevich, N., and Verkhovskaya, M. (2016) Resting state of respiratory Complex I from *Escherichia coli*, *FEBS Lett.* 590, 1570-1575.
- (14) Belevich N, B. G., Chen Z., Sinha S.C., Verkhovskaya M. (2017) Activation of respiratory Complex I from *Escherichia coli* studied by fluorescent probes, *Heliyon* 3, e00224.
- (15) Kotlyar, A. B., and Vinogradov, A. D. (1990) Slow active inactive transition of the mitochondrial NADH-ubiquinone reductase, *Biochim. Biophys. Acta* 1019, 151-158.
- (16) Euro, L., Belevich, G., Wikström, M., and Verkhovskaya, M. (2009) High affinity cation-binding sites in Complex I from *Escherichia coli*, *Biochim. Biophys. Acta* 1787, 1024-1028.
- (17) Wiedenmann, A., Dimroth, P., and von Ballmoos, C. (2008) $\Delta\psi$ and ΔpH are equivalent driving forces for proton transport through isolated F-0 complexes of ATP synthases, *Biochim. Biophys. Acta* 1777, 1301-1310.
- (18) Verkhovskaya, M., Knuuti, J., and Wikström, M. (2011) Role of Ca^{2+} in structure and function of Complex I from *Escherichia coli*, *Biochim. Biophys. Acta* 1807, 36-41.
- (19) Belevich, G., Euro, L., Wikström, M., and Verkhovskaya, M. (2007) Role of the conserved arginine 274 and histidine 224 and 228 residues in the NuoCD subunit of complex I from *Escherichia coli*, *Biochemistry* 46, 526-533.
- (20) Vinogradov, A. D., and Grivennikova, V. G. (2001) The mitochondrial complex I: Progress in understanding of catalytic properties, *IUBMB Life* 52, 129-134.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(21) Shinkarev, V. P., and Wraight, C. A. (1997) The interaction of quinone and detergent with reaction centers of purple bacteria .1. Slow quinone exchange between reaction center micelles and pure detergent micelles, *Biophys. J.* 72, 2304-2319.

(22) Grivennikova, V. G., Kapustin, A. N., and Vinogradov, A. D. (2001) Catalytic activity of NADH-ubiquinone oxidoreductase (Complex I) in intact mitochondria - Evidence for the slow active/inactive transition, *J. Biol. Chem.* 276, 9038-9044.

(23) Knuuti, J., Belevich, G., Sharma, V., Bloch, D.A., Verkhovskaya, M. (2013) A single amino acid residue controls ROS production in the respiratory Complex I from Escherichia coli, *Mol. Microbiol.* 90, 1190-1200

Activation of proton translocation by respiratory Complex I

Nikolai Belevich, Christoph von Ballmoos, Marina Verkhovskaya