

RESEARCH ARTICLE

Open Access



Death at no cost? Persons with no health insurance claims in the last year of life in Switzerland

Radoslaw Panczak^{1*} , Viktor von Wyl², Oliver Reich^{3,4}, Xhyljeta Luta¹, Maud Maessen^{1,5}, Andreas E. Stuck⁶, Claudia Berlin¹, Kurt Schmidlin¹, David C. Goodman^{1,7}, Matthias Egger¹, Kerri Clough-Gorr^{1,8} and Marcel Zwahlen¹

Abstract

Background: Lack of health insurance claims (HIC) in the last year of life might indicate suboptimal end-of-life care, but reasons for no HIC are not fully understood because information on causes of death is often missing. We investigated association of no HIC with characteristics of individuals and their place of residence.

Methods: We analysed HIC of persons who died between 2008 and 2010, which were obtained from six providers of mandatory Swiss health insurance. We probabilistically linked these persons to death certificates to get cause of death information and analysed data using sex-stratified, multivariable logistic regression. Supplementary analyses looked at selected subgroups of persons according to the primary cause of death.

Results: The study population included 113,277 persons (46% males). Among these persons, 1199 (proportion 0.022, 95% CI: 0.021–0.024) males and 803 (0.013, 95% CI: 0.012–0.014) females had no HIC during the last year of life. We found sociodemographic and health differentials in the lack of HIC at the last year of life among these 2002 persons. The likelihood of having no HIC decreased steeply with older age. Those who died of cancer were more likely to have HIC (adjusted odds ratio for males 0.17, 95% CI: 0.13–0.22; females 0.19, 95% CI: 0.12–0.28) whereas those dying of mental and behavioural disorders (AOR males 1.83, 95% CI: 1.42–2.37; females 1.65, 95% CI: 1.27–2.14), and males dying of suicide (AOR 2.15, 95% CI: 1.72–2.69) and accidents (AOR 2.41, 95% CI: 1.96–2.97) were more likely to have none. Single, widowed, and divorced persons also were more likely to have no HIC (AORs in range of 1.29–1.80). There was little or no association between the lack of HIC and characteristics of region of residence. Patterns of no HIC differed across main causes of death. Associations with age and civil status differed in particular for persons who died of cancer, suicide, accidents and assaults, and mental and behavioural disorders.

Conclusions: Particular groups might be more likely to not seek care or not report health insurance costs to insurers. Researchers should be aware of this aspect of health insurance data and account for persons who lack HIC.

Keywords: End-of-life, Delivery of health care, Health care cost, Health insurance, Switzerland

Background

Health insurance claims (HIC) offer cost-effective research potential for studying large populations to track determinants and variations in the use of healthcare [1, 2]. HIC are particularly important in Switzerland, where population-wide data about healthcare use, particularly relating to cost and end-of-life (EOL) care, are scarce or fragmented [3].

Swiss HIC data inform aspects of healthcare delivery such as EOL cost trajectories [4], the burden of schizophrenia [5], and potentially inappropriate medications [6], just to name a few. Yet despite its strengths, HIC data lack information that otherwise could improve their usefulness in health services research. Information that is important in, for example, EOL studies, on cause of death, is often not readily available [7].

Swiss residents enjoy one of the best performing healthcare systems and have one of the highest life expectancies in the world. At the same time, this system

* Correspondence: r.panczak@gmail.com

¹Institute of Social and Preventive Medicine, University of Bern, Finkenhubelweg 11, 3012 Bern, Switzerland

Full list of author information is available at the end of the article



is characterized by high costs and complexity that make it difficult to manage and change. With large choice and wide supply of services, individuals might find it hard to find optimal solutions [8]. Similarly, high spending might not necessarily mean high quality; in 2010 the Economist Intelligence Unit ranked Switzerland 30th out of 40 in quality of end-of-life care [9].

Healthcare expenditures tend to rise, often sharply, near the end of life [4, 10]. Studies often use development of cost over a certain period or aggregate overall cost within a certain period prior to death [11]. With advances in healthcare in general, and the growing intensity of EOL care in particular, these costs tend to be substantial. However, some small proportion of persons die with either no healthcare use at all, or at least no HIC. Lack of HIC in the last year of life might indicate suboptimal EOL care, but the reasons for no HIC are not fully understood. We therefore investigated associations between the lack of end-of-life health insurance claims and characteristics of individuals and where they live.

Methods

Study design & data sources

This study used routinely collected Swiss HIC data in a retrospective, secondary analysis of data that are described in detail elsewhere [12]. To summarize, data from six of the 10 largest insurers operating in the Swiss market were pooled and used to track healthcare use over the last year of life of adults who died between 2008 and 2010 [13]. The insurance providers delivered data on sex, date of birth, date of death, place of residence (community or postcode), and complete records of HIC of policyholders. Based on the communities in which they resided, we deterministically linked data on level of urbanization, language region, and neighbourhood socioeconomic status.

Using dates of birth and death, sex, and place of residence, we probabilistically linked the insured persons file to the death certificates in the Swiss Federal Statistical Office's database (see [12] for details on linkage procedure and results). In addition to causes of death, this data linkage also provided information about civil status and nationality.

Study setting


Basic health insurance, which covers all services related to illness and pregnancy deemed medically necessary and cost-appropriate, is mandatory for and offered to all Swiss residents with no prior checks or restrictions [14]. Eighty-one private insurers (at the time closest to the end of the study period, 10 August 2011) offered the mandatory basic insurance package. Insurers vary greatly in size and coverage from the 858,005 insured by CSS

(which provided data for this study) to 170 persons insured by Krankenkasse Zeneggen [13]. The mean number of those insured by all providers is 96,045 (standard deviation 167,746).

Residents choose a deductible in a range of 300–2500 Swiss Francs (1 CHF = 0.85 Euro = 1.01 US\$, as of 25 December 2017); higher deductibles and managed care plans lower the cost of premiums. Social assistance subsidizes premiums for low-income persons. Hospital claims are delivered directly to insurers, whereas a majority of other services is paid by individuals and later reimbursed by insurers after deductibles have been met. Individuals can voluntarily supplement the basic insurance package with private insurance to add further provider and treatment choices (e.g., complementary medicine, dental care) and cover additional benefits (e.g., a private room during a hospital stay). A separate mandatory insurance system covers HIC that are accident related [4, 7, 8, 12, 15].

Conceptual framework

We hypothesized that having no health insurance claims in the last year of life may occur in two main ways: either a person used no healthcare, or healthcare was delivered but no HIC were filed (Fig. 1). In the first case, a person could have died having had no need of healthcare, having had all healthcare needs met by the family, having refused treatment, having been incapable of finding or paying for healthcare, having been undertreated, or a person could have died suddenly with no possibility of medical care. In the second case, care was in fact delivered but for one reason or another information on its cost never reached the insurer. This could happen,



for example, in situations in which healthcare was paid entirely out-of-pocket or by supplemental healthcare insurance, the cost of care did not reach the level of the relevant deductible and information about that cost did not reach the insurer, or a patient or caregiver was not willing or capable of handling the HIC documents.

Analyses

Guided by previous work [12], we stratified analyses by sex. In the last year of life, the absence of reimbursed HIC as opposed to having some HIC was a binary outcome of the analyses. We calculated frequencies of persons with and without HIC and proportions of persons without HIC across covariates. We used logistic regression with robust standard errors that adjusted for clustering of decedents within regions of residence [16]. Multivariable models included age in 5-year bands, nationality (Swiss or foreigner, including unknown), and civil status (single, married, widowed, divorced), and cause of death, which was coded according to the 10th revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10): cardiovascular diseases (CVD, all I codes), cancer (all C codes), mental and behavioural disorders (all F codes), diseases of the nervous system (all G codes), respiratory diseases (all J codes), diseases of the digestive system (all K codes) accidents and assaults (all V, all W and X00-X59, X85-X99, Y00-Y09, Y85-Y86 codes), suicide (X60-X84 codes), and other (remaining codes). Characteristics of regions of residence included level of urbanization (urban, periurban, rural), language region (German, French, Italian) and quintiles of median area-based socioeconomic position (Swiss-SEP) index [17]. We conducted supplementary analyses by main causes of death to explore whether the associations differed across these strata. These analyses further aggregated age categories in order to have a sufficient number of events across groups.

Results

The probabilistic linkage to the death certificate database had a 95.6% success rate; unlinked individuals were similar to the linked ones and were excluded. The study population consisted of 113,277 persons (Table 1) who comprised 61.3% of those who died in Switzerland between 2008 and 2010. The sex, age, nationality, civil status, level of urbanization, language region, Swiss-SEP and, most importantly, cause of death distributions were almost identical to overall mortality in that time period [12].

One thousand one hundred ninety nine males (0.022, 95% confidence interval: 0.021–0.024) and 803 females (0.013, 95% CI: 0.012–0.014) did not have any reimbursed HIC in the last year of life. The proportion of

persons with no HIC decreased sharply with age, particularly among males; for example, 0.194 (95% CI: 0.151–0.238) of men who were 19–25 at death had no claim, as opposed to approximately 0.010 (95% CI: 0.007–0.013) of those 76 and older (Fig. 2, left panel). More males dying of accidents and assaults (0.095, 95% CI: 0.083–0.107) and suicide (0.118, 95% CI: 0.101–0.135) had no HIC, whereas the proportion of persons who died of cancer and had no HIC was low (0.005, 95% CI: 0.004–0.006 for men; 0.004, 95% CI: 0.003–0.005 for women). Slightly more foreign, single, or divorced males had no HIC.

A strong, negative age gradient remained in the multivariable logistic regression model (Fig. 2, right panel; see Additional file 1 for exact estimates of AOR); the adjusted odds ratio (AOR) for the youngest males was 2.63 (95% CI: 1.79–3.86, compared to persons 61–65 years old), whereas for the oldest it was 0.24 (95% CI: 0.17–0.35). In comparison to persons dying from CVD, cancer patients were unlikely to have no HIC (AOR 0.17, 95% CI: 0.13–0.22 for males; 0.19, 95% CI: 0.12–0.28 for females) with weaker effects for young males dying of respiratory and digestive organs diseases. On the other hand, males dying from accidents or assaults (AOR 2.41 95% CI: 1.96–2.97) or suicide (AOR 2.15 95% CI: 1.72–2.69) had higher probability of not having any HIC. For both sexes, those who died of mental and behavioural disorders as well as single, widowed, and divorced persons were more likely to have no HIC. There was little or no association with place of residence apart from a lowered AOR for females in the Italian speaking part of Switzerland.

We found different patterns of association across selected main causes of death among those having no HIC [see Additional file 2]. Persons who died of CVD resembled the overall findings. Persons who died of accidents and assaults showed an association mainly with age. Associations with age and civil status varied the most. For example, no association with age was observed among either persons who died of cancer or females who died of mental and behavioural disorders and suicide. Neither did civil status play a role for males who died of mental and behavioural disorders and suicide.

Discussion

Principal findings

We found demographic, health, and socioeconomic differentials in the lack of health insurance claims, and possibly costs, in the last year of life. Several groups of patients identified by sex, age, civil status, and cause of death had a higher probability of not having HIC. Region of residence had little effect, and associations with age and civil status varied for certain causes of death.

Table 1 Study population. Distribution of persons with and without health insurance claims and proportion (and 95% confidence interval) across analysed variables. Attribution of causes of death follows ICD-10 coding

Category	Males					Females				
	HIC exist		No HIC		Proportion no HIC (95% CI)	HIC exist		No HIC		Proportion no HIC (95% CI)
	No.	Col %	No.	Col %		No.	Col %	No.	Col %	
Age group										
19–25	253	0%	61	5%	0.194 (0.151–0.238)	105	0%	8	1%	0.071 (0.024–0.118)
26–30	214	0%	40	3%	0.157 (0.113–0.202)	126	0%	3	0%	0.023 (0.000–0.049)
31–35	262	1%	38	3%	0.127 (0.089–0.164)	153	0%	9	1%	0.056 (0.020–0.091)
36–40	384	1%	48	4%	0.111 (0.081–0.141)	252	0%	4	0%	0.016 (0.000–0.031)
41–45	671	1%	80	7%	0.107 (0.084–0.129)	391	1%	10	1%	0.025 (0.010–0.040)
46–50	1111	2%	93	8%	0.077 (0.062–0.092)	702	1%	14	2%	0.020 (0.009–0.030)
51–55	1606	3%	107	9%	0.062 (0.051–0.074)	1048	2%	27	3%	0.025 (0.016–0.034)
56–60	2411	5%	123	10%	0.049 (0.040–0.057)	1468	2%	23	3%	0.015 (0.009–0.022)
61–65	3672	7%	105	9%	0.028 (0.023–0.033)	2191	4%	32	4%	0.014 (0.009–0.019)
66–70	4584	9%	107	9%	0.023 (0.019–0.027)	2801	5%	41	5%	0.014 (0.010–0.019)
71–75	5713	11%	85	7%	0.015 (0.012–0.018)	4153	7%	52	6%	0.012 (0.009–0.016)
76–80	7925	15%	77	6%	0.010 (0.007–0.012)	6607	11%	84	10%	0.013 (0.010–0.015)
81–85	9496	18%	89	7%	0.009 (0.007–0.011)	11,268	19%	125	16%	0.011 (0.009–0.013)
86–90	8404	16%	92	8%	0.011 (0.009–0.013)	13,527	23%	159	20%	0.012 (0.010–0.013)
91+	5411	10%	54	5%	0.010 (0.007–0.013)	14,366	24%	212	26%	0.015 (0.013–0.016)
Nationality										
Swiss	46,768	90%	1013	84%	0.021 (0.020–0.022)	55,671	94%	748	93%	0.013 (0.012–0.014)
Foreigner	5349	10%	186	16%	0.034 (0.029–0.038)	3487	6%	55	7%	0.016 (0.011–0.020)
Civil status										
Single	6247	12%	380	32%	0.057 (0.052–0.063)	6978	12%	134	17%	0.019 (0.016–0.022)
Married	30,452	58%	476	40%	0.015 (0.014–0.017)	13,064	22%	131	16%	0.010 (0.008–0.012)
Widowed	10,796	21%	138	12%	0.013 (0.011–0.015)	33,875	57%	425	53%	0.012 (0.011–0.014)
Divorced	4622	9%	205	17%	0.042 (0.037–0.048)	5241	9%	113	14%	0.021 (0.017–0.025)
Cause of death										
CVD	17,398	33%	353	29%	0.020 (0.018–0.022)	22,858	39%	321	40%	0.014 (0.012–0.015)
Cancer	16,107	31%	76	6%	0.005 (0.004–0.006)	13,250	22%	48	6%	0.004 (0.003–0.005)
Mental & behavioural disorders	2313	4%	87	7%	0.036 (0.029–0.044)	4670	8%	110	14%	0.023 (0.019–0.027)
Nervous system	2178	4%	40	3%	0.018 (0.012–0.024)	3256	6%	66	8%	0.020 (0.015–0.025)
Respiratory	3637	7%	39	3%	0.011 (0.007–0.014)	3360	6%	41	5%	0.012 (0.008–0.016)
Digestive	2088	4%	35	3%	0.016 (0.011–0.022)	2419	4%	34	4%	0.014 (0.009–0.018)
Accidents	1980	4%	208	17%	0.095 (0.083–0.107)	1955	3%	33	4%	0.017 (0.011–0.022)
Suicide	1213	2%	162	14%	0.118 (0.101–0.135)	581	1%	15	2%	0.025 (0.013–0.038)
Other	5203	10%	199	17%	0.037 (0.032–0.042)	6809	12%	135	17%	0.019 (0.016–0.023)
Urbanicity										
Urban	16,086	31%	392	33%	0.024 (0.021–0.026)	20,503	35%	349	43%	0.017 (0.015–0.018)
Peri-urban	22,725	44%	528	44%	0.023 (0.021–0.025)	24,819	42%	324	40%	0.013 (0.011–0.014)
Rural	13,306	26%	279	23%	0.021 (0.018–0.023)	13,836	23%	130	16%	0.009 (0.008–0.011)
Language region										
German	36,616	70%	877	73%	0.023 (0.022–0.025)	42,034	71%	548	68%	0.013 (0.012–0.014)
French	12,857	25%	268	22%	0.020 (0.018–0.023)	13,905	24%	237	30%	0.017 (0.015–0.019)

Table 1 Study population. Distribution of persons with and without health insurance claims and proportion (and 95% confidence interval) across analysed variables. Attribution of causes of death follows ICD-10 coding (Continued)

Category	Males					Females				
	HIC exist		No HIC		Proportion no HIC (95% CI)	HIC exist		No HIC		Proportion no HIC (95% CI)
	No.	Col %	No.	Col %		No.	Col %	No.	Col %	
Italian	2644	5%	54	5%	0.020 (0.015–0.025)	3219	5%	18	2%	0.006 (0.003–0.008)
Swiss-SEP quintile										
1st (lowest)	3807	7%	81	7%	0.021 (0.016–0.025)	3847	7%	40	5%	0.010 (0.007–0.013)
2nd	11,480	22%	233	19%	0.020 (0.017–0.022)	12,323	21%	139	17%	0.011 (0.009–0.013)
3rd	14,256	27%	312	26%	0.021 (0.019–0.024)	16,128	27%	194	24%	0.012 (0.010–0.014)
4th	17,455	33%	451	38%	0.025 (0.023–0.027)	21,196	36%	334	42%	0.016 (0.014–0.017)
5th (highest)	5119	10%	122	10%	0.023 (0.019–0.027)	5664	10%	96	12%	0.017 (0.013–0.020)
Total	52,117	100%	1199	100%	0.022 (0.021–0.024)	59,158	100%	803	100%	0.013 (0.012–0.014)

Abbreviations: Col, column; HIC, health insurance claims; CI, confidence interval; Mental & behav., mental and behavioural disorders; Swiss-SEP, Swiss neighbourhood index of socioeconomic position [13]

Strengths

This is the first study to the authors' knowledge to have looked at the lack of mandatory health insurance (MHI) reimbursed healthcare in the last year of life. We used a large, diverse, and representative database of HIC augmented by probabilistic linkage to a database of causes of death and regional characteristics [12]. MHI covers the entire range of providers, including hospital and ambulatory care, medication, and nursing home medical costs.

Relation to other studies

Unsurprisingly, patterns identified in this study reflect other findings on overall cost of care [12]. For example, persons who died of cancer had higher costs and lower probability of having had no HIC whereas for younger, widowed, and divorced persons the opposite was true. Men and unmarried persons in the U.S. were also found less likely to receive care in the last year of life [18]. Being male, younger, healthier, and living alone was also found to be associated with higher failure to pay insurance premiums in Switzerland [19], and could partly support our findings of either delayed healthcare or lack of healthcare needs in groups with these attributes. Reich et al. showed that persons of lower socioeconomic status, who receive social assistance, generally have higher intensity and cost of healthcare [20], which parallels our weak association of no HIC among individuals from high Swiss-SEP regions. Similarly as in the case of mortality, the use of ecological instead of individual SEP may be a reason for weaker association [17]. The Reich et al. study also identified higher rates of social assistance among those suffering from psychological disorders and psychoses (identified using pharmaceutical cost groups) [20], which supports the idea that these groups might be more economically vulnerable. Mental illness has been associated with problems in paying medical bills [21] and forgone medical and prescription care [22],

and these findings parallel results of this study. Finally, a review of international data (excluding Switzerland) estimated that around 23% of persons had no contact with primary care in the last year of life, with lower figures for women than men and for older than younger persons [23]. Our estimates indicate lower proportions, potentially suggesting higher use of healthcare; however, we included any MHI claim, which might lead to an overestimate.

Implications

Our results suggest that persons with mental and behavioural disorders, those who are prone to suicide, and persons who are unmarried might be more likely to be unable to identify health needs, fail to seek needed healthcare, or to some degree be less able to handle healthcare-related administrative tasks. Married persons, in contrast, might be more likely to have their HIC submitted after death by a spouse. Future research should therefore explore why healthcare is not utilized by particular groups. Researchers and policy makers also should be aware that analyses based only on persons having HIC might have a differential bias that misses certain groups; studies of healthcare costs might choose to log-transform the outcomes and exclude persons with zero cost [24]. Although the lack of cost seems to be rare phenomenon in EOL studies, other aspects of healthcare use might be more affected.

Limitations

As is also true of other HIC-based studies, this analysis had limited access to individual-level characteristics that could potentially explain observed patterns. For instance, morbidity, functional status, patient preferences, chosen deductibles, and individual socioeconomic position are not available in HIC data yet could influence lack of HIC and should be explored. An estimated 2–3% of all claim invoices are paid by patients directly and never reach their insurers [7]. This could potentially have had




Fig. 2 Proportions (left panel) and adjusted odds ratios (AOR, right panel) and their 95% confidence intervals (CI) of lack of health insurance claims. AORs from sex-stratified, multivariable logistic models with robust standard errors. Lack of CI in the left panel indicates very narrow CI. Lack of CI in the right panel indicates reference category (for instance CVD). Dashed lines in the left panel represent sex-specific means. Abbreviations: CVD, cardiovascular diseases; Mental & behav., Mental and behavioural disorders; Nat., nationality; Civ., civil status at the time of death; Urb., level of urbanization; Lan., language region; SSEP, Swiss neighbourhood index of socioeconomic position [13] (in quintiles). Attribution of causes of death follows ICD-10 coding

an impact on our results, particularly among persons with high deductibles or those dying from causes of death not associated with frequent or expensive health-care use. Switzerland has a relatively high share of out-of-pocket payments in the last year of life [10], and persons with high deductibles are known to incur lower costs [25]. Though the last year of life is used frequently [9], it is still an arbitrary time frame [26]. Our previous

analyses indicate that it performs similarly to the last 3 months of life when analysing costs [12].

Conclusions

Particular groups might be more likely to not seek care or not report health insurance costs to insurers. Researchers should be aware of this aspect of health insurance data and account for persons who lack HIC.

Additional files

Additional file 1: Adjusted odds ratios (AOR, right panel) and their 95% confidence intervals (CI) of lack of health insurance claims (as presented on Fig. 2). AORs from sex-stratified, multivariable logistic models with robust standard errors. Abbreviations: CVD, cardiovascular diseases; Mental & behave., Mental and behavioural disorders; Nat., nationality; Civ., civil status at the time of death; Urb., level of urbanization; Lan., language region; Swiss-SEP, Swiss neighbourhood index of socioeconomic position [13]. Attribution of causes of death follows ICD-10 coding. (XLS 33 kb)

Additional file 2: Adjusted odds ratios (AOR) and their 95% confidence intervals (CI) of lack of health insurance claims across selected main causes of death. AORs from sex-stratified, multivariable logistic models with robust standard errors. Lack of CI indicates reference category (for instance CVD). There were no events (having no HIC) among females aged 19–50 dying of mental and behavioural disorders. Abbreviations: CVD, cardiovascular diseases; Mental & behav., Mental and behavioural disorders; Nat., nationality; Civ., civil status at the time of death; Urb., level of urbanization; Lan., language region; Swiss-SEP, Swiss neighbourhood index of socioeconomic position [13]. Attribution of causes of death follows ICD-10 coding. (PDF 82 kb)

Abbreviations

AOR: Adjusted odds ratio; CI: Confidence interval; CVD: Cardiovascular diseases; EOL: End of life; HIC: Health insurance claims; ICD-10: 10th revision of the International statistical classification of diseases and related health problems; MHI: Mandatory health insurance; Swiss-SEP: Swiss neighbourhood index of socioeconomic position

Acknowledgements

We are extremely grateful to the insurance companies that were part of the study: CSS, Groupe Mutuel, Helsana, Sanitas, SWICA, and Visana. They provided us with the data used in the study and offered helpful comments on data management. Initial funding and data were obtained by the late André Busato, who passed away on 12 November 2013. We thank Christopher Ritter for his editorial contributions.

Funding

This work was supported by Swiss National Science Foundation funding ('NRP 67 End of Life' project number: 139333; grant number 406740_139333) and by a joint grant of FMH Swiss Medical Association, KKA Konferenz der Kantonalen Ärztgesellschaften, and Newindex AG. The funding bodies influenced neither the study design; the collection, analysis, and interpretation of data; the writing of the report; nor the decision to submit the article for publication.

Availability of data and materials

This study relied upon access to claims data granted by six insurance companies, which own the data. Both privacy protections and contractual agreements with the data providers prohibit us from sharing the data.

Authors' contributions

RP & MZ jointly designed the study. RP & MZ jointly managed and analysed the data and interpreted results. KS performed record linkage. RP wrote the first draft of the manuscript. VwW, OR, XL, MM, AES, CB, KS, DCG, ME, KMCG commented on and critically reviewed the drafted manuscript. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Ethical approval was obtained from the Ethics Committee of the Canton of Bern. Consent to participate was not needed as this retrospective study used routinely collected data.

Consent for publication

Not applicable.

Competing interests

All authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Institute of Social and Preventive Medicine, University of Bern, Finkenhubelweg 11, 3012 Bern, Switzerland. ²Epidemiology, Biostatistics & Prevention Institute, University of Zürich, Hirschengraben 84, 8001 Zurich, Switzerland. ³Department of Health Sciences, Helsana Insurance Group, Palmstrasse 26b, 8401 Winterthur, Switzerland. ⁴SWICA Gesundheitsorganisation, sante24, Winterthur, Switzerland. ⁵University Center for Palliative Care, Inselspital, Bern University Hospital, University of Bern, Freiburgstrasse 28, 3010 Bern, Switzerland. ⁶Department of Geriatrics, Inselspital, Bern University Hospital, and University of Bern, Freiburgstrasse, 3010 Bern, Switzerland. ⁷The Dartmouth Institute of Health Policy & Clinical Practice, Lebanon, NH, USA. ⁸Section of Geriatrics, Boston University Medical Center, Boston, MA, USA.

Received: 26 October 2016 Accepted: 7 March 2018

Published online: 14 March 2018

References

- Riley GF. Administrative and claims records as sources of health care cost data. *Med Care*. 2009;47:551–5.
- Davies JM, Gao W, Sleeman KE, Lindsey K, Murtagh FE, Teno JM, et al. Using routine data to improve palliative and end of life care. *BMJ Support Palliat Care*. 2016;6:257–62.
- Berlin C, Panczak R, Egger M. Versorgungsforschung mit Routinedaten in der Schweiz : eine Herausforderung. *Schweizerische Ärztezeitung*. 2014;95:1892–5.
- von Wyl V, Telser H, Weber A, Fischer B, Beck K. Cost trajectories from the final life year reveal intensity of end-of-life care and can help to guide palliative care interventions. *BMJ Support Palliat Care*. Published Online First: 15 October 2015. <https://doi.org/10.1136/bmjspcare-2014-000784>.
- Pletscher M, Mattli R, Von Wyl A, Reich O, Wieser S. The societal costs of schizophrenia in Switzerland. *J Ment Health Policy Econ*. 2015;18:93–103.
- Blozik E, Rapold R, Reich O. Prescription of potentially inappropriate medication in older persons in Switzerland: does the dispensing channel make a difference? *Risk Manag Healthc Policy*. 2015;8:73–80.
- Reich O, Signorell A, Busato A. Place of death and health care utilization for people in the last 6 months of life in Switzerland: a retrospective analysis using administrative data. *BMC Health Serv Res*. 2013;13:116.
- Billi-Andorno N, Zeltner T. Individual responsibility and community solidarity - the Swiss health care system. *N Engl J Med*. 2015;373:2193–7.
- Economist Intelligence Unit. The quality of death: Ranking end-of-life care across the world [Internet]. 2010. Available from: http://www.lienfoundation.org/sites/default/files/qod_index_2.pdf
- Penders YW, Rietjens J, Albers G, Croezen S, Van den Block L. Differences in out-of-pocket costs of healthcare in the last year of life of older people in 13 European countries. *Palliat Med*. 2016;31(1):42–52.
- Langton JM, Blanch B, Drew AK, Haas M, Ingham JM, Pearson S-A. Retrospective studies of end-of-life resource utilization and costs in cancer care using health administrative data: a systematic review. *Palliat Med*. 2014;28:1167–96.
- Panczak R, Luta X, Maessen M, Stuck AE, Berlin C, Schmidlin K, et al. Regional variation of cost of Care in the Last 12 months of life in Switzerland small-area analysis using insurance claims data. *Med Care*. 2017; 55:155–63.
- Federal Office of Public Health FOPH. Statistik der obligatorischen Krankenversicherung 2010 [statistics on compulsory health insurance]. 2010. Available from: <https://www.bag.admin.ch/bag/de/home/service/zahlen-fakten/statistiken-zur-krankenversicherung/statistik-der-obligatorischen-krankenversicherung.html>
- Thomson S, Busse R, Crivelli L, Van de Ven W, Van de Voorde C. Statutory health insurance competition in Europe: a four-country comparison. *Health Policy*. 2013;109:209–25.
- De Pietro C, Camenzind P, Sturmy I, Crivelli L, Edwards-Garavoglia S, Spranger A, et al. Switzerland: Health System Review. *Heal Syst Transit*. 2015;17:1–288.
- Teno JM, Gozalo PL, Bynum JPW, Leland NE, Miller SC, Morden NE, et al. Change in end-of-life care for Medicare beneficiaries. Site of death, place of care, and health care transitions in 2000, 2005, and 2009. *JAMA*. 2013;309:470–7.

17. Panczak R, Galobardes B, Voorpostel M, Spoerri A, Zwahlen M, Egger M. A Swiss neighbourhood index of socioeconomic position: development and association with mortality. *J Epidemiol Community Health*. 2012;66:1129–36.
18. Wachterman MW, Sommers BD. The impact of gender and marital status on end-of-life care: evidence from the National Mortality Follow-Back Survey. *J Palliat Med*. 2006;9:343–53.
19. von Wyl V, Beck K. Failure to pay for social health insurance premiums: acts of protest or desperation? *Prev Med Rep*. 2015;2:127–33.
20. Reich O, Wolffers F, Signorell A, Blozik E. Health care utilization and expenditures in persons receiving social assistance in 2012 evidence from Switzerland. *Glob J Health Sci*. 2014;7:1–11.
21. Herman PM, Rissi JJ, Walsh ME. Health insurance status, medical debt, and their impact on access to care in Arizona. *Am J Public Health*. 2011;101:1437–43.
22. Baughman KR, Burke RC, Hewit MS, Sudano JJ, Meeker J, Hull SK. Associations between difficulty paying medical bills and forgone medical and prescription drug care. *Popul Health Manag*. 2015;0:1–9.
23. Luoma JB, Martin CE, Pearson JL. Contact with mental health and primary care providers before suicide: a review of the evidence. *Am J Psychiatry*. 2002;159:909–16.
24. Manning WG, Mullahy J. Estimating log models: to transform or not to transform? *J Health Econ*. 2001;20:461–94.
25. Bähler C, Huber CA, Brüngger B, Reich O. Multimorbidity, health care utilization and costs in an elderly community-dwelling population: a claims data based observational study. *BMC Health Serv Res*. 2015;15:23.
26. Bach PB, Schrag D, Begg CB. Resurrecting treatment histories of dead patients. A study design that should be laid to rest. *JAMA*. 2004;292:2765–70.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

