Annihilators in JB-algebras

By M. BATTAGLIA

Universität Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland

(Received 9 October 1989; revised 11 January 1990)

Abstract

Orthogonality is defined for all elements in a JB-algebra and Topping's results on annihilators in JW-algebras are generalized to the context of JB- and JBW-algebras. A pair (a, b) of elements in a JB-algebra A is said to be orthogonal provided that $a^2 \circ b$ equals zero. It is shown that this relation is symmetric. The annihilator $S^\perp$ of a subset S of A is defined to be the set of elements a in A such that, for all elements s in S , the pair (s, a) is orthogonal. It is shown that the annihilators are closed quadratic ideals and, if A is a JBW-algebra, a subset I of A is a w^* -closed quadratic ideal if and only if I coincides with its biannihilator $I^{\perp\perp}$. Moreover, in a JBW-algebra A the formation of the annihilator of a w^* -closed quadratic ideal is an orthocomplementation on the complete lattice of w^* -closed quadratic ideals which makes it into a complete orthomodular lattice. Further results establish a connection between ideals, central idempotents and annihilators in JBW-algebras.

1. Introduction

Topping[7] studied the w^* -closed Jordan algebras of bounded self-adjoint operators, the so-called JW-algebras. A generalization of these objects was given by Alfsen, Shultz and Størmer[1] by introducing JB- and JBW-algebras.

It is the aim of the present paper to generalize Topping's theory on annihilators in JW-algebras to the context of JB- and JBW-algebras. Annihilators in JW-algebras are defined in terms of the underlying associative operator product and therefore there is no obvious extension of this definition to JB-algebras.

Let A be a JB-algebra. A pair (a, b) of elements in A is said to be orthogonal provided that $a^2 \circ b$ equals zero. The relation of orthogonality is symmetric: this is shown in Section 3.

In Section 4 the annihilator $S^\perp$ of a subset S of the JB-algebra A is defined as the set of elements a in A such that, for all elements s in S , the pair (s, a) is orthogonal. It is shown that the annihilator of a general subset S of a JB-algebra A is a closed quadratic ideal.

In Section 5 it is shown that a subset I of a JBW-algebra A is a w^* -closed quadratic ideal if and only if I coincides with its biannihilator $I^{\perp\perp}$. It also follows that the formation of the annihilator of a w^* -closed quadratic ideal in a JBW-algebra A is an orthocomplementation on the complete lattice of w^* -closed quadratic ideals which makes it into a complete orthomodular lattice. Further results establish a connection between ideals, central idempotents and annihilators in JBW-algebras.

2. Preliminaries

A (real) *Jordan algebra* A is a real vector space together with a bilinear mapping $\circ: A^2 \rightarrow A$, called the *Jordan product*, satisfying, for all elements a and b in A ,

$$(i) \ a \circ b = b \circ a, \quad (ii) \ (a^2 \circ b) \circ a = a^2 \circ (b \circ a).$$

In a real associative algebra $(A, \cdot)$ a canonical Jordan product can be defined, for all elements a and b in A , by

$$a \circ b := \frac{1}{2}(a \cdot b + b \cdot a).$$

A Jordan algebra which is isomorphic to a subalgebra of a Jordan algebra formed in this way is said to be special.

Let A be a Jordan algebra and let a be an element in A . Then a linear mapping $U_a: A \rightarrow A$ is defined, for all elements b in A , by

$$U_a b := 2(a \circ b) \circ a - a^2 \circ b.$$

A subspace I of a Jordan algebra A is said to be an ideal, resp. a quadratic ideal, provided that $I \circ A \subseteq I$, resp. $U_I A \subseteq I$. An element a in a Jordan algebra A is called central provided that, for all elements b and c in A , $(a \circ b) \circ c$ equals $a \circ (b \circ c)$. The set $Z(A)$ of all central elements in A is called the centre of A . Notice that $Z(A)$ is a subalgebra.

A Jordan algebra A with unit e together with a complete norm $\|\cdot\|$ such that, for all elements a and b in A ,

$$(i) \ \|a \circ b\| \leq \|a\| \|b\|, \quad (ii) \ \|a\|^2 \leq \|a^2 + b^2\|$$

is called a *JB-algebra*. Notice that in a JB-algebra the Jordan product is jointly continuous. The set A^2 of all squares of elements in a JB-algebra A is a proper convex cone. Therefore A^2 induces an ordering $\leq$ on A . For all elements a in a JB-algebra A , the mapping $U_a: A \rightarrow A$ is *positive*, i.e. $U_a A^2 \subseteq A^2$.

A JB-algebra A which is a Banach space dual is said to be a *JBW-algebra*. Notice that in a JBW-algebra the Jordan product is separately w^* -continuous and that a w^* -closed subalgebra of a JBW-algebra has a unit.

An element p in a JB-algebra A is called *idempotent* if $p^2 = p$. Let $U(A)$ denote the collection of idempotents in A . With respect to the ordering $\leq$ and the orthocomplementation defined, for all idempotents p in A , by $p' := e - p$, $U(A)$ becomes an orthomodular poset with 0 as the least and e as the greatest element. Let p and q be idempotents in A . Then $p \leq q'$ if and only if $p \circ q$ equals zero and $p \leq q$ if and only if $p \circ q$ equals p if and only if $U_q U_p$ equals U_p . In a JBW-algebra A the triple $(U(A), \leq, ')$ is a complete orthomodular lattice.

Let a be an element in a JBW-algebra A . Then there exists a least idempotent $r(a)$ in A such that $r(a) \circ a$ equals a . The element $r(a)$ is called the *support idempotent* of a . Notice that $r(a)$ is the unit in the smallest w^* -closed subalgebra $W(a)$ of A containing a . For every idempotent p in a JBW-algebra A there exists a least central idempotent $c(p)$ in A majorizing p . The element $c(p)$ is called the *central support* of p .

Edwards[2] has shown that every w^* -closed quadratic ideal in a JBW-algebra A is of the form $U_p A$, for a suitable idempotent p in A , and conversely, for an

idempotent p in A , $U_p A$ is a w^* -closed quadratic ideal. Moreover, $U_p A$ is an ideal if and only if p is central.

For the details of the theory of JB- and JBW-algebras the reader is referred to [1, 2, 3, 4, 6].

3. Orthogonality in JB-algebras

Let A be a JB-algebra. A pair (a, b) of elements in A is said to be *orthogonal*, denoted by $a \perp b$, if $a^2 \circ b = 0$. Hence a pair (p, q) of idempotents is orthogonal if and only if $p \leq q'$.

Notice that, for elements a and b in a JB-algebra A , $a \circ b = 0$ does not imply $a^2 \circ b = 0$. In the JB-algebra M_3^s , i.e. the Jordan algebra of hermitian 3×3 matrices with Cayley number entries, consider the elements

$$a = \begin{pmatrix} 0 & i & 0 \\ -i & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad b = \begin{pmatrix} 0 & -k & 0 \\ k & 0 & k \\ 0 & -k & 0 \end{pmatrix}.$$

Since $ik = -j$, $ki = j$ and $i^2 = -1$, $a \circ b = 0$ but $a^2 \circ b = 2b \neq 0$.

It is the aim of this section to show that orthogonality is a symmetric relation and that $a \perp b$ entails $a^n \circ b = 0$ for all natural numbers n .

LEMMA 3.1. *Let A be a JB-algebra. Let (a, b) be an orthogonal pair of elements in A . Then $a^2 \circ b^2$ is a positive element in A .*

Proof. By [5], theorem I-10, the smallest subalgebra B of A containing a^2 and b is special, i.e. there is an isomorphism j from B into an associative algebra $(J, \cdot)$ equipped with the canonical Jordan product. Then

$$j(a)^2 \cdot j(b) + j(b) \cdot j(a)^2 = 0.$$

Multiplying the equation on the left and right, respectively, by $j(b)$ in the underlying associative product we obtain

$$j(a)^2 \cdot j(b)^2 = j(b)^2 \cdot j(a)^2.$$

Therefore a^2 and b^2 generate an associative subalgebra in A . By the continuity of the Jordan product, the smallest closed subalgebra C of A containing a^2 and b^2 is associative as well. Since a^2 and b^2 are positive elements in A , there exist, by [2], lemma 2.1, elements c and d in C such that $a^2 = c^2$ and $b^2 = d^2$. By the associativity of C , we have $(c \circ d)^2 = a^2 \circ b^2$.

LEMMA 3.2. *Let A be a JB-algebra. Let (a, b) be an orthogonal pair of elements in A . Then*

$$a^2 \circ b^2 = U_b a^2 = U_a b^2 = U_a b = a \circ b = 0.$$

Proof. From $a \perp b$ it follows that $U_b a^2 = -a^2 \circ b^2$. Then, by Lemma 3.1 and the positivity of the operator U_b , both $U_b a^2$ and $a^2 \circ b^2$ are zero. The remaining assertions follow from the identities

$$(U_a b^2)^2 = U_a U_b U_b a^2, \quad (U_a b)^2 = U_a U_b a^2, \\ (a \circ b)^2 = \frac{1}{2} b \circ U_a b + \frac{1}{4} U_a b^2 + \frac{1}{4} U_b a^2.$$

COROLLARY 3.3. *Let A be a JB-algebra and let a and b be elements in A . Then the following conditions are equivalent:*

- (i) $a \perp b$; (ii) $a \circ b = 0$ and $a^2 \circ b = 0$; (iii) $a^n \circ b = 0$ for all $n \in \mathbb{N}$.

Proof. (i) $\Rightarrow$ (ii). See Lemma 3.2.

(ii) $\Rightarrow$ (iii). By [5], theorem I-10, the smallest subalgebra B of A containing a and b is special, i.e. there is an isomorphism j from B into an associative algebra $(J, \cdot)$ equipped with the canonical Jordan product. Suppose now that $a^{n-1} \circ b = 0$. Then

$$j(a)^{n-1} \cdot j(b) + j(b) \cdot j(a)^{n-1} = 0.$$

Multiplying the equation on the left and right, respectively, by $j(a)$ in the underlying associative product and adding the results we get

$$j(a)^n \cdot j(b) + j(b) \cdot j(a)^n + j(a)^{n-1} \cdot j(b) \cdot j(a) + j(a) \cdot j(b) \cdot j(a)^{n-1} = 0.$$

From this and the assumption it follows that $a^n \circ b = 0$.

(iii) $\Rightarrow$ (i). This is obvious.

We now proceed to the main result of this section.

PROPOSITION 3.4. *Let A be a JB-algebra. Let a and b be elements in A . Then $a \perp b$ if and only if $b \perp a$.*

Proof. Suppose that $a \perp b$. By Lemma 3.2 and the positivity of U_a and U_b ,

$$0 \leq U_a b^4 \leq \|b^2\| U_a b^2 = 0.$$

Therefore $U_a b^4$ equals zero and the assertion now follows from Lemma 3.2 and the identity

$$(a \circ b^2)^2 = \frac{1}{4} U_a b^4 + \frac{1}{4} U_{b^2} a^2 + \frac{1}{2} b^2 \circ U_a b^2.$$

4. Annihilators in JB-algebras

Let A be a JB-algebra and let S be a subset of A . The *annihilator* $S^\perp$ of S is defined to be

$$S^\perp := \{a \in A; s \perp a \text{ for all } s \in S\}.$$

The set $S^{\perp\perp}$ is referred to as the *biannihilator* of S .

LEMMA 4.1. *Let A be a JB-algebra, let S and T be subsets of A and let $(S_i)_{i \in I}$ be a family of subsets of A . Then*

- (i) $S \subseteq T$ implies $T^\perp \subseteq S^\perp$; (ii) $S \subseteq S^{\perp\perp}$;
 (iii) $(\bigcup_{i \in I} S_i)^\perp = \bigcap_{i \in I} (S_i)^\perp$; (iv) $S^{\perp\perp\perp} = S^\perp$;
 (v) $S^\perp$ is a closed subspace of A ; (vi) $S^\perp = (\text{cl } S)^\perp = (\text{lin } S)^\perp$.

Proof. (i), (ii), (iii) and (iv). These are immediate.

(v) By bilinearity and continuity of the Jordan product, $S^\perp$ is a closed subspace.

(vi) By (v), $\text{cl } S$ and $\text{lin } S$ all are contained in $S^{\perp\perp}$ and the assertion follows, by (i).

THEOREM 4.2. *Let A be a JB-algebra and let S be a subset of A . Then the annihilator $S^\perp$ of S is a closed quadratic ideal.*

Proof. Let a, b and c be elements in A such that the pair (a, b) is orthogonal. By the positivity of the operators U_a , U_b and U_c and Lemma 3.2, it follows that

$$0 \leq U_a U_b U_c b^2 \leq \|b^2\| U_a U_b c^2 \leq \|b^2\| \|c^2\| U_a b^2 = 0.$$

Therefore $U_a U_b U_c b^2 = 0$. Since $0 \leq a^4 \leq \|a^2\| a^2$, we conclude, by Lemma 3.2, that

$$0 \leq U_b a^4 \leq \|a^2\| U_b a^2 = 0.$$

Therefore $U_b a^4 = 0$. By Lemma 3.2 and the identity

$$(a^2 \circ U_b c)^2 = \frac{1}{4} U_a U_b U_c b^2 + \frac{1}{4} U_b U_c U_b a^4 + \frac{1}{2} a^2 \circ U_b U_c U_b a^2,$$

$a^2 \circ U_b c$ equals zero. Therefore by Lemma 4.1 (v), the assertion follows.

PROPOSITION 4.3. *Let A be a JB-algebra and let p and q be idempotents in A . Then*

(i) $\{p\}^\perp = U_p A = (U_p A)^\perp$, and (ii) $p \leq q$ if and only if $\{p\}^\perp \supseteq \{q\}^\perp$.

Proof. (i) Let a be an element in $U_p A$ and let b be an element in $U_{p'} A$. Then, by (2.66) and (2.69) from [4], we have $a^2 \circ b = 0$. Therefore $U_{p'} A \subseteq (U_p A)^\perp$. By Lemma 4.1 (i), we have $(U_p A)^\perp \subseteq \{p\}^\perp$. Now let $b \in \{p\}^\perp$; then $p' \circ b = b$. Therefore $b \in U_{p'} A$.

(ii) Since $p \leq q$ if and only if $q' \leq p'$, (ii) follows from (i) and $U_{q'} A \subseteq U_{p'} A$.

5. Annihilators in JBW-algebras

In JBW-algebras Lemma 4.1 (vi) and Theorem 4.2 can be improved.

LEMMA 5.1. *Let A be a JBW-algebra and let S be a subset of A . Then*

(i) $S^\perp$ is a w^* -closed quadratic ideal in A , and

(ii) $S^\perp = (\text{cl } S)^\perp = (w^*\text{-cl } S)^\perp = (\text{lin } S)^\perp$.

Proof. By Theorem 4.2 and separate w^* -continuity of the Jordan product, (i) follows. The proof of (ii) is similar to that of Lemma 4.1 (vi).

We are now in a position to identify the subsets S of a JBW-algebra which coincide with their biannihilator $S^{\perp\perp}$.

THEOREM 5.2. *Let A be a JBW-algebra. A subset S of A is a w^* -closed quadratic ideal if and only if S coincides with its biannihilator $S^{\perp\perp}$.*

Proof. By Lemma 5.1, a subset which coincides with its biannihilator is a w^* -closed quadratic ideal. Since every w^* -closed quadratic ideal is of the form $U_p A$ for a suitable idempotent p in A , by [2], theorem 2.3, it follows from Proposition 4.3 that it coincides with its biannihilator.

COROLLARY 5.3. *Let A be a JBW-algebra, let $U(A)$ be the collection of idempotents in A and let $\mathcal{T}(A)$ be the collection of w^* -closed quadratic ideals in A . Let p be an idempotent in A and define $\phi(p) := U_p A$. Then the mapping $\phi: U(A) \rightarrow \mathcal{T}(A)$ is an ortho-order isomorphism from the complete orthomodular lattice $(U(A), \leq, ')$ onto $(\mathcal{T}(A), \subseteq, ^\perp)$.*

Proof. By Theorem 5.2, Proposition 4.3 and [2], corollary 2.5, the assertion follows.

In order to identify the annihilator of a general subset of a JBW-algebra we begin by examining annihilators of singleton sets.

LEMMA 5.4. *Let A be a JBW-algebra and let a and b be elements in A . Then the following conditions are pairwise equivalent: (i) $a \perp b$; (ii) $a \perp r(b)$; (iii) $r(a) \perp r(b)$.*

Proof. (i) $\Rightarrow$ (ii) $\Rightarrow$ (iii). Since $W(b)$ coincides with the w^* -closure of the linear hull of the set $\{b^n; n \in \mathbb{N}\}$ and since the Jordan product is separately w^* -continuous, this follows from Corollary 3.3.

(iii) $\Rightarrow$ (i). If $r(a) \circ r(b)$ equals zero then $r(a) \leq r(b)'$. Therefore $p \leq r(b)'$ for all idempotents p in $W(a)$, and it follows that $r(b) \leq p'$. Hence $p \circ q = 0$ for all idempotents q in $W(b)$ and all idempotents p in $W(a)$, and, by spectral theory, $a \perp b$.

COROLLARY 5.5. *Let A be a JBW-algebra and let a be an element in A . Then $r(a)$ is the unique idempotent in A such that $\{a\}^\perp = \{r(a)\}^\perp$.*

Proof. By Lemma 5.4, $a \perp b$ if and only if $r(a) \perp b$. Hence $\{a\}^\perp = \{r(a)\}^\perp$. Uniqueness follows from Corollary 5.3.

THEOREM 5.6. *Let A be a JBW-algebra and let S be a non-empty subset of A . Let $W(S)$ denote the smallest w^* -closed subalgebra of A containing S and let p be the unit of $W(S)$. Then $S^\perp = U_p A$.*

Proof. By Lemma 4.1 (iii), Corollary 5.5, Proposition 4.3 and Corollary 5.3,

$$S^\perp = \bigcap_{a \in S} \{a\}^\perp = \bigcap_{a \in S} \{r(a)\}^\perp = \bigcap_{a \in S} U_{r(a)'} A = U_r A,$$

where r denotes $\bigvee_{a \in S} r(a)$. It remains to prove that $r = p$. Clearly $r \leq p$. Conversely, $r(a) \leq r$, for all elements a in S . Therefore $W(S) \subseteq U_r A$ and $p \leq r$ follows.

We present two applications of the previously developed theory.

COROLLARY 5.7. *Let A be a JBW-algebra, let S be a subset of A and let $W(S)$ denote the smallest w^* -closed subalgebra of A containing S . Then*

$$S^\perp = \{a \in A; W(S) \circ a = \{0\}\}.$$

Proof. Let p be the unit of $W(S)$. Then $W(S) \subseteq U_p A$ and, by Theorem 5.6, $S^\perp$ equals $U_p A$. Hence, by [4], theorem 2.6.5, we have $W(S) \circ S^\perp = \{0\}$. Conversely, if $W(S) \circ a = \{0\}$ then $b^2 \circ a = 0$ for all b in S .

COROLLARY 5.8. *Let A be a JBW-algebra and let a and b be elements in A . Then $r(a+b) \leq r(a) \vee r(b)$.*

Proof. From $r(a) \leq r(a) \vee r(b)$, $r(b) \leq r(a) \vee r(b)$, Proposition 4.3 and Corollary 5.5 it follows that $\{r(a) \vee r(b)\}^\perp \subseteq \{a\}^\perp \cap \{b\}^\perp$. Therefore, for all z in $\{r(a) \vee r(b)\}^\perp$, both $a \circ z^2$ and $b \circ z^2$ equal zero. Hence $(a+b) \circ z^2 = 0$ and

$$z \in \{a+b\}^\perp = \{r(a+b)\}^\perp,$$

by Corollary 5.5. Therefore $\{r(a) \vee r(b)\}^\perp \subseteq \{r(a+b)\}^\perp$ and an application of Proposition 4.3 proves the claim.

The final results generalize [7], proposition 5 and section 7.

PROPOSITION 5.9. *Let A be a JBW-algebra and let I be an ideal in A . Then there exists a central idempotent f in A such that $I^\perp = U_f A$.*

Proof. By the separate w^* -continuity of the Jordan product, $w^*\text{-cl } I$ is an ideal. By [2], theorem 2.3, there is a central idempotent f in A such that $w^*\text{-cl } I = U_f A$. Then f' is a central idempotent in A and $I^\perp = U_{f'} A$, by Theorem 5.6 and Lemma 5.1.

PROPOSITION 5.10. *Let A be a JBW-algebra and let a be an element in A . Let (a) be the smallest ideal containing a and let $r(a)$ be the support idempotent of a . Then the central support $c(r(a))$ of $r(a)$ is the unique (central) idempotent such that $\{c(r(a))\}^\perp = (a)^\perp$.*

Proof. By Proposition 5.9, there is a central idempotent f in A such that

$$(a)^\perp = \{f\}^\perp = U_{f'}A.$$

Hence $f \circ a = a$. Therefore $c(r(a)) \leq f$. Since $U_{c(r(a))}A$ is an ideal containing a , it follows that

$$U_{c(r(a))}A = (U_{c(r(a))}A)^\perp \subseteq (a)^\perp = U_{f'}A.$$

Therefore $c(r(a))' \leq f'$.

COROLLARY 5.11. *Let A be a JBW-algebra and let p and q be idempotents in A . Let $c(p)$ and $c(q)$ be the central supports of p and q , respectively. Let (p) and (q) be the smallest ideals containing p and q , respectively. Then the following conditions are equivalent:*

- (i) $c(p) \perp c(q)$; (ii) $c(p) \perp q$; (iii) $p \perp c(q)$; (iv) $q \in (p)^\perp$; (v) $p \in (q)^\perp$.

Proof. (i) $\Leftrightarrow$ (ii) $\Leftrightarrow$ (iii). These are immediate.

(ii) $\Leftrightarrow$ (iv). Since $U_{c(p)}A$ is an ideal containing p , $c(p) \perp q$ entails

$$q \in U_{c(p)}A = (U_{c(p)}A)^\perp \subseteq (p)^\perp.$$

Conversely, let q be an element in $(p)^\perp$. Then, by Proposition 5.10, q is contained in $\{c(p)\}^\perp$. Therefore $c(p) \perp q$.

(iii) $\Leftrightarrow$ (v). The proof is similar to the one above.

PROPOSITION 5.12. *Let A be a JBW-algebra with centre $Z(A)$ and let S be a subset of $Z(A)$. Then $S^\perp$ is a w^* -closed ideal in A .*

Proof. By Lemma 5.1, $S^\perp$ is a w^* -closed subspace of A . Let $a \in A$ and $b \in S^\perp$. It is enough to show $c \perp a \circ b$, for each element c in S . Since c is central, c^2 also is central and it follows that

$$c^2 \circ (b \circ a) = (c^2 \circ b) \circ a = 0.$$

The author wishes to thank G. T. Rüttimann for encouragement and valuable discussions and the Swiss National Science Foundation for supporting this research.

REFERENCES

- [1] E. M. ALFSEN, F. W. SCHULTZ and E. STØRMER. A Gelfand-Neumark theorem for Jordan algebras. *Adv. in Math.* **28** (1978), 11–56.
- [2] C. M. EDWARDS. Ideal theory in JB-Algebras. *J. London Math. Soc.* (2) **16** (1977), 507–513.
- [3] C. M. EDWARDS. On the facial structure of a JB-algebra. *J. London Math. Soc.* (2) **19** (1979), 335–344.
- [4] H. HANCHE-OLSEN and E. STØRMER. *Jordan Operator Algebras*. Monographs Stud. Math. no. 21 (Pitman, 1984).
- [5] N. JACOBSON. *Structure and representations of Jordan algebras*. Amer. Math. Soc. Colloq. Publ. no. 39 (American Mathematical Society, 1968).
- [6] F. W. SHULTZ. On normed Jordan algebras which are Banach dual spaces. *J. Funct. Anal.* **31** (1979), 360–376.
- [7] D. M. TOPPING. *Jordan algebras of self-adjoint operators*. Memoirs Amer. Math. Soc. no. 53 (American Mathematical Society, 1965).