DEGENERATIONS FOR SELF-INJECTIVE ALGEBRAS OF TREECLASS D_n

ROBERT AEHLE

ABSTRACT

Let Λ be a connected representation finite selfinjective algebra. According to G. Zwara the partial orders $\leq_{\text{ext}}$ and $\leq_{\text{deg}}$ on the isomorphism classes of d -dimensional Λ -modules are equivalent if and only if the stable Auslander–Reiten quiver Γ_Λ of Λ is not isomorphic to $\mathbb{Z}D_{3m}/\tau^{2m-1}$ for all $m \geq 2$. The paper describes all minimal degenerations $M \leq_{\text{deg}} N$ with $M \not\leq_{\text{ext}} N$ in the case when $\Gamma_\Lambda \cong \mathbb{Z}D_{3m}/\tau^{2m-1}$ for some $m \geq 2$.

1. Introduction

1.1. The affine variety $\text{mod}_d \Lambda$

Let k be an algebraically closed field and Λ be a finite dimensional associative k -algebra with unit. We denote by $\text{mod} \Lambda$ the category of finitely generated Λ -left-modules. A d -dimensional Λ -module M is the vectorspace k^d together with a multiplication by Λ from the left.

Now let $\lambda_1 = 1, \lambda_2, \dots, \lambda_n$ be a k -basis of Λ . Then $\lambda_i \lambda_j = \sum_l a_{ij}^l \lambda_l$ for $i, j = 1, \dots, n$ with the structure constants $a_{ij}^l \in k$. The multiplication of M by λ_i induces an endomorphism of k^d which we can represent by a $d \times d$ matrix over k with respect to the standard basis of k^d . Thus M corresponds to a unique n -tuple of matrices $m = (E, m_2, \dots, m_n) \in (\text{Mat}_{d \times d}(k))^n$, where E denotes the identity matrix, and such an n -tuple m with $m_1 = E$ corresponds to a d -dimensional Λ -module if and only if it satisfies the equations $m_i m_j = \sum_l a_{ij}^l m_l$ for $i, j = 1, \dots, n$. We denote the set of all n -tuples corresponding to a d -dimensional Λ -module by $\text{mod}_d \Lambda$ and we will identify the module with its n -tuple. For each i with $1 \leq i \leq n$ let X^i denote the matrix $(x_{\mu\nu}^i)_{\mu, \nu=1, \dots, d}$. Then $\text{mod}_d \Lambda$ is the zero set of the ideal $I \subset k[x_{\mu\nu}^\xi]$ ($\mu, \nu = 1, \dots, d; \xi = 1, \dots, n$), where I is generated by the components of the matrices $X^i X^j - \sum_l a_{ij}^l X^l$ for $i, j = 1, \dots, n$. This gives $\text{mod}_d \Lambda$ the structure of an affine variety, which does not have to be irreducible.

The general linear group $\text{GL}_d(k)$ acts on $\text{mod}_d \Lambda$ by conjugation, that is to say $g \cdot (m_1, \dots, m_n) = (gm_1 g^{-1}, \dots, gm_n g^{-1})$ for $g \in \text{GL}_d(k)$ and $(m_1, \dots, m_n) \in \text{mod}_d \Lambda$. The orbits under this action are the isomorphism classes of d -dimensional Λ -modules (see [7]). This definition of $\text{mod}_d \Lambda$ depends on the chosen basis of Λ only up to a $\text{GL}_d(k)$ equivariant isomorphism of affine varieties.

1.2. Partial orders on isomorphism classes of $\text{mod}_d \Lambda$

A module N is called a degeneration of M (in symbols $M \leq_{\text{deg}} N$) if N belongs to the Zariski closure of the $\text{GL}_d(k)$ -orbit of M in $\text{mod}_d \Lambda$. Since orbits are irreducible

Received 19 February 2001; revised 17 October 2001.

2000 *Mathematics Subject Classification* 14L30, 16G70.

and open in their closure, this defines a partial order on the set of isomorphism classes of d -dimensional Λ -modules. It is an interesting problem to express the partial order $\leq_{\deg}$ in algebraic terms. There are several articles in this direction, including works by S. Abeasis and A. del Fra [1], K. Bongartz [4, 5], C. Riedtmann [11] and G. Zwara [12, 14], connecting $\leq_{\deg}$ to other partial orders on the isomorphism classes of d -dimensional Λ -modules.

In [15] Zwara gives an alternative description of $\leq_{\deg}$, that is to say $M \leq_{\deg} N$ if and only if there exists a short exact sequence

$$0 \longrightarrow S \longrightarrow S \oplus M \longrightarrow N \longrightarrow 0 \quad (1)$$

for some Λ -module S .

We are concerned with two other partial orders on the isomorphism classes of d -dimensional Λ -modules. The partial order $\leq_{\text{ext}}$ is the transitive closure of the relation $M \leq_{\text{ext}} N$ if there exists a short exact sequence

$$0 \longrightarrow N_1 \longrightarrow M \longrightarrow N_2 \longrightarrow 0 \quad (2)$$

with $N \cong N_1 \oplus N_2$. We take the pullback of the sequence (2) with the canonical projection $N \longrightarrow N_2$ according to the isomorphism $N \cong N_1 \oplus N_2$. This results in a sequence as in (1) with $S = N_1$, so $\leq_{\text{ext}}$ implies $\leq_{\deg}$.

The hom order $\leq$ is the partial order given by $M \leq N$ if and only if

$$[M, X] \leq [N, X]$$

for every Λ -module X , where $[U, V] := \dim_k \text{Hom}_{\Lambda}(U, V)$ for Λ -modules U and V . It follows immediately from (1) and the left-exactness of $\text{Hom}_{\Lambda}(_, X)$ that $\leq_{\deg}$ implies $\leq$. The reverse implication is not true in general. However it holds for representation finite algebras (see [14]) and tame concealed algebras (see [4]).

1.3. Statement of the theorem

We define the Auslander–Reiten quiver Γ_{Λ} of Λ as the quiver whose vertices are representatives of the isomorphism classes of indecomposable Λ -modules. There is an arrow $x \longrightarrow y$ between the vertices x and y if there exists an irreducible morphism from a Λ -module represented by x to one of y . This definition coincides with the usual one in the representation finite case (see [3]) and is appropriate for our consideration.

We denote by τ the Auslander–Reiten translation. It is a bijection from the isomorphism classes of indecomposable non-projective Λ -modules to the isomorphism classes of indecomposable non-injective Λ -modules.

The stable Auslander–Reiten quiver Γ_{Λ}^s of Λ is the full subquiver of Γ_{Λ} containing all the vertices x for which $\tau^n(x)$ is defined for all $n \in \mathbb{Z}$.

Let Λ be connected and selfinjective of finite representation type. C. Riedtmann showed in [8] that the stable Auslander–Reiten quiver Γ_{Λ}^s of Λ is isomorphic to $\mathbb{Z}\Delta/G$ where Δ is one of the Dynkin diagrams A_n, D_n, E_6, E_7, E_8 and G is an admissible automorphism group of $\mathbb{Z}\Delta$. Using the results in [10], [9] and [6] about the category of modules over representation finite selfinjective algebras, G. Zwara showed in [13] that the partial orders $\leq_{\text{ext}}$ and $\leq_{\deg}$ coincide if and only if $\Gamma_{\Lambda}^s \not\cong \mathbb{Z}D_{3m}/\tau^{2m-1}$ for all $m \geq 2$.

We want to investigate the difference between the partial orders $\leq_{\deg}$ and $\leq_{\text{ext}}$ in those exceptional cases. In particular, we want to describe the minimal degenerations

FIGURE 1. $\mathbb{Z}D_{3m}/\tau^{2m-1}$.

$M \leq_{\text{deg}} N$ with $M \not\leq_{\text{ext}} N$ for a connected selfinjective algebra of finite representation type with stable Auslander–Reiten quiver $\mathbb{Z}D_{3m}/\tau^{2m-1}$. A degeneration $M \leq_{\text{deg}} N$ is called minimal if it is a proper degeneration, that is to say $M \not\cong N$, and if there exists no module P with $M \not\cong P \not\cong N$ and $M \leq_{\text{deg}} P \leq_{\text{deg}} N$. It is an interesting question how complicated minimal degenerations are. Some results concerning the complexity of degenerations can be found in [2].

G. Zwara proved in [15, Theorem 4] that for a minimal degeneration $M \leq_{\text{deg}} N$ with $M \not\leq_{\text{ext}} N$ there exist decompositions $M \cong M' \oplus W$ and $N \cong N' \oplus W$ such that N' is indecomposable and $M' \leq_{\text{deg}} N'$ is a minimal degeneration. Therefore it is enough to concentrate on degenerations to indecomposables.

The stable translation quiver $\mathbb{Z}D_{3m}$ has the vertices c_j^i where $i \in \{1, \dots, 3m\}$ and $j \in \mathbb{Z}$. There are arrows $c_j^i \rightarrow c_j^{i+1}$ and $c_j^{i+1} \rightarrow c_{j+1}^i$ for $1 \leq i \leq 3m-2$ and arrows $c_j^{3m-2} \rightarrow c_j^{3m}$ and $c_j^{3m} \rightarrow c_{j+1}^{3m-2}$. The translation is given by $\tau(c_j^i) = c_{j-1}^i$. Thus the vertices c_j^i and c_{j+2m-1}^i are identified in the quotient $\mathbb{Z}D_{3m}/\tau^{2m-1}$. In Figure 1 the stable Auslander–Reiten quiver $\mathbb{Z}D_{3m}/\tau^{2m-1}$ is drawn for m even. Every letter refers to the vertex at its left and the thick diagonal lines indicate the $(2m-1)$ -period of the translation τ .

THEOREM 1.1. *Let Λ be a connected and selfinjective algebra of finite representation type whose stable Auslander–Reiten quiver is isomorphic to $\mathbb{Z}D_{3m}/\tau^{2m-1}$. There exists a proper degeneration $M \leq_{\text{deg}} N$ to the indecomposable Λ -module N if and only if N corresponds to a vertex c_l^s with $m+1 \leq s \leq 2m$. Moreover, the module M is determined by N up to isomorphism.*

2. Preliminaries

We want to represent pairs of modules (M, N) in terms of $\mathbb{Z}$ -valued difference-functions on the set of isomorphism classes of indecomposable modules and to characterize those functions corresponding to pairs (M, N) with $M \leq_{\text{deg}} N$. This will enable us to give a combinatorial proof of the theorem.

We say that the modules M and N are disjoint if they have no common direct summand. We denote by $\bar{M}$ the isomorphism class of the module M and by $\mathcal{S}$ the set of ordered pairs $(\bar{M}, \bar{N})$ such that M and N are disjoint. To every pair $(\bar{M}, \bar{N})$ we associate the function $\delta_{M,N}$ given by $\delta_{M,N}(X) = [N, X] - [M, X]$.

Let $\mu(X, A)$ be the multiplicity of the indecomposable direct summand X in the direct sum decomposition of A . In particular $A \cong \bigoplus_{X: X \text{ indec}} X^{\mu(X,A)}$.

$$\begin{array}{ccc}
& \oplus_{\bar{X}; X \text{ indec.}} \mathbb{Z} \bar{X} & \\
\alpha \nearrow & & \searrow \beta \\
\mathcal{S} & \xrightarrow{\gamma} & (\oplus_{\bar{Y}; Y \text{ indec.}} \mathbb{Z} \bar{Y})^*
\end{array}$$

FIGURE 2.

Let $(\oplus_{\bar{Y}; Y \text{ indec.}} \mathbb{Z} \bar{Y})^* = \text{Hom}_{\mathbb{Z}}(\oplus_{\bar{Y}; Y \text{ indec.}} \mathbb{Z} \bar{Y}, \mathbb{Z})$. We consider the diagram in Figure 2 where α, β and γ are given by

$$\begin{aligned}
\alpha(\bar{M}, \bar{N}) &= \sum_{\bar{X}; X \text{ indec.}} (\mu(X, N) - \mu(X, M)) \bar{X}, \\
\gamma(\bar{M}, \bar{N}) &= \delta_{M, N},
\end{aligned}$$

$$\beta \left(\sum_{\bar{X}; X \text{ indec.}} \lambda_{\bar{X}} \bar{X} \right) = \sum_{\bar{X}; X \text{ indec.}} \lambda_{\bar{X}} [X, \quad].$$

The diagram commutes since

$$\beta \circ \alpha(\bar{M}, \bar{N}) = \sum_{\bar{X}} (\mu(X, N) - \mu(X, M)) [X, \quad] = [N, \quad] - [M, \quad] = \delta_{M, N}.$$

Obviously α is a bijection and β is $\mathbb{Z}$ -linear.

LEMMA 2.1. *If Λ is of finite representation type then β is an isomorphism.*

Proof. The map β is $\mathbb{Z}$ -linear between free $\mathbb{Z}$ -modules of the same finite rank. Thus it suffices to show that β is surjective. For each indecomposable module X we consider the exact sequence

$$X \longrightarrow E'_X \longrightarrow \tau^{-1}X \longrightarrow 0$$

which is the Auslander–Reiten sequence starting in X if X is not injective. Otherwise we set $E'_X = X/\text{soc}(X)$ and $\tau^{-1}X = 0$. The functor $\text{Hom}_{\Lambda}(\quad, Y)$ induces the exact sequence

$$0 \longrightarrow \text{Hom}_{\Lambda}(\tau^{-1}X, Y) \longrightarrow \text{Hom}_{\Lambda}(E'_X, Y) \longrightarrow \text{Hom}_{\Lambda}(X, Y) \longrightarrow k^{\mu(X, Y)} \longrightarrow 0$$

of k -vectorspaces. Thus for every indecomposable module Y we have

$$([X, \quad] + [\tau^{-1}X, \quad] - [E'_X, \quad])(Y) = \begin{cases} 1 & \text{if } Y \cong X \\ 0 & \text{otherwise,} \end{cases}$$

showing that β is surjective. □

We want to describe the inverse of β . For each indecomposable module X we consider the exact sequence

$$0 \longrightarrow \tau X \longrightarrow E_X \longrightarrow X$$

which is the Auslander–Reiten sequence ending in X if X is not projective. Otherwise we set $E_X = \text{rad}(X)$ and $\tau X = 0$. Then β^{-1} is given by

$$\beta^{-1}(\delta) = \sum_{\bar{X}; X \text{ indec.}} (\delta(X) + \delta(\tau X) - \delta(E_X)) \bar{X}.$$

Thus, if Λ is of finite representation type, γ is bijective and we write $\gamma^{-1}(\delta) = (\bar{M}_\delta, \bar{N}_\delta) \in \mathcal{S}$. Then

$$\begin{aligned} \sum_{\bar{X}; X \text{ indec}} (\delta(X) + \delta(\tau X) - \delta(E_X)) \bar{X} &= \beta^{-1}(\delta) = \alpha \circ \gamma^{-1}(\delta) \\ &= \sum_{\bar{X}; X \text{ indec}} (\mu(X, N_\delta) - \mu(X, M_\delta)) \bar{X} \end{aligned}$$

and in consequence

$$\delta(X) + \delta(\tau X) - \delta(E_X) = \mu(X, N_\delta) - \mu(X, M_\delta) \quad (3)$$

for every indecomposable module X and every $\delta \in (\bigoplus_{\bar{Y}; Y \text{ indec}} \mathbb{Z}\bar{Y})^*$.

Let $\mathcal{S}' \subset \mathcal{S}$ be the subset containing all pairs $(\bar{M}, \bar{N})$ with $M \leq_{\deg} N$.

LEMMA 2.2. *If Λ is of finite representation type then γ restricts to a bijection between $\mathcal{S}'$ and the set of non-negative functions $\delta \in (\bigoplus_{\bar{Y}; Y \text{ indec}} \mathbb{Z}\bar{Y})^*$ such that $\delta(I) = 0$ for every injective module I .*

Proof. Zwara showed in [14] that the partial orders $\leq_{\deg}$ and $\leq$ coincide for representation finite algebras. Hence $\gamma(\bar{M}, \bar{N})$ is a non-negative function for every $(\bar{M}, \bar{N}) \in \mathcal{S}'$. If I is an injective module then $[N, I] = [M, I]$ holds in consequence of the exactness of $\text{Hom}_\Lambda(-, I)$ and (1). On the other hand, let $\delta \in (\bigoplus_{\bar{Y}; Y \text{ indec}} \mathbb{Z}\bar{Y})^*$ be non-negative such that $\delta(I) = 0$ for every injective module. We have to show that $\dim_k N_\delta = \dim_k M_\delta$ holds. We consider the injective module $\text{Hom}_k(\Lambda_\Lambda, k)$, where Λ_Λ denotes Λ as Λ -right module. Then the adjoint isomorphism gives $[A, \text{Hom}_k(\Lambda_\Lambda, k)] = \dim_k \text{Hom}_k(A, k) = \dim_k A$ for every Λ -module A . In particular $\dim_k N_\delta = \dim_k M_\delta$. $\square$

Let $\mathcal{S}'_N = \{(\bar{X}, \bar{Y}) \in \mathcal{S}' \mid \bar{Y} = \bar{N}\}$. As a consequence of Lemma 2.2 and (3) we can describe all isomorphism classes of modules degenerating to an indecomposable.

LEMMA 2.3. *Let Λ be of finite representation type and N be indecomposable. Then $\mathcal{S}'_N$ is mapped bijectively by γ to the set of non-negative functions $\delta \in (\bigoplus_{\bar{Y}; Y \text{ indec}} \mathbb{Z}\bar{Y})^*$ with $\delta(I) = 0$ for every injective module I , and satisfying*

$$\delta(X) + \delta(\tau X) - \delta(E_X) \begin{cases} = 1 & \text{if } X \cong N \\ \leq 0 & \text{otherwise} \end{cases} \quad (4)$$

for every indecomposable X .

Note that for $(\bar{M}, \bar{N}) = \gamma^{-1}(\delta)$ we have then

$$M \cong \bigoplus_{\bar{X}; X \not\cong N} X^{-(\delta(X) + \delta(\tau X) - \delta(E_X))}, \quad (5)$$

where the direct sum is taken over all isomorphism classes of indecomposable Λ -modules except that of N .

3. Proof of Theorem 1.1

Let Λ be a selfinjective finite dimensional k -algebra of finite representation type with stable Auslander–Reiten quiver isomorphic to $\mathbb{Z}D_{3m}/\tau^{2m-1}$. If $M \leq_{\deg} N$ is a

proper degeneration to the indecomposable N , then M and N are disjoint. If N were projective then the sequence (1) would split in contradiction to $M \not\cong N$. Thus N corresponds to a vertex c_l^s in the stable Auslander–Reiten quiver of Λ .

In subsection 3.1 we will characterize a function $\delta \in \gamma(\mathcal{S}'_N)$ describing a proper degeneration to N as the unique solution of a linear system depending on two vertices. One of these vertices is the vertex corresponding to N . In Subsection 3.2 we analyse this linear system. We will show that if this linear system has a solution in the natural numbers, then this solution is uniquely determined by the vertex c_l^s and $m+1 \leq s \leq 2m$ holds. It follows then from Lemma 2.3 that there exists up to isomorphism at most one module M degenerating to N . Finally we give, for the indecomposable module N corresponding to the vertex c_l^s with $m+1 \leq s \leq 2m$, a non-negative function δ which satisfies (4) for every indecomposable. Thus by Lemma 2.3 there exists a proper degeneration $M \leq_{\text{deg}} N$.

By reindexing the stable Auslander–Reiten quiver $\mathbb{Z}D_{3m}/\tau^{2m-1}$ we can assume that $l = 1$. From now on N always corresponds to the vertex c_1^s .

3.1. Characterization of δ by a linear system

We denote by $p := 2m - 1$ the period of the Auslander–Reiten translation τ and by $h := 3m - 1$ the ‘height’ of the Dynkin diagram D_{3m} .

Let us fix an element $\delta \in \gamma(\mathcal{S}'_N)$. We set

$$a_j^i := \begin{cases} 0 & \text{if } i = 0 \\ \delta(c_j^i) & \text{if } 1 \leq i \leq h-1 \\ b_j^h + b_j^{h+1} & \text{if } i = h, \end{cases}$$

where $b_j^i := \delta(c_j^i)$ for $h \leq i \leq h+1$.

Note that all the integers a_j^i and b_j^i are non-negative. We consider for each vertex c_j^i the Auslander–Reiten sequence ending in c_j^i . Since $\delta(I) = 0$ if I is an injective Λ -module we obtain the following set of inequalities from Lemma 2.3.

If $1 \leq i \leq h-1$

$$a_j^i + a_{j-1}^i - a_{j-1}^{i+1} - a_j^{i-1} \begin{cases} = 1 & \text{if } c_j^i = c_1^s \\ \leq 0 & \text{otherwise.} \end{cases} \quad (6)$$

If $h \leq i \leq h+1$

$$b_j^i + b_{j-1}^i - a_j^{h-1} \begin{cases} = 1 & \text{if } c_j^i = c_1^s \\ \leq 0 & \text{otherwise.} \end{cases} \quad (7)$$

These inequalities are the key to proving the theorem. First we derive some information on the τ -orbits in $\mathbb{Z}D_{3m}/\tau^{2m-1}$. We sum up the δ -values along each τ -orbit and set

$$a_i = \sum_{j=1}^p a_j^i \quad \text{for } 0 \leq i \leq h,$$

$$b_i = \sum_{j=1}^p b_j^i \quad \text{for } h \leq i \leq h+1.$$

Then by definition $a_h = b_h + b_{h+1}$ and $a_0 = 0$. For each fixed i we add up the inequalities of (6) and (7) respectively and we obtain

$$2a_i \leq a_{i+1} + a_{i-1} + \delta_{i,s} \quad \text{for } 1 \leq i \leq h-1, \quad (8)$$

$$2b_i \leq a_{h-1} + \delta_{i,s} \quad \text{for } h \leq i \leq h+1. \quad (9)$$

Here $\delta_{i,s}$ denotes the Kronecker symbol. By definition and the inequality (9) we get $2a_h = 2b_h + 2b_{h+1} \leq 2a_{h-1} + 1$. Hence

$$a_h \leq a_{h-1}. \quad (10)$$

REMARK 3.1. From $\delta(N) + \delta(\tau N) - \delta(E_N) = 1$ it follows immediately that $a_s \geq \delta(N) + \delta(\tau N) > 0$ if $s \leq h-1$ and $a_h > 0$ if $s \geq h$.

The following lemma implies that the case $s \geq h$ does not occur. In view of Figure 1 this means that the vertex c_1^s is not one of the somehow exceptional vertices on the upper boundary.

LEMMA 3.2. *It holds that $s \leq h-1$ and there exists an integer t with $2 \leq t \leq s$ such that*

$$t = 2b_h = 2b_{h+1} \quad \text{and} \quad a_i = \begin{cases} 0 & \text{if } 0 \leq i \leq s-t \\ i - (s-t) & \text{if } s-t \leq i \leq s \\ t & \text{if } s \leq i \leq h. \end{cases}$$

In particular $a_s = t$ is an even integer.

Proof. The inequalities in (8) are equivalent to

$$a_i - a_{i-1} \leq a_{i+1} - a_i + \delta_{i,s} \quad (11)$$

for $1 \leq i \leq h-1$. Suppose that $s \geq h$. It follows from (11) and (10) that

$$0 \leq a_1 = a_1 - a_0 \leq \dots \leq a_h - a_{h-1} \leq 0.$$

This implies that $a_i = 0$ for all $i \in \{1, \dots, h\}$ in contradiction to $a_h > 0$ by Remark 3.1. Hence $s \leq h-1$.

Again by (11) and (10) we obtain the following chain of inequalities:

$$\begin{aligned} 0 \leq a_1 = a_1 - a_0 &\leq a_2 - a_1 \leq \dots \leq a_s - a_{s-1} \\ &\leq a_{s+1} - a_s + 1 \leq \dots \leq a_h - a_{h-1} + 1 \leq 1. \end{aligned}$$

If $a_s - a_{s-1} = 0$ then $a_1 = a_2 = \dots = a_s = 0$ in contradiction to $a_s > 0$ by Remark 3.1. Hence there is an integer t with $0 < t \leq s$ such that

$$\begin{aligned} 0 &= a_1 - a_0 = \dots = a_{s-t} - a_{s-t-1}, \\ 1 &= a_{s-t+1} - a_{s-t} = \dots = a_s - a_{s-1} \\ &= a_{s+1} - a_s + 1 = \dots = a_h - a_{h-1} + 1. \end{aligned}$$

Our claim for the a_i is an easy consequence. In particular we have $a_{h-1} = a_h = b_h + b_{h+1}$, but $2b_{h+1} \leq a_{h-1}$ and $2b_h \leq a_{h-1}$ by (9). Hence we see that $t = a_{h-1} = 2b_h = 2b_{h+1}$ is an even integer. $\square$

As an immediate consequence of Lemma 3.2 we note that

$$2a_i - a_{i+1} - a_{i-1} = \begin{cases} 1 & \text{if } i = s \\ -1 & \text{if } i = s - t \\ 0 & \text{otherwise} \end{cases} \quad (12)$$

for $1 \leq i \leq h - 1$.

We want to describe the non-negative integers a_j^i and b_j^i and hence the function δ as the unique solution of a linear system. By Lemma 3.2 we have $1 = a_{s-t+1} = \sum_j \delta(c_j^{s-t+1})$, so there exists exactly one vertex c_{u-1}^{s-t+1} with $\delta(c_{u-1}^{s-t+1}) = a_{u-1}^{s-t+1} = 1$. Note that the index u is only determined modulo p . In the sequel let $\tilde{u}$ be the representative of u with $1 \leq \tilde{u} \leq p$.

If $s > t$ the Auslander–Reiten sequence ending in c_u^{s-t} gives rise to the equation

$$a_u^{s-t} + a_{u-1}^{s-t} - a_{u-1}^{s-t+1} - a_u^{s-t-1} = -1 \quad (13)$$

because $a_i = 0$ for $i \leq s - t$, by Lemma 3.2.

We consider the following linear system which depends on the positions of the vertices c_1^s and c_{u-1}^{s-t+1} or equivalently on the integers s, t and $\tilde{u}$. The lower index is taken to be in $\mathbb{Z}/p\mathbb{Z}$.

$$x_j^0 = 0, \quad x_j^1 = \begin{cases} 1 & \text{if } c_j^1 = c_{u-1}^{s-t+1} \\ 0 & \text{otherwise,} \end{cases} \quad x_j^h = y_j^h + y_j^{h+1}. \quad (14)$$

If $1 \leq i \leq h - 1$

$$x_j^i + x_{j-1}^i - x_{j-1}^{i+1} - x_j^{i-1} = \begin{cases} 1 & \text{if } c_j^i = c_1^s \\ -1 & \text{if } c_j^i = c_u^{s-t} \\ 0 & \text{otherwise.} \end{cases} \quad (15)$$

If $h \leq i \leq h + 1$

$$y_j^i + y_{j-1}^i = x_j^{h-1}. \quad (16)$$

LEMMA 3.3. *If x_j^i, y_j^i is a rational solution of the linear system (14)–(16) then $y_j^{h+1} = y_j^h$ holds for all j .*

Proof. Suppose that there is j_0 with $y_{j_0}^{h+1} > y_{j_0}^h$. Since $y_j^{h+1} + y_{j+1}^{h+1} = x_{j+1}^{h-1} = y_j^h + y_{j+1}^h$ by (16) we have $-y_{j_0+1}^{h+1} > -y_{j_0+1}^h$ and successively

$$-y_{j_0}^{h+1} = (-1)^p y_{j_0+p}^{h+1} > (-1)^p y_{j_0+p}^h = -y_{j_0}^h, \quad \text{as } p \text{ is odd,}$$

in contradiction to $y_{j_0}^{h+1} > y_{j_0}^h$. $\square$

To any integer solution x_j^i, y_j^i of this linear system we can associate a function $\delta' \in (\bigoplus_{\bar{Y}; Y \text{ indec}} \mathbb{Z}\bar{Y})^*$ by setting $\delta'(c_j^i) = x_j^i$ for $1 \leq i \leq h - 1$, $\delta'(c_j^i) = y_j^i$ for $h \leq i \leq h + 1$ and $\delta'(I) = 0$ for every injective module I . Under the same conditions we will speak of a function $\delta' \in (\bigoplus_{\bar{Y}; Y \text{ indec}} \mathbb{Z}\bar{Y})^*$ as a solution of the linear system (14)–(16).

LEMMA 3.4. *The unique solution of the linear system (14)–(16) is given by $x_j^i = a_j^i$ and $y_j^i = b_j^i$. In particular $a_j^h = 2b_j^h$ is an even integer for all j .*

Proof. With respect to uniqueness, it is obvious that the values of x_j^i are determined by the equations in (14) and (15). The values y_j^i are given by $y_j^i = x_j^h/2$ by (14) and Lemma 3.3.

The equations in (14) are obviously satisfied by a_j^i and b_j^i .

Because of (13) and $a_1^s + a_0^s - a_0^{s+1} - a_1^{s-1} = 1$ it remains for (15) to show that

$$a_j^i + a_{j-1}^i - a_{j-1}^{i+1} - a_j^{i-1} = 0$$

for $c_u^{s-t} \neq c_j^i \neq c_1^s$. By (12) we have

$$0 = 2a_i - a_{i+1} - a_{i-1} = \sum_j (a_j^i + a_{j-1}^i - a_{j-1}^{i+1} - a_j^{i-1})$$

for $s \neq i \neq s-t$ and

$$0 = 2a_{s-t} - a_{s-t+1} - a_{s-t-1} + 1 = \sum_{j \neq u} (a_j^{s-t} + a_{j-1}^{s-t} - a_{j-1}^{s-t+1} - a_j^{s-t-1})$$

for $i = s-t$ and

$$0 = 2a_s - a_{s+1} - a_{s-1} - 1 = \sum_{j \neq 1} (a_j^s + a_{j-1}^s - a_{j-1}^{s+1} - a_j^{s-1})$$

for $i = s$. However because of (6) each of the summands on the right-hand side is less than or equal to zero. Thus each summand on the right-hand side is zero.

Concerning (16) we remark that for $h \leq i \leq h+1$ we have

$$0 = 2b_i - a_{h-1} = \sum_j (b_j^i + b_{j-2}^i - a_{j-1}^{h-1})$$

by Lemma 3.2. Since each summand on the right-hand side is less than or equal to zero by (7) it is zero.

Finally we have $a_j^h = 2b_j^h$ by (14) and Lemma 3.3. $\square$

This means that every $\delta \in \gamma(\mathcal{S}'_N)$ is the unique solution of a linear system which itself depends on some vertex c_{u-1}^{s-t+1} . In the next section we will show that c_{u-1}^{s-t+1} is uniquely determined by c_1^s .

3.2. Computation of c_{u-1}^{s-t+1}

Note that the vertices c_1^s and c_{u-1}^{s-t+1} have to be positioned in $\mathbb{Z}D_{3m}/\tau^{2m-1}$ in such a way that the unique solution of the linear system (14)–(16) takes values in $\mathbb{N}$. We will show that this condition implies that $m+1 \leq s \leq 2m$ and determines c_{u-1}^{s-t+1} as c_{s-m}^{2m-s+1} .

It is possible to solve this problem directly by examining the linear system (14)–(16), but this procedure is rather complicated. Therefore we use a different method. We will ignore the equations of (16) and only use that a_h^j is even. Furthermore we use a covering technique to simplify the computations.

To this purpose we consider the stable translation-quiver $\mathbb{Z}A_h$ with vertices d_j^i where $1 \leq i \leq h$ and $j \in \mathbb{Z}$. There are arrows from d_j^i to d_j^{i+1} and from d_j^{i+1} to d_{j+1}^i for $1 \leq i \leq h-1$. The translation τ is given by $\tau(d_j^i) = d_{j-1}^i$. In particular we are interested in two quotients of $\mathbb{Z}A_h$, namely $Q_1 := \mathbb{Z}A_h/\tau^p$ and $Q_2 := \mathbb{Z}A_h/\tau^{2p}$.

FIGURE 3. Q_2 .

Now let $s', t', u' \in \mathbb{N}$ with $2 \leq t' \leq s' \leq h-1$ and $1 \leq u' \leq p$. Let $\delta_r : \{\text{vertices of } Q_r\} \rightarrow \mathbb{Z}$ for $r = 1, 2$ be a function satisfying

$$\delta_r(d_j^1) = \begin{cases} 1 & \text{if } d_j^1 = d_{u'-1}^{s'-t'+1} \\ 0 & \text{otherwise,} \end{cases} \quad (17)$$

$$\delta_r(d_j^1) + \delta_r(d_{j-1}^1) - \delta_r(d_{j-1}^2) = \begin{cases} -1 & \text{if } d_j^1 = d_{u'}^{s'-t'} \\ 0 & \text{otherwise,} \end{cases} \quad (18)$$

$$\delta_r(d_j^i) + \delta_r(d_{j-1}^i) - \delta_r(d_{j-1}^{i+1}) - \delta_r(d_j^{i-1}) = \begin{cases} 1 & \text{if } d_j^i = d_1^{s'} \\ -1 & \text{if } d_j^i = d_{u'}^{s'-t'} \\ 0 & \text{otherwise,} \end{cases} \quad (19)$$

for $2 \leq i \leq h-1$.

The values of $\delta_r(d_j^1)$ are determined by condition (17). The values of $\delta_r(d_j^i)$ with $i \geq 2$ are determined by the values of $\delta_r(d_j^{i'})$ with $i' < i$ because of conditions (18) and (19). Thus there exists exactly one such function δ_r .

The function δ_2 is easy to calculate.

LEMMA 3.5. *If $u' - t' \leq 1$ then the function δ_2 is given by $\delta_2(d_j^i) = 1$ if the vertex d_j^i lies in the shaded area (including the boundary) of Figure 3 and $\delta_2(d_j^i) = 0$ otherwise.*

Proof. It is straightforward to check that Equations (17), (18) and (19) hold. $\square$

We define the function $\delta'_1 : \{\text{vertices of } Q_1\} \rightarrow \mathbb{Z}$ by $\delta'_1(d_j^i) = \delta_2(d_j^i) + \delta_2(d_{j+p}^i)$. Of course δ'_1 satisfies Equations (17), (18) and (19), from which we see that $\delta'_1 = \delta_1$ and consequently

$$\delta_1(d_j^i) = \delta_2(d_j^i) + \delta_2(d_{j+p}^i). \quad (20)$$

We note the following.

LEMMA 3.6. *If $u' - t' \leq 1$ then $\delta_1(d_j^i)$ is an even integer if and only if $\delta_2(d_j^i) = \delta_2(d_{j+p}^i)$.*

Proof. Since δ_2 takes only values in $\{0, 1\}$ by Lemma 3.5 the claim follows from (20). $\square$

Let us now consider again our function $\delta \in \gamma(\mathcal{S}'_N)$. This function induces a function $\bar{\delta} : \{\text{vertices of } Q_1\} \rightarrow \mathbb{Z}$ by $\bar{\delta}(d_j^i) = a_j^i$. We set $s' = s$, $t' = t$ and $u' = \tilde{u}$.

Then the function $\bar{\delta}$ satisfies Equations (17), (18) and (19) by Lemma 3.4 and consequently $\bar{\delta} = \delta_1$.

LEMMA 3.7. *If $\delta \in \gamma(\mathcal{S}'_N)$ is the unique solution of the linear system (14)–(16) then we have $\tilde{u} - t \leq 1$.*

Proof. Suppose the lemma is false. Then $1 < \tilde{u} - t \leq \tilde{u} \leq p$ and in consequence none of the integers $2p, 1, p, p+1$ is congruent modulo $2p$ to an integer in $\{\tilde{u} - t, \tilde{u} - t + 1, \dots, \tilde{u} - 1\}$. By (17)–(19) and since $2 \leq t < p$ we have

$$\delta_2(d_j^s) = \begin{cases} 1 & \text{if } d_j^s = d_{j'}^s \text{ with } j' \in \{\tilde{u} - t, \tilde{u} - t + 1, \dots, \tilde{u} - 1\} \\ 0 & \text{otherwise.} \end{cases}$$

Hence by (20) and Remark 3.1

$$\begin{aligned} 0 &= \delta_2(d_1^s) + \delta_2(d_{1+p}^s) + \delta_2(d_{2p}^s) + \delta_2(d_p^s) \\ &= \bar{\delta}(d_1^s) + \bar{\delta}(d_p^s) \\ &= \delta(N) + \delta(\tau N) \\ &\geq 1 \end{aligned}$$

which is obviously a contradiction. $\square$

We are able now to determine $\tilde{u}$ and t by means of s .

LEMMA 3.8. *If $\delta \in \gamma(\mathcal{S}'_N)$ solves the linear system (14)–(16) then $c_{u-1}^{s-t+1} = c_{s-m}^{2m-s+1}$ and $m+1 \leq s \leq 2m$ holds. In particular $t = 2(s-m)$ and $\tilde{u} = t/2 + 1$.*

Proof. By Lemma 3.7 we have $\tilde{u} - t \leq 1$. Therefore we can apply Lemma 3.5 to describe δ_2 . On the other hand, we know from Lemma 3.4 that $\delta_1(d_j^h) = \bar{\delta}(d_j^h) = a_j^h$ is always an even integer. In view of Lemma 3.6 this means that $\delta_2(d_j^h) = \delta_2(d_{j+p}^h)$ for all j . Thus we have (see Figure 3)

$$\begin{aligned} \tilde{u} - 1 - 1 &\equiv -(h-s) - (\tilde{u} - t - (h-s)) \pmod{2p}, \\ -(h-s) + p &\equiv \tilde{u} - 1 \pmod{2p}. \end{aligned}$$

Substituting p by $2m-1$ and h by $3m-1$ we calculate that $\tilde{u} = t/2 + 1$ and $t = 2(s-m)$. Since $2 \leq t \leq s$ by Lemma 3.2 we get $2 \leq 2(s-m) \leq s$ which is equivalent to $m+1 \leq s \leq 2m$. $\square$

Hence the existence of a proper degeneration to the indecomposable N corresponding to the vertex c_1^s implies that $m+1 \leq s \leq 2m$. Furthermore, if $M \leq_{\text{deg}} N$ is a proper degeneration to N , then $\gamma((\bar{M}, \bar{N})) \in (\bigoplus_{\bar{Y}: Y \text{ indec}} \mathbb{Z}\bar{Y})^*$ is the unique solution of the linear system (14)–(16) with $c_{u-1}^{s-t+1} = c_{s-m}^{2m-s+1}$, by Lemma 3.4 and Lemma 3.8. Hence there exists up to isomorphism at most one module M degenerating to N .

On the other hand, let $m + 1 \leq s \leq 2m$ and N correspond to the vertex c_1^s . We set $s' = s$, $t' = 2(s - m)$ and $u' = t'/2 + 1$ and define δ_2 according to (17), (18) and (19). Since $u' - t' \leq 1$ the function δ_2 is described by Lemma 3.5. In consequence we have $\delta_2(d_j^h) = \delta_2(d_{j+p}^h)$. We define δ by

$$\delta(c_j^i) = \begin{cases} \delta_2(d_j^i) + \delta_2(d_{j+p}^i) & \text{if } 1 \leq i \leq h - 1 \\ 1/2(\delta_2(d_j^h) + \delta_2(d_{j+p}^h)) & \text{if } h \leq i \leq h + 1. \end{cases}$$

Then δ takes value in $\mathbb{N}$ and solves the linear system (14)–(16). Indeed (14) is a consequence of (17), (15) of (18) and (19), and (16) can be checked easily using Lemma 3.5. Thus $\delta \in \gamma(\mathcal{S}'_N)$ which means in view of Lemma 2.3 that there exists a proper degeneration to N . This completes the proof of Theorem 1.1.

Note that the function δ constructed above for the indecomposable module N corresponding to the vertex c_1^s describes the module M degenerating to N in the following way. Let $P_1, \dots, P_m$ be representatives of the isomorphism classes of projective indecomposable Λ -modules. By Lemma 3.4 and (5)

$$M \cong M_1 \oplus \bigoplus_{i=1}^m P_i^{\delta(\text{rad } P_i)}$$

where M_1 corresponds to the vertex c_{s-m+1}^{2m-s} if $s < 2m$ and $M_1 = 0$ if $s = 2m$.

Acknowledgements. The results presented in this paper form a part of the author's doctoral dissertation written under the supervision of Professor C. Riedtmann. The author thanks C. Riedtmann and G. Zwara for helpful discussions and gratefully acknowledges support from the Schweizerischer Nationalfonds.

References

1. S. ABEASIS and A. DEL FRA, 'Degenerations for the representations of a quiver of type A_m ', *J. Algebra* 93 (1985) 376–412.
2. R. AEHLE, C. RIEDTMANN and G. ZWARA, 'Complexity of degenerations of modules', *Comment. Math. Helv.* 76 (2001) 781–803.
3. M. AUSLANDER, I. REITEN and S. O. SMALØ, *Representation theory of Artin algebras*, Cambridge Studies in Advanced Mathematics 36 (Cambridge University Press, 1995).
4. K. BONGARTZ, 'Degenerations for representations of tame quivers', *Ann. Sci. École Norm. Sup.* 28 (1995) 647–668.
5. K. BONGARTZ, 'On degenerations and extensions of finite dimensional modules', *Adv. Math.* 121 (1996) 245–287.
6. O. BRETSCHER, C. LÄSER and C. RIEDTMANN, 'Selfinjective and simply connected algebras', *Manuscripta Math.* 36 (1981) 253–307.
7. H. KRAFT, 'Geometric methods in representation theory', *Representations of algebras*, Lecture Notes in Mathematics 944 (Springer, 1982) 180–258.
8. C. RIEDTMANN, 'Algebren, Darstellungsköcher und zurück', *Comment. Math. Helv.* 55 (1980) 199–224.
9. C. RIEDTMANN, 'Representation-finite selfinjective algebras of class A_n ', *Representation theory II*, Lecture Notes in Mathematics 832 (Springer, 1980) 449–520.
10. C. RIEDTMANN, 'Representation-finite selfinjective algebras of class D_n ', *Compositio Math.* 49 (1983) 231–282.
11. C. RIEDTMANN, 'Degenerations for representations of quivers with relations', *Ann. Sci. École Norm. Sup.* 4 (1986) 275–301.
12. G. ZWARA, 'Degenerations for modules over representation-finite biserial algebras', *J. Algebra* 198 (1997) 563–581.
13. G. ZWARA, 'Degenerations for modules over representation-finite selfinjective algebras', *Colloq. Math.* 75 (1998) 91–95.

14. G. ZWARA, 'Degenerations for modules over representation-finite algebras', *Proc. Amer. Math. Soc.* 127 (1999) 1313–1322.
15. G. ZWARA, 'Degenerations of finite dimensional modules are given by extensions', *Compositio Math.* 121 (2000) 205–218.

Mathematisches Institut
Universität Bern
Sidlerstrasse 5
3012 Bern
Switzerland
aehle@math-stat.unibe.ch