

Astrid Konrad
Michael Mähler
Stephan Arni
Beatrice Flogerzi
Sonja Klingelhöfer
Frank Seibold

Ameliorative effect of IDS 30, a stinging nettle leaf extract, on chronic colitis

Accepted: 25 May 2004
Published online: 25 August 2004
© Springer-Verlag 2004

A. Konrad · S. Arni · B. Flogerzi ·
F. Seibold (✉)
Division of Gastroenterology,
Inselspital, University Hospital,
University of Bern,
Freiburgstrasse, 3010 Berne, Switzerland
e-mail: frank.seibold@insel.ch
Tel.: +41-31-6328025
Fax: +41-31-6329765

M. Mähler
Central Animal Facility and Institute
for Laboratory Animal Science,
Medical School Hannover,
30623 Hannover, Germany

S. Klingelhöfer
Strathmann AG,
Hamburg, Germany

Abstract *Background and aims:* Anti-TNF- α antibodies are very effective in the treatment of acute Crohn's disease, but are limited by the decline of their effectiveness after repeated applications. The stinging nettle leaf extract, IDS 30, is an adjuvant remedy in rheumatic diseases dependent on a cytokine suppressive effect. We investigated the effect of IDS 30 on disease activity of murine colitis in different models. *Methods:* C3H.IL-10-/- and BALB/c mice with colitis induced by dextran sodium sulphate (DSS) were treated with either IDS 30 or water. Mice were monitored for clinical signs of colitis. Inflammation was scored histologically, and faecal IL-1 β and mucosal cytokines were measured by ELISA. Mononuclear cell proliferation of spleen and Peyer's patches were quantified by ³H-thymidine. *Results:* Mice with

chronic DSS colitis or IL-10-/- mice treated with IDS 30 clinically and histologically revealed significantly ($p < 0.05$) fewer signs of colitis than untreated animals. Furthermore, faecal IL-1 β and mucosal TNF- α concentrations were significantly lower ($p < 0.05$) in treated mice. Mononuclear cell proliferation after stimulation with lipopolysaccharide was significantly ($p < 0.001$) reduced in mice treated with IDS 30. *Conclusions:* The long-term use of IDS 30 is effective in the prevention of chronic murine colitis. This effect seems to be due to a decrease in the Th1 response and may be a new therapeutic option for prolonging remission in inflammatory bowel disease.

Keywords Inflammatory bowel disease · IDS 30 · Murine colitis · TNF- α

Introduction

Until now, the pathogenesis of inflammatory bowel disease (IBD) is not totally understood. However, in Crohn's disease (CD), genetic, immunological and environmental factors clearly play an important role. One important concept is the imbalance between T-helper (Th) cell subsets with an overreaction of Th1 cells. Various pro-inflammatory cytokines (especially TNF- α) are secreted by the mucosa of the intestinal tract during the disease [1]. Furthermore, the cells producing TNF- α , as well as the TNF- α concentration in stools, were enhanced [2, 3]. Based on these results, agents have been developed to

suppress TNF- α as a new approach in the therapy of CD. An efficient therapeutic strategy for patients with CD is the intravenous application of anti-TNF- α antibodies (infliximab). Infliximab has been shown to be effective in the treatment of refractory luminal CD, as well as in the treatment of fistulae [4, 5]. Some patients, however, are refractory to immune suppressive treatment and need repeated infusions of infliximab for maintenance of remission [6]. In addition to the high costs of this therapy, some patients develop side effects such as infusion reactions or infectious complications [7–9]. Finally, a loss of effectiveness has been observed after repeated applications [10]. Another anti-TNF- α approach is the therapy with

soluble receptors (e.g. etanercept). This recombinant receptor/Fc fusion protein binds to human TNF- α , but a placebo-controlled trial revealed no significant clinical benefit in CD [11]. In the search for further anti-TNF- α remedies, a few open-label trials have been published using the teratogen thalidomide, which inhibits the production of TNF- α [12–14]. This medication, however, is rejected by the majority of patients because of its side effects. Therefore, it is necessary to evaluate further strategies leading to a modulation of cytokine expression such as novel anti-TNF- α strategies [15, 16].

In this paper we investigate the effects of IDS 30, a stinging nettle leaf extract that is a well-known adjuvant drug in the continuous therapy of rheumatic diseases. Stinging nettle leaf extract was able to reduce pain in the joints and 26% of patients were able to withdraw their non-steroidal anti-rheumatic (NSAR) medications. IDS 30 is a standardised extract of stinging nettle leaves that potently suppresses TNF- α and IL-1 β [17, 18]. The effects are at least in part due to an inhibitory effect on NF- κ B. Inhibition of NF- κ B was not mediated by a direct modification of DNA binding, but rather by preventing degradation of its inhibitory subunit I κ B- α [19]. Furthermore, it has been shown that these extracts mediate a switch in cytokine patterns derived from T-helper cells [20]. IDS 30 (Strathmann AG; Hamburg, Germany) is a specially developed lipophilic stinging nettle leaf extract with high concentration of 13-*S*-hydroxy-9*Z*, 11*E*, 15*Z*-octadecatrienoic acid (13-HOTE). This oxylipin has been identified as a cytokine-suppressive agent in stinging nettle leaves.

In CD, the therapeutic options for maintenance therapy are still limited and most of them are characterised by potential side effects. Thus, we investigated the preventive and therapeutic effect of the orally applied anti-TNF- α active medication IDS 30 on murine colitis. The following experimental IBD models were used: interleukin-10 gene-deficient mice (IL-10 $^{-/-}$) developing a spontaneous colitis, and when orally administered with the dextran sodium sulphate salt (DSS induces acute and chronic colitis) [21]. These models are widely accepted as a reproducible mucosal colonic inflammation in mice [22].

Materials and methods

Materials

The stinging nettle leaf extract IDS 30 (Hox alpha) was kindly provided by Strathmann AG, Germany. The drug was administered at a final concentration of 0.5 mg/ml in drinking water. The daily uptake of IDS 30 in mice corresponded to the dose used in humans for the treatment of rheumatoid arthritis.

Animals

Forty-two BALB/c mice were obtained from Harlan (Horst, The Netherlands). Sixteen homozygous 4-week-old interleukin 10 gene-deficient mice (IL-10 $^{-/-}$), generated on a C3H/HeJBir genetic background (C3H.IL-10 $^{-/-}$), were housed and bred under specific pathogen-free conditions (individually-ventilated cages, IVC). All provisions for the facility were sterilised by autoclaving. All animal experiments were approved by the Swiss authorities.

Five 8-week-old BALB/c mice were treated with 3.5% DSS salt (ICN Biomedicals; Aurora, OH, USA) for 5 days to induce an acute chemical colitis. Treatment with IDS 30 was started for the 4 days following the induction of colitis, and the mice were sacrificed on day 9.

Eight 8-week-old BALB/c mice were treated with the prepared IDS 30 drinking solution or water for 7 days, and another set of eight mice received water as a control. On day 8, 3.5% DSS was administered for 5 days to induce an acute chemical colitis. Treatment with IDS 30 was continued for 4 days following the induction of colitis, and the mice were sacrificed on day 16.

The chronic DSS colitis was induced by three cycles of 3% DSS for 4 days. The DSS phase was interrupted by periods of 1 week's treatment with IDS 30 (or water) each time, and a final treatment with IDS 30 (or water) for the last week. Again, eight mice were treated with IDS and eight mice with water as a control in these experiments.

Eight 4-week-old C3H.IL-10 $^{-/-}$ mice without clinical signs of colitis received either the water with 0.5 mg/ml IDS 30 or water alone for 10 weeks, and were sacrificed on day 70.

Monitoring

The drinking volume, weight and clinical signs of colitis for each mouse were monitored four times a week (redness of the anus, ulcerations of the anus, weight loss, diarrhoea and bloody diarrhoea).

Histological injury grading

All mice were sacrificed by CO₂ inhalation. Longitudinal halves of colons and caeca were taken in their entirety; the length was measured and fixed in 4% phosphate-buffered formalin. Samples were embedded in paraffin in toto and sectioned at 5 μ m. Sections were stained with H&E and the slides were scored double-blinded by two different experts (M.M. and B.F.) as previously described [23]. Briefly, the histological score ranged from grade 0 to 3 (Table 1). The intestinal inflammation of each mouse was scored at the proximal, middle and distal colon, and the scores for each section were computed, resulting in a range from 0 to 9. Furthermore, the intestinal inflammation from the caecum was scored separately and ranged from 0 to 3.

Faecal sample preparation

Faecal samples were collected and weighed. Samples were homogenised in 3 ml/0.1 g stool weight in a solution of 1 mg/ml soy trypsin inhibitor and 1 mg/ml PMSF (phenylmethylsulphonyl fluoride from Sigma; Buchs, Switzerland) in PBS (phosphate-buffered saline, Sigma). Supernatants were collected after centrifugation at 10,000 g for 15 min, sterile filtered and stored at -70°C prior to testing. IL-1 β concentrations were measured by sandwich ELISA using a commercial test system (R&D Systems; Abingdon, UK).

Table 1 Histopathological score

Score	Characteristics
Grade 0	No change from normal tissue
Grade 1	Mild mucosal hyperplasia; inflammation and/or fibrosis in the lamina propria mucosae; occasional erosions and/or small ulcers (involving up to a total of 20 crypts)
Grade 2	Moderate mucosal hyperplasia; multifocal areas of inflammation and/or fibrosis extending into the tunica submucosa and sometimes transmural; crypt abscesses; erosions and ulcers (involving up to a total of 40 crypts)
Grade 3	Severe mucosal hyperplasia; diffuse areas of inflammation and/or fibrosis extending into the tunica muscularis; crypt abscesses; erosions and ulcers (involving more than a total of 40 crypts)

Mucosal cytokine secretion

Longitudinal halves of colons were removed from the control and treated animals, put in cold Hanks-balanced salt solution without calcium and magnesium (HBSS, Life Technologies; Basel, Switzerland), cut into 2- to 4-mm squares and resuspended in complete Roswell Park Memorial Institute (RPMI) medium. Mucosal pieces were then incubated at 37°C in 5% CO₂ in the presence or absence of 2 µg/ml lipopolysaccharide (LPS from *E. coli*, Serotype 055:B5, Sigma, Switzerland). After 6 h, supernatants were collected for measurement of TNF-α and IL-1β and stored at -70°C prior to testing. Cytokine secretion was measured using commercial ELISAs (BD Biosciences Pharmingen; Basel, Switzerland; R&D Systems).

Isolation of mononuclear cells

Peyer's patches and spleens were taken, placed in HBSS, washed, strained through a mesh sieve (40-µm pores) and centrifuged for 8 min at 1,800 rpm. The supernatant was removed and the spleen pellet was resuspended in erythrocyte lysis buffer (ACK-buffer) and cells washed in HBSS three times before reconstitution to 2×10⁶ cells/ml in complete RPMI medium, supplemented with 5% heat-inactivated foetal bovine serum and 100 U/ml antibiotic/antimycotic solution (Life Technologies).

Proliferation assay for mononuclear cells

Mononuclear cells were cultured in complete RPMI medium. Mononuclear cells (2×10⁶/ml) were co-cultivated either with 1–10 µg/ml LPS, 3 µg/ml of the cell mitogen Concanavalin A (Con A, Böhringer; Mannheim, Germany) or 5 µg/ml anti-CD3 (Pharmingen), and incubated for 4 days. Proliferation was determined by ³H-thymidine incorporation (Amersham; Little Chalfont, UK). Cells were pulsed with ³H-thymidine for the final 18 h and harvested (Skatron; Lier, Norway). Incorporated ³H-thymidine was quantified by scintillation counting (Beckmann; Fullerton, CA, USA).

Statistical analysis

The results were expressed as mean ± standard error of the mean. The significance of the difference in means was determined by the unpaired, two-tailed Student's *t*-test (GraphPadPrism 3.0).

Results

Ameliorative effect of the stinging nettle leaf extract IDS 30 on disease activity of chronic DSS colitis

The clinical effects of a daily oral treatment with IDS 30 were compared with those in untreated controls in a DSS-induced chronic and acute colitis model. Since a therapeutic effect of IDS 30 in rheumatic diseases was only seen after several weeks of treatment, we investigated the chronic DSS colitis model. As shown in Fig. 1a, in the chronic DSS model, no significant difference in weight over the time of the experiment was observed when comparing mice treated with IDS 30 with untreated mice. However, during the DSS cycles, the loss of weight was earlier and more severe in the control compared with the IDS 30 treatment group with increasing evidence in the repeated DSS phases (not significant). Only the water-treated group (25%) and not the mice treated with IDS 30 showed more frequently clinical signs of colitis (redness and ulcerations of the anus, bloody diarrhoea; *p*<0.05). These observations were supported by the significantly reduced colon length in the water-treated group compared with mice receiving IDS 30 (7.3 vs. 8.2 cm, *p*<0.05), and were confirmed by a non-significant, more severe histological score in untreated mice. The mice treated with IDS 30 were scored with 4.25±0.25 for the colon, whereas the control group averaged over 4.75±0.25 (*p*=0.20). The caeca were scored with 2.00±0.0 vs. 2.5±0.5 (Table 2).

Amelioration of disease in IL-10-/- mice

In a further experiment on chronic murine colitis over a 10-week period, we applied IDS 30 to 4-week-old C3H.IL-10-/- mice while a control group was treated with water as a placebo. We found a parallel gain in weight in both groups (Fig. 1b). In 29% of the placebo group, we observed clinical signs of colitis, such as redness of the anus and perianal ulcerations; the same observations were made in only 14% of the mice treated with IDS 30. Significant discrepancies were measured in the colon length (8.4 vs. 7.1 cm; *p*<0.05) and histological injury grading (*p*<0.05). The colons of mice treated with IDS 30 were scored with 2.5±0.5, whereas the control




Fig. 1 **a** Increase in the weight of BALB/c mice and C3H/He JBir.IL-10^{-/-} mice treated with IDS 30 compared with those treated with water. In the chronic dextran sodium sulphate (DSS) model, colitis was induced after 1 week's pre-treatment with IDS 30 by three cycles of 3% DSS for 4 days interrupted by a period of 1 week's treatment with IDS 30 (or water). **b** Increased weight of the C3H.IL-10^{-/-} mice treated for 10 weeks with IDS 30

versus water. In the acute DSS model, **c** BALB/c mice were treated with IDS 30 after colitis had been established or **d** mice were pre-treated with IDS 30 or water for 1 week before the induction of colitis with DSS. Afterwards, 3.5% DSS was applied for 5 days, and the treatment was continued for 4 days (**c**, **d**). Note that $n=5$ BALB/c mice in experiment (**c**), $n=8$ BALB/c mice per group in **a**, **d**, and $n=8$ IL-10 knockout mice per group in **b**

Table 2 Colon length and histological score of treated and untreated animals. DSS dextran sodium sulphate

	IDS 30-treated mice; colon length in cm (mean \pm SEM)	Water-treated mice; colon length in cm (mean \pm SEM)	IDS 30-treated mice; score/colon (mean \pm SEM)	Water-treated mice; score/colon (mean \pm SEM)	IDS 30-treated mice; score/caecum (mean \pm SEM)	Water-treated mice score/caecum (mean \pm SEM)
Chronic DSS model (BALB/c mice)	8.2 \pm 0.4*	7.3 \pm 0.2	4.25 \pm 0.25	4.75 \pm 0.25	2.00 \pm 0.0	2.5 \pm 0.5
C3H.IL-10 ^{-/-} mice	8.4 \pm 0.3*	7.1 \pm 0.3	2.5 \pm 0.5*	4.75 \pm 0.25	2.00 \pm 0.0*	2.75 \pm 0.25
Acute DSS model (BALB/c mice) therapy	7.0 \pm 0.2	6.8 \pm 0.4	3.6 \pm 0.2	4.00 \pm 0.0	2.4 \pm 0.25	2.6 \pm 0.25
Acute DSS model (BALB/c mice) prevention	7.1 \pm 0.3	6.9 \pm 0.3	3.25 \pm 0.25	4.00 \pm 0.0	2.25 \pm 0.25	2.75 \pm 0.25

* $p < 0.05$

group averaged over 4.75 \pm 0.25. Caeca were scored with 2.0 \pm 0.0 vs. 2.75 \pm 0.25 (Fig. 2; Table 2).

IDS 30 is not effective in the prevention or amelioration of acute colitis

After the demonstration of the beneficial effect of IDS 30 on chronic murine colitis, we investigated the effect of

this component in the acute DSS model. We observed a similar progression in weight in the groups treated and untreated with IDS 30 (Fig. 1c). Neither the clinical signs of colitis nor the colon length differed significantly in either group (7.0 cm [IDS 30] vs. 6.8 cm [control]; Table 2). We found a non-significant, more severe histological score in untreated mice. The colon scores were 3.6 \pm 0.2 in the IDS 30 group vs. 4.00 \pm 0.0 in the water group, and 2.4 \pm 0.25 vs. 2.6 \pm 0.25 for the caecum.




Fig. 2 Histological sections from C3H.IL-10 knockout mice (H&E, $\times 10$). The histology of mice treated with IDS 30 was scored with 2.50 ± 0.50 , whereas the control group averaged over 4.75 ± 0.25 . **a** Proximal colon of an IL-10-deficient mouse treated with IDS 30 with normal appearance. **b** Proximal colon of a water-treated IL-10-deficient mouse showing severe inflammatory cell infiltration in the lamina propria and submucosa ($n=8$ mice per group)

When mice were pre-treated for 1 week with IDS 30 or water prior to the DSS application, we found less weight loss in the mice treated with IDS 30 during the DSS phase (not significant) than those in the water-treated group (Fig. 1d). There was no difference in the drinking volume between the two groups. At the end of the DSS feeding, we observed clinical signs of colitis (redness of the anus, diarrhoea and bloody diarrhoea) less frequently in the group treated with IDS 30 than in the control group (38 vs. 50%). Neither the clinical signs of colitis nor the colon length differed significantly in either group (7.1 cm vs. 6.9 cm [pre-treatment regimen]; Table 2). We found a non-significant, more severe histological score in untreated mice. The animals pre-treated with IDS 30 were scored 3.25 ± 0.25 vs. 4.00 ± 0.0 in the water group for the colon, and 2.25 ± 0.25 vs. 2.75 ± 0.25 for the caecum (Table 2).

Treatment with IDS 30 reduces faecal IL-1 β concentration

To verify the disease activity during murine colitis, stool samples were collected at different times, and IL-1 β was measured using ELISA. We observed similar concentrations in BALB/c- and in C3H.IL-10 $^{-/-}$ mice prior to treatment (Fig. 3a–c). In the chronic DSS model, there were non-significantly higher IL-1 β concentrations in untreated than in treated mice after a period of 14 days (mean_{water} 82 pg/ml, mean_{IDS 30} 8 pg/ml; $p=0.17$; Fig. 3b). After more than 5 weeks of treatment, we observed a significant difference between the group treated with IDS 30 and the untreated group (mean_{water} 634 pg/ml, mean_{IDS 30} 229 pg/ml; $p=0.043$; Fig. 3a). In C3H.IL10 $^{-/-}$ mice we detected significantly ($p=0.02$) higher levels of IL-1 β at day 70 in faecal samples of the untreated group than in the group of mice that received IDS 30 (mean_{water} 472 pg/ml, mean_{IDS 30} 51 pg/ml; Fig. 3b).

In the acute DSS model, higher concentrations of IL-1 β at day 11 were measured in the placebo group (mean_{water} 260 pg/ml) during acute colitis than in the group treated with IDS 30 (mean_{IDS 30} 220 pg/ml; not significant; $p=0.25$; Fig. 3c).

Reduction of mucosal TNF- α and IL-1 β production after IDS 30 treatment in BALB/c mice

To investigate the influence of IDS 30 on intestinal cytokine production, colons of treated and untreated BALB/c mice from the chronic and acute DSS models were removed to obtain mucosal organ cultures that were cultured in the presence or absence of 2 $\mu\text{g/ml}$ LPS. In the chronic DSS model, there were significantly higher levels of TNF- α in mucosal supernatants of colon ($p=0.035$) of placebo than in animals treated with IDS 30 after stimulation with LPS (Fig. 4a). Without stimulation, lower concentrations of the pro-inflammatory cytokine TNF- α were detected in supernatants of mucosal organ cultures from mice of the acute DSS model receiving IDS 30 than in untreated animals. However, after stimulation with LPS, no difference was observed between the two groups (Fig. 4b). The investigation of mucosal IL-1 β concentrations revealed no differences between treated and untreated animals under baseline conditions. However, the mice treated with IDS 30 in the presence of LPS had significantly lower levels of IL-1 β in comparison to the water-treated group of mice. These observations were made in the chronic DSS model ($p=0.044$) as well as in the acute DSS model ($p=0.042$; Fig. 4c, d).




Fig. 3 Stool samples for IL-1 β determination were collected prior, during and after treatment with IDS 30 or water. IL-1 β concentrations were determined by ELISA. **a** In the chronic DSS model, the IL-1 β levels were significantly lower in the mice treated with IDS 30 at the end of the experiment. **b** IL-1 β concentrations in

faecal samples from untreated C3H.IL-10^{-/-} mice compared with IDS 30-treated animals ($*p < 0.05$). **c** IL-1 β concentrations in faecal samples from IDS 30 and water-treated BALB/c mice prior treatment, in the DSS phase and 5 days after the DSS application in the acute DSS model ($n = 8$ mice per group)

Proliferation of mononuclear cell

Splenic mononuclear cells from BALB/c mice of the chronic DSS model treated with IDS 30 proliferated fewer splenocytes after stimulation with LPS compared with untreated mice (Fig. 5a). We found a significant difference after stimulation with 1 $\mu\text{g/ml}$ ($*p = 0.0272$) and 10 $\mu\text{g/ml}$ LPS ($**p = 0.0038$). Similar results were obtained in Peyer's patches after stimulation with 1 $\mu\text{g/ml}$ ($***p = 0.0007$) and 10 $\mu\text{g/ml}$ LPS ($**p = 0.0019$). Comparing T-cells of BALB/c mice treated with IDS 30 with those of untreated BALB/c mice, we found no difference in the proliferative response after stimulation with either anti-CD3 or Con A (data not shown). In order to investigate whether the IDS 30 itself has any influence on the proliferative response, we stimulated mononuclear cells from IDS 30-treated and -untreated BALB/c and C3H.IL-10^{-/-} mice using different concentrations (1–10 mg/ml) of IDS 30. No proliferation could be detected (data not shown).

Discussion

Inflammatory bowel disease and rheumatoid arthritis have partially common pathomechanisms and several medical therapies such as steroids and infliximab are effective in

both diseases. IDS 30 is approved as an adjuvant therapy of rheumatic diseases in Germany. This paper has described the biological efficacy of IDS 30 on colitis in three different experimental mouse models for IBD. We found a substantial effect of IDS 30 in two models of chronic colitis. The limited effect of IDS 30 in the treatment of acute colitis may be due to the short duration of treatment with this substance. This effect was also seen in clinical studies with patients suffering from arthritis in which the clinical benefit was detectable after a duration of treatment of several weeks. The clinical observations in this study were confirmed by the measurement of IL-1 β in stools, a widely used disease activity marker for murine colitis [24]. We found significantly higher IL-1 β concentrations in untreated animals than in treated ones.

Several mechanisms of the stinging nettle leaf extract, IDS 30, have been discussed. This includes the suppression of cytokine production via an inhibition of NF- κ B activation by IDS 30 [19]. In IBD models, the inhibition of NF- κ B or the pro-inflammatory cytokine TNF- α by an antisense oligonucleotide strategy, led to an improvement in the disease [25, 26]. The effect of IDS 30 on NF- κ B in the context of IBD has not been investigated so far. Furthermore, IDS 30 leads to a suppression of matrix metalloproteinases as shown in chondrocytes *in vivo*. It is known that the elevation of these metalloproteinases plays a certain role in IBD [27, 28], but the effect of IDS 30 in




Fig. 4 a, b Mucosal TNF- α and **c, d** IL-1 β secretion in BALB/c mice at the end of the IDS 30 DSS treatment. **b** Under basal conditions, lower concentrations of the pro-inflammatory cytokine TNF- α were detected in mucosal organ cultures from DSS-treated mice, which received IDS 30 vs. placebo (acute DSS model). **a** After stimulation with 2 μ g/ml lipopolysaccharide (LPS), significantly higher levels of TNF- α were detected in the supernatants of

the colon of placebo than in IDS 30-treated animals in the chronic DSS model. **c, d** For the IL-1 β secretions, no difference was observed under baseline conditions, whereas in the presence of LPS, significantly lower levels of IL-1 β were observed in IDS 30-treated mice from the chronic as well as from the acute DSS model ($n=8$ mice per group). *Significant difference ($p < 0.05$) between the groups

this context remains to be evaluated. A third important mechanism of IDS 30 may be the decrease in the secretion of pro-inflammatory cytokine TNF- α in the intestinal mucosa. An imbalance of a Th1 and Th2 reaction with a preponderance of a Th1 reaction in CD is a widely accepted pathogenic concept. Madsen et al. have shown that colitis in IL-10 $^{-/-}$ mice is associated with high levels of mucosal TNF- α . When the mice were housed under germ-free conditions, the disease was prevented and the mucosal TNF- α ranged within normal levels [29]. These findings suggest that the IL-10 $^{-/-}$ colitis is a result of a Th1-triggered immune response to the mucosal microflora. Furthermore, the clinical data on the effect of an anti-TNF therapy support the importance of TNF- α in the pathogenesis of CD [30, 31]. In another study, it was shown that stinging nettle leaf extract is able to inhibit a Th1 reaction in vitro. The production of IL-2 and IFN- α proteins as well as mRNA by mononuclear cells was inhibited in a dose-dependent manner [20]. Furthermore, in vitro data on macrophages showed that TNF- α and IL-1 β secretion after LPS stimulation were reduced by the addition of a stinging nettle leaf extract of up to 99% in a dose-dependent manner [18]. This study clearly demonstrates the downward regulatory effect of IDS 30 on the Th1-triggered immune response in BALB/c as well as in C3H/HeJBir.IL-10 $^{-/-}$ mice after stimulation with LPS. We detected a significant decrease in IL-1 β as well as less

TNF- α secretion in the supernatants of the intestinal mucosa in treated animals. In vitro, this effect is mediated by a reduced maturation of dendritic cells and their expression of CD83 and CD86, leading to a reduced T-cell response. One of the most potent inducers of pro-inflammatory cytokines is LPS, the most common pathogen-associated molecular pattern. LPS isolated from *E. coli* is well characterised, it signals through the toll-like receptor complex and induces Th cells to secrete high levels of pro-inflammatory cytokines [32]. In this study, the mononuclear cells of spleens and Peyer's patches from mice treated with IDS 30 proliferated less after stimulation with LPS from *E. coli* than those from water-treated mice. This indicates the suppression of a Th1-type reaction by IDS 30. Our in vivo models clearly show a reduced mucosal TNF- α and IL-1 β secretion triggered by the application of IDS 30 during long-term treatment. In humans, increased secretion of IL-1 β and TNF- α is commonly used as a predictive marker for acute relapse. Schreiber et al. demonstrated the positive correlation between high secretion of the pro-inflammatory cytokines TNF- α or IL-1 β from lamina propria mononuclear cells and relapse in patients with CD [33].

Until now, the therapeutic goal in the therapy of IBD is the remission of clinical symptoms as well as the healing of the mucosal inflammation [34]. The therapeutic tools to prolong remission and to reduce mucosal inflammation




Fig. 5 Effect of IDS 30 on the proliferative response of mononuclear cells in treated and untreated BALB/c mice from the chronic DSS model. Mononuclear cells from the spleen and Peyer's patches were isolated from mice treated with IDS-30 or water and stimulated with LPS. The proliferative response was measured by ^3H -thymidine incorporation. Results are expressed as a mean (cpm) + SEM. **a** Mononuclear cells of spleens from BALB/c mice treated with IDS 30 proliferated less than splenocytes from untreated animals after stimulation with LPS. This difference is significant for the tested doses of LPS stimulation (1 $\mu\text{g}/\text{ml}$ and 10 $\mu\text{g}/\text{ml}$ LPS). **b** After stimulation, Peyer's patches with LPS (1 $\mu\text{g}/\text{ml}$ and 10 $\mu\text{g}/\text{ml}$ LPS) mononuclear cells from IDS 30-treated animals proliferated less than those from untreated animals (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$)

are still limited. In CD, the most efficient therapies comprise immune suppressive therapies such as azathioprine, 6-mercaptopurine and methotrexate. These substances may maintain about 60% of the patients in remission [35]. The repeated application of the anti-TNF antibody infliximab is mainly limited by its costs, the

intravenous application, infection complications and the possible loss of efficacy following repeated applications. The need for further anti-TNF- α drugs is undisputed, as shown in many recently published experimental studies [26, 36]. Our data show the TNF- α -suppressive effect of IDS 30. However, it could not be compared with infliximab in terms of efficacy and time of response. In arthritis patients, though, this medication is well tolerated. Our data were much more impressive in the chronic murine colitis models than in the acute colitis model. Apparently, IDS 30 needs several days for a clinical effect to be seen. This is supported by the fact that the LPS-stimulated TNF- α release decreased during the IDS 30 treatment. The basal TNF- α was not affected by the treatment with IDS 30, as already described in another study of TNF- α suppression [26]. In that study, the antisense TNF- α suppression was effective in the acute DSS model, but only if higher concentrations than in the chronic colitis model were used. Another explanation might be the different pathophysiological effects caused by acute versus chronic DSS application. Acute DSS colitis is independent of lymphocytes, mainly triggered by toxic effects of DSS on epithelial cells of the colon and can be induced in SCID-mice, whereas in chronic DSS colitis, immunologic mechanisms are involved [37]. Our data support the hypothesis of the beneficial effect by TNF- α inhibition on preventing disease in different models of experimental IBD.

In conclusion, the use of IDS 30 in the experimental treatment of IBD is effective in the prevention and amelioration of chronic murine colitis in DSS-treated BALB/c and in C3H.IL-10-/- mice. This is the first study to demonstrate the beneficial immunomodulatory effect of IDS 30 on experimental IBD in vivo. Therefore, IDS 30 appears to be a new candidate for maintenance therapy in IBD, which needs to be shown in controlled studies with patients.

Acknowledgements This study was supported by the Swiss National Science Foundation (SNF 31-59031.99), by a grant to Astrid Konrad from the Deutsche Forschungsgemeinschaft (KO-2228/2-1) and by Strathmann AG, Hamburg, Germany.

References

1. Van Deventer SH (1997) Tumour necrosis factor and Crohn's disease. *Gut* 40:443-448
2. Breese EJ, Michie CA, Nicholls SW, Murch SH, Williams CD, Domizio P et al (1994) Tumor necrosis factor alpha-producing cells in the intestinal mucosa of children with inflammatory bowel disease. *Gastroenterology* 106:1455-1466
3. Braegger CP, Nicholls SW, Murch SH, Stephens S, MacDonald TT (1992) Tumour necrosis factor alpha in stool as a marker of intestinal inflammation. *Lancet* 339:89-91
4. Targan SR, Hanauer SB, van Deventer SJ, Mayer L, Present DH, Braakman T et al (1997) A short-term study of chimeric monoclonal antibody cA2 to tumor necrosis factor alpha for Crohn's disease. Crohn's disease cA2 Study group. *N Engl J Med* 337:1029-1035

5. Present DH, Rutgeerts P, Targan S, Hanauer SB, Mayer L, van Hogezaand RA et al (1999) Infliximab for the treatment of fistulas in patients with Crohn's disease. *N Engl J Med* 340:1398–1405
6. Van Deventer SJH (2002) Anti-tumour necrosis factor therapy in Crohn's disease: where are we now? *Gut* 51:362–363
7. Noman M, Baert F, D'Haens G et al (2001) HACA formation after infliximab (remicade) treatment in Crohn's disease is clearly associated with infusion reactions. *Gastroenterology* 120:A69
8. Baert F, Noman M, Vermeire S, Van Assche G, Haens G, Carbonez A et al (2003) Influence of immunogenicity on the long-term efficacy of infliximab in Crohn's disease. *N Engl J Med* 348:601–608
9. Keane J, Gershon S, Wise RP, Mirabile-Levens E, Kasznica J, Schwiertman WD et al (2001) Tuberculosis associated with infliximab, a tumor necrosis factor- α neutralizing agent. *N Engl J Med* 345:1098–1104
10. Hanauer SB, Feagan BG, Lichtenstein GR, Mayer LF, Schreiber S, Colombel JF et al (2002) Maintenance infliximab for Crohn's disease: the ACCENT I randomised trial. *Lancet* 359:1541–1549
11. Sandborn WJ, Hanauer SB, Katz S, Safdi M, Wolf DG, Baerg RD et al (2001) Etanercept for active Crohn's disease: a randomized, double-blind, placebo-controlled trial. *Gastroenterology* 121:1088–1094
12. Vasilias EA, Kam LY, Abreu-Martin MT, Hassard PV, Papadakis KA, Yang H et al (1999) An open-label pilot study of low-dose thalidomide in chronically active, steroid-dependent Crohn's disease. *Gastroenterology* 117:1278–1287
13. Bauditz J, Wedel S, Lochs H (2002) Thalidomide reduces tumour necrosis factor alpha and interleukin 12 production in patients with chronic active Crohn's disease. *Gut* 50:196–200
14. Sabate JM, Villarejo J, Lemann M, Bonnet J, Allez M, Modigliani R (2002) An open-label study of thalidomide for maintenance therapy in responders to infliximab in chronically active and fistulizing refractory Crohn's disease. *Aliment Pharmacol Ther* 16:1117–1124
15. Podolsky DK (2002) Inflammatory bowel disease. *N Engl J Med* 347:417–429
16. Rutgeerts P (2002) A critical assessment of new therapies in inflammatory bowel disease. *J Gastroenterol Hepatol* 17:S176–S185
17. Teucher T, Obertreis B, Ruttkowski T, Schmitz H (1996) Cytokine secretion in whole blood of healthy subjects following oral administration of *Urtica dioica* L. plant extract. *Arzneimittelforschung* 46:906–910
18. Obertreis B, Ruttkowski T, Teucher T, Behnke B, Schmitz H (1996) Ex-vivo in-vitro inhibition of lipopolysaccharide stimulated tumor necrosis factor-alpha and interleukin-1 beta secretion in human whole blood by extractum urtica dioicae foliorum. *Arzneimittelforschung* 46:389–394
19. Riehemann K, Behnke B, Schulze-Osthoff K (1999) Plant extracts from stinging nettle (*Urtica dioica*), an anti-rheumatic remedy, inhibit the proinflammatory transcription factor NF-kappaB. *FEBS Lett* 442:89–94
20. Klingelhofer S, Obertreis B, Quast S, Behnke B (1999) Antirheumatic effect of IDS 23, a stinging nettle leaf extract, on in vitro expression of T helper cytokines. *J Rheumatol* 26:2517–2522
21. Cooper HS, Murthy SN, Shah RS, Sedergran DJ (1993) Clinicopathologic study of dextran sulphate sodium experimental murine colitis. *Lab Invest* 69:238–249
22. Elson CO, Sartor RB, Tennyson GS, Riddell RH (1995) Experimental models of inflammatory bowel disease. *Gastroenterology* 109:1344–1367
23. Konrad A, Mähler M, Flogerzi B, Varga L, Kalousek M, Lange J et al (2003) Amelioration of murine colitis by feeding a solution of lysed *E. coli*. *Scand J Gastroenterol* 38:172–179
24. Lindsay JO, Ciesielki CJ, Scheinin T, Hodgson HJ, Brennan FM (2001) The prevention and treatment of murine colitis using gene therapy with adenoviral vectors encoding IL-10. *J Immunol* 157:7625–7633
25. Neurath MF, Petterson S, Meyer zum Buschenfelde KH, Strober W (1996) Local administration of antisense phosphorothioate oligonucleotides to the p65 subunit of NF-kappaB abrogates established experimental colitis in mice. *Nat Med* 2:998–1004
26. Myers KJ, Murthy S, Flanagan A, Witchell DR, Butler M, Murray S et al (2003) Antisense oligonucleotide blockade of tumor necrosis factor- α in two murine models of colitis. *J Pharmacol Exp Ther* 304:411–424
27. Schulze-Tanzil G, de Sousa P, Behnke B, Klingelhofer S, Scheid A, Shakibaei M (2002) Effects of the antirheumatic remedy hox alpha—a new stinging nettle leaf extract—on matrix metalloproteinases in human chondrocytes in vitro. *Histol Histopathol* 17:477–485
28. Broer J, Behnke B (2002) Immunosuppressant effect of IDS 30, a stinging nettle leaf extract, on myeloid dendritic cells in vitro. *J Rheumatol* 29:659–666
29. Madsen KL, Doyle JS, Jewell LD, Tavernini MM, Fedorak RN (1999) *Lactobacillus* species prevents colitis in interleukin 10 gene-deficient mice. *Gastroenterology* 116:1107–1114
30. Schreiber S, Campieri M, Colombel JF, van Deventer SJH, Feagan B, Fedorak R et al (2001) Use of anti-tumour necrosis factor agents in inflammatory bowel disease. European guidelines for 2001–2003. *Int J Colorectal Dis* 16:1–11
31. D'Haens G (2003) Anti-TNF therapy for Crohn's disease. *Curr Pharm Des* 9:289–294
32. Netea MG, van Deuren M, Kullberg BJ, Cavaillon JM, Van der Meer JW (2002) Does the shape of lipid A determine the interaction of LPS with toll-like receptors? *Trends Immunol* 23:135–139
33. Schreiber S, Nikolaus S, Hampe J, Hämling J, Koop I, Groessner B et al (1999) Tumour necrosis factor α and interleukin 1β in relapse of Crohn's disease. *Lancet* 353:459–461
34. Arnott IDR, Watts D, Ghosh S (2002) Review article: is clinical remission the optimum therapeutic goal in the treatment of Crohn's disease? *Aliment Pharmacol Ther* 16:857–867
35. Feagan BG, Fedorak RN, Irvine EJ, Wild G, Sutherland L, Steinhart AH et al (2000) A comparison of methotrexate with placebo for the maintenance of remission in Crohn's disease. North American Crohn's Study Group Investigators. *N Engl J Med* 342:1627–1632
36. Loher F, Schmall K, Freytag P, Landauer N, Hallwachs R, Bauer C et al (2003) The specific type 4 phosphodiesterase inhibitor mesopram alleviates experimental colitis in mice. *J Pharmacol Exp Ther* 305:549–556
37. Dieleman LA, Ridwan BU, Tennyson GS, Beagley KW, Bucy RP, Elson CO (1994) Dextran sulfate sodium-induced colitis occurs in severe combined immunodeficient mice. *Gastroenterology* 107:1643–1652