

# Accepted Manuscript

Vertical Right Axillary Mini-Thoracotomy for Correction of Ventricular Septal Defects and Complete Atrioventricular Septal Defects

Paul Philipp Heinisch, Marc Wildbolz, Maria Julia Beck, Maris Bartkevics, Brigitta Gahl, Balthasar Eberle, Gabor Erdoes, Hans-Joerg Jenni, Florian Schoenhoff, Jean-Pierre Pfammatter, Thierry Carrel, Alexander Kadner

PII: S0003-4975(18)30730-6

DOI: [10.1016/j.athoracsur.2018.05.003](https://doi.org/10.1016/j.athoracsur.2018.05.003)

Reference: ATS 31650

To appear in: *The Annals of Thoracic Surgery*


Received Date: 29 December 2017

Revised Date: 28 April 2018

Accepted Date: 3 May 2018

Please cite this article as: Heinisch PP, Wildbolz M, Beck MJ, Bartkevics M, Gahl B, Eberle B, Erdoes G, Jenni HJ, Schoenhoff F, Pfammatter JP, Carrel T, Kadner A, Vertical Right Axillary Mini-Thoracotomy for Correction of Ventricular Septal Defects and Complete Atrioventricular Septal Defects, *The Annals of Thoracic Surgery* (2018), doi: 10.1016/j.athoracsur.2018.05.003.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



# Vertical Right Axillary Mini-Thoracotomy for Correction of Ventricular Septal Defects and Complete Atrioventricular Septal Defects

Running Head: Mini thoracotomy for VSD and CAVSD

Paul Philipp Heinisch<sup>1</sup>, Marc Wildbolz<sup>2</sup>, Maria Julia Beck<sup>1</sup>, Maris Bartkevics<sup>1</sup>, Brigitta Gahl<sup>1</sup>, Balthasar Eberle<sup>3</sup>, Gabor Erdoes<sup>3</sup>, Hans-Joerg Jenni<sup>1</sup>, Florian Schoenhoff<sup>1</sup>, Jean-Pierre Pfammatter<sup>2</sup>, Thierry Carrel<sup>1</sup>, Alexander Kadner<sup>1</sup>

<sup>1</sup> Centre for Congenital Heart Disease, Department of Cardiovascular Surgery, Inselspital, University Hospital, University Bern, Bern, Switzerland

<sup>2</sup> Centre for Congenital Heart Disease, Department of Cardiology, Inselspital, University Hospital, University Bern, Bern, Switzerland

<sup>3</sup> Department of Anaesthesiology and Pain Therapy, Inselspital, University Hospital, University Bern, Bern, Switzerland

**Key Words:** Vertical right axillary mini-thoracotomy, ventricular septal defect, complete atrioventricular septal defects

*Corresponding author:*

Alexander Kadner, MD

Centre for Congenital Heart Disease

Department of Cardiovascular Surgery

Inselspital, University Hospital Bern

Freiburgstrasse 18

CH - 3010 Berne, Switzerland

Email: alexander.kadner@insel.ch

**Abstract**

**Background.** Vertical-right-axillary-mini-thoracotomy (VRAMT) is the standard approach for correction of atrial septal defect (ASD) and partial atrioventricular septal defects (PAVSD) at our institution. This observational single-centre study compares our initial results with the VRAMT approach for the repair of ventricular septal defects (VSD) and complete atrioventricular septal defects (CAVSD) in infants and children to an approach using standard median sternotomy (MS).

**Methods.** Perioperative course of patients undergoing VSD and CAVSD correction through either a VRAMT or a MS were analysed retrospectively. The surgical technique for the VRAMT involved a 4-5 cm vertical incision in the right axillary fold.

**Results.** Of 84 patients, 25 patients (VSD, n=15; CAVSD, n=10) underwent correction through a VRAMT approach, whereas 59 patients (VSD, n=35; CAVSD, n=24) had repair through MS. VSD and CAVSD-groups were comparable with respect to age and weight. No significant differences were observed for aortic cross-clamp duration, intensive care unit stay, hospital stay and echocardiographic follow-up. There was no need for any conversion from VRAMT to MS in any case. No wound infection nor thoracic deformities were observed in both groups.

**Conclusions.** VRAMT can be considered as a safe and effective approach for the repair of VSD and CAVSD in selected patient groups, and the outcome data appears comparable to those of median sternotomy.

Evolving surgical techniques and medical care have led to a significant decrease of the mortality and complication rate following congenital heart surgery during the past decades. A complete surgical repair allows both a normal life expectancy and excellent quality of life in the majority of children. At the same time, there is a quest for less invasive and aesthetically superior surgical access [1-3]. Median sternotomy (MS) has been the standard surgical approach for heart surgery, providing optimal exposure of all cardiac structures and allowing easy accessibility for central arterial and venous cannulation and installation of cardio-pulmonary-bypass (CPB). However, median sternotomy is invasive and results in a scar that is well visible on the chest of growing children, which can be potentially stigmatising and trigger long-lasting psychological distress [4-6].

Consequently, several alternative surgical approaches have been proposed like right anterolateral thoracotomy [7,8], posterolateral thoracotomy [9,10], and partial sternotomy [3,11]. They have however been concluded as suboptimal, sometimes because of visibility of the scar, sometimes because of long term consequences like thorax deformity or, especially such as the anterolateral-thoracotomy when performed in pre-pubescent females, impairment of breast development [12,13].

The axillar region represents an interesting surgical access to the thoracic cavity due to particular anatomical properties: The skin is flexible, muscles are relatively rare and can therefore be spared, it is far away from breast tissue in girls, and the resulting scar is hidden under the resting arm [6]. As other studies have already examined, the right axillary access has proven sufficiently exposure of the right atrium and right heart structures, as needed for instance for ASD, VSD and CAVSD. [4]. Furthermore, an eventual scar in the right axilla has shown to be fairly unsuspecting for heart disease, since the anatomical correlation to the heart in the right hemi-thorax is not given [6].

Several groups have already reported their experience with this approach for the correction of atrial-septal-defect (ASD), ventricular-septal-defect (VSD), and partial-atrioventricular-septal-defects (PAVSD) [6,14-18]. Correction of complete-atrioventricular-septal-defects (CAVSD) through a vertical-right-axillary-mini-thoracotomy (VRAMT) has not yet been described before.

The objective of this study is to analyse outcomes and surgical results of VRAMT for correction of VSD and CAVSD in comparison to repair through a MS.

## Patients and Methods

### *Patient selection and data collection*

An observational retrospective single-centre study was conducted using perioperative clinical patient data retrieved from clinical records, surgery and follow-up reports. The study was approved by the Cantonal-Ethics-Committee (approval no. 2016-01484). Patients who underwent corrective heart surgery by VRAMT for either a perimembranous VSD or a CAVSD (Rastelli type A), starting from January 2012 until August 2016 were included and compared to patients who underwent correction via MS. All surgical procedures either VRAMT or MS for the correction of VSD or CAVSD were performed by the same surgeon (AK). To ensure comparability between study groups, patients with a body weight >10 kg were excluded. An observational design was used conforming to the STROBE statement [19]. All data was gathered in a standardized database using the Research-Electronic-Data-Capture (REDCap) system. Patient characteristics, procedural data and outcomes are shown in Tables 1-4.

### *Patient selection for VRAMT*

Since several years, VRAMT has been the standard approach for repair of secundum ASD, sinus venosus defect, partial-anomalous-pulmonary-venous-return, partial atrioventricular septal defects, permanent pacemaker and ICD-implantation in paediatric patients at our institution. Given the growing experience with this cosmetically appealing and less invasive approach, and its comparable results, our group extended its use, since 2012, to perimembranous VSD repair. Since 2014 we also used the VRAMT approach for elective correction of CAVSD with a Rastelli classification type A [20].

### *Statistical analysis*

Variables were presented as numbers with % or as median with interquartile ranges. Differences were investigated using Fisher's exact test for four field tables,  $\chi^2$  test for tables with more than four fields and Kruskal-Wallis test for continuous variables and were presented with 95% confidence interval (CI). All p values and confidence intervals were two sided. All calculations were performed using Stata 12 (College Station, Texas).

### *Follow up*

The follow-up included ambulatory visits with clinical examination and echocardiography by senior paediatric cardiologists at one, three and six months postoperatively, and yearly thereafter. All follow-up data until 31<sup>st</sup> of August 2017 were included in the analyses. Follow-up variables are presented as numbers with percent or as mean  $\pm$  standard deviation (SD). For the purpose of this study all echocardiographic data at discharge and the most recent follow-up are included.

### *Definition of major adverse cardiac and cerebrovascular events*

Major adverse cardiac and cerebrovascular events (MACCE) was defined as one of these events: sudden cardiac death, fatal or non-fatal myocardial-infarction (MI), arrhythmias needing intervention with permanent pacemaker, stroke, or renal failure requiring prolonged dialysis. MI criteria for this study were defined according to the *2012 Third Universal Definition of Myocardial Infarction by the European Society of Cardiology Guidelines*. elevation of cardiac biomarker values  $>10 \times 99^{\text{th}}$  percentile URL in patients with normal baseline troponin values. In addition, either (i) new pathological Q waves or new LBBB, or (ii) angiographic documented new native coronary artery occlusion, or (iii) imaging evidence of new loss of viable myocardium or new regional wall motion abnormality [21].

### *Surgical technique*

Patients received general anaesthesia after induction with sevoflurane, sufentanil and rocuronium. For tracheal intubation, a single-lumen paediatric endotracheal tube was used. Intraoperative monitoring consisted of American-Society-of-Anaesthesiologists standard monitoring, invasive arterial blood pressure via a left-sided radial artery or left-sided femoral artery catheter, central-venous-pressure via a jugular venous catheter, nasopharyngeal temperature, urine output, transesophageal-echocardiography (TEE) and cerebral-near-infrared-spectroscopy (NIRS). Patients undergoing correction of VSD and CAVSD through a VRAMT were placed in a left decubitus position (Figure 1). Prior to positioning, the anterior, posterior axillary line and 4th intercostal space was outlined in supine position with arms in adduction. Following lateral positioning of the child the arm was elevated to expose the axillary region.

A vertical incision of 4-5 cm parallel to the right anterior axillary fold was performed overlying the 4<sup>th</sup> or 5<sup>th</sup>-intercostal-space. The subcutaneous tissue was generously mobilized. the serratus anterior muscle

overlying the 4<sup>th</sup>-intercostal-space was identified and the lateral border of the pectoral muscle slightly mobilized. Care was taken to avoid injury to the long thoracic nerve and artery lying posteriorly on the serratus anterior muscle. Thoracotomy was performed at the superior margin of the 5<sup>th</sup>-rib in the 4th intercostal space. A retractor by *Fehling Surgical Instruments* (V-CUT Titanium MSX-1 for 3-7 kg Bodyweight or MSX-2 for 7-15kg Bodyweight) was used for VRAMT. The pericardium was opened 1- 2 cm anterior to the phrenic nerve (Figure 2 A.). Stay sutures were placed along both the margins of the pericardium and fixed to the surrounding drapes to retain the lungs.

CPB was established following cannulation of the ascending aorta and both vena cava. The inferior-vena-cava was either cannulated directly or in patients > 5kg a percutaneous-venous-cannula (BIO-MEDICUS, Medtronic, Pediatric-Venous-Cannulae) was inserted in the Seldinger-technique into the femoral vein and a LV-vent (Figure 2.B). CPB was conducted in mild-hypothermia, the aorta was cross-clamped directly and antegrade-cardioplegia was administered by using a low-dose (1.5 ml/kg), single-shot crystalloid solution (Cardioplexol™) [22]. The surgical repair of VSD and CAVSD through VRAMT was performed using standard technique and is shown in Figure 3. Deairing was performed via the LV-vent and the aortic root cardioplegia needle and removed after successful de-airing and echocardiographic confirmation of good left-ventricular function. During rewarming epicardial temporary-pacing-electrodes were placed to the right atrium and ventricle. Following decannulation and haemostasis the pericardium was closed except a distal opening of about 1-2 cm for drainage of pericardial fluid into the right pleura. A subpleural-paravertebral-catheter was placed covering two segments each cranial and caudal to the VRAMT interspace for postoperative pain management at a period of 72 hours, using a continuous infusion of bupivacaine 0.125% with adrenaline 5 mcg/ml. Two chest drains were inserted, and the chest was closed with absorbable suture material (Figure 4 A.).

## Results

### *Perioperative Data*

#### *VSD patients*

There were 50 patients included in the VSD group, n= 35 operated through median sternotomy (MS), and n= 15 patients through vertical right axillary mini-thoracotomy (VRAMT). Median age was 4.1 (3.5 - 5.7) months in the MS-VSD group and 4.3 (3.3 - 6.3) months in the VRAMT-VSD group (p= 0.841).

While 31% of patients were female in the MS-VSD group, the VRAMT-VSD group held a majority, 60%, of female patients. Patients were comparable regarding bodyweight ( $p= 0.525$ ) and height ( $p= 0.688$ ). Baseline data and additional diagnosis of VSD patients are shown in Table 1.

#### *CAVSD patients*

From a total of 34 patients with a CAVSD Rastelli type A, 24 were operated through MS and 10 patients through VRAMT. The median-age was 3.7 (3.1 - 7.8) months in the MS-group and 6.1 (4.5 - 10.5) months in the VRAMT-group. CAVSD patients were comparable for bodyweight ( $p= 0.174$ ) and height ( $p= 0.219$ ). Non-cardiac diagnosis included Trisomy 21 for 17 patients in the MS-group and 4 patients in the VRAMT-group. Table 2 shows the preoperative demographics of CAVSD patients.

#### *Intra- and postoperative Assessment*

None of the patients with VRAMT had to be converted to another surgical access. There were two postoperative deaths, one in the MS-VSD-group (Trisomy 18 with severe co-morbidities) and one in the MS-CAVSD-group (pulmonary hypertension crisis and resuscitation).

#### *VSD patients*

The intra- and postoperative data of VSD patients is listed in table 3. The median operating duration ( $p= 0.597$ ) and aortic-cross-clamp-times ( $p= 0.182$ ) were comparable between the groups. Cardiopulmonary bypass time was longer in the VRAMT groups ( $p= 0.015$ ). The median ventilation-time resulted in 43 (22 - 82) hours for patients in the MS-group and 24 (17 - 47) hours in the VRAMT-group ( $p= 0.043$ ).

#### *CAVSD patients*

The operating times for the VRAMT were significantly shorter ( $p= 0.003$ ), whereas central CPB-time ( $p= 0.597$ ) and aortic-cross-clamp-time ( $p= 0.241$ ) of the patients in the MS and VRAMT-group were comparable. Postoperative rhythms in the MS-group included one persistent AV-Block III° with the

indication for a permanent pacemaker. Table 4 shows the intra- and postoperative data of CAVSD patients.

#### *Echocardiographic Follow-up*

##### *VSD patients*

For VSD patients the mean follow-up was complete with a time of 8.1 months ( $\pm 6.9$ ) in the VRAMT-group and 52.3 months ( $\pm 25.7$ ) in the MS group. No significant residual defects were found in patients of both groups. No patient required a reoperation. At last follow-up, there were four patients with trivial residual VSDs, two in the MS-VSD-group, and two in the VRAMT-VSD-group respectively. In clinical examination, no wound complications or thoracic deformity has been found during the follow-up.

##### *CAVSD patients*

Mean follow-up time in the CAVSD-group was complete with a mean follow-up time of 8.6 months ( $\pm 10.3$ ) in the VRAMT-group and 52.9 months ( $\pm 28.9$ ) in the MS group. No patient required a reoperation during the follow-up period. At last follow-up, trivial residual VSDs were found in two patients from the MS-CAVSD-group.

Left AV-valve insufficiency was found in 17 patients (77%) in the MS-CAVSD-group (trivial  $n=15$ , mild to moderate  $n=2$ ) vs. seven patients (70%) in the VRAMT-CAVSD-group (Trivial  $n=5$ , mild to moderate  $n=2$ ), respectively. In the MS-CAVSD-group 15 patients (68%) presented with trivial and one patient (5%) with mild to moderate right AV-valve insufficiency, while in the VRAMT-CAVSD-group six patients (60%) with trivial right AV-valve insufficiency. In clinical examination, no wound complications or thoracic deformity has been found during the follow-up.

#### **Comment**

The use of less invasive lateral thoracotomy in paediatric patients for the correction of congenital heart defects has been reported as feasible and safe in various publications [4,5,18,23,24]. The vertical right axillary mini-thoracotomy aims to provide superior cosmetic results without compromising the surgical

outcome when employed for congenital heart defects, such as atrial septal defects (ASD), ventricular septal defect (VSD), and partial atrio-ventricular septal defects (PAVSD).

In our institution, the surgical correction of perimembranous ventricular septal defects through a right-axillary-mini-thoracotomy was started in 2012. Due to the growing experience and surgical results, and the high satisfaction of the children and their parents, as well as cardiologists, we extended this approach to the correction of “simple” CAVSD with a Rastelli type A. To our knowledge, there are no reports on the application of a VRAMT approach for the correction of CAVSD.

As far as aortic-cross-clamp-times, cardiopulmonary-bypass-time and operation-time are concerned, VRAMT procedures are comparable to standard MS surgery, indicating an adequate exposure and access to the anatomical structures. Furthermore, VRAMT procedures can be performed using standard instruments for access and the correction itself. These findings confirm previously reported advantages and results by other groups [4,18,23-25].

As an advantage of our approach patients weighing < 5kg were operated with central arterial and venous cannulation avoiding potential peripheral vascular complications. In patients weighing > 5 kg peripheral percutaneous-femoral-venous-cannulation was performed. No vascular complications were encountered. The use of femoral artery cannulation for CPB has led to vascular complications in other studies and should therefore be performed only with limited indication and meticulous technique [4,25].

The echocardiographic follow-up revealed a moderate AV-valve insufficiency in two patients for the MS-CAVSD-group (8%) and also in two patients for the VRAMT-CAVSD-group (20%). Furthermore, no wound complications or development of thorax deformities were observed. Long-term follow-up revealed comparable surgical results for VRAMT in comparison to MS for VSD patients. There was no need for re-operations in the current follow-up period for VSD and CAVSD patients. The long term outcome after primary repair of CAVSD remains strongly influenced by the function of the left atrioventricular valve, including in particular the risk for reoperation [26]. Regarding reinterventions after VRAMT-CAVSD repair, our group would aim towards a re-thoracotomy in these patients. However, we have to admit, that the experience is limited to only two cases necessitating reoperation due to endocarditis after PAVSD repair and ASD repair. Both patients had an uneventful perioperative course and good surgical results.

In the current presented study collective there were two deaths in the MS-groups, one in the MS-VSD-group and one in the MS-CAVSD-group. The patient in the MS-VSD suffered from Trisomy 18 and severe neurological and respiratory insufficiency. Following assessment of the situation of the patient, an interdisciplinary team advised against corrective cardiac surgery, nevertheless the surgical intervention was the explicit wish of the patient's parents.

The intraoperative course was uneventful, however the patient developed progressive global hemodynamic and respiratory insufficiency starting at postop day 12. Four days later the interdisciplinary team together with the parents decided for limited palliative therapy.

The patient in the MS-CAVSD-group was diagnosed with Trisomy 21. Surgery was performed on an elective schedule with uneventful intraoperative course. During the first postoperative day, the patient was resuscitated because of a severe pulmonary hypertension crisis. ECMO implantation was performed. Unfortunately, the patient died on the 5<sup>th</sup> postoperative day due to neurological complications and multi-organ failure. Overall the rate of MACE revealed no significant difference between the MS and VRAMT-groups.

In conclusion, this study confirms previous reports which described VRAMT as a feasible, efficient and safe approach for surgical repair of perimembranous VSD. Cardiopulmonary bypass can be performed without peripheral arterial cannulation. Furthermore, this approach can be applied to the correction of CAVSD with Rastelli type A. Considering the beneficial aesthetic factor of this less invasive approach and at the comparable surgical results in the long-term, VRAMT can be considered as an equal surgical approach for a broader range of congenital heart defects.

#### *Study limitations*

We acknowledge some limitations of our study. It was a retrospective observational analysis and therefore cause and effect are hard to establish.

**List of Abbreviations**

ASD - atrial septal defect  
BMI - body mass index  
CAVSD - complete atrioventricular septal defects  
CD – chest drains  
CI - confidence interval  
CPB - cardio-pulmonary-bypass  
CL - cardioplegia line  
ICD - implantable cardioverter defibrillator  
LBBB – left bundle branch block  
MACCE - major adverse cardiac and cerebrovascular events  
MI - myocardial infarction  
PAVSD - partial atrioventricular septal defects  
PE - pacing electrodes  
PFO - persistent foramen ovale  
PLSVC - persistent left superior vena cava,  
MS - median sternotomy  
NIRS - cerebral-near-infrared-spectroscopy  
REDCap - research electronic data capture  
SD - standard deviation  
SVC - superior vena cava  
TEE - transoesophageal-echocardiography  
URL – upper reference limit  
VRAMT - vertical-right-axillary-mini-thoracotomy  
VSD - ventricular septal defects

## Figure Legends

Figure 1. Placement of patients in a left decubitus position. The marked anterior axillary line and the 4<sup>th</sup> intercostal space serve as a guiding for the axillary incision.

Figure 2. (A) Vertical skin incision over the region of the 4<sup>th</sup> – 5<sup>th</sup> ICS with a length of 3 - 5 cm. Mobilisation of subcutaneous tissue and sparing of muscles. Visualization of the right atrium (arrow) and superior vena cava (SVC). (B) Direct access cannulation with aortic cannulation (Ao), venous cannulation of the superior vena cava (SVC) and cardioplegia line (CL). The inferior vena cava was cannulated using percutaneous technique.

Figure 3. (A) Surgical field after atriotomy for the correction of a perimembranous ventricular septal defect (VSD). (B) Closure of VSD with a xeno-pericardial patch (arrow).

Figure 4. (A) Initial postoperative access site after closed vertical skin incision. Chest drains (CD) in the right pleural cavity and placement of atrial/ventricular pacing electrodes (PE). (B) Long-term cosmetic result after VSD closure through VRAMT.

Table 1. Preoperative Data of VSD Patients.

Demographics VSD-cohort	MS N = 35	VRAMT N = 15	Difference and 95% confidence interval	p Value
Age [months]	4.1 (3.5; 5.7)	4.3 (3.3; 6.3)	-0.2 (-2.5; 2.2)	0.841
Premature Birth (SSW < 37)	6 (17%)	2 (13%)	4% (-19%; 27%)	1.000
Gender (Female)	11 (31%)	9 (60%)	-29% (-58%; 1%)	0.114
Weight (kilograms)	5.0 (4.4; 6.2)	5.7 (4.3; 6.8)	-0.3 (-1.2; 0.5)	0.525
Height (cm)	61.0 (58.0; 65.0)	60.0 (58.0; 69.0)	-1.1 (-5.2; 3.0)	0.688
BMI	13.7 (12.8; 14.3)	13.5 (12.2; 15.8)	-0.4 (-1.4; 0.7)	0.604
Body Surface Area [m <sup>2</sup> ]	0.3 (0.3; 0.3)	0.3 (0.3; 0.3)	-0.0 (-0.0; 0.0)	0.539
<b>Additional cardiac diagnosis</b>				
ASD Type II	17 (49%)	9 (60%)	-11% (-43%; 20%)	0.545
PLSVC	2 (6%)	1 (7%)	-1% (-16%; 14%)	1.000
PFO	7 (20%)	0 (0%)	20% (-1%; 41%)	0.087
<b>Non-cardiac diagnosis</b>				
Genetic	12 (34%)	2 (13%)	21% (-7%; 49%)	0.179
Trisomy 18	1 (3%)	0 (0%)	3% (-6%; 12%)	1.000
Trisomy 21	9 (26%)	2 (13%)	12% (-14%; 38%)	0.468
Di George-Syndrome (22q11-Syndrom)	1 (3%)	0 (0%)	3% (-6%; 12%)	1.000
Neurology	2 (6%)	0 (0%)	6% (-7%; 18%)	0.519
Metabolic	2 (6%)	1 (4%)	-1% (-16%; 14%)	1.000
Urogenital	1 (3%)	0 (0%)	3% (-6%; 12%)	1.000
Visceral	1 (3%)	0 (0%)	3% (-6%; 12%)	1.000

BMI: body mass index, ASD: atrial septal defect, PLSVC: persistent left superior vena cava, PFO: persistent foramen ovale. Variables are presented as numbers with % or as median with interquartile ranges.

Table 2. Preoperative Data of CAVSD Patients.

Demographics CAVSD-cohort	MS N = 24	VRAMT N = 10	Difference and 95% confidence interval	p Value
<b>Demographics</b>				
Age [months]	3.7 (3.1; 7.8)	6.1 (4.5; 10.5)	5.7 (-16.8; 28.1)	0.121
Premature Birth	5 (21%)	0 (0%)	21% (-6%; 48%)	0.291
Gender (Female)	13 (54%)	5 (50%)	4% (-35%; 43%)	1.000
Weight (kilograms)	4.9 (4.4; 6.9)	5.8 (4.6; 8.0)	0.6 (-3.6; 4.7)	0.174
Height (cm)	59.0 (56.0; 65.5)	67.0 (57.8; 68.5)	0.8 (-13.4; 15.0)	0.219
BMI	14.2 (13.2; 15.4)	14.4 (13.3; 17.1)	-0.6 (-2.2; 0.9)	0.484
Body Surface Area [m <sup>2</sup> ]	0.3 (0.3; 0.3)	0.3 (0.3; 0.4)	0.0 (-0.1; 0.2)	0.212
<b>Additional cardiac diagnosis</b>				
PLSVC	1 (4%)	0 (0%)	4% (-9%; 17%)	1.000
PFO	2 (8%)	0 (0%)	8% (-10%; 27%)	1.000
<b>Non-cardiac diagnosis</b>				
Trisomy 21	17 (71%)	4 (40%)	31% (-6%; 68%)	0.130
Neurology	2 (8%)	1 (10%)	-2% (-24%; 21%)	1.000
Metabolic	1 (4%)	2 (20%)	-16% (-37%; 6%)	0.201
Urogenital	1 (4%)	0 (0%)	4% (-9%; 17%)	1.000
Visceral	4 (17%)	0 (0%)	17% (-8%; 41%)	0.296

BMI: body mass index, PLSVC: persistent left superior vena cava, PFO: persistent foramen ovale.

Variables are presented as numbers with % or as median with interquartile ranges.

Table 3. Intra- and postoperative Data of VSD Patients.

Intraoperative Data	MS N = 35	VRAMT N = 15	Difference and 95% confidence interval	p Value
OR-time [min]	155 (140; 180)	155 (130; 180)	3.0 (-17.4; 23.3)	0.597
CPB-time [min]	63 (56; 77)	77 (63; 90)	-17.5 (-28.2; -6.9)	<b>0.015</b>
Aortic-cross-clamp-time [min]	36 (29; 41)	41 (33; 51)	-5.0 (-10.8; 0.9)	0.182
Postoperative Data	MS N = 35	VRAMT N = 15	Difference and 95% confidence interval	p Value
<b>Postoperative</b>				
Fast-Track: Extubation in OR	0 (0%)	1 (7%)	-7% (-15%; 2%)	0.300
ICU Stay [h]	96.0 (72.0; 166.0)	72.0 (48.0; 164.0)	30.0 (-14.2; 74.3)	0.212
Ventilation [h]	43.0 (22.0; 82.0)	24.0 (17.0; 47.0)	30.3 (-0.8; 61.4)	<b>0.043</b>
<b>Hospital Stay [in d]</b>	10.0 (8.0; 13.0)	9.0 (8.0; 12.0)	7.4 (-10.4; 25.1)	0.727
<b>Mortality</b>				
Mortality	1 (3%)	0 (0%)	3% (-6%; 12%)	1.000

OR: operation room, ICU: intensive care unit, CPB: cardio-pulmonary-bypass. Variables are presented as numbers with % or as median with interquartile ranges

Table 4. Intra- and postoperative Data of CAVSD Patients.

Intraoperative Data	MS N = 24	VRAMT N = 10	Difference and 95% confidence interval	p Value
OR-time [min]	225 (195; 269)	189 (171; 198)	52.8 (20.0; 85.6)	<b>0.003</b>
CPB-time [min]	106 (84; 144)	106 (93; 119)	8.0 (-13.8; 29.8)	0.597
Aortic-cross-clamp-time [min]	68 (55; 78)	64.0 (48; 72)	8.6 (-2.7; 19.9)	0.241

Postoperative Data	MS N = 24	VRAMT N = 10	Difference and 95% confidence interval	p Value
<b>ICU</b>				
Fast-Track: Extubation in OR	1 (4%)	0 (0%)	4% (-9%; 17%)	1.000
ICU Stay [h]	151.5 (92.3; 229.8)	126.0 (112.5; 192.0)	48.3 (-74.9; 171.5)	0.431
Ventilation [h]	71.5 (31.0; 100.8)	32.5 (23.8; 54.0)	43.4 (-1.3; 88.1)	0.059
<b>Hospital Stay [in d]</b>	14.0 (8.0; 29.0)	12.5 (8.0; 20.0)	6.3 (-6.4; 18.9)	0.681
<b>MACCE</b>				
Renal failure	1 (4%)	0 (0%)	4% (-9%; 17%)	1.000
Stroke	1 (4%)	0 (0%)	4% (-9%; 17%)	1.000
Pacemaker Implantation	1 (4%)	0 (0%)	4% (-9%; 17%)	1.000
<b>Mortality</b>				
Mortality	1 (4%)	0 (0%)	4% (-9%; 17%)	1.000

OR: operation room, ICU: intensive care unit, CPB: cardio-pulmonary-bypass MACCE: major adverse cardiac and cerebrovascular events. Variables are presented as numbers with % or as median with interquartile ranges.

## References

- [1] Hopkins RA, Bert AA, Buchholz B, Guarino K, Meyers M. Surgical patch closure of atrial septal defects. *Ats* 2004;77:2144–9–authorreply2149–50.
- [2] Khan JH, McElhinney DB, Reddy VM, Hanley FL. Repair of secundum atrial septal defect: limiting the incision without sacrificing exposure. *Ats* 1998;66:1433–5.
- [3] Nicholson IA, Bichell DP, Bacha EA, del Nido PJ. Minimal sternotomy approach for congenital heart operations. *Ats* 2001;71:469–72.
- [4] Dave HH, Comber M, Solinger T, Bettex D, Dodge-Khatami A, Prêtre R. Mid-term results of right axillary incision for the repair of a wide range of congenital cardiac defects. *Eur J Cardiothorac Surg* 2009;35:864–9–discussion869–70.
- [5] Prêtre R, Kadner A, Dave H, Dodge-Khatami A, Bettex D, Berger F. Right axillary incision: A cosmetically superior approach to repair a wide range of congenital cardiac defects. *J Thorac Cardiovasc Surg* 2005;130:277–81.
- [6] Yan L, Zhou Z-C, Li H-P, Lin M, Wang H-T, Zhao Z-W, et al. Right vertical infra-axillary mini-incision for repair of simple congenital heart defects: a matched-pair analysis. *Eur J Cardiothorac Surg* 2013;43:136–41.
- [7] Rosengart TK, Stark JF. Repair of atrial septal defect through a right thoracotomy. *Ann Thorac Surg* 1993;55:1138–40.
- [8] Mishaly D, Ghosh P, Preisman S. Minimally invasive congenital cardiac surgery through right anterior minithoracotomy approach. *Ann Thorac Surg* 2008;85:831–5.
- [9] Yoshimura N, Yamaguchi M, Oshima Y, Oka S, Ootaki Y, Yoshida M. Repair of atrial septal defect through a right posterolateral thoracotomy: a cosmetic approach for female patients. *Ats* 2001;72:2103–5.
- [10] Houyel L, Petit J, Planché C, Sousa-Uva M, Roussin R, Belli E, et al. [Right postero-lateral thoracotomy for open heart surgery in infants and children. Indications and results]. *Arch Mal Coeur Vaiss* 1999;92:641–6.
- [11] Seipelt RG, Popov A, Danner B, Paul T, Tirilomis T, Schoendube FA, et al. Minimally invasive partial inferior sternotomy for congenital heart defects in children. *J Cardiovasc Surg (Torino)* 2010;51:929–33.
- [12] Bleiziffer S, Schreiber C, Burgkart R, Regenfelder F, Kostolny M, Libera P, et al. The influence of right anterolateral thoracotomy in prepubescent female patients on late breast development and on the incidence of scoliosis. *J Thorac Cardiovasc Surg* 2004;127:1474–80.

- [13] Cherup LL, Siewers RD, Futrell JW. Breast and pectoral muscle maldevelopment after anterolateral and posterolateral thoracotomies in children. *Ats* 1986;41:492–7.
- [14] Wang Q, Li Q, Zhang J, Wu Z, Zhou Q, Wang D-J. Ventricular septal defects closure using a minimal right vertical infraaxillary thoracotomy: seven-year experience in 274 patients. *Ann Thorac Surg* 2010;89:552–5.
- [15] An G, Zhang H, Zheng S, Wang W, Ma L. Mid-term outcomes of Common Congenital Heart Defects Corrected through a Right Subaxillary Thoracotomy. *Heart Lung Circ* 2016;0.
- [16] An G, Zhang H, Zheng S, Wang W, Wu Q, Xing Q. Minimally invasive surgical closure for doubly committed subarterial ventricular septal defects through a right subaxillary thoracotomy. *Interact Cardiovasc Thorac Surg* 2016;23:924–8.
- [17] Hu C-X, Tan J, Chen S, Ding H, Xu Z-W. Comparison of clinical outcomes and postoperative recovery between two open heart surgeries: minimally invasive right subaxillary vertical thoracomy and traditional median sternotomy. *Asian Pac J Trop Med* 2014;7:625–9.
- [18] Dodge-Khatami A, Salazar JD. Right Axillary Thoracotomy for Transatrial Repair of Congenital Heart Defects: VSD, Partial AV Canal With Mitral Cleft, PAPVR or Warden, Cor Triatriatum, and ASD. *Operative Techniques in Thoracic and Cardiovascular Surgery* 2015;20:384–401.
- [19] Bastuji-Garin S, Sbidian E, Gaudy-Marqueste C, Ferrat E, Roujeau J-C, Richard M-A, et al. Impact of STROBE Statement Publication on Quality of Observational Study Reporting: Interrupted Time Series versus Before-After Analysis. *PLoS ONE* 2013;8:e64733.
- [20] Ghosh PK, Unger F. *Cardiac Reconstructions*. Berlin, Heidelberg: Springer Science & Business Media; 2012.
- [21] Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, White HD, Joint ESC/ACCF/AHA/WHF Task Force for the Universal Definition of Myocardial Infarction, et al. Third universal definition of myocardial infarction. *Circulation*, vol. 126, American Heart Association, Inc; 2012, pp. 2020–35.
- [22] Kaireit K, Deen J, Vernieuwe L, de Bruyn A, Kalantary S, Rodrigus I. Cardioplexol, a new cardioplegic solution for elective CABG. *J Cardiothorac Surg* 2013;8:P120.
- [23] Kadner A, Dodge-Khatami A, Dave H, Knirsch W, Bettex D, Prêtre R. Closure of restrictive ventricular septal defects through a right axillary thoracotomy. *Heart Surg Forum* 2006;9:E836–9.
- [24] Fonseca LD, Silva JPD, Turquetto AL, Franchi SM, Cascudo CM, Castro RM, et al. Horizontal right Axillary minithoracotomy: Aesthetic and effective option for ventricular and atrial septal defect repair in infants and toddlers. *Revista Brasileira De Cirurgia Cardiovascular* 2014.
- [25] Yang X, Wang D, Wu Q. Repair of Partial Atrioventricular Septal Defect through a Minimal Right Vertical Infra-Axillary Thoracotomy. *J Card Surg* 2003;18:262–4.

- [26] Bové T, Strubbe I, Vandekerckhove K, Panzer J, De Groote K, De Wolf D, et al. Surgical repair of atrioventricular septal defects: incidence and mode of failure of the left atrioventricular valve. *Interact Cardiovasc Thorac Surg* 2018;150:369.



