



# Whole-Genome Sequence of the First Extended-Spectrum $\beta$ -Lactamase-Producing Strain of *Salmonella enterica* subsp. *enterica* Serovar Napoli

Mathieu Clément,<sup>a,b</sup> Alban Ramette,<sup>a</sup> Odette J. Bernasconi,<sup>a,b</sup> Luigi Principe,<sup>c</sup> Francesco Luzzaro,<sup>c</sup>  Andrea Endimiani<sup>a</sup>

<sup>a</sup>Institute for Infectious Diseases, University of Bern, Bern, Switzerland

<sup>b</sup>Graduate School of Cellular and Biomedical Sciences, University of Bern, Bern, Switzerland

<sup>c</sup>Clinical Microbiology and Virology Unit, A. Manzoni Hospital, Lecco, Italy

**ABSTRACT** We report here the whole-genome sequence of the first extended-spectrum  $\beta$ -lactamase (ESBL)-producing strain of *Salmonella enterica* subsp. *enterica* serovar Napoli, LC0541/17, isolated from the stools of an ambulatory pediatric patient in northern Italy. The strain was of sequence type 474 (ST474) and possessed a 90-kb Inc11 ST49 plasmid carrying the *bla*<sub>CTX-M-1</sub> ESBL gene.

An increase in infections and outbreaks due to strains of *Salmonella enterica* subsp. *enterica* serovar Napoli has been recorded in Europe (1–3). This serovar has great propensity to cause bacteremia and is associated with hospitalization (4). However, only one draft genome sequence (GenBank accession number [LFIH00000000](https://doi.org/10.1128/MRA.00973-18)) is available (5), which is from strain SN310 belonging to sequence type 474 (ST474) and is resistant to aminoglycosides and responsible for a multischool outbreak in Milan, Italy (1). To our knowledge, *Salmonella* Napoli strains resistant to third-generation cephalosporins have not yet been reported.

In June 2017, an ~1-year-old girl living close to Milan was referred to the pediatrician because of diarrhea and abdominal pain. *Salmonella* Napoli strain LC0541/17 was isolated from the patient's stools. Susceptibility tests indicated that LC0541/17 was a suspected ESBL producer. Here, we describe the whole-genome sequence (WGS) of LC0541/17.

Genomic DNA was extracted from fresh overnight colonies grown on MacConkey agar at 36°C using the QIAamp DNA minikit (Qiagen). Both the MinION device (Oxford Nanopore Technologies) with the 1D library kit and the MiSeq platform (Illumina, Inc., done by GATC Biotech AG) with 2 × 150-bp reads were used to perform WGS. Adapters from the Nanopore and Illumina reads were removed using Porechop 0.2.3 (<https://github.com/rrwick/Porechop>) and Trimmomatic 0.33 (6), respectively. A total of 39,878 of the longest and best-quality Nanopore reads were extracted until reaching 500,001,524 bp (*N*<sub>50</sub>, 12,177 bp; maximum, 58,306 bp) using Filtlong 0.2.0 (<https://github.com/rrwick/Filtlong>), and contigs were assembled using Canu 1.5 (7). Contigs were circularized using Circlator 1.5.5 (8). The assembly was polished with Illumina-trimmed reads using Pilon 1.22, and annotation was performed using Prokka 1.13 (9, 10). Strain and plasmid STs were obtained using mlst 2.10 (<https://github.com/tseemann/mlst>), which relies on the PubMLST database (<https://pubmlst.org/>). Plasmid and antimicrobial resistance genes (ARGs) were found using ABRicate 0.8 (<https://github.com/tseemann/abricate>), which uses ResFinder and PlasmidFinder databases and pathogenicity islands using SPIFinder (<https://cge.cbs.dtu.dk/services/SPIFinder/>). Virulence gene profiles (VPs) were assessed using BLASTn. All software was run with default or advised parameters.

Two circular contigs were obtained, one chromosome and one plasmid (pLC0541\_17).

Received 11 July 2018 Accepted 16 August 2018 Published 13 September 2018

**Citation** Clément M, Ramette A, Bernasconi OJ, Principe L, Luzzaro F, Endimiani A. 2018. Whole-genome sequence of the first extended-spectrum  $\beta$ -lactamase-producing strain of *Salmonella enterica* subsp. *enterica* serovar Napoli. Microbiol Resour Announc 7:e00973-18. <https://doi.org/10.1128/MRA.00973-18>.

**Editor** Julie C. Dunning Hotopp, University of Maryland School of Medicine

**Copyright** © 2018 Clément et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Andrea Endimiani, [aendimiani@gmail.com](mailto:aendimiani@gmail.com).

The chromosome size was 4,679,033 bp, with a GC content of 52.2%. Annotation indicated 4,555 coding sequences (CDSs) (not including ARGs), 22 rRNAs, 84 tRNAs, and 1 transfer-messenger RNA (tmRNA). LC0541/17 was of ST474, and five pathogenicity islands were detected, of which three were also observed in strain SN310. Ten *Salmonella* Napoli VPs were previously characterized (2). LC0541/17 was found to belong to the VP1 profile [*ssaQ mgtC spi\_4D(siiD) sopB sopE1 bcfC*], which is the most prevalent (36%) among isolates from humans, animals, food, or the environment (2).

The size of pLC0541\_17 was 90,558 bp, with a GC content of 49.1%. It was an IncI1 ST49 plasmid with 102 CDSs carrying the *bla*<sub>CTX-M-1</sub> gene but no other ARGs. The CTX-M-1 group ESBLs are the most prevalent in Italy (11); however, only several 80- to 100-kb IncI1 ST49 CTX-M-1 plasmids have been reported in Denmark in *Escherichia coli* of either human, pig, or cattle origin (12). In particular, GenBank database comparison (BLASTn) revealed that pLC0541\_17 was similar (99% identity, 99 to 100% coverage) to two of these plasmids (GenBank accession numbers [KJ563250](#) and [KM052220](#)) and another from France (GenBank accession number [LT985288](#)). As for these three plasmids, pLC0541\_17 carried the *bla*<sub>CTX-M-1</sub> gene inserted into the following genetic environment: *pilJ orf477 bla*<sub>CTX-M-1</sub> *ISecpB1 pill*.

The emergence of ESBL-producing *Salmonella* Napoli isolates is alarming. Public health institutions should coordinate surveillance programs to monitor the spread of these life-threatening pathogens.

**Data availability.** The sequences of the chromosome and plasmid are deposited in GenBank under accession numbers [CP030838](#) and [CP030839](#), respectively. Raw data are available in the Sequence Read Archive under accession number [SRP156370](#).

## ACKNOWLEDGMENTS

This work was supported by NRP-72, National Research Programme “Antimicrobial Resistance” (Swiss National Science Foundation; grant 177378 to A.E. and A.R.). M.C. is a Ph.D. student (2018 to 2021) supported by grant NRP-72. O.J.B. is a Ph.D. student (2015 to 2018) supported by the Hans Sigrist Foundation, Bern, Switzerland.

## REFERENCES

- Huedo P, Gori M, Amato E, Bianchi R, Valerio E, Magnoli L, Pontello M. 2016. A multischool outbreak due to *Salmonella enterica* serovar Napoli associated with elevated rates of hospitalizations and bacteremia, Milan, Italy, 2014. *Foodborne Pathog Dis* 13:417–422. <https://doi.org/10.1089/fpd.2015.2091>.
- Graziani C, Busani L, Dionisi AM, Caprioli A, Ivarsson S, Hedenström I, Luzzi I. 2011. Virulotyping of *Salmonella enterica* serovar Napoli strains isolated in Italy from human and nonhuman sources. *Foodborne Pathog Dis* 8:997–1003. <https://doi.org/10.1089/fpd.2010.0833>.
- Sabbatucci M, Dionisi AM, Pezzotti P, Lucarelli C, Barco L, Mancin M, Luzzi I. 2018. Molecular and epidemiologic analysis of reemergent *Salmonella enterica* serovar Napoli, Italy, 2011–2015. *Emerg Infect Dis* 24:562–565. <https://doi.org/10.3201/eid2403.171178>.
- Huedo P, Gori M, Zolin A, Amato E, Ciceri G, Bossi A, Pontello M. 2017. *Salmonella enterica* serotype Napoli is the first cause of invasive nontyphoidal salmonellosis in Lombardy, Italy (2010–2014), and belongs to Typhi subclade. *Foodborne Pathog Dis* 14:148–151. <https://doi.org/10.1089/fpd.2016.2206>.
- Huedo P, Gori M, Scaltriti E, Morganti M, Casadei G, Amato E, Pontello M. 2015. Draft genome sequence of *Salmonella enterica* subsp. *enterica* serovar Napoli strain SN310, cause of a multischool outbreak in Milan, Italy, in 2014. *Genome Announc* 3:e01044–15. <https://doi.org/10.1128/genomeA.01044-15>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res* 27:722–736. <https://doi.org/10.1101/gr.215087.116>.
- Hunt M, Silva ND, Otto TD, Parkhill J, Keane JA, Harris SR. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biol* 16:294. <https://doi.org/10.1186/s13059-015-0849-0>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
- Giani T, Antonelli A, Caltagirone M, Mauri C, Nicchi J, Arena F, Nucleo E, Bracco S, Pantosti A, Luzzaro F, Pagani L, Rossolini GM. 2017. Evolving  $\beta$ -lactamase epidemiology in *Enterobacteriaceae* from Italian nationwide surveillance, October 2013: KPC-carbapenemase spreading among outpatients. *Euro Surveill* 22:pil=30583. <https://www.eurosurveillance.org/content/10.2807/1560-7917.ES.2017.22.31.30583>.
- Jakobsen L, Bortolaia V, Bielak E, Moodley A, Olsen SS, Hansen DS, Frimodt-Møller N, Guardabassi L, Hasman H. 2015. Limited similarity between plasmids encoding CTX-M-1  $\beta$ -lactamase in *Escherichia coli* from humans, pigs, cattle, organic poultry layers and horses in Denmark. *J Glob Antimicrob Resist* 3:132–136. <https://doi.org/10.1016/j.jgar.2015.03.009>.