Outcomes of Emergency Abdominal Surgery in Octogenarians:

A single-center analysis

Joël L. Lavanchy^a, Melanie M. Holzgang^a, Tobias Haltmeier^a, Daniel Candinas^a, Beat Schnüriger^a

^a Department of Visceral Surgery and Medicine, Inselspital, Bern University Hospital,
University of Bern, Switzerland

DOI: <https://doi.org/10.1016/j.amjsurg.2018.11.023>

Correspondence to:

Beat Schnüriger, MD

Department of Visceral Surgery and Medicine

Inselspital, Bern University Hospital

3010 Bern, Switzerland

beat.schnuriger@gmail.com

+41 (0)31 632 74 26

Presented at the Annual Swiss Trauma and Acute Care Surgery Day, March 16-17

2018, Bern, Switzerland, at the European Congress for Trauma & Emergency Surgery,

May 6-8 2018, Valencia, Spain and at the Annual Meeting of the Swiss Society of

Surgery, May 16-18 2018, Basel, Switzerland

Short summary

Octogenarians undergoing emergency abdominal surgery had a mortality rate of 16.4%. Mortality was independently predicted by age, ASA score $\geq$ 4, mesenteric ischemia and ICU admission. Therefore, the indication for emergency abdominal surgery should be assessed cautiously, including patients' and relatives' wishes, surgeons, intensivists, anesthesiologists, and nursing staff.

Keywords:

Emergency abdominal surgery; octogenarians; mortality; early mortality; loss of independency

Abbreviations

ASA: American society of Anesthesiologists

BMI: body mass index

CCI: Charlson Comorbidity Index

EAS: emergency abdominal surgery

HLOS: hospital length of stay

ICU intensive care unit

ICU LOS: intensive care unit length of stay

Introduction

As life expectancy is increasing worldwide, societies are challenged to adapt health care systems to the aging population.¹ There is an ongoing discussion whether the evolution of medicine and health technology will be able to compensate for the increasing age-related burden of disease.² Adding to the problem, increasing life expectancy actually is one of the main drivers for increasing health care expenditures.³ Therefore the effectiveness of medical treatments in patients with extensive treatment needs, such as surgery in the elderly,⁴ is of utmost importance for a fair allocation of resources.

In a large retrospective cohort study including 1.8 million individuals, over 18% of patients underwent surgery in the last month of their life.⁵ Among other factors, higher age has been shown to be associated with deficits in surgical decision making.⁶ Multidimensional patient-centred approaches⁷ and best communication practices^{8, 9} have been discussed to facilitate decision making in elderly patients with surgical emergencies. Recent studies investigated outcomes of patients >80 years of age undergoing cardiovascular or neurosurgical interventions.¹⁰⁻¹² However, literature on outcomes of octogenarians undergoing emergency abdominal surgery (EAS) is scarce.¹³ Therefore, this study opted to assess the overall and early in-hospital mortality rate, loss of independence, and factors associated with these outcomes in octogenarians undergoing EAS.

Material and methods

This study is reported in accordance with the STROBE (Strengthening the Reporting of Observational studies in Epidemiology) statement¹⁴ and was approved by the cantonal ethics committee of Bern, Switzerland (KEK 2017-01284).

Study design

This is a single centre retrospective observational study including patients ≥ 80 years of age that underwent non-trauma emergency abdominal surgery at the Bern University Hospital from 01/12/2011 to 31/12/2016. Patients with malignant disease and those who underwent non-operative treatment for acute abdominal disease were excluded (Figure 1).

Irrespective of age, the treating surgical team assessed all patients that potentially require EAS. The goals of care were discussed with the patients and their relatives. Non-curative treatment was initiated only if the patients specifically expressed their wish for palliative care.

Data were extracted from the Bern University Hospital administrative database and electronic patient charts. The following variables were collected: age, sex, body mass index (BMI), vital signs and laboratory values at admission, American Society of Anesthesiologists (ASA) score,¹⁵ Charlson Comorbidity Index (CCI),¹⁶ diagnoses, type of surgery (open, laparoscopic), intensive care unit length of stay (ICU LOS), hospital length of stay (HLOS), nursing effort (hours of nursing per hospital day), complications according

to Dindo Clavien classification,¹⁷ origin of admittance/discharge (home, nursing facility, other hospital), and in-hospital mortality.

The main outcome was overall in-hospital mortality. Secondary outcomes included early mortality and loss of independence. Early mortality was defined as death ≤ 7 days from surgery. Loss of independence was defined as being admitted from home and being discharged to a nursing facility or other hospital.

The association of clinically important variables, including patient and treatment characteristics, on outcomes was assessed in univariable and multivariable regression analysis.

Statistical analysis

Normality of distribution was assessed using histograms and the Shapiro-Wilk test. Results were reported as numbers and percentages or medians and interquartile ranges, as appropriate. Categorical variables were compared using Chi-Square or Fisher's exact test. Students' t test or Mann-Whitney U test were used for analysis of continuous variables.

The impact of clinical variables on in-hospital mortality and loss of independency was assessed in univariable analysis. Clinical variables with a p -value ≤ 0.1 in univariable analysis were then included in a multivariable regression model in order to assess independency. Subsequently, forward logistic regression was performed to create an independent predictive model for mortality and loss of independency, respectively. Mortality rates were shown as Kaplan Meier curves. P -values ≤ 0.05 were considered

statistically significant. Statistical analysis was performed using SPSS Statistics version 25 (IBM Corporation, Armonk, New York).

Results

Within the 61-months study period, 752 octogenarians with acute abdominal disease were screened. After the exclusion of patients treated non-operatively and those with malignant disease, a total of 140 octogenarians undergoing EAS were included in the analysis (*Figure 1*).

Median age was 83.9 (IQR 81.9-87.2) years, 55.7% (n=78) were female, and the median BMI was 24.5 (17.1-45.7) kg/m² (*Table 1*). An ASA score ≥ 4 was found in 40.7% (n=57) and 47.2 % (n=66) had a CCI ≥ 3 . Patients were treated for cholecystitis (27.1%, n=38), small or large bowel obstruction (22.1%, n=31), hollow viscus perforation (16.4%, n=23), perforated diverticulitis (12.9%, n=18), mesenteric ischemia (10.0%, n=14), incarcerated hernia (9.3%, n=13), and appendicitis (2.1%, n=3). A total of 37 patients (26.4%) were operated laparoscopically, 73.6% underwent laparotomy (n=103). The postoperative ICU-admission rate was 47.9%. Median HLOS and ICULOS was days 9.0 (6.0-15.8) and 2.0 (1.0-4.0) days, respectively. A total of 39 octogenarians (27.9 %) revealed postoperative complications Dindo Clavien ≥ 3 .

Overall mortality

Overall in-hospital mortality was 16.4% (n=23). *Table 1* outlines the demographics and characteristics of the study population stratified by mortality. When comparing survivors and non-survivors, a statistically significant age difference (83.6 vs. 86.4 years, $p=0.024$) was found. Moreover, ASA scores ≥ 4 were significantly less frequent (32.5% vs. 82.6%, $p<0.001$) and the ICU admission rate was significantly lower (41.0% vs. 82.6%, $p<0.001$)

in survivors than in non-survivors. On the other hand, survivors were significantly more often admitted from home compared to non-survivors (64.1% vs. 34.8%, $p=0.010$). Regarding abdominal disease, acute cholecystitis was more frequent (32.5% vs. 0.0%, $p=0.001$) and mesenteric ischemia less frequent (3.4% vs. 43.5%, $p<0.001$) in survivors compared to non-survivors. Of note, survivors were more frequently operated laparoscopically compared to the non-survivors (29.9% vs. 8.7% $p=0.039$). Multivariable forward regression analysis revealed following independent predictors for overall mortality: age, ASA scores ≥ 4 , mesenteric ischemia and ICU admission (*Table 2*). The logistic regression model fit the data well (Nagelkerkes' $R^2=0.575$)

Early mortality

Fourteen of 23 non-survivors died within 7 days and were defined as early mortalities. Ultimately, multi-organ failure ($n=7$, 50%), postoperative withdrawal of care ($n=5$, 36%), cardiac arrest ($n=1$, 7%), and respiratory insufficiency ($n=1$, 7%) led to the early deaths. The reasons for postoperative withdrawal of care were as following: patients' will ($n=2$), relatives' wishes ($n=3$). The demographics of this subgroup of patients are presented in *Table 3*. Figures 2 A-C are showing the Kaplan Meier survival curves stratified by the categorical independent predictors for mortality. Of note, Octogenarians with an ASA score ≥ 4 or with mesenteric ischemia showed earlier mortality (*Figure 2 A and 2 B*).

Loss of independency

A total 76 of the 117 patients (54.3%) that survived their hospital stay were admitted from home. Of these, 29 patients (38.2%) were subsequently discharged to a nursing facility or another hospital and were defined as patients with loss of independency. *Table 4* outlines the demographics and characteristics of the study population stratified by loss of independency. Patients with loss of independency showed significantly higher temperatures (37.2 vs. 36.8 °C, $p=0.040$) and respiratory frequencies (22 vs. 18, $p=0.007$) on admission, had a higher nursing intensity during their hospital stay (8.9 vs. 7.1 hours per day, $p=0.002$) and were significantly more often admitted to the ICU (48.3% vs. 23.9%, $p=0.044$) compared to patients without loss of independency. In multivariable logistic regression analysis the worst lactate level within 24 hours from admission was independently associated with a loss of independency (Nagelkerkes' $R^2=0.322$).

Discussion

In the current study, a total of 140 octogenarians undergoing EAS were retrospectively assessed. The overall in-hospital mortality was 16.4%. Age, ASA scores ≥ 4 , mesenteric ischemia and ICU admission were independently associated with mortality. Of those that survived and were admitted from home, loss of independency was seen in 38.2% of patients. The worst lactate level within 24 hours of admission was independently associated with a loss of independency.

Overall Mortality

The current population revealed a 16% mortality rate, which is in accordance with two currently available studies showing a 15 to 33% mortality rate in an emergency surgery population ≥ 80 years.^{18, 19} In the current study, acute mesenteric ischemia was identified as a strong independent predictor for mortality, whereas acute cholecystitis was associated with decreased mortality. This should be taken into account when therapeutic decisions are made in this age group. Furthermore, the severity of pathophysiological derangements as indicated by an ASA score ≥ 4 or postoperative ICU admission, was independently associated with significantly higher mortality.

In the current study, ICU admission was an independent predictor for mortality. The pathophysiologic derangement should carefully be assessed before admitting a patient to the ICU. ICU admission may lead to overtreatment. Of note, a large trial including patients aged ≥ 75 years showed a significantly higher rate of ICU admission and consecutively

increased in-hospital mortality in the group of patients that were preoperatively systematically evaluated by intensivists.²⁰

Surprisingly, in the current series, the CCI did not predict mortality. This might be due to equally distributed co-morbidities in surviving and non-surviving octogenarians. Similarly, a previous study showed that comorbidities do not accurately predict mortality in octogenarians undergoing EAS. However, in this study detailed information on abdominal pathologies were not reported.¹⁹

Age was independently associated with mortality. This underlines the fact that age persists to be a strong predictor for worse outcomes even in octogenarians. This is of particular relevance in the context of Switzerland's high life expectancy, which is the second largest in the world.¹ A similar observation has been made in a Finnish population, however, in patients aged ≥ 65 years.²¹

Although not an independent predictor for mortality, preoperative independent living seems to influence outcomes of octogenarians undergoing EAS. In the current univariate analysis, octogenarians coming from home survived significantly more often. This has also been demonstrated in another study including patients aged ≥ 85 years, although this study analysed a mixed population of patients undergoing non-emergency and emergency abdominal surgery.²² Similarly, an increased mortality of patients ≥ 65 years of age undergoing emergent and non-emergent abdominal surgery has been demonstrated for nursing home residents compared to patients admitted from home.²³

Early mortality

Ten percent of octogenarians died within 7 days of surgery and were defined as early mortalities. This subgroup of patients was separately analysed in order to investigate ethically questionable surgical interventions. When further investigating the ultimate reasons for the mortalities, it was found that over a third of early mortalities or a fifth of overall deaths were due to withdrawal of care. This observation emphasizes the importance to elaborate an interdisciplinary treatment plan in accordance with the patients' will and relatives' wishes before surgery is performed. Here a clear communication containing the actual diagnosis, the pre-existing health conditions, and the risks and benefits of surgery is required to avoid unrealistic expectations by patients and families.⁸ There are different frailty scores available, that possibly help to further improve decision making, however, they need to be carefully validated in this subgroup of very old patients.²⁴⁻²⁶

Loss of independency

To our knowledge, this is the first study assessing the predictors for loss of independency of octogenarians underdoing EAS. In this study, the highest lactate level within 24 hours after admission was independently associated with loss of independency. Lactate level is a known predictor for mortality in EAS of elderly patients aged over 70 years²⁷ and a well-known surrogate marker for the severity of sepsis.²⁸ However, it has never been shown as a predictor for loss of independency. In the current study, higher lactate levels as a predictor for loss of independency rather reflect the overall severity of disease than a direct effect of lactate on the loss of independency.

Limitations

Major limitation of this study is its retrospective design. Therefore, the dataset is limited to the variables available from the institutional electronic charts. In the literature there are multiple specific frailty indices described, however, in clinical practice these need to be applied prospectively.²⁴⁻²⁶ Furthermore, the mortality after hospital discharge could not be obtained and therefore a long-term survival analysis was not available. Moreover, the heterogeneity of the patient population and the relatively small number of patients does not allow for stratification and subgroup analysis. Therefore, the conclusions were drawn cautiously.

Conclusions

In octogenarians undergoing EAS, age, ASA score ≥ 4 , mesenteric ischemia and ICU admission was independently associated with mortality. One third of early mortality was due to withdrawal of care. Therefore, to prevent ethically questionable acute surgical interventions, the deliberation of treatment options and the goals of care should involve emergency physicians, surgeons, anaesthesiologists, and intensivists. Furthermore, the treatment goals needs to be in consistency with the patients' will or with the relatives in their role as proxy decision makers for the patient.

Acknowledgements

The Authors Joël Lavanchy, Melanie Holzgang, Tobias Haltmeier, Daniel Candinas and Beat Schnüriger have no conflict of interest or financial ties to declare.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

1. World health statistics 2016: monitoring health for the SDGs, sustainable development goals. Geneva, Switzerland 2016.
2. Christensen K, Doblhammer G, Rau R, Vaupel JW. Ageing populations: the challenges ahead. *Lancet*. 2009;374(9696):1196-208. [https://doi.org/10.1016/S0140-6736\(09\)61460-4](https://doi.org/10.1016/S0140-6736(09)61460-4).
3. de Meijer C, Wouterse B, Polder J, Koopmanschap M. The effect of population aging on health expenditure growth: a critical review. *Eur J Ageing*. 2013;10(4):353-61. <https://doi.org/10.1007/s10433-013-0280-x>.
4. Regenbogen SE, Gust C, Birkmeyer JD. Hospital surgical volume and cost of inpatient surgery in the elderly. *J Am Coll Surg*. 2012;215(6):758-65. <https://doi.org/10.1016/j.jamcollsurg.2012.07.011>.
5. Kwok AC, Semel ME, Lipsitz SR, et al. The intensity and variation of surgical care at the end of life: a retrospective cohort study. *Lancet*. 2011;378(9800):1408-13. [https://doi.org/10.1016/S0140-6736\(11\)61268-3](https://doi.org/10.1016/S0140-6736(11)61268-3).
6. Cooper Z, Hevelone N, Sarhan M, et al. Identifying Patient Characteristics Associated With Deficits in Surgical Decision Making. *J Patient Saf*. 2016. <https://doi.org/10.1097/PTS.0000000000000323>.
7. Glance LG, Osler TM, Neuman MD. Redesigning surgical decision making for high-risk patients. *N Engl J Med*. 2014;370(15):1379-81. <https://doi.org/10.1056/NEJMp1315538>.
8. Cooper Z, Koritsanszky LA, Cauley CE, et al. Recommendations for Best Communication Practices to Facilitate Goal-concordant Care for Seriously Ill Older Patients With Emergency Surgical Conditions. *Ann Surg*. 2016;263(1):1-6. <https://doi.org/10.1097/SLA.0000000000001491>.
9. Cauley CE, Block SD, Koritsanszky LA, et al. Surgeons' Perspectives on Avoiding Nonbeneficial Treatments in Seriously Ill Older Patients with Surgical Emergencies: A Qualitative Study. *J Palliat Med*. 2016;19(5):529-37. <https://doi.org/10.1089/jpm.2015.0450>.

10. Kawahito K, Kimura N, Yamaguchi A, et al. Early and Late Surgical Outcomes of Acute Type A Aortic Dissection in Octogenarians. *Ann Thorac Surg*. 2017.
<https://doi.org/10.1016/j.athoracsur.2017.06.057>.
11. Kuo K, Shah P, Hiebert B, et al. Predictors of survival, functional survival, and hospital readmission in octogenarians after surgical aortic valve replacement. *J Thorac Cardiovasc Surg*. 2017;154(5):1544-53 e1. <https://doi.org/10.1016/j.jtcvs.2017.05.047>.
12. Steinberger J, Bronheim RS, Vempati P, et al. Morbidity and Mortality of Meningioma Resection Increases in Octogenarians. *World Neurosurg*. 2017.
<https://doi.org/10.1016/j.wneu.2017.09.021>.
13. Fukuda N, Wada J, Niki M, et al. Factors predicting mortality in emergency abdominal surgery in the elderly. *World J Emerg Surg*. 2012;7(1):12.
<https://doi.org/10.1186/1749-7922-7-12>.
14. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet*. 2007;370(9596):1453-7. [https://doi.org/10.1016/S0140-6736\(07\)61602-X](https://doi.org/10.1016/S0140-6736(07)61602-X).
15. Saklad M. Grading of patients for surgical procedures. *Anesthesiology*. 1941;2(3):281-4.
16. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373-83.
17. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg*. 2004;240(2):205-13. <https://doi.org/10.1097/01.sla.0000133083.54934.ae>.
18. Ford PN, Thomas I, Cook TM, et al. Determinants of outcome in critically ill octogenarians after surgery: an observational study. *Br J Anaesth*. 2007;99(6):824-9.
<https://doi.org/10.1093/bja/aem307>.
19. Rubinfeld I, Thomas C, Berry S, et al. Octogenarian abdominal surgical emergencies: not so grim a problem with the acute care surgery model? *J Trauma*. 2009;67(5):983-9. <https://doi.org/10.1097/TA.0b013e3181ad6690>.
20. Guidet B, Leblanc G, Simon T, et al. Effect of Systematic Intensive Care Unit Triage on Long-term Mortality Among Critically Ill Elderly Patients in France: A

Randomized Clinical Trial. *JAMA*. 2017;318(15):1450-9.

<https://doi.org/10.1001/jama.2017.13889>.

21. Ukkonen M, Kivivuori A, Rantanen T, Paaianen H. Emergency Abdominal Operations in the Elderly: A Multivariate Regression Analysis of 430 Consecutive Patients with Acute Abdomen. *World J Surg*. 2015;39(12):2854-61.

<https://doi.org/10.1007/s00268-015-3207-1>.

22. Mirbagheri N, Dark JG, Watters DA. How do patients aged 85 and older fare with abdominal surgery? *J Am Geriatr Soc*. 2010;58(1):104-8. <https://doi.org/10.1111/j.1532-5415.2009.02612.x>.

23. Finlayson E, Wang L, Landefeld CS, Dudley RA. Major abdominal surgery in nursing home residents: a national study. *Ann Surg*. 2011;254(6):921-6.

<https://doi.org/10.1097/SLA.0b013e3182383a78>.

24. Rockwood K, Stadnyk K, MacKnight C, et al. A brief clinical instrument to classify frailty in elderly people. *Lancet*. 1999;353(9148):205-6. [https://doi.org/10.1016/S0140-6736\(98\)04402-X](https://doi.org/10.1016/S0140-6736(98)04402-X).

25. Rockwood K, Song X, MacKnight C, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ*. 2005;173(5):489-95.

<https://doi.org/10.1503/cmaj.050051>.

26. Rolfson DB, Majumdar SR, Tsuyuki RT, et al. Validity and reliability of the Edmonton Frail Scale. *Age Ageing*. 2006;35(5):526-9.

<https://doi.org/10.1093/ageing/afl041>.

27. Sharrock AE, McLachlan J, Chambers R, et al. Emergency Abdominal Surgery in the Elderly: Can We Predict Mortality? *World J Surg*. 2017;41(2):402-9.

<https://doi.org/10.1007/s00268-016-3751-3>.

28. Cheng HH, Chen FC, Change MW, et al. Difference between elderly and non-elderly patients in using serum lactate level to predict mortality caused by sepsis in the emergency department. *Medicine (Baltimore)*. 2018;97(13):e0209.

<https://doi.org/10.1097/MD.00000000000010209>.

Table 1: Patient's baseline characteristics and comparison of survivors with non-survivors

	Overall (N=140)	Survivors (n=117)	Non-survivors (n=23)	p value
Age, y, median (IQR)	83.9 (81.9-87.2)	83.6 (81.6-87.1)	86.4 (83.3-88.3)	0.024*
Sex, female/male, n (%)	78/62 (55.7/44.3)	67/50 (57.3/42.7)	11/12 (47.8/52.2)	0.493†
BMI, kg/m ² , median (IQR)	24.5 (22.4-27.7)	24.5 (22.5-28.0)	23.9 (21.4-26.2)	0.569*
ASA score, n (%)				
I	-	-	-	<0.001†
II	7 (5.0)	7 (6.0)	-	
III	76 (54.3)	72 (61.5)	4 (17.4)	
IV	49 (35.0)	37 (31.6)	12 (52.2)	
V	8 (5.7)	1 (0.9)	7 (30.4)	
Charlson Comorbidity Index, n (%)				
0	23 (16.4)	19 (16.2)	4 (17.4)	0.931†
1	20 (14.3)	17 (14.5)	3 (13.0)	
2	31 (22.1)	27 (23.1)	4 (17.4)	
3	25 (17.9)	20 (17.1)	5 (21.7)	
4	21 (15.0)	18 (15.4)	3 (13.0)	
≥5	20 (14.3)	16 (13.7)	4 (17.4)	
Diagnosis, n (%)				
Cholecystitis	38 (27.1)	38 (32.5)	-	0.001†
Small or large bowel obstruction	31 (22.1)	27 (23.1)	4 (17.4)	0.784†
Hollow viscus perforation	23 (16.4)	19 (16.2)	4 (17.4)	1.000†
Diverticulitis	18 (12.9)	14 (12.0)	4 (17.4)	0.498†
Mesenteric ischemia	14 (10.0)	4 (3.4)	10 (43.5)	<0.001†
Hernia	13 (9.3)	12 (10.3)	1 (4.3)	0.694†
Appendicitis	3 (2.1)	3 (2.6)	-	1.000†
Vital signs at admission, median (IQR)				
Temperature, °C	36.8 (36.5-37.3)	36.9 (36.5-37.4)	36.7 (36.3-37.1)	0.155*
Respiratory frequency, /min.	21 (17-24)	21 (17-24)	22 (19-27)	0.437*
Heart rate, /min.	87 (75-102)	85 (75-99)	96 (79-114)	0.103‡
Systolic blood pressure, mmHg	134 (117-159)	135 (117-159)	130 (109-159)	0.869*
Laboratory values at admission, median (IQR)				
White blood cell count, G/l	12.5 (9.4-16.1)	12.2 (9.2-14.8)	16.1 (11.1-19.5)	0.024*
CRP, mg/l	47 (10-222)	57 (11-240)	27 (5-162)	0.031‡
Creatinine, mmol/l	100 (77-137)	98 (75-131)	115 (94-182)	0.103*
Creatinine clearance,	51 (35-72)	55 (36-72)	38 (28-48)	0.047*
Lactate at admission, mmol/l	1.7 (1.2-2.6)	1.6 (1.2-2.5)	2.2 (1.5-4.4)	0.028‡
Lactate, worst within first 24 h, mmol/l	2.0 (1.2-2.9)	1.8 (1.2-2.7)	3.3 (2.0-6.4)	0.001*
pH	7.40 (7.32-7.43)	7.40 (7.32-7.43)	7.41 (7.33-7.44)	0.727*
Sepsis at admission, n (%)	85 (60.7)	74 (63.2)	11 (47.8)	0.242†
Nursing effort, h per day, median (IQR)	9.2 (7.1-13.0)	8.8 (6.9-11.5)	15.5 (10.9-23.1)	0.215‡
Admission from, n (%)				
Home	83 (59.3)	75 (64.1)	8 (34.8)	0.010†
Nursing home, rehab.	11 (7.9)	10 (8.5)	1 (4.3)	
Other Hospital	46 (32.9)	32 (27.4)	14 (60.9)	
Operation technique, n (%)				
Laparotomy	117 (83.6)	82 (70.1)	21 (91.3)	0.039†
Laparoscopy	23 (16.4)	35 (29.9)	2 (8.7)	
ICU admission, n (%)	67 (49.7)	48 (41.0)	19 (82.6)	<0.001†

IQR: interquartile range; BMI: body mass index; ASA score: American Society of Anesthesiology score; ICU: intensive care unit; * Mann-Whitney-U test; † Fisher's exact test; ‡ Students' t test.

Table 2: Multivariable analysis of outcomes

	In hospital mortality		
	OR	95% CI	<i>p</i> value
Age, y	1.24	1.04-1.47	0.015
ASA score ≥ 4	11.15	2.39-52.02	0.002
Mesenteric ischemia	52.60	8.93-309.94	<0.001
ICU admission	9.23	1.74-49.04	0.009
	Loss of independency		
	OR	95% CI	<i>p</i> value
Lactate, worst within first 24 h, mmol/l	2.36	1.09-5.10	0.029
Nursing intensity, h per day	1.16	0.97-1.39	0.103

Forward stepwise logistic regression analysis.

ASA score: American Society of Anesthesiology score; ICU: intensive care unit; OR: Odds ratio; CI: confidence interval

Table 3: Characteristics of patients with withdrawal of care

Patient	Age (years)	Sex	ASA score	Sepsis at admission	Diagnosis	Intervention	Time of withdrawal of care
1	91.5	Male	5	Yes	Colonic perforation (splenic flexure)	Subtotal colectomy, end ileostomy	POD 1 (by relatives)
2	88.3	Female	4	Yes	Mesenteric ischemia	Thrombectomy SMA, segmental small bowel resection	POD 1 (by patient)
3	95.6	Female	5	No	Small bowel obstruction	Adhesiolysis	POD 4 (by relatives)
4	86.4	Male	4	No	Small bowel obstruction	Adhesiolysis	POD 4 (by relatives)
5	88.1	Female	4	Yes	Colonic perforation (sigmoid colon)	Rectosigmoidectomy, end colostomy	POD 6 (by patient)

ASA score: American Society of Anesthesiology score; SMA: superior mesenteric artery; POD: Postoperative day

Table 4: Comparison of patients stratified by loss of independency

	No loss of independency (n=47)	Loss of independency (n=29)	p value
Age, y, median (IQR)	84.0 (81.9-87.1)	83.9 (81.2-86.9)	0.724*
Sex, female/male, n (%)	28/19 (59.6/40.4)	15/14 (51.7/48.3)	0.634†
BMI, kg/m ² , median (IQR)	24.7 (22.3-27.7)	24.0 (22.7-27.1)	0.817*
ASA score, n (%)			
I	-	-	0.166†
II	5 (10.6)	1 (3.4)	
III	33 (70.2)	17 (58.6)	
IV	9 (19.1)	11 (37.9)	
V	-	-	
Charlson Comorbidity Index, n (%)			
0	8 (17.0)	6 (20.7)	0.237†
1	8 (17.0)	2 (6.9)	
2	13 (27.7)	8 (27.6)	
3	7 (14.9)	6 (20.7)	
4	8 (17.0)	1 (3.4)	
≥5	3 (6.4)	6 (20.7)	
Diagnosis, n (%)			
Cholecystitis	18 (38.3)	7 (24.1)	0.222†
Small or large bowel obstruction	8 (17.0)	9 (31.0)	0.169†
Hollow viscus perforation	6 (12.8)	5 (17.2)	0.739†
Diverticulitis	4 (8.5)	2 (6.9)	1.000†
Mesenteric ischemia	-	3 (10.3)	0.052†
Hernia	8 (17.0)	3 (10.3)	0.517†
Appendicitis	3 (6.4)	-	0.283†
Vital signs at admission, median (IQR)			
Temperature, °C	36.8 (36.3-37.1)	37.1 (36.8-37.5)	0.045*
Respiratory frequency, /min.	18.5 (15-23)	22.0 (20-28)	0.010*
Heart rate, /min.	81.5 (71-96)	85.0 (75-99)	0.526*
Systolic blood pressure, mmHg	133.5 (120- 160)	144.0 (127-162)	0.334*
Laboratory values at admission, median (IQR)			
White blood cell count, G/l	11.9 (8.8-15.1)	13.2 (10.2-14.8)	0.521*
CRP, mg/l	45 (8-165)	29 (6-213)	0.710*
Creatinine, mmol/l	91 (73-117)	105 (78-127)	0.171*
Creatinine clearance,	61 (42-76)	53 (38-70)	0.382*
Lactate at admission, mmol/l	1.5 (1.1-2.3)	1.6 (1.2-3.1)	0.170*
Lactate, worst within first 24 h, mmol/l	1.5 (1.1-2.3)	2.0 (1.2-3.1)	0.112*
pH	7.40 (7.33-7.42)	7.40 (7.35-7.44)	0.531*
Sepsis at admission, n (%)	26 (55.3)	19 (65.5)	0.473†
Nursing effort, h per day, median (IQR)	7.0 (5.0-8.9)	8.9 (7.2-11.4)	0.002*
Operation technique, n (%)			
Laparotomy	31 (66.0)	21 (72.4)	0.619†
Laparoscopy	16 (34.0)	8 (27.6)	
ICU admission, n (%)	11 (23.4)	14 (48.3)	0.043†
ICU LOS, median (IQR)	8.0 (6.0-13.0)	13.0 (9.0-20.0)	0.291*

IQR: interquartile range; BMI: body mass index; ASA score: American Society of Anesthesiology score; ICU: intensive care unit; * Mann-Whitney-U test; † Fisher's exact test.

Figure 1: Flow chart of the study outline

Figure 2: Kaplan-Meier survival curves stratified by ASA score (Figure 2 A), presence of mesenteric ischemia (Figure 2 B) and ICU admission (Figure 2 C)