



27 **Reducing sodium intake during childhood is recommended for the primordial prevention of**  
28 **hypertension, and cardiovascular diseases (CVD). There is however no estimation of the effect**  
29 **of sustained reduction on sodium intake beginning in childhood on CVD later in life. Based on a**  
30 **simple impact model, we estimated that a sodium intake reduction between 1 and 2 g per day in**  
31 **childhood could reduce coronary diseases by 3-6%, strokes by 3-5%, heart failures by 4-9%,**  
32 **and CVD mortality by 4-9% later in life.**

33 Worldwide, cardiovascular diseases (CVD) cause 17.6 million deaths per year, and half would be  
34 attributable to high blood pressure (BP) (1, 2). Since experimental studies have shown that high  
35 sodium intake increases BP (3), a key public health strategy to reduce the burden of high BP, and its  
36 associated consequences, is to reduce dietary sodium intake at the population level (4). Sodium intake  
37 is high in most populations: The average sodium intake among adults has been estimated be around 4  
38 g/day worldwide (corresponding to 10 g of salt) (5), with the majority of the world population having  
39 intakes above the maximum of 2 g/day recommended by the WHO (6). Because high BP has its roots  
40 in childhood and tracks to adulthood (7), it has been advocated that primordial prevention beginning  
41 as early as from the first years of life can have a large potential to reduce the global burden of high BP  
42 (8). Further, as dietary preferences and habits are largely ingrained during childhood, high sodium  
43 intake in childhood could lead to high sodium intake in adulthood, with associated higher BP levels  
44 and increased risk of CVD.

45 To the best of our knowledge, there is however no estimation of the long-term effect of sustained  
46 reduction on sodium intake beginning in childhood on CVD later in life. While several cohorts and  
47 trials have investigated the associations between sodium intake in childhood and BP later in childhood  
48 and in early adulthood (9, 10), no study has assessed the effect of reduced salt intake in childhood on  
49 BP after 25 years of age and associated CVD risk in adulthood. Using a simple impact model, we  
50 therefore indirectly estimated the percentage of CVD events and deaths prevented in adulthood by  
51 reducing sodium intake during childhood at the population level.

52 Assuming a causal relationship between sodium intake and BP during childhood, we modeled the  
 53 effect of a reduction of sodium intake on BP and CVD incidence and mortality in adulthood. First, we  
 54 chose two different levels of sodium intake reductions during childhood, i.e., 1 and 2 grams/day  
 55 respectively. Second, we calculated the change in systolic BP in childhood caused each sodium intake  
 56 reduction, using the findings from a recent systematic review on the effect of sodium intake on BP  
 57 during childhood (11). Third, we extrapolated this systolic BP change in childhood to a change in  
 58 systolic BP in adulthood, assuming that, a change in 1 mmHg in childhood would lead to a change in  
 59 2 mmHg in adulthood, based on the mean difference between the highest and lowest blood pressure  
 60 trajectories reported in a cohort study (12). Fourth, we estimated the effect of this change in BP in  
 61 adulthood on the risk of CVD later in life, using risk estimates from a recent systematic review (13).  
 62 The equations used were the following:

63 1)  $\Delta_{SBP\ child} [mmHg] = \Delta_{SBP/Na\ child} [mmHg/g\ sodium/d] \times \Delta_{Na\ child} [g/d]$

64 2)  $\Delta_{SBP\ adult} [mmHg] = 2 \times \Delta_{SBP\ child} [mmHg]$

65 3)  $\Delta_{CVD} (\%) = \Delta_{Risk} [\%/mmHg] \times \Delta_{SBP\ adult} [mmHg]$

66 where  $\Delta_{SBP\ child}$  is the expected change in systolic BP in children;  $\Delta_{SBP/Na\ child}$  is the change in systolic  
 67 BP expected for one gram change in sodium intake in children, i.e., 0.8 mmHg/g (11);  $\Delta_{Na\ child}$  is the  
 68 reduction in sodium intake per day in children, i.e., 1 or 2 g;  $\Delta_{SBP\ adult}$  is the expected change in  
 69 systolic BP in adults;  $\Delta_{CVD}$  is the relative reduction in fatal and non-fatal events of coronary heart  
 70 disease, stroke, and heart failure, and CVD mortality;  $\Delta_{Risk}$  is the percentage reduction in coronary  
 71 heart disease, stroke, heart failure and CVD mortality per mmHg systolic BP (i.e. 1.7%, 2.7%, 2.8%,  
 72 and 1.3% respectively (13)). To estimate the uncertainty around the estimates, we used the 95%  
 73 confidence intervals (CI) of the expected change in systolic BP per gram change in sodium intake in  
 74 children, i.e., 0.4 to 1.3 mmHg/g (11).

75 The results are shown in **Table 1**. We estimated that a reduction of sodium intake by 1 g per day from  
 76 childhood to adulthood would reduce BP during childhood by 0.8 mmHg, and during adulthood by

77 1.6 mmHg. Due to this reduction in BP in adults, 3% of coronary heart diseases, 3% of strokes, 4% of  
78 heart failures, and 4% of CVD deaths could be prevented. With a sodium reduction of 2 g per day, up  
79 to 6% of coronary heart disease, 5% of stroke, 9% of heart failure, and 9% of CVD deaths could be  
80 prevented.

81 Our simple model however has several limitations. First, the effect of sodium reduction in childhood  
82 was likely underestimated due to regression dilution bias in the available estimates due to  
83 measurement errors in sodium intake (14). Second, our analysis assumed a simple linear effect of  
84 sodium intake on BP in childhood, although it has been suggested that the dose-response relationship  
85 between sodium intake and BP could be J- or U- shaped (15, 16). Depending on the level of sodium  
86 intake, this could result in over- or under-estimation of the actual effect in some segments of the  
87 population. Third, we assumed that there was a causal relationship between sodium intake and BP and  
88 that there were no negative consequences of reducing sodium intake irrespective of the baseline  
89 sodium intake level and hypertension status of the individuals (17). Fourth, we assumed a difference  
90 in systolic BP in childhood of 1 mmHg resulted in a change of 2 mmHg in adulthood, although blood  
91 trajectories are much more complex and depend on other factors, such as overweight and smoking  
92 (12). Fifth, we did not consider the effect of a salt intake reduction on diastolic BP. Finally, the  
93 uncertainty around the estimates of the percentage of CVD events and deaths prevented is likely  
94 greater than reported, as our impact model does not account for several sources of uncertainties.

95 Nevertheless, this modeling study helps evaluate the impact of early life CVD preventive population  
96 based approach. Other studies have shown that population approaches to prevent CVD potentially can  
97 prevent more CVD events and deaths than targeted approaches. For instance, a modelling study using  
98 data from the Framingham Heart Study and the NHANES II found that a reduction of 2 mmHg  
99 diastolic BP at the population level could reduce the percentage of coronary heart diseases by 6%  
100 (18), similarly to what we have found in our analysis. By comparison, a targeted approach where  
101 individuals with diastolic BP above 95 mmHg are given anti-hypertension medication has been  
102 estimated to prevent 4% of the coronary heart disease events (18).

103 While limiting sodium intake during childhood would have a small impact from a clinical point of  
104 view, our analysis suggests that it could have a substantial impact on the burden of CVD later in life.  
105 Sodium intake can be limited during childhood by having more low-salt food options and reducing the  
106 amount of salt added during cooking and at the table both at home and at school. Primordial  
107 prevention of high blood pressure starting early in life, through a reduction in sodium intake in  
108 childhood, has the potential to have a substantial impact on the risk of CVD later in life.

109

110 **References**

- 111 1. Global Burden of Disease Causes of Death Collaborators. Global, regional, and national age-  
112 sex specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the Global Burden  
113 of Disease Study 2016. *Lancet*. 2017;390(10100):1151-210.
- 114 2. Global Burden of Disease Risk Factors Collaborators. Global, regional, and national  
115 comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks  
116 or clusters of risks, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016.  
117 *Lancet*. 2017;390(10100):1345-422.
- 118 3. Aburto NJ, Ziolkovska A, Hooper L, Elliott P, Cappuccio FP, Meerpohl JJ. Effect of lower  
119 sodium intake on health: systematic review and meta-analyses. *BMJ*. 2013;346:f1326.
- 120 4. World Health Organization. Global action plan for the prevention and control of  
121 noncommunicable diseases 2013-2020. Geneva: World Health Organization; 2013. Available from:  
122 <http://www.who.int/nmh/publications/ncd-action-plan/en/>. Accessed on 29 September 2016.
- 123 5. Powles J, Fahimi S, Micha R, Khatibzadeh S, Shi P, Ezzati M, et al. Global, regional and  
124 national sodium intakes in 1990 and 2010: a systematic analysis of 24 h urinary sodium excretion and  
125 dietary surveys worldwide. *BMJ Open*. 2013;3(12):e003733.
- 126 6. World Health Organization. Guideline: Sodium intake for adults and children. Geneva: World  
127 Health Organization (WHO); 2012. Available from:  
128 [http://www.who.int/nutrition/publications/guidelines/sodium\\_intake\\_printversion.pdf](http://www.who.int/nutrition/publications/guidelines/sodium_intake_printversion.pdf). Accessed on 10  
129 June 2017.
- 130 7. Chen X, Wang Y. Tracking of blood pressure from childhood to adulthood: a systematic  
131 review and meta-regression analysis. *Circulation*. 2008;117(25):3171-80.
- 132 8. Olsen MH, Angell SY, Asma S, Boutouyrie P, Burger D, Chirinos JA, et al. A call to action  
133 and a lifecourse strategy to address the global burden of raised blood pressure on current and future  
134 generations: the Lancet Commission on hypertension. *Lancet*. 2016;388(10060):2665-712.

- 135 9. Geleijnse JM, Hofman A, Witteman JC, Hazebroek AA, Valkenburg HA, Grobbee DE. Long-  
136 term effects of neonatal sodium restriction on blood pressure. *Hypertension*. 1997;29(4):913-7.
- 137 10. Krupp D, Shi L, Egert S, Wudy SA, Remer T. Prospective relevance of fruit and vegetable  
138 consumption and salt intake during adolescence for blood pressure in young adulthood. *Eur J Nutr*.  
139 2015;54(8):1269-79.
- 140 11. Leyvraz M, Chatelan A, da Costa BR, Taffe P, Paradis G, Bovet P, et al. Sodium intake and  
141 blood pressure in children and adolescents: A systematic review and meta-analysis of experimental  
142 and observational studies. *Int J Epidemiol*. 2018(dyy121):1-15.
- 143 12. Theodore RF, Broadbent J, Nagin D, Ambler A, Hogan S, Ramrakha S, et al. Childhood to  
144 Early-Midlife Systolic Blood Pressure Trajectories: Early-Life Predictors, Effect Modifiers, and Adult  
145 Cardiovascular Outcomes. *Hypertension*. 2015;66(6):1108-15.
- 146 13. Etehad D, Emdin CA, Kiran A, Anderson SG, Callender T, Emberson J, et al. Blood pressure  
147 lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis.  
148 *Lancet*. 2016;387(10022):957-67.
- 149 14. Hutcheon JA, Chiolero A, Hanley JA. Random measurement error and regression dilution  
150 bias. *BMJ*. 2010;340:c2289.
- 151 15. Jackson SL, Cogswell ME, Zhao L, Terry AL, Wang CY, Wright J, et al. Association  
152 between urinary sodium and potassium excretion and blood pressure among adults in the United  
153 States: National Health and Nutrition Examination Survey, 2014. *Circulation*. 2018;137(3):237-46.
- 154 16. Mente A, O'Donnell MJ, Rangarajan S, McQueen MJ, Poirier P, Wielgosz A, et al.  
155 Association of urinary sodium and potassium excretion with blood pressure. *N Engl J Med*.  
156 2014;371(7):601-11.
- 157 17. DiNicolantonio JJ, O'Keefe JH. Salt and hypertension: what do we know? *Curr Opin Cardiol*.  
158 2018;33(4):377-81.
- 159 18. Cook NR, Cohen J, Hebert PR, Taylor JO, Hennekens CH. Implications of small reductions  
160 in diastolic blood pressure for primary prevention. *Arch Intern Med*. 1995;155(7):701-9.





**Table 1.** Effect of reducing sodium intake by 1 and 2 g/day in childhood on blood pressure in childhood and on CVD risk and mortality in adulthood.

<b>Reduction in sodium (g Na/d)</b>	<b>Equivalent in salt (g NaCl/d)</b>	<b>Change in systolic blood pressure in children (mmHg) (11)</b>	<b>Change in systolic blood pressure in adult (mmHg) (12)</b>	<b>Percentage of coronary heart diseases prevented (13)</b>	<b>Percentage of strokes prevented (13)</b>	<b>Percentage of heart failures prevented (13)</b>	<b>Percentage of CVD deaths prevented (13)</b>
-1 g	-2.5 g	-0.8 (-0.4, -1.3) <sup>1</sup>	-1.6 (-0.8, -2.6) <sup>2</sup>	-3% (-2%, -5%) <sup>2</sup>	-3% (-1%, -4%) <sup>2</sup>	-4% (-2%, -7%) <sup>2</sup>	-4% (-2%, -7%) <sup>2</sup>
-2 g	-5 g	-1.6 (-0.8, -2.6) <sup>1</sup>	-3.2 (-1.6, -5.2) <sup>2</sup>	-6% (-3%, -10%) <sup>2</sup>	-5% (-3%, -9%) <sup>2</sup>	-9% (-4%, -14%) <sup>2</sup>	-9% (-4%, -15%) <sup>2</sup>

Abbreviations: CVD: cardiovascular diseases; Na: sodium; NaCl: salt. <sup>1</sup> 95% confidence interval; <sup>2</sup> lower and upper estimates.

## Summary Table

### What is known about topic

- Reducing sodium intake during childhood is advocated for the primordial prevention of hypertension, and cardiovascular diseases.
- There is no estimation of the effect of sustained reduction on sodium intake beginning in childhood on CVD later in life.

### What this study adds

- This modeling study helps evaluate the impact of sodium reduction in childhood for the primordial prevention of cardiovascular diseases incidence and mortality later in life.
- Our findings suggest that a sodium intake reduction in childhood could prevent a substantial proportion of cardiovascular diseases later in life.