

Advances in therapeutic options for portal hypertension

Marina Vilaseca, Sergi Guixé-Muntet, Anabel Fernández-Iglesias and Jordi Gracia-Sancho 

Ther Adv Gastroenterol

2018, Vol. 11: 1–19

DOI: 10.1177/
1756284818811294

© The Author(s), 2018.
Article reuse guidelines:
sagepub.com/journals-
permissions

Abstract: Portal hypertension represents one of the major clinical consequences of chronic liver disease, having a deep impact on patients' prognosis and survival. Its pathophysiology defines a pathological increase in the intrahepatic vascular resistance as the primary factor in its development, being subsequently aggravated by a paradoxical increase in portal blood inflow. Although extensive preclinical and clinical research in the field has been developed in recent decades, no effective treatment targeting its primary mechanism has been defined. The present review critically summarizes the current knowledge in portal hypertension therapeutics, focusing on those strategies driven by the disease pathophysiology and underlying cellular mechanisms.

Keywords: cirrhosis, hepatic hemodynamic, HVPG, portal pressure, sinusoid

Received: 20 July 2018; revised manuscript accepted: 15 October 2018.

Introduction

Advanced chronic liver disease (CLD) represents the fourth cause of death due to noncommunicable diseases worldwide. Its high mortality is mainly due to long-time asymptomatic stages during disease progression and, thus, is diagnosed after appearance of portal hypertension (PH) and its clinical complications such as hepatic encephalopathy, variceal hemorrhage or ascites. In fact, PH is the major driver of the complications of CLD, the measurement of hepatic venous pressure gradient (HVPG) being a predictor of the risk of clinical decompensation and survival in compensated cirrhotic patients.^{1,2}

Clinically significant PH is defined as an HVPG of 10 mmHg or greater and depends on changes in portal blood flow or in intrahepatic vascular resistance (IHVR). During cirrhosis development, elevation in IHVR due to deregulation of hepatic cells and liver fibrosis is the primary factor involved in increasing portal pressure (PP) leading to secondary splanchnic vasodilation and extrahepatic shunt formation, which further aggravate PH and liver dysfunction. Notably, current treatments do not target the main

underlying mechanism, but consist in extrahepatic vasoconstrictors [i.e. nonselective beta blockers (NSBB) and somatostatin] aimed at ameliorating PH by reducing the hyperdynamic circulation, or in strategies focused on the prevention of PH-derived complications such as infections or encephalopathy. In the specific case of NSBB, it should be noted that carvedilol may also exert intrahepatic vasodilatation due to its anti- α 1-adrenergic properties.³ Despite the extended use of these drugs as the most effective treatment for PH, a recent analysis based on over 100,000 cases showed that 30-day mortality following a decompensation in cirrhosis remains equal, or even higher, than that observed 10 years ago,⁴ strongly suggesting there is a need for new therapeutic approaches for PH and CLD.

In this review we summarize the potential therapeutic strategies for the treatment of PH, which importantly derive from preclinical knowledge and mainly target the increased IHVR. These approaches consist of pharmacological, cell-derived or lifestyle interventions, as shown below.

Correspondence to:
Jordi Gracia-Sancho
Liver Vascular Biology
Research Group,
Barcelona Hepatic
Hemodynamic Laboratory,
IDIBAPS Biomedical
Research Institute,
CIBEREHD, Rosselló 149,
4th floor, 08036 Barcelona,
Spain
jordi.gracia@idibaps.org

Marina Vilaseca
Anabel Fernández-Iglesias
Hepatic Hemodynamic
Laboratory, IDIBAPS
Biomedical Research
Institute, Barcelona, Spain
Sergi Guixé-Muntet
Department of Biomedical
Research, University of
Bern, Bern, Switzerland






Figure 1. Schematic representation of different therapeutic approaches for portal hypertension focused on their liver vasomodulation effect.

Drugs in bold denote currently ongoing RCT and known mechanisms of action are indicated in parenthesis. APAP, acetaminophen; COX, cyclooxygenase; ECA, enzyme converter of angiotensin; ERβ, estrogen receptor beta; FXR, farnesoid X receptor; OCA, obeticholic acid; 5-MTHF, 5 methylfolate; eNOS, endothelial nitric oxide synthase; ET, endothelin receptor; GTP, guanosine triphosphate; cGMP, cyclic guanosine monophosphate; KLF2, Kruppel-like factor 2; NO, nitric oxide; sGC, soluble guanylate cyclase; L-arg, L-arginine; LTC, leukotriene; MAS, Mas receptor; PDE, phosphodiesterase; NCX, NO-donor compound derived from ursodeoxycholic acid; JNK, Jun N-terminal kinases; PLA2, phospholipase A2; RhoK, Rho kinase; RAA, renin-angiotensin-aldosterone; RCT, randomized controlled trial; TXA2, thromboxane A2.

Vasoprotective strategies

CLD is characterized by persistent injury of the liver due to different etiologies, including alcoholic and nonalcoholic steatohepatitis (NASH) or viral infections. This in turn leads to alterations of the hepatic microvasculature, which in combination with fibrosis constitute the main components of increased IHVR. This scenario of vascular alterations has led to the assessment of different vasoprotective strategies for the treatment of intrahepatic-derived PH, mainly vasomodulators, anticoagulants and antiangiogenic approaches.

Vasomodulators

One of the causes of the abovementioned microvascular alterations during CLD is a shift/alteration in the synthesis of vasoactive molecules in the liver, where vasoconstrictors prevail over vasodilators. Thus, several vasomodulator strategies aimed at enhancing intrahepatic vasodilation and normalizing excessive vasoconstriction have been recently proposed (Figure 1).

Inhibition of vasoconstriction. One of the most potent vasoconstrictors that contributes to microvascular dysfunction in CLD is endothelin. This molecule is synthesized by the activated endothelium and acts as a ligand for the endothelin receptors (ET) on hepatic stellate cells (HSCs), which are the final effectors of vasoconstriction. Indeed, the use of antagonists targeting ET (either ET_A, ET_B or both) showed improvements in the hepatic sinusoid and in liver fibrosis in preclinical models^{5,6} and decreased PP in a small clinical study.⁷ However, its possible beneficial effects were not further validated in human randomized-controlled trials (RCTs).^{8,9} Currently ongoing human RCTs evaluating the unspecific antagonist macitentan¹⁰ and ambrisentan¹¹ will provide essential information to support or discard this therapeutic strategy in cirrhosis and PH.

Urotensin is another vasoconstrictor playing a role in PH, known to correlate with HVP levels.¹² Similar to ET, blockage of the urotensin II receptor with the antagonist palosuran decreases




Figure 2. Schematic representation of different therapeutic approaches for portal hypertension based on their effect on apoptosis, oxidative stress or inflammation as main features of chronic liver damage.

Known mechanisms of action are indicated in parenthesis.

ASK1, apoptosis signal-regulating kinase; CeO₂, cerium oxide; EET, epoxyeicosatrienoic acids; GLP-1R, glucagon-like peptide 1 receptor; JAK2, janus kinase 2; NADPH, nicotinamide adenine dinucleotide phosphate; NO, nitric oxide; rMn, Recombinant Manganese; ROS, reactive oxygen species; SOD, superoxide dismutase.

PP in experimental models.¹³ Nevertheless, there are no current studies on cirrhotic patients in order to validate the translatability of these results to the bedside.

Eicosanoids represent another important family of vasoconstrictors derived from the conversion of arachidonic acid. Indeed, cyclooxygenase (COX)-derived prostanoids (a subclass of eicosanoids) are overexpressed in the cirrhotic liver.^{14,15} In this regard, several drugs have been used to chronically inhibit either their synthesis (i.e. by COX inhibition with nitroflurbiprofen¹⁶ or celecoxib^{17,18}) or to block their receptor (e.g. thromboxane A2 receptor with terutroban¹⁹). Although these strategies decreased PP in preclinical models of cirrhosis, there is only one ongoing RCT with the thromboxane receptor antagonist ifetroban.²⁰ Similar strategies aimed at reducing arachidonic acid metabolites focused on inhibiting epoxygenases using MS-PPOH²¹ and cysteinyl leukotrienes with montelukast.²²

In addition to the increased synthesis of vasoconstrictors, the cirrhotic liver is also hypersensitive to these molecules. Thus, research aimed to inhibit the effect of not only hepatic vasoconstrictors, but also systemic ones. In this regard, the renin-angiotensin-aldosterone (RAA) system is

a hormone system that controls blood pressure and fluid balance, playing also an important role regulating hepatic hemodynamics. Importantly, angiotensin II is a potent vasoconstrictor peptide with hepatic effects that favors the development of PH.²³ Several strategies have been developed to reduce its effects on the liver: angiotensin-converting-enzyme inhibitors (captopril) and angiotensin receptor blockers (losartan, candesartan, irbesartan) have been tested in preclinical and clinical settings and showed beneficial effects on reducing PH but mainly in compensated patients with Child A cirrhosis.²⁴ Given the potential of these drugs, some strategies aimed at focusing on the RAA system downstream pathway. Indeed, the Janus kinase inhibitor AG490 showed a PP-lowering effect additive to propranolol in preclinical models,^{25–27} while the Rho-kinase inhibitors Y-27632²⁸ and fasudil²⁹ showed comparable beneficial effects. Indeed, RhoA modulation using sodium ferulate³⁰ or diarylpropionitrile³¹ showed beneficial effects in preclinical PH, and interestingly, pointed out the relevance of the latter (which is an estrogen-receptor β agonist) in postmenopausal women. On the other hand, the receptor Mas also represents an alternative downstream target of the RAA pathway. Its activation showed promising effects in experimental models of cirrhosis, also

reducing PP and liver fibrosis.³² Nevertheless, another study blocking the Mas receptor with A779 led to the same decrease in PP,³³ leaving the Mas receptor role in a paradigm.

Vasopressin, also named antidiuretic hormone, is a hormone that contributes to the regulation of water and electrolyte homeostasis and constricts arterioles, increasing vascular resistance and blood pressure. The blockage of vasopressin using conivaptan has been recently tested in an RCT and showed no beneficial effects on PP.³⁴ Indeed, previous data support the concept that vaptans may not only be ineffective but also have negative secondary effects in terms of morbidity and mortality in patients with cirrhosis.^{35,36} Partial vasopressin agonists in the mesentery may have better projection. Indeed, a recent study demonstrated beneficial effects of FE 204038 reducing PP in cirrhotic rats,³⁷ and a recently started RCT evaluates the compound FE 204205 in cirrhotic patients with PH.³⁸

Promotion of vasodilation. Nitric oxide (NO) is a potent vasodilator synthesized by the NO synthase (NOS) that regulates vascular tone and inflammation; however, it becomes markedly deregulated in the cirrhotic liver. Several therapeutic approaches focused at increasing its levels, either by enhancing NO production (NOS-dependent strategies), or by providing NO as an exogenous source or decreasing NO degradation (NOS-independent strategies).

Delivering NO to precise cellular locations, like the liver sinusoid, represents a challenge. However, nanoparticle delivery has lately become an important and emerging field. In fact, recent studies using NO-releasing nanoparticles coated with vitamin A, which is mainly stored in HSCs in the liver, reduced PP in a preclinical cirrhotic model.³⁹ Similar effects were observed when NO synthesis was enhanced: the NOS cofactor tetrahydrobiopterin (sapropterin) and the NOS transcription enhancer (AVE 9488) decreased PP in cirrhotic animal models,^{40,41} nevertheless no benefits were observed in an RCT evaluating the first.⁴² In this same strategy, a currently ongoing RCT will describe the effectiveness of serelaxin, a drug that has been used for the treatment of acute heart failure due its capability to activate the relaxin receptor and NOS.⁴³

The soluble guanylate cyclase (sGC) is the only known receptor for NO and represents its

downstream mediator. When sGC is activated by NO, it induces the formation of the intermediate cyclic guanosine monophosphate (cGMP), which confers potent vasodilation. In this regard, the stimulation of the nonoxidized form of sGC or the oxidized and free-sGC form may represent novel approaches for activating cGMP production and subsequently ameliorate the vasoconstrictor environment. In fact, a recent study using the nonoxidized and free-sGC stimulator riociguat showed improved liver fibrosis and PP in cirrhotic animals.⁴⁴ However, no studies have evaluated the potential of activators of the oxidized form of sGC for the treatment of PH. Considering that oxidative stress plays a key role in cirrhosis and PH, and that sGC activators are able to generate cGMP even in these detrimental conditions,⁴⁵ their usefulness as a therapeutic option for PH deserves further investigation.

In addition to sGC strategies, there are other approaches aimed at potentiating the effects of cGMP by preventing its degradation by phosphodiesterase-5 (PDE-5). Indeed, different PDE-5 inhibitors showed lowering effects on PP in cirrhotic patients.⁴⁶⁻⁴⁸

Transcriptional modulators. In addition to specific modulators of vasoactive molecules, other strategies rely on the modulation of transcription factors that regulate the expression of many of these proteins. For example, farnesoid X receptor (FXR) is a bile-acid-responsive transcription factor highly expressed in the liver that controls the expression of many genes involved in metabolic regulation, hepatic fibrosis, vascular homeostasis and inflammation, and thus it has been considered a promising therapeutic target in CLD. Obeticholic acid (OCA), a semisynthetic FXR agonist, has been tested in preclinical models of cirrhosis showing beneficial effects on PH by reducing the IHVR.⁴⁹ Indeed, different ongoing clinical trials are investigating its effects in advanced CLD,⁵⁰⁻⁵² including patients with NASH cirrhosis.^{53,54} This drug, however, has raised some cases of overdose-related toxicity in patients with severe or even mild liver dysfunction, so its dosage needs to be carefully adjusted.⁵⁵ In addition to OCA, new nonsteroidal FXR agonists have been developed to evade enterohepatic recirculation. These drugs (e.g. PX201606, GS-9674), have been tested in animals with CLD showing increased endothelial NOS expression, reduced inflammatory response and improved liver injury.^{56,57} The effects of these newly

formulated compounds on advanced CLD and PH require further investigation.

On the other hand, statins are lipid-lowering drugs that have shown strong hepatosinusoidal protective effects in preclinical models of CLD, ultimately leading to PP decrease.^{58–60} Underlying mechanisms of statins mostly rely on their capability to induce the expression of the transcription factor Kruppel-like factor 2 that leads to liver sinusoidal endothelial cell (LSEC) phenotype amelioration, HSC deactivation, and global improvement in hepatic injury.^{61–66} Encouraged by the preclinical results, several clinical studies have tested statins in PH-cirrhotic patients. Recent analyses have shown that statins are associated with decreased risk of decompensation, death and hepatocellular carcinoma development.^{67,68} Moreover, simvastatin has shown beneficial effects when administered alone⁶⁹ or combined with beta blockers.⁷⁰ Future and ongoing RCTs will extend our knowledge regarding the use of statins for the treatment of PH.⁷¹

Anticoagulants

It is widely accepted that cirrhotic patients have an imbalance in anti- and procoagulant factors, leading to both increased risk of bleedings and microthrombosis in the liver, which may in turn increase liver hypoxia and inflammation.⁷² Therefore, different anticoagulants have been lately tested in the setting of CLD with success in the prevention of hepatic fibrosis.⁷³ Indeed, the direct antithrombin enoxaparin has been shown to reduce PP and liver fibrosis in experimental CLD,⁷⁴ confirming at the bench side its clinical benefits on cirrhosis decompensation and survival.⁷⁵ A different anticoagulant strategy is the direct inhibitor of factor Xa rivaroxaban, which has been shown to reduce liver microthrombosis, HSC activation and PP in experimental models of cirrhosis.⁷⁶ Moreover, it is currently being tested in RCTs on patients with cirrhosis⁷⁷ or with portal-vein thrombosis.⁷⁸ Additional, ongoing RCTs will provide more data on the use of anticoagulants for CLD-patients.⁷⁹

Antiangiogenics

Angiogenesis is an important feature in the development of PH, being triggered by hypoxia and inflammation which are present during hepatic fibrogenesis.⁸⁰

Vascular endothelial growth factor (VEGF) is the main proangiogenic factor promoting neovascularization in the mesentery and contributes to the development of increased splanchnic vasodilation. Therefore, different studies have focused on blocking VEGF receptors (VEGFR) to decrease splanchnic circulation and improve PP. Initially, monoclonal antibodies and specific inhibitors of VEGFR2 were used in PH animals showing improvement in portosystemic collateral vessel formation but showing no effect on PP.^{81,82} Subsequent preclinical studies investigated oral tyrosine kinase inhibitors that block VEGFR2, such as sorafenib,^{83,84} brivanib^{84,85} and regorafenib⁸⁶ and showed improved PP and systemic shunting in cirrhotic and noncirrhotic rats with PH. Indeed, tyrosine kinase inhibitors have also been shown to decrease liver fibrosis.⁸⁷ These beneficial preclinical results were confirmed in a small clinical trial where sorafenib significantly improved PH in patients with liver cirrhosis and hepatocellular carcinoma.⁸⁸

Apart from VEGF, other molecules such as pigment epithelium-derived factor (PEDF) or vashoin-1 have been shown as potent endogenous inhibitors of angiogenesis. In fact, overexpression by adenovirus of PEDF or vashoin-1 resulted in decreased mesenteric angiogenesis, portosystemic shunting, PH and liver fibrosis.^{89,90} Nevertheless, no current RCTs are being performed following this approach. It is important to highlight that angiogenesis is not only deleterious but also required for repairment of hepatic tissue during liver fibrosis resolution, therefore total angiogenesis blockade may not be a realistic therapeutic approach.

Cell death, oxidative stress and inflammation

Prevention of cell death

The main factor in the progression of cirrhosis is chronic liver injury. This is a scenario of persistent cell death (due to different etiologies), which leads to excessive inflammation and associated cytotoxicity, and oxidative stress. This sequence of events has led to the hypothesis that pharmacological inhibition of cell death may represent a therapeutic option for the treatment of PH and CLD (Figure 2).

Emricasan is an orally active pan-caspase inhibitor that has proven promising potential in the setting

of CLD. Indeed, a preclinical study presented as a meeting abstract describes that CCl₄-cirrhotic rats receiving emricasan displayed improved PP and IHVR compared with vehicle-treated rats, in addition to improved liver function and microcirculation, and these effects were accompanied by improved LSEC and HSC phenotype and reduced inflammation.⁹¹ More recently, another study in bile-duct-ligated (BDL) mice reported similar effects of this caspase inhibitor.⁹² At the bedside, emricasan did not show positive effects in acute-on-chronic liver failure.⁹³ However, a proof-of-concept clinical study suggests that emricasan ameliorates PP in patients with compensated cirrhosis and severe PH.⁹⁴ The completion of a bigger ongoing clinical trial⁹⁵ may confirm these promising results in the near future.

These findings are in agreement with other studies assessing the use of caspase inhibitors in chronic liver diseases, such as NASH (reviewed elsewhere⁹⁶). Of note, inhibition of the apoptosis signal-regulating kinase 1 is another potential candidate strategy in CLD, as it has shown significant results in reducing the fibrosis stage of NASH patients.⁹⁷

Despite the exciting results, there is still controversy on the role of cell-death modulation in CLD. Indeed, a recent study suggests that the use of caspase inhibitors would not avoid cell death, as hepatocytes may still die from necrosis.⁹⁸ On the other hand, some vasoprotective drugs modulate hepatic fibrosis at least in part due to selective apoptosis of HSCs,⁶⁵ while other studies suggest that treatment with low doses of an apoptotic agent (gliotoxin) may achieve similar effects without affecting hepatocyte viability.⁹⁹

Future studies will clarify the usefulness of cell-death inhibition in a context where the injury is not present anymore, such as the regression of CLD after the removal of the hepatic injury.

Antioxidants

During liver damage, increased inflammation, mitochondrial damage and enzymatic alterations [i.e. xanthine and nicotinamide adenine dinucleotide phosphate (NADPH) oxidases] induce oxidative stress. Antioxidants are molecules that compensate this oxidative stress by promoting the conversion of reactive oxygen species (ROS) into less reactive molecules, either by direct enzymatic

activity or by transcriptional/post-transcriptional modulation of other enzymes. However, cirrhotic livers are known to produce reduced levels of antioxidants compared with control livers, thus further aggravating the increased oxidative stress in this setting.^{100,101} For this reason, several studies aimed to develop pharmacological antioxidant-enhancing strategies for CLD. One of these is mitoquinone, a mitochondria-targeted antioxidant with oral administration approved for human use. This drug has already proved positive effects in preventing oxidative damage in models of liver diseases such as NASH or ischemia/reperfusion.^{102,103} In a recent study, treatment with mitoquinone reduced PP and IHVR in two different preclinical models of CLD [CCl₄ and thioacetamide (TAA)] by reducing oxidative stress and fibrosis.¹⁰¹ Importantly, these antifibrotic effects were also validated in precision-cut liver slices from human tissue, suggesting potential applicability in clinical practice. Moreover, a recent abstract reports similar effects of this drug in the BDL rat model.¹⁰⁴

Apocynin is a natural antioxidant of vegetal origin which has a specific inhibitory effect on NADPH oxidase and thus has been studied for its antioxidant properties.^{105–107} In this regard, a recent study assessed the effects of apocynin in a model of mild fibrosis.¹⁰⁸ Although authors described amelioration of markers of liver damage, oxidative stress, inflammation and fibrosis, these effects were not observed in a previous study using animals with advanced CLD probably due to a progressive reduction in NADPH oxidase activity during cirrhosis development.¹⁰⁹ This advocates for the use of NADPH oxidase inhibitors during progression but not at advanced stages of CLD. Indeed, a study using a new generation of NADPH oxidase inhibitor demonstrated anti-inflammatory and hepatoprotective effects in a preclinical model of NASH without significant fibrosis,¹¹⁰ altogether encouraging future studies to ascertain which would be the appropriate therapeutic window for this strategy in cirrhosis.

In addition to their protective effects against oxidative stress, antioxidants may also play an important role in ameliorating microvascular dysfunction, as they prevent the scavenging of NO by the superoxide anion (O₂⁻).¹⁰⁰ In this direction, back in 2006, Hernández-Guerra and colleagues assessed whether acute administration of ascorbic acid would prevent the increase in HVPG associated with the

defective ability of the cirrhotic endothelium to accommodate the postprandial increase in portal blood flow. Indeed, patients receiving vitamin C displayed lower HVPG after a meal compared with placebo, suggesting improved microvascular function.¹¹¹ Interestingly, a more recent publication demonstrated that consumption of dark chocolate had similar effects on the postprandial increase of HVPG.¹¹² Since then, several studies have addressed the role of antioxidant therapy in the improvement of NO bioavailability. In this regard, the modulation of the activity of the antioxidant enzyme superoxide dismutase (SOD) has proven promising effects in animal models of CLD.^{113,114} In a study from 2011, administration of tempol, a SOD analog, improved the IHVR of cirrhotic rats in an NO-dependent manner.¹¹⁴ Similarly, a more recent study assessed the administration of a recombinant form of human manganese SOD in two different animal models of CLD resulting in improved liver hemodynamics, as well as reduced oxidative stress and liver fibrosis.¹¹⁵ Resveratrol is an antioxidant found in the skin of grapes that activates the histone deacetylase Sirt1, which regulates several vasoprotective pathways leading to NO overexpression.¹¹⁶ Recent studies have reported hepatic antifibrotic effects of resveratrol in models of mild liver injury linked to its antioxidant properties^{117–119} while its administration to rats with severe PH reduced PP and hepatic fibrosis, and ameliorated hepatic microvascular dysfunction.¹²⁰

Strategies targeting inflammation

Anti-inflammatory strategies. Rapamycin is an immunosuppressor that negatively regulates the mammalian target of rapamycin (mTOR) complexes, which play a central role in the regulation of many of the cellular pathways including proliferation, inflammation, apoptosis and autophagy. In the field of CLD, rapamycin has shown the ability to reduce the PP in rats with PH due to its intrahepatic^{121,122} and extrahepatic effects.^{123,124}

The Janus kinase 2 protein (JAK2) is involved in the toll-like receptor (TLR) signaling in response to lipopolysaccharide, and upregulates profibrotic and inflammatory genes. As commented above, AG490 is a specific inhibitor of JAK2 but it can also regulate the TLR signaling pathway. Recently, its potent antifibrotic and anti-inflammatory effects were proven in livers of BDL-cirrhotic rats, partially normalizing PH. Furthermore, the same study reported additional extrahepatic

hemodynamic amelioration in the partial portal-vein ligation model,²⁶ while combination with NSBB achieved additive effects.²⁷

Epoxyeicosatrienoic acids (EETs) are products of the arachidonic acid metabolism with anti-inflammatory properties.¹²⁵ These molecules are further metabolized by the soluble epoxide hydrolase (sEH), thus inhibition of this enzyme represents a good strategy in maintaining or increasing EET levels. Indeed, the anti-inflammatory effects of the sEH inhibitor t-TUCB have been recently validated in steatotic livers.¹²⁶ In addition to these effects, EETs have also been described to show vasoactive properties,¹²⁷ thus suggesting additional potential in liver vascular diseases. Indeed, recent studies reported that t-TUCB treatment ameliorates PH in CCl₄-cirrhotic rats, altogether improving fibrogenesis and endothelial dysfunction^{128,129} while additionally ameliorating extrahepatic circulation.¹³⁰ Although inflammatory markers were also downregulated by this inhibitor, whether the hemodynamic effects were additive or directly mediated by inflammation was not assessed. Despite these promising reports, there is some controversy regarding the role of EETs in cirrhosis, as another study reported opposite effects in the splanchnic territory.²¹

In addition to conventional anti-inflammatory pharmacological strategies, engineered nanoparticles represent an alternative therapeutic option with targetable delivery. Cerium oxide nanoparticles have been shown to behave as anti-inflammatory agents, with no exception in the liver. Indeed, these nanoparticles prevented macrophage infiltration and overexpression of hepatic inflammatory genes, causing a reduction in oxidative stress and improving PP in rats with mild PH.¹³¹

Antidiabetic drugs. As expected from drugs aimed at normalizing glucose levels, antidiabetic treatments usually target glucose and lipid transport, as well as metabolism. However, these drugs have also been reported to have vasoprotective effects associated with NO bioavailability, inflammation and oxidative stress.^{132–134} In consequence, these beneficial effects have been assessed in CLD. One such pharmacological strategy is use of liraglutide. This glucagon-like peptide 1 (GLP-1) analog has shown the ability to ameliorate PH and liver microvascular function in cirrhotic animals by reducing HSC activation and the levels of inflammatory markers independently of the

GLP-1 receptor.¹³⁵ In this same study, the effects of liraglutide were validated in different human models *in vitro*, thus shortening the gap between the preclinical study and clinical trials. Indeed, liraglutide has shown antifibrotic effects also in NASH patients,¹³⁶ thus representing a promising drug with high probabilities of success as a treatment for PH and CLD. Similarly, the antidiabetic drug metformin has shown improvement in liver hemodynamic and fibrosis, which were accompanied by a reduction in inflammation and oxidative stress.¹³⁷ Interestingly, the observed hemodynamic effects were complementary to the effects of NSBB. Lastly, fenofibrate, another antidiabetic drug with described positive vascular effects, also improved PP in cirrhotic rats in addition to hepatic fibrosis and microvascular function.¹³⁸

An additional strategy to these drugs is the use of antibodies that specifically block the interaction between metabolic hormones and their ligands. Leptin, the satiety hormone, is overexpressed in obese people, who may suffer from leptin resistance. Furthermore, this hormone has been implicated in fibrosis and is highly expressed in cirrhotic patients, probably due to inflammation.^{139,140} Interestingly, the use of a competitive antibody against leptin receptor proved to be effective in ameliorating PP and IHVR in cirrhotic rats, suggesting that leptin is not only a marker in CLD but is also a targetable effector.¹⁴¹

Microbiota modulators. Intestinal bacteria are known to modulate splanchnic flow and thus participate in the PH-derived hyperdynamic syndrome. In addition, during cirrhosis, increased gut permeability allows translocation of bacteria or their products into the bloodstream, causing general and hepatic inflammation.^{142,143} For this reason, different clinical trials have assessed the use of antibiotics (rifaximin,^{144–146} norfloxacin¹⁴⁷) as prophylactic intervention against general infections in decompensated cirrhotic patients, which overall improved their outcome.

However, recent research suggests that the composition of the microbiota population may also play a role in gut permeability and, interestingly, in liver hemodynamics. In this regard, oral administration of the probiotic VSL#3 (a mix of four *Lactobacillus* species) prevented the increase in the systemic levels of inflammatory cytokines and vasodilators in a rat model of BDL cirrhosis¹⁴⁸ and ameliorated liver function and pro-inflammatory markers in cirrhotic

patients compared with the placebo group.¹⁴⁹ *Bifidobacterium pseudocatenulatum* CECT7765 supplementation had similar anti-inflammatory features. On one hand, liver or blood macrophages isolated from cirrhotic animals and patients (respectively) switched to an anti-inflammatory M2 phenotype when treated *in vitro* with *B. pseudocatenulatum*¹⁵⁰ while on the other hand, oral administration of the bifidobacteria to BDL-cirrhotic rats improved their hepatic hemodynamic parameters, liver damage and markers of inflammation *in vivo*.¹⁵¹

These findings suggest that not only direct administration of bacteria but also diets that modify the gut microbiota could have therapeutic value in CLD and PH. Indeed, a recent clinical trial reported that cohorts of healthy controls and patients with cirrhosis have different microbiota populations, that microbiota diversity negatively correlates with the degree of cirrhosis, and that cirrhotic patients with diets that ensure higher microbiota diversity had lower risk of hospitalization.¹⁵² Dietary goods associated with microbiota diversity included fermented milk products, coffee, tea, dark chocolate and vegetables, while consumption of carbonated drinks (including caffeinated ones) was predictor of poor diversity. Interestingly, most of the microbiota-beneficial products are rich in polyphenols (antioxidants), thus holding a potential combined effect to that.

Strategies targeting fibrogenesis

As stated above, it is well known that chronic liver damage in combination with paracrine signaling from the activated endothelium lead to the activation of HSCs, promoting fibrogenesis.^{63,153}

Although most of the vasoprotective strategies described above ultimately lead to significant amelioration in fibrosis, there are a few pharmacological approaches that directly target the process of fiber formation/stabilization. One of these is simtuzumab, a monoclonal antibody against lysyl oxidase homologue 2 (LOXL2). LOXL2 is a protein that participates in the cross-linking of collagens and elastin. This protein is highly expressed in fibrotic livers (unlike control livers) and its inhibition with antibodies has shown positive results in preventing and reducing liver fibrosis in preclinical models.¹⁵⁴ Despite these promising results, simtuzumab proved to be ineffective in a recent phase II RCT in NASH-cirrhotic patients,¹⁵⁵ highlighting

the controversy that is often found in the translation of some therapies from the laboratory to the clinic.

Cell therapy

The lack of pharmacologic therapeutic options to ameliorate PH and cirrhosis has prompted the use of regenerative therapies or cell-derived interventions. In 2006, Terai and colleagues performed the first clinical trial studying the autologous bone marrow cell infusion (ABMi) in decompensated cirrhotic patients, showing a decrease in Child-Pugh score as well as better liver function.¹⁵⁶ Latterly, infusion of mesenchymal stem cells (MSCs) from bone marrow or from umbilical cord to patients with liver cirrhosis demonstrated amelioration in hepatic fibrosis and function with increased serum albumin levels and with a good safety profile.¹⁵⁷ However, the last investigations in the field are focused on less invasive therapies using induced pluripotent stem cells (iPS).¹⁵⁸ iPS have been proposed as a new therapeutic option to replace damaged hepatocytes by healthy hepatocyte-like cells in CLD.¹⁵⁹ Nevertheless, hepatocyte-like cells induced by reprogramming methods remain immature without the specific hepatocyte characteristics such as detoxification or production of albumin and urea.¹⁶⁰ Along this theme, transplantation of human fetal hepatocytes¹⁶¹ was evaluated in patients with end-stage CLD; although the small population size of the study was an important limitation for obtaining a robust conclusion, model for end-stage liver disease (MELD) score and portosystemic encephalopathy episodes were ameliorated in transplanted patients. Even though several clinical trials evaluating stem cell therapies in CLD are still ongoing,¹⁶²⁻¹⁶⁴ none of them specifically performs HVPG measurements; therefore, these RCTs will not provide new results about the effect of stem cells in PH.

On the other hand, preclinical studies have evaluated the effectiveness of cell-derived therapy ameliorating hepatic fibrosis, inflammation and liver hemodynamics using different sources and potency of stem cells in fibrotic and cirrhotic rodents. Specifically, human placenta-derived stem cells (PDSCs) are considered a promising therapeutic option for their proliferation and differentiation capacities, as well as their low immunogenicity potential.¹⁶⁵ In this sense, the anti-inflammatory and regenerative effects of

chorionic plate-derived MSC transplantation in CCl₄-injured rats¹⁶⁶ and the amelioration in fibrosis and cirrhosis progression in fibrotic rodents receiving human amniotic membrane-derived mesenchymal¹⁶⁷ or epithelial stem cells¹⁶⁸ has been reported. In addition, transplantation of human amniotic MSCs in TAA-induced cirrhotic mice was able to restore the hepatic function and improve liver fibrosis;¹⁶⁹ importantly, this study suggested the PDSCs transplantation as a better therapeutic option in comparison with adipose tissue-derived MSCs.

Regarding the effects of stem cells on PH and hepatic microcirculation, much less is known. Brückner and colleagues suggested that transplantation of hepatocyte-like cells *in vitro* differentiated from rat adipose tissue MSCs into CCl₄-cirrhotic rats led to an amelioration in PH after 3 weeks of treatment without changes in liver dysfunction or fibrosis.¹⁷⁰ However, an ongoing preclinical study presented as a meeting abstract has shown the beneficial effects of transplantation of human amniotic stem cells (mesenchymal or epithelial) in CCl₄-induced cirrhotic rats with PH, improving HSC and LSEC phenotype, inflammation and liver function, resulting in an amelioration in PH and microvascular dysfunction.¹⁷¹

The promising results of cell therapy in preclinical models of CLD and PH support future and ongoing clinical trials using this strategy.

Lifestyle and dietary interventions

The basis of most of the abovementioned strategies is to improve deregulated processes in CLD (vasoconstriction, oxidative stress, etc.) with drugs or other molecules that specifically target these molecular pathways. Alternatively, wider effects on these pathological processes can also be achieved by modification in lifestyle; mainly diet and exercise.

Dietary approaches

The effects of certain diet components (dark chocolate, coffee/tea, fermented milk) on gut microbiota and its implications in CLD have been described above.^{112,152,172} In summary, consumption of these products ensures a proper diversity in the gut microbiota while some of them also possess antioxidant properties, thus ameliorating liver damage and hemodynamics in CLD. In addition

to these, the following dietary supplementations may have the potential to ameliorate PH.

Caffeine intake is extended worldwide, either in the form of coffee, tea or carbonated drinks. Independently of its effects on the gut microbiota,¹⁷³ caffeine consumption (in the form of coffee) has been associated with reduced liver fibrosis in precirrhotic patients.¹⁷⁴ Although there are no reports of its effects in PH in humans, prophylactic but also therapeutic treatment of BDL-cirrhotic rats with caffeine or caffeinated coffee ameliorated PP, liver inflammation and fibrosis, while also improving the extrahepatic vasculature.^{175,176}

Taurine is an amino-sulfonic acid ubiquitously expressed in mammals with many pleiotropic effects¹⁷⁷ and is also a component of the so-labeled 'energy drinks'. It has been reported that taurine deficiency leads to CLD in preclinical models,¹⁷⁸ suggesting a protective role in cirrhosis. Indeed, oral taurine administration has been proven to ameliorate PP due to reduction in fibrosis and systemic vasodilation in a rat model of mild cirrhosis¹⁷⁹ while more recently, a clinical trial in a small cohort of patients with clinically significant PH (HVPG >12 mmHg) also reported reduction in PP.¹⁸⁰ Thus, consumption of low carbonated 'energy drinks' (which are rich in taurine, and also contain caffeine) may have a positive impact in PH.

Finally, curcumin is a dietary product most common in Asian countries but extended worldwide that possesses anti-inflammatory, antiangiogenic and antiproliferative properties.¹⁸¹ Indeed, its administration to BDL-cirrhotic rats decreased hepatic fibrosis and ameliorated liver endothelial phenotype while inducing splanchnic vasoconstriction, which led to a reduction in PP.¹⁸²

Lifestyle interventions

In addition to nutritional strategies, nonsedentary lifestyle exerts beneficial effects on CLD complications such as PH. 'The SportDiet study' demonstrated that moderate exercise in combination with a controlled diet reduced PP and body weight in overweight patients with compensated cirrhosis and PH.¹⁸³ Moreover, another RCT evaluating the effects of exercise and diet intervention in patients with cirrhosis and PH showed an improvement in HVPG determined before and postintervention.¹⁸⁴ Moreover, the impact of

exercise therapy on cirrhosis and its complications has been extensively studied in patients^{185,186} as well as in preclinical models.¹⁸⁷

Conclusion

The knowledge about the pathophysiology of CLD has improved dramatically in the last few years. Consequently, many preclinical studies reported novel treatments with promising improvements in PH and its complications that, importantly, act at different levels of the disease (either improving the microcirculation or the liver's response to damage). Indeed, some of these proposed treatments are currently being tested in RCTs and could become new options for the treatment of CLD in the near future.

However, it seems obvious that the rate of success of the preclinical strategies at the bedside is still far from optimal. In this regard, new studies on the mechanisms of the disease (still to be fully understood) will provide new insights on novel targetable molecular pathways and metabolites. Nevertheless, we believe that in addition to the important research on drug development and efficacy, it is time to start wondering why these therapies fail in translation. What are the differences between preclinical models and current patients that make these therapies ineffective? Should the preclinical models integrate important variables present in humans, like comorbidities, epigenetics, aging, etc.? Are therapeutic windows different in humans? If so, could we 'desensitize' the human liver to these therapeutic windows? In our opinion, this second line of research could widen the current bottleneck in the translation of pharmacological strategies and reopen the box of failed therapeutic options in search of second chances.

Acknowledgement

MV and SG-M contributed equally to this article.

Funding

SG-M has a postdoctoral fellowship from the Stiftung für Leberkrankheiten and the Asociación Española Para el Estudio del Hígado. AF-I has a Sara Borrell contract from the Instituto de Salud Carlos III (CD15/00050). JG-S has received continued funding from the Instituto de Salud Carlos III (currently, FIS PI17/00012), the Spanish Ministry of Science, Innovation and Universities, Centro de Investigación Biomédica

en Red en Enfermedades Hepáticas y Digestivas (CIBEREHD), the European Union Funds FEDER ‘una manera de hacer Europa’ and the Stiftung für Leberkrankheiten.

Conflict of interest statement

The authors declare that there is no conflict of interest.

ORCID iD

Jordi Gracia-Sancho  <https://orcid.org/0000-0001-7736-4089>

References

- Bosch J, Abraldes JG, Berzigotti A, *et al.* The clinical use of HVPG measurements in chronic liver disease. *Nat Rev Gastroenterol Hepatol* 2009; 6: 573–582.
- Ripoll C, Groszmann R, Garcia-Tsao G, *et al.* Hepatic venous pressure gradient predicts clinical decompensation in patients with compensated cirrhosis. *Gastroenterology* 2007; 133: 481–488.
- Li T, Ke W, Sun P, *et al.* Carvedilol for portal hypertension in cirrhosis: systematic review with meta-analysis. *BMJ Open* 2016; 6: e010902.
- Kanwal F, Tansel A, Kramer JR, *et al.* Trends in 30-day and 1-year mortality among patients hospitalized with cirrhosis from 2004 to 2013. *Am J Gastroenterol* 2017; 112: 1287–1297.
- Watanabe N, Takashimizu S, Nishizaki Y, *et al.* An endothelin A receptor antagonist induces dilatation of sinusoidal endothelial fenestrae: implications for endothelin-1 in hepatic microcirculation. *J Gastroenterol* 2007; 42: 775–782.
- Feng H-Q, Weymouth ND and Rockey DC. Endothelin antagonism in portal hypertensive mice: implications for endothelin receptor-specific signaling in liver disease. *AJP Gastrointest Liver Physiol* 2009; 297: G27–G33.
- Zipprich A, Schenkel E, Gittinger F, *et al.* Selective endothelin-A blockade decreases portal pressure in patients with cirrhosis: a pilot study combining a local intraarterial and systemic administration. *J Hepatol* 2016; 64: S247.
- Tripathi D, Theraponos G, Ferguson JW, *et al.* Endothelin-1 contributes to maintenance of systemic but not portal haemodynamics in patients with early cirrhosis: a randomised controlled trial. *Gut* 2006; 55: 1290–1295.
- Lebrec D, Bosch J, Jalan R, *et al.* Hemodynamics and pharmacokinetics of tezosentan, a dual endothelin receptor antagonist, in patients with cirrhosis. *Eur J Clin Pharmacol* 2012; 68: 533–541.
- EU Clinical Trials Register, www.clinicaltrialsregister.eu/ctr-search/trial/2014-004624-21/DE#P. (accessed 17 July 2018)
- EU Clinical Trials Register, www.clinicaltrialsregister.eu/ctr-search/trial/2011-001139-22/AT. (accessed 17 July 2018)
- Heller J, Schepke M, Neef M, *et al.* Increased urotensin II plasma levels in patients with cirrhosis and portal hypertension. *J Hepatol* 2002; 37: 767–772.
- Trebicka J, Leifeld L, Hennenberg M, *et al.* Hemodynamic effects of urotensin II and its specific receptor antagonist palosuran in cirrhotic rats. *Hepatology* 2008; 47: 1264–1276.
- Gracia-Sancho J, Maeso-Díaz R, Fernández-Iglesias A, *et al.* New cellular and molecular targets for the treatment of portal hypertension. *Hepatol Int* 2015; 9: 183–191.
- Gracia-Sancho J, Laviña B, Rodríguez-Vilarrupla A, *et al.* Enhanced vasoconstrictor prostanoid production by sinusoidal endothelial cells increases portal perfusion pressure in cirrhotic rat livers. *J Hepatol* 2007; 47: 220–227.
- Laleman W, Van Landeghem L, Van der Elst I, *et al.* Nitroflurbiprofen, a nitric oxide-releasing cyclooxygenase inhibitor, improves cirrhotic portal hypertension in rats. *Gastroenterology* 2007; 132: 709–719.
- Gao J-H, Wen S-L, Yang W-J, *et al.* Celecoxib ameliorates portal hypertension of the cirrhotic rats through the dual inhibitory effects on the intrahepatic fibrosis and angiogenesis. *PLoS One* 2013; 8: e69309.
- Gao J-H, Wen S-L, Feng S, *et al.* Celecoxib and octreotide synergistically ameliorate portal hypertension via inhibition of angiogenesis in cirrhotic rats. *Angiogenesis* 2016; 19: 501–511.
- Rosado E, Rodríguez-Vilarrupla A, Gracia-Sancho J, *et al.* Terutroban, a TP-receptor antagonist, reduces portal pressure in cirrhotic rats. *Hepatology* 2013; 58: 1424–1435.
- US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT02802228>. (accessed 17 July 2018)
- Di Pascoli M, Zampieri F, Verardo A, *et al.* Inhibition of epoxyeicosatrienoic acid production in rats with cirrhosis has beneficial

- effects on portal hypertension by reducing splanchnic vasodilation. *Hepatology* 2016; 64: 923–930.
22. Steib CJ, Bilzer M, Op den Winkel M, *et al.* Treatment with the leukotriene inhibitor montelukast for 10 days attenuates portal hypertension in rat liver cirrhosis. *Hepatology* 2010; 51: 2086–2096.
 23. Bolognesi M, Di Pascoli M, Verardo A, *et al.* Splanchnic vasodilation and hyperdynamic circulatory syndrome in cirrhosis. *World J Gastroenterol* 2014; 20: 2555–2563.
 24. Tandon P, Abraldes JG, Berzigotti A, *et al.* Renin–angiotensin–aldosterone inhibitors in the reduction of portal pressure: a systematic review and meta-analysis. *J Hepatol* 2010; 53: 273–282.
 25. Granzow M, Schierwagen R, Klein S, *et al.* Angiotensin-II type 1 receptor-mediated Janus kinase 2 activation induces liver fibrosis. *Hepatology* 2014; 60: 334–348.
 26. Wang D, Yin J, Dong R, *et al.* Inhibition of Janus kinase-2 signalling pathway ameliorates portal hypertensive syndrome in partial portal hypertensive and liver cirrhosis rats. *Dig Liver Dis* 2015; 47: 315–323.
 27. Wang D, Wang Q, Yin J, *et al.* Combined administration of propranolol + AG490 offers better effects on portal hypertensive rats with cirrhosis. *J Gastroenterol Hepatol* 2016; 31: 1037–1044.
 28. Klein S, Van Beuge MM, Granzow M, *et al.* HSC-specific inhibition of Rho-kinase reduces portal pressure in cirrhotic rats without major systemic effects. *J Hepatol* 2012; 57: 1220–1227.
 29. Fukuda T, Narahara Y, Kanazawa H, *et al.* Effects of fasudil on the portal and systemic hemodynamics of patients with cirrhosis. *J Gastroenterol Hepatol* 2014; 29: 325–329.
 30. Wei L, Yang J, Wang M, *et al.* Sodium ferulate lowers portal pressure in rats with secondary biliary cirrhosis through the RhoA/Rho-kinase signaling pathway: a preliminary study. *Int J Mol Med* 2014; 34: 1257–1267.
 31. Zhang C-G, Zhang B, Deng W-S, *et al.* Role of estrogen receptor β selective agonist in ameliorating portal hypertension in rats with CCl₄-induced liver cirrhosis. *World J Gastroenterol* 2016; 22: 4484–4500.
 32. Klein S, Herath CB, Schierwagen R, *et al.* Hemodynamic effects of the non-peptidic angiotensin-(1-7) agonist AVE0991 in liver cirrhosis. *PLoS One* 2015; 10: e0138732.
 33. Grace JA, Klein S, Herath CB, *et al.* Activation of the MAS receptor by angiotensin-(1-7) in the renin-angiotensin system mediates mesenteric vasodilatation in cirrhosis. *Gastroenterology* 2013; 145: 874–884.e5.
 34. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT00592475>. (accessed 17 July 2018)
 35. Wong F, Bernardi M, Horsmans Y, *et al.* Effects of satavaptan, an oral vasopressin V₂ receptor agonist, on management of ascites and morbidity in liver cirrhosis in a long-term, placebo-controlled study. *J Hepatol* 2009; 50: S42–S43.
 36. European Association for the Study of the Liver. EASL clinical practice guidelines on the management of ascites, spontaneous bacterial peritonitis, and hepatorenal syndrome in cirrhosis. *J Hepatol* 2010; 53: 397–417.
 37. Fernández-Varo G, Oró D, Cable EE, *et al.* Vasopressin 1a receptor partial agonism increases sodium excretion and reduces portal hypertension and ascites in cirrhotic rats. *Hepatology* 2016; 63: 207–216.
 38. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT02929407>. (accessed 17 July 2018)
 39. Duong HTT, Dong Z, Su L, *et al.* The use of nanoparticles to deliver nitric oxide to hepatic stellate cells for treating liver fibrosis and portal hypertension. *Small* 2015; 11: 2291–2304.
 40. Matei V, Rodríguez-Vilarrupla A, Deulofeu R, *et al.* The eNOS cofactor tetrahydrobiopterin improves endothelial dysfunction in livers of rats with CCl₄ cirrhosis. *Hepatology* 2006; 44: 44–52.
 41. Biecker E, Trebicka J, Kang A, *et al.* Treatment of bile duct-ligated rats with the nitric oxide synthase transcription enhancer AVE 9488 ameliorates portal hypertension. *Liver Int* 2008; 28: 331–338.
 42. Reverter E, Mesonero F, Seijo S, *et al.* Effects of sapropterin on portal and systemic hemodynamics in patients with cirrhosis and portal hypertension: a bicentric double-blind placebo-controlled study. *Am J Gastroenterol* 2015; 110: 985–992.
 43. EU Clinical Trials Register, www.clinicaltrialsregister.eu/ctr-search/trial/2015-004031-12/GB. (accessed 17 July 2018)

44. Schwabl P, Brusilovskaya K, Supper P, *et al.* The soluble guanylate cyclase stimulator riociguat reduces fibrogenesis and portal pressure in cirrhotic rats. *Sci Rep* 2018; 8: 9372.
45. Buys ES, Zimmer DP, Chickering J, *et al.* Discovery and development of next generation sGC stimulators with diverse multidimensional pharmacology and broad therapeutic potential. *Nitric Oxide* 2018; 78: 72–80.
46. Lee K-C, Yang Y-Y, Wang Y-W, *et al.* Acute administration of sildenafil enhances hepatic cyclic guanosine monophosphate production and reduces hepatic sinusoid resistance in cirrhotic patients. *Hepatol Res* 2008; 38: 1186–1193.
47. Deibert P, Schumacher Y-O, Ruecker G, *et al.* Effect of vardenafil, an inhibitor of phosphodiesterase-5, on portal haemodynamics in normal and cirrhotic liver – results of a pilot study. *Aliment Pharmacol Ther* 2006; 23: 121–128.
48. Deibert P, Lazaro A, Stankovic Z, *et al.* Beneficial long term effect of a phosphodiesterase-5-inhibitor in cirrhotic portal hypertension: a case report with 8 years follow-up. *World J Gastroenterol* 2018; 24: 438–444.
49. Verbeke L, Farre R, Trebicka J, *et al.* Obeticholic acid, a farnesoid X receptor agonist, improves portal hypertension by two distinct pathways in cirrhotic rats. *Hepatology* 2014; 59: 2286–2298.
50. EU Clinical Trials Register, www.clinicaltrialsregister.eu/ctr-search/trial/2016-002965-67/NL. (accessed 17 July 2018)
51. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT01473524>. (accessed 17 July 2018)
52. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT02308111>. (accessed 17 July 2018)
53. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT02548351>. (accessed 17 July 2018)
54. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT03439254>. (accessed 17 July 2018)
55. US Food and Drug Administration. FDA Drug Safety Communication: FDA warns about serious liver injury with Ocaliva (obeticholic acid) for rare chronic liver disease, (publish date 9 September 2017). <https://www.fda.gov/Drugs/DrugSafety/ucm576656.htm> (accessed 17 July 2018)
56. Baghdasaryan A, Claudel T, Gumhold J, *et al.* Dual farnesoid X receptor/TGR5 agonist INT-767 reduces liver injury in the Mdr2-/- (Abcb4-/-) mouse cholangiopathy model by promoting biliary HCO₃⁻ output. *Hepatology* 2011; 54: 1303–1312.
57. Zhang S, Wang J, Liu Q, *et al.* Farnesoid X receptor agonist WAY-362450 attenuates liver inflammation and fibrosis in murine model of non-alcoholic steatohepatitis. *J Hepatol* 2009; 51: 380–388.
58. Trebicka J, Hennenberg M, Laleman W, *et al.* Atorvastatin lowers portal pressure in cirrhotic rats by inhibition of RhoA/Rho-kinase and activation of endothelial nitric oxide synthase. *Hepatology* 2007; 46: 242–253.
59. Abraldes JG, Rodríguez-Vilarrupla A, Graupera M, *et al.* Simvastatin treatment improves liver sinusoidal endothelial dysfunction in CCl4 cirrhotic rats. *J Hepatol* 2007; 46: 1040–1046.
60. Meireles CZ, Pasarin M, Lozano JJ, *et al.* Simvastatin attenuates liver injury in rodents with biliary cirrhosis submitted to hemorrhage/resuscitation. *Shock* 2017; 47: 370–377.
61. Trebicka J, Hennenberg M, Odenthal M, *et al.* Atorvastatin attenuates hepatic fibrosis in rats after bile duct ligation via decreased turnover of hepatic stellate cells. *J Hepatol* 2010; 53: 702–712.
62. Klein S, Klösel J, Schierwagen R, *et al.* Atorvastatin inhibits proliferation and apoptosis, but induces senescence in hepatic myofibroblasts and thereby attenuates hepatic fibrosis in rats. *Lab. Invest* 2012; 92: 1440–1450.
63. Marrone G, Russo L, Rosado E, *et al.* The transcription factor KLF2 mediates hepatic endothelial protection and paracrine endothelial-stellate cell deactivation induced by statins. *J Hepatol* 2013; 58: 98–103.
64. Miao Q, Zeng X, Ma G, *et al.* Simvastatin suppresses the proangiogenic microenvironment of human hepatic stellate cells via the Kruppel-like factor 2 pathway. *Rev Esp Enferm Dig* 2015; 107: 63–71.
65. Marrone G, Maeso-Díaz R, García-Cardena G, *et al.* KLF2 exerts antifibrotic and vasoprotective effects in cirrhotic rat livers: behind the molecular mechanisms of statins. *Gut* 2015; 64: 1434–1443.
66. Rodríguez S, Raurell I, Torres-Arauz M, *et al.* A nitric oxide-donating statin decreases portal pressure with a better toxicity profile than

- conventional statins in cirrhotic rats. *Sci Rep* 2017; 7: 40461.
67. Mohanty A, Tate JP and Garcia-Tsao G. Statins are associated with a decreased risk of decompensation and death in veterans with hepatitis c-related compensated cirrhosis. *Gastroenterology* 2016; 150: 430–440.e1.
 68. Simon TG, Bonilla H, Yan P, *et al.* Atorvastatin and fluvastatin are associated with dose-dependent reductions in cirrhosis and hepatocellular carcinoma, among patients with hepatitis C virus: results from ERCHIVES. *Hepatology* 2016; 64: 47–57.
 69. Pollo-Flores P, Soldan M, Santos UC, *et al.* Three months of simvastatin therapy vs. placebo for severe portal hypertension in cirrhosis: a randomized controlled trial. *Dig Liver Dis* 2015; 47: 957–963.
 70. Abraldes JG, Villanueva C, Aracil C, *et al.* Addition of simvastatin to standard therapy for the prevention of variceal rebleeding does not reduce rebleeding but increases survival in patients with cirrhosis. *Gastroenterology* 2016; 150: 1160–1170.e3.
 71. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT02465645>. (accessed 17 July 2018)
 72. Tacke F, Luedde T and Trautwein C. Inflammatory pathways in liver homeostasis and liver injury. *Clin Rev Allergy Immunol* 2009; 36: 4–12.
 73. Jairath V and Burroughs AK. Anticoagulation in patients with liver cirrhosis: complication or therapeutic opportunity? *Gut* 2013; 62: 479–482.
 74. Cerini F, Vilaseca M, Lafoz E, *et al.* Enoxaparin reduces hepatic vascular resistance and portal pressure in cirrhotic rats. *J Hepatol* 2016; 64: 834–842.
 75. Villa E, Cammà C, Marietta M, *et al.* Enoxaparin prevents portal vein thrombosis and liver decompensation in patients with advanced cirrhosis. *Gastroenterology* 2012; 143: 1253–1260.e4.
 76. Vilaseca M, García-Calderó H, Lafoz E, *et al.* The anticoagulant rivaroxaban lowers portal hypertension in cirrhotic rats mainly by deactivating hepatic stellate cells. *Hepatology* 2017; 65: 2031–2044.
 77. EU Clinical Trials Register, www.clinicaltrialsregister.eu/ctr-search/trial/2014-005523-27/ES. (accessed 17 July 2018)
 78. EU Clinical Trials Register, www.clinicaltrialsregister.eu/ctr-search/trial/2016-003240-37/ES. (accessed 17 July 2018)
 79. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT02247414>. (accessed 17 July 2018)
 80. Corpechot C, Barbu V, Wendum D, *et al.* Hypoxia-induced VEGF and collagen I expressions are associated with angiogenesis and fibrogenesis in experimental cirrhosis. *Hepatology* 2002; 35: 1010–1021.
 81. Fernandez M, Vizzutti F, Garcia-Pagan JC, *et al.* Anti-VEGF receptor-2 monoclonal antibody prevents portal-systemic collateral vessel formation in portal hypertensive mice. *Gastroenterology* 2004; 126: 886–894.
 82. Fernandez M, Mejias M, Angermayr B, *et al.* Inhibition of VEGF receptor-2 decreases the development of hyperdynamic splanchnic circulation and portal-systemic collateral vessels in portal hypertensive rats. *J Hepatol* 2005; 43: 98–103.
 83. Reiberger T, Angermayr B, Schwabl P, *et al.* Sorafenib attenuates the portal hypertensive syndrome in partial portal vein ligated rats. *J Hepatol* 2009; 51: 865–873.
 84. Yang Y-Y, Liu R-S, Lee P-C, *et al.* Anti-VEGFR agents ameliorate hepatic venous dysregulation/microcirculatory dysfunction, splanchnic venous pooling and ascites of NASH-cirrhotic rat. *Liver Int* 2014; 34: 521–534.
 85. Lin H-C, Huang Y-T, Yang Y-Y, *et al.* Beneficial effects of dual vascular endothelial growth factor receptor/fibroblast growth factor receptor inhibitor brivanib alaninate in cirrhotic portal hypertensive rats. *J Gastroenterol Hepatol* 2014; 29: 1073–1082.
 86. Uschner FE, Nikolova I, Klein S, *et al.* The multikinase inhibitor regorafenib improves portal hypertension in presence and absence of cirrhosis in rats. *J Hepatol* 2016; 64: S134–S135.
 87. Qu K, Huang Z, Lin T, *et al.* New insight into the anti-liver fibrosis effect of multitargeted tyrosine kinase inhibitors: from molecular target to clinical trials. *Front Pharmacol* 2015; 6: 300.
 88. Pinter M, Sieghart W, Reiberger T, *et al.* The effects of sorafenib on the portal hypertensive syndrome in patients with liver cirrhosis and hepatocellular carcinoma—a pilot study. *Aliment Pharmacol Ther* 2012; 35: 83–91.

89. Mejias M, Coch L, Berzigotti A, *et al.* Antiangiogenic and antifibrogenic activity of pigment epithelium-derived factor (PEDF) in bile duct-ligated portal hypertensive rats. *Gut* 2015; 64: 657–666.
90. Coch L, Mejias M, Berzigotti A, *et al.* Disruption of negative feedback loop between vasohibin-1 and vascular endothelial growth factor decreases portal pressure, angiogenesis, and fibrosis in cirrhotic rats. *Hepatology* 2014; 60: 633–647.
91. Gracia-Sancho J, Contreras PC, Vila S, *et al.* The pan caspase inhibitor emricasan improves the hepatic microcirculatory dysfunction of CCl₄-cirrhotic rats leading to portal hypertension amelioration and cirrhosis regression. *Hepatology* 2016; 64: 811–1050.
92. Eguchi A, Koyama Y, Wree A, *et al.* Emricasan, a pan-caspase inhibitor, improves survival and portal hypertension in a murine model of common bile-duct ligation. *J Mol Med* 2018; 96: 575–583.
93. Mehta G, Rousell S, Burgess G, *et al.* A placebo-controlled, multicenter, double-blind, phase 2 randomized trial of the pan-caspase inhibitor emricasan in patients with acutely decompensated cirrhosis. *J Clin Exp Hepatol* 2018; 8: 224–234.
94. Garcia-Tsao G, Fuchs M, Shiffman M, *et al.* Emricasan (IDN-6556) lowers portal pressure in patients with compensated cirrhosis and severe portal hypertension. *Hepatology*. Epub ahead of print 31 July 2018. DOI: 10.1002/hep.30199.
95. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT02960204>. (accessed 17 July 2018)
96. Wilson CH and Kumar S. Caspases in metabolic disease and their therapeutic potential. *Cell Death Differ* 2018; 25: 1010–1024.
97. Loomba R, Lawitz E, Mantry PS, *et al.* The ASK1 inhibitor selonsertib in patients with nonalcoholic steatohepatitis: a randomized, phase 2 trial. *Hepatology* 2018; 67: 549–559.
98. Ni H-M, McGill MR, Chao X, *et al.* Caspase inhibition prevents tumor necrosis factor- α -induced apoptosis and promotes necrotic cell death in mouse hepatocytes in vivo and in vitro. *Am J Pathol* 2016; 186: 2623–2636.
99. Wright MC, Issa R, Smart DE, *et al.* Gliotoxin stimulates the apoptosis of human and rat hepatic stellate cells and enhances the resolution of liver fibrosis in rats. *Gastroenterology* 2001; 121: 685–698.
100. Gracia-Sancho J, Laviña B, Rodríguez-Vilarrupla A, *et al.* Increased oxidative stress in cirrhotic rat livers: A potential mechanism contributing to reduced nitric oxide bioavailability. *Hepatology* 2007; 47: 1248–1256.
101. Vilaseca M, García-Calderó H, Lafoz E, *et al.* Mitochondria-targeted antioxidant mitoquinone deactivates human and rat hepatic stellate cells and reduces portal hypertension in cirrhotic rats. *Liver Int* 2017; 37: 1002–1012.
102. Chacko BK, Srivastava A, Johnson MS, *et al.* Mitochondria-targeted ubiquinone (MitoQ) decreases ethanol-dependent micro and macro hepatosteatosis. *Hepatology* 2011; 54: 153–163.
103. Mukhopadhyay P, Horváth B, Zsengeller Z, *et al.* Mitochondrial reactive oxygen species generation triggers inflammatory response and tissue injury associated with hepatic ischemia-reperfusion: therapeutic potential of mitochondrially targeted antioxidants. *Free Radic Biol Med* 2012; 53: 1123–1138.
104. Turkseven S, Bolognesi M, Brocca A, *et al.* In cirrhotic rats, mitochondria-targeted antioxidant mitoquinone attenuates liver inflammation and fibrosis by modulating oxidative stress and mitophagy. *J Hepatol* 2018; 68: S365–S604.
105. Petrônio MS, Zeraik ML, Fonseca LM, *et al.* Apocynin: chemical and biophysical properties of a NADPH oxidase inhibitor. *Molecules* 2013; 18: 2821–2839.
106. Castro MC, Francini F, Schinella G, *et al.* Apocynin administration prevents the changes induced by a fructose-rich diet on rat liver metabolism and the antioxidant system. *Clin Sci (Lond)* 2012; 123: 681–692.
107. Fan R, Shan X, Qian H, *et al.* Protective effect of apocynin in an established alcoholic steatohepatitis rat model. *Immunopharmacol Immunotoxicol* 2012; 34: 633–638.
108. Rahman MM, Muse AY, Khan DMIO, *et al.* Apocynin prevented inflammation and oxidative stress in carbon tetra chloride induced hepatic dysfunction in rats. *Biomed Pharmacother* 2017; 92: 421–428.
109. Gracia-Sancho J, Laviña B, Rodríguez-Vilarrupla A, *et al.* Evidence against a role for NADPH oxidase modulating hepatic vascular tone in cirrhosis. *Gastroenterology* 2007; 133: 959–966.

110. Bettaieb A, Jiang JX, Sasaki Y, *et al.* Hepatocyte nicotinamide adenine dinucleotide phosphate reduced oxidase 4 regulates stress signaling, fibrosis, and insulin sensitivity during development of steatohepatitis in mice. *Gastroenterology* 2015; 149: 468–480.e10.
111. Hernández-Guerra M, García-Pagán JC, Turnes J, *et al.* Ascorbic acid improves the intrahepatic endothelial dysfunction of patients with cirrhosis and portal hypertension. *Hepatology* 2006; 43: 485–491.
112. De Gottardi A, Berzigotti A, Seijo S, *et al.* Postprandial effects of dark chocolate on portal hypertension in patients with cirrhosis: results of a phase 2, double-blind, randomized controlled trial. *Am J Clin Nutr* 2012; 96: 584–590.
113. Lavina B, Gracia-Sancho J, Rodríguez-Vilarrupla A, *et al.* Superoxide dismutase gene transfer reduces portal pressure in CCl₄ cirrhotic rats with portal hypertension. *Gut* 2009; 58: 118–125.
114. García-Calderó H, Rodríguez-Vilarrupla A, Gracia-Sancho J, *et al.* Tempol administration, a superoxide dismutase mimetic, reduces hepatic vascular resistance and portal pressure in cirrhotic rats. *J Hepatol* 2011; 54: 660–665.
115. Guillaume M, Rodríguez-Vilarrupla A, Gracia-Sancho J, *et al.* Recombinant human manganese superoxide dismutase reduces liver fibrosis and portal pressure in CCl₄-cirrhotic rats. *J Hepatol* 2013; 58: 240–246.
116. Gracia-Sancho J, Villarreal G, Zhang Y, *et al.* Activation of SIRT1 by resveratrol induces KLF2 expression conferring an endothelial vasoprotective phenotype. *Cardiovasc Res* 2010; 85: 514–519.
117. Chan C-C, Lee K-C, Huang Y-H, *et al.* Regulation by resveratrol of the cellular factors mediating liver damage and regeneration after acute toxic liver injury. *J Gastroenterol Hepatol* 2014; 29: 603–613.
118. Kessoku T, Imajo K, Honda Y, *et al.* Resveratrol ameliorates fibrosis and inflammation in a mouse model of nonalcoholic steatohepatitis. *Sci Rep* 2016; 6: 22251.
119. Szkudelski T and Szkudelska K. Potential of resveratrol in mitigating metabolic disturbances induced by ethanol. *Biomed Pharmacother* 2018; 101: 579–584.
120. Di Pascoli M, Diví M, Rodríguez-Vilarrupla A, *et al.* Resveratrol improves intrahepatic endothelial dysfunction and reduces hepatic fibrosis and portal pressure in cirrhotic rats. *J Hepatol* 2013; 58: 904–910.
121. Wang W, Yan J, Wang H, *et al.* Rapamycin ameliorates inflammation and fibrosis in the early phase of cirrhotic portal hypertension in rats through inhibition of mTORC1 but not mTORC2. *PLoS One* 2014; 9: e83908.
122. Biecker E, De Gottardi A, Neef M, *et al.* Long-term treatment of bile duct-ligated rats with rapamycin (sirolimus) significantly attenuates liver fibrosis: analysis of the underlying mechanisms. *J Pharmacol Exp Ther* 2005; 313: 952–961.
123. Mejias M, Garcia-Pras E, Gallego J, *et al.* Relevance of the mTOR signaling pathway in the pathophysiology of splenomegaly in rats with chronic portal hypertension. *J Hepatol* 2010; 52: 529–539.
124. Wu K-C, Huang H-C, Chang T, *et al.* Effect of sirolimus on liver cirrhosis and hepatic encephalopathy of common bile duct-ligated rats. *Eur J Pharmacol* 2018; 824: 133–139.
125. Spector AA, Fang X, Snyder GD, *et al.* Epoxyeicosatrienoic acids (EETs): metabolism and biochemical function. *Prog Lipid Res* 2004; 43: 55–90.
126. López-Vicario C, Alcaraz-Quiles J, García-Alonso V, *et al.* Inhibition of soluble epoxide hydrolase modulates inflammation and autophagy in obese adipose tissue and liver: role for omega-3 epoxides. *Proc Natl Acad Sci. U S A* 2015; 112: 536–541.
127. Khan AH, Falck JR, Manthali VL, *et al.* Epoxyeicosatrienoic acid analog attenuates angiotensin II hypertension and kidney injury. *Front Pharmacol* 2014; 5: 216.
128. Deng W, Zhu Y, Lin J, *et al.* Inhibition of soluble epoxide hydrolase lowers portal hypertension in cirrhotic rats by ameliorating endothelial dysfunction and liver fibrosis. *Prostaglandins Other Lipid Mediat* 2017; 131: 67–74.
129. Zhang C-H, Zheng L, Gui L, *et al.* Soluble epoxide hydrolase inhibition with t-TUCB alleviates liver fibrosis and portal pressure in carbon tetrachloride-induced cirrhosis in rats. *Clin Res Hepatol Gastroenterol* 2018; 42: 118–125.
130. Huang Y, Qin J, Sun D, *et al.* Inhibition of soluble epoxide hydrolase reduces portal pressure by protecting mesenteric artery myogenic responses in cirrhotic rats. *Prostaglandins Other Lipid Mediat* 2017; 131: 17–24.

131. Oró D, Yudina T, Fernández-Varo G, *et al.* Cerium oxide nanoparticles reduce steatosis, portal hypertension and display anti-inflammatory properties in rats with liver fibrosis. *J Hepatol* 2016; 64: 691–698.
132. Shiraki A, Oyama J, Komoda H, *et al.* The glucagon-like peptide 1 analog liraglutide reduces TNF- α -induced oxidative stress and inflammation in endothelial cells. *Atherosclerosis* 2012; 221: 375–382.
133. Saisho Y. Metformin and inflammation: its potential beyond glucose-lowering effect. *Endocr Metab Immune Disord Drug Targets* 2015; 15: 196–205.
134. Stolarz AJ, Farris RA, Wiley CA, *et al.* Fenofibrate attenuates neutrophilic inflammation in airway epithelia: potential drug repurposing for cystic fibrosis. *Clin Transl Sci* 2015; 8: 696–701.
135. De Mesquita FC, Guixé-Muntet S, Fernández-Iglesias A, *et al.* Liraglutide improves liver microvascular dysfunction in cirrhosis: evidence from translational studies. *Sci Rep* 2017; 7: 3255.
136. Armstrong MJ, Gaunt P, Aithal GP, *et al.* Liraglutide safety and efficacy in patients with non-alcoholic steatohepatitis (LEAN): a multicentre, double-blind, randomised, placebo-controlled phase 2 study. *Lancet* 2016; 387: 679–690.
137. Tripathi DM, Erice E, Lafoz E, *et al.* Metformin reduces hepatic resistance and portal pressure in cirrhotic rats. *Am J Physiol Gastrointest Liver Physiol* 2015; 309: G301–G309.
138. Rodríguez-Vilarrupla A, Laviña B, García-Calderó H, *et al.* PPAR α activation improves endothelial dysfunction and reduces fibrosis and portal pressure in cirrhotic rats. *J Hepatol* 2012; 56: 1033–1039.
139. Zhang W, Niu M, Yan K, *et al.* Stat3 pathway correlates with the roles of leptin in mouse liver fibrosis and sterol regulatory element binding protein-1c expression of rat hepatic stellate cells. *Int J Biochem Cell Biol* 2013; 45: 736–744.
140. Zhai X, Yan K, Fan J, *et al.* The β -catenin pathway contributes to the effects of leptin on SREBP-1c expression in rat hepatic stellate cells and liver fibrosis. *Br J Pharmacol* 2013; 169: 197–212.
141. Delgado MG, Gracia-Sancho J, Marrone G, *et al.* Leptin receptor blockade reduces intrahepatic vascular resistance and portal pressure in an experimental model of rat liver cirrhosis. *Am J Physiol Gastrointest Liver Physiol* 2013; 305: G496–G502.
142. Zhao T-Y, Su L-P, Ma C-Y, *et al.* IGF-1 decreases portal vein endotoxin via regulating intestinal tight junctions and plays a role in attenuating portal hypertension of cirrhotic rats. *BMC Gastroenterol* 2015; 15: 77.
143. Gómez-Hurtado I, Such J and Francés R. Microbioma y traslocación bacteriana en la cirrosis. *Gastroenterol Hepatol* 2016; 39: 687–696.
144. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT02508623>. (accessed 17 July 2018)
145. EU Clinical Trials Register, www.clinicaltrialsregister.eu/ctr-search/trial/2012-002890-71/DK. (accessed 17 July 2018)
146. EU Clinical Trials Register, www.clinicaltrialsregister.eu/ctr-search/trial/2013-004708-20/GB. (accessed 17 July 2018)
147. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT01134692>. (accessed 17 July 2018)
148. Rincón D, Vaquero J, Hernando A, *et al.* Oral probiotic VSL#3 attenuates the circulatory disturbances of patients with cirrhosis and ascites. *Liver Int* 2014; 34: 1504–1512.
149. Dhiman RK, Rana B, Agrawal S, *et al.* Probiotic VSL#3 reduces liver disease severity and hospitalization in patients with cirrhosis: a randomized, controlled trial. *Gastroenterology* 2014; 147: 1327–1337.e3.
150. Moratalla A, Caparrós E, Juanola O, *et al.* Bifidobacterium pseudocatenulatum CECT7765 induces an M2 anti-inflammatory transition in macrophages from patients with cirrhosis. *J Hepatol* 2016; 64: 135–45.
151. Gómez-Hurtado I, Zapater P, Portune K, *et al.* Improved hemodynamic and liver function in portal hypertensive cirrhotic rats after administration of B. pseudocatenulatum CECT 7765. *Eur J Nutr*. Epub ahead of print 10 May 2018. DOI: 10.1007/s00394-018-1709-y.
152. Bajaj JS, Idilman R, Mabudian L, *et al.* Diet affects gut microbiota and modulates hospitalization risk differentially in an international cirrhosis cohort. *Hepatology* 2018; 68: 234–247.
153. Marrone G, Shah VH and Gracia-Sancho J. Sinusoidal communication in liver fibrosis and regeneration. *J Hepatol* 2016; 65: 608–617.

154. Ikenaga N, Peng Z-W, Vaid KA, *et al.* Selective targeting of lysyl oxidase-like 2 (LOXL2) suppresses hepatic fibrosis progression and accelerates its reversal. *Gut* 2017; 66: 1697–1708.
155. Harrison SA, Abdelmalek MF, Caldwell S, *et al.* Simtuzumab is ineffective for patients with bridging fibrosis or compensated cirrhosis caused by nonalcoholic steatohepatitis. *Gastroenterology* 2018; 155: 1140–1153.
156. Terai S, Ishikawa T, Omori K, *et al.* Improved liver function in patients with liver cirrhosis after autologous bone marrow cell infusion therapy. *Stem Cells* 2006; 24: 2292–2298.
157. Lyra AC, Soares MBP, da Silva LFM, *et al.* Infusion of autologous bone marrow mononuclear cells through hepatic artery results in a short-term improvement of liver function in patients with chronic liver disease: a pilot randomized controlled study. *Eur J Gastroenterol Hepatol* 2010; 22: 33–42.
158. Hannoun Z, Steichen C, Dianat N, *et al.* The potential of induced pluripotent stem cell derived hepatocytes. *J Hepatol* 2016; 65: 182–199.
159. Nagamoto Y, Takayama K, Ohashi K, *et al.* Transplantation of a human iPSC-derived hepatocyte sheet increases survival in mice with acute liver failure. *J Hepatol* 2016; 64: 1068–1075.
160. Yu Y, Liu H, Ikeda Y, *et al.* Hepatocyte-like cells differentiated from human induced pluripotent stem cells: relevance to cellular therapies. *Stem Cell Res* 2012; 9: 196–207.
161. Pietrosi G, Vizzini G, Gerlach J, *et al.* Phases I–II matched case-control study of human fetal liver cell transplantation for treatment of chronic liver disease. *Cell Transplant* 2015; 24: 1627–1638.
162. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT01560845>. (accessed 17 July 2018)
163. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT03209986>. (accessed 17 July 2018)
164. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT01573923>. (accessed 17 July 2018)
165. Parolini O, Alviano F, Bagnara GP, *et al.* Concise review: isolation and characterization of cells from human term placenta: outcome of the first international workshop on placenta derived stem cells. *Stem Cells* 2008; 26: 300–311.
166. Jung J, Choi JH, Lee Y, *et al.* Human placenta-derived mesenchymal stem cells promote hepatic regeneration in ccl4 -injured rat liver model via increased autophagic mechanism. *Stem Cells* 2013; 31: 1584–1596.
167. Zhang D, Jiang M and Miao D. Transplanted human amniotic membrane-derived mesenchymal stem cells ameliorate carbon tetrachloride-induced liver cirrhosis in mouse. *PLoS One* 2011; 6: e16789.
168. Manuelpillai U, Lourensz D, Vaghjiani V, *et al.* Human amniotic epithelial cell transplantation induces markers of alternative macrophage activation and reduces established hepatic fibrosis. *PLoS One* 2012; 7: e38631.
169. Lee P-H, Tu C-T, Hsiao C-C, *et al.* Antifibrotic activity of human placental amnion membrane-derived CD34+ mesenchymal stem/progenitor cell transplantation in mice with thioacetamide-induced liver injury. *Stem Cells Transl Med* 2016; 5: 1473–1484.
170. Brückner S, Zipprich A, Hempel M, *et al.* Improvement of portal venous pressure in cirrhotic rat livers by systemic treatment with adipose tissue-derived mesenchymal stromal cells. *Cytotherapy* 2017; 19: 1462–1473.
171. Fernández-Iglesias A, Pietrosi G, Pampalone M, *et al.* Stem cells as a new therapeutic strategy for portal hypertension and cirrhosis. *J Hepatol* 2018; 68: S8–S9.
172. Gracia-Sancho J and Bosch J. Is it tea time for portal hypertension? *Clin Sci* 2014; 126: 631–632.
173. Cowan TE, Palmnäs MSA, Yang J, *et al.* Chronic coffee consumption in the diet-induced obese rat: impact on gut microbiota and serum metabolomics. *J Nutr Biochem* 2014; 25: 489–495.
174. Modi AA, Feld JJ, Park Y, *et al.* Increased caffeine consumption is associated with reduced hepatic fibrosis. *Hepatology* 2010; 51: 201–209.
175. Hsu S-J, Lee F-Y, Wang S-S, *et al.* Caffeine ameliorates hemodynamic derangements and portosystemic collaterals in cirrhotic rats. *Hepatology* 2015; 61: 1672–1684.
176. Arauz J, Zarco N, Hernández-Aquino E, *et al.* Coffee consumption prevents fibrosis in a rat model that mimics secondary biliary cirrhosis in humans. *Nutr Res* 2017; 40: 65–74.

177. Huxtable RJ. Physiological actions of taurine. *Physiol Rev* 1992; 72: 101–163.
178. Warskulat U, Borsch E, Reinehr R, *et al.* Chronic liver disease is triggered by taurine transporter knockout in the mouse. *FASEB J* 2006; 20: 574–576.
179. Liang J, Deng X, Lin Z-X, *et al.* Attenuation of portal hypertension by natural taurine in rats with liver cirrhosis. *World J Gastroenterol* 2009; 15: 4529–4537.
180. Schwarzer R, Kivaranovic D, Mandorfer M, *et al.* Randomised clinical study: the effects of oral taurine 6g/day vs placebo on portal hypertension. *Aliment Pharmacol Ther* 2018; 47: 86–94.
181. Bachmeier B, Killian P and Melchart D. The role of curcumin in prevention and management of metastatic disease. *Int J Mol Sci* 2018; 19: 1716.
182. Hsu S-J, Lee J-Y, Lin T-Y, *et al.* The beneficial effects of curcumin in cirrhotic rats with portal hypertension. *Biosci Rep* 2017; 37: BSR20171015.
183. Berzigotti A, Albillos A, Villanueva C, *et al.* Effects of an intensive lifestyle intervention program on portal hypertension in patients with cirrhosis and obesity: the SportDiet study. *Hepatology* 2017; 65: 1293–1305.
184. Macías-Rodríguez RU, Ilarraza-Lomelí H, Ruiz-Margáin A, *et al.* Changes in hepatic venous pressure gradient induced by physical exercise in cirrhosis: results of a pilot randomized open clinical trial. *Clin Transl Gastroenterol* 2016; 7: e180.
185. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT02776553>. (accessed 17 July 2018)
186. US National Library of Medicine. ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT02949505>. (accessed 17 July 2018)
187. Lafoz E, Vilaseca M, García-Calderó H, *et al.* Impacto del ejercicio con y sin dieta en la regresión de la cirrosis experimental en ratas tras la detención del agente etiológico hepatotóxico. *Gastroenterol Hepatol* 2018; 41: 24.

Visit SAGE journals online
[journals.sagepub.com/
home/tag](https://journals.sagepub.com/home/tag)

 SAGE journals