



## Projection theorems in hyperbolic space

ZOLTÁN M. BALOGH AND ANNINA ISELI 

**Abstract.** We establish Marstrand-type projection theorems for orthogonal projections along geodesics onto  $m$ -dimensional subspaces of the hyperbolic  $n$ -space by a geometric argument. Moreover, we obtain a Besicovitch–Federer type characterization of purely unrectifiable sets in terms of these hyperbolic orthogonal projections.

**Mathematics Subject Classification.** 28A78, 53C22.

**Keywords.** Hausdorff dimension, Projections, Hyperbolic space.

**1. Introduction.** Marstrand’s theorem [20] states that given a Borel set  $A$  in  $\mathbb{R}^2$ , for almost every line  $L$  the orthogonal projection of  $A$  onto  $L$  is a set of Hausdorff dimension equal to the minimum of 1 and the Hausdorff dimension of  $A$ . This result has marked the start of a large sequence of results in the same spirit. In particular, Marstrand’s theorem has been sharpened and generalized to higher dimensions by Kaufman [19], Falconer [9], and Mattila [22]. Also, similar problems have been studied in various other settings such as Heisenberg groups [1, 2, 15] and normed spaces [3, 17], as well as for radial projections in [25], different notions of measure and dimension [10, 11, 14], and restricted families of projections [8, 12, 26] (and references therein). In this paper, we prove Marstrand-type projection theorems as well as a Besicovitch–Federer-type projection theorem (i.e. a characterization of purely unrectifiable sets in terms of projections) for orthogonal projection along geodesics in the hyperbolic  $n$ -space. In particular, we generalize previous results of the authors [4] to higher dimensions. An extended introduction to the topic is provided in this previous work. For a more exhaustive background on projection theorems in various settings, we recommend the recent survey article [21] and the references therein.

---

This research was supported by the Swiss National Science Foundation Grant Nr. 00020 165507.

By  $\mathbb{H}^n$  denote the hyperbolic  $n$ -space and by  $d$  the hyperbolic metric on  $\mathbb{H}^n$ . Fix a base point  $p \in \mathbb{H}^n$  and identify the tangent plane  $T_p\mathbb{H}^n$  with  $\mathbb{R}^n$ . Now, consider the exponential mapping  $\exp_p : \mathbb{R}^n \rightarrow \mathbb{H}^n$  at  $p$ . Note that for every  $m$ -plane  $V$  (i.e.  $m$ -dimensional linear subspace of  $\mathbb{R}^n$ ) the image  $\exp_p(V)$  is a geodesically convex  $m$ -dimensional submanifold of  $\mathbb{H}^n$  that is isometric to  $\mathbb{H}^m$ . Since  $\mathbb{H}^n$  is a simply connected Riemannian manifold of constant sectional curvature equal to  $-1$ , for all  $x \in \mathbb{H}^n$ , there exists a unique point  $q \in \exp_p(V)$  such that

$$\text{dist}(x, \exp_p(V)) = d(x, q).$$

Define the projection of  $\mathbb{H}^n$  onto the hyperbolic  $m$ -plane  $\exp_p(V)$  by

$$P_V : \mathbb{H}^n \rightarrow \exp_p(V), \quad P_V(x) = q.$$

As standard arguments show (see Proposition 2.4 in [7]), for all  $x \in \mathbb{H}^n$  and all  $m$ -planes  $V$ , the geodesic segment  $[x, P_V(x)]$  intersects  $\exp_p(V)$  orthogonally in the point  $P_V(x)$ . Therefore, we will refer to the collection of mappings  $P_V : \mathbb{H}^n \rightarrow \exp_p(V)$ , for  $m$ -planes  $V$ , as the family of orthogonal projections (along geodesics) onto  $m$ -planes in  $\mathbb{H}^n$ .

It is known that the projections  $P_V : \mathbb{H}^n \rightarrow \exp_p(V)$  are 1-Lipschitz (i.e. distance non-increasing) with respect to the hyperbolic metric  $d$ , and hence  $\dim P_V(A) \leq \dim A$ , for all sets  $A \subseteq \mathbb{H}^n$  and all  $m$ -planes  $V$ . Moreover, the facts that  $P_V A \subset V$  and  $\dim V = m$  imply that  $\dim P_V A \leq m$  for all  $m$ -planes  $V$ . This yields the same upper bound

$$\dim P_V A \leq \min\{m, \dim A\}$$

as in the Euclidean setting. It is therefore a natural question whether the generic lower bounds for  $\dim P_V A$  is the same as well, i.e. whether Marstrand-type projection theorems generalize to the hyperbolic setting.

We call the family of all  $m$ -planes  $V$  in  $\mathbb{R}^n$  the Grassmannian of  $m$ -planes (in  $\mathbb{R}^n$ ) which we denote by  $G(n, m)$ . The Grassmannian  $G(n, m)$  carries a natural measure  $\sigma_{n,m}$  that is induced by the Haar measure on  $O(n)$  via the group action of  $O(n)$  on  $G(n, m)$ ; see [23, Chapter 3]. Moreover, the Grassmannian can be smoothly parametrized by local charts in  $\mathbb{R}^K$ , where  $K = (n - m)m$ ; see [18, Section 2.3]. This yields a notion of zero sets for the  $s$ -dimensional Hausdorff measure  $\mathcal{H}^s$ ,  $s > 0$ , and of Hausdorff dimension  $\dim$  of subset of  $G(n, m)$ .

The following Marstrand-type theorem is a main result of this paper. It can be considered an analog of results in Euclidean space due to Marstrand [20], Kaufman [19], Falconer [9], Mattila [22], and Peres–Schlag [27].

**Theorem 1.** *For the family of orthogonal projections  $P_V : \mathbb{H}^n \rightarrow \exp_p(V)$ ,  $V \in G(n, m)$ , onto  $m$ -planes in  $\mathbb{H}^n$  and for all Borel sets  $A \subseteq \mathbb{H}^n$ , the following hold.*

- (1) *If  $\dim A \leq m$ , then*
  - (a)  *$\dim(P_V A) = \dim A$  for  $\sigma_{n,m}$ -a.e.  $V \in G(n, m)$ ,*
  - (b) *For  $0 < \alpha \leq \dim A$ ,*  

$$\dim(\{V \in G(n, m) : \dim(P_V A) < \alpha\}) \leq (n - m - 1)m + \alpha.$$

- (2) If  $\dim A > m$ , then
- (a)  $\mathcal{H}^m(P_V A) > 0$  for  $\sigma_{n,m}$ -a.e.  $V \in G(n, m)$ ,
  - (b)  $\dim(\{V \in G(n, m) : \mathcal{H}^m(P_V A) = 0\}) \leq (n - m)m + m - \dim A$ .
- (3) If  $\dim A > 2m$ , then
- (a)  $P_V A$  has non-empty interior in  $V$  for  $\sigma_{n,m}$ -a.e.  $V \in G(n, m)$ ,
  - (b)  $\dim(\{V \in G(n, m) : (P_V A)^\circ \neq \emptyset\}) \leq (n - m)m + 2m - \dim A$ .

We will prove Theorem 1 by a comparison argument. Namely, we will define a self-map of the unit ball that by conjugation transforms hyperbolic orthogonal projections (displayed in the Poincaré model) into Euclidean orthogonal projections; see Section 2. The same arguments will allow us to establish a Besicovitch–Federer-type [6, 13] characterization of purely  $m$ -unrectifiable subsets of  $\mathbb{H}^n$ . Recall that a subset  $A$  of a metric space  $X$  is called  $m$ -rectifiable if there exist at most countably many Lipschitz mappings  $f_i : \mathbb{R}^m \rightarrow X$  such that

$$\mathcal{H}^m\left(A \setminus \bigcup f_i(\mathbb{R}^m)\right) = 0.$$

On the other hand, a set  $F \subseteq X$  is called purely  $m$ -unrectifiable if  $\mathcal{H}^m(F \cap A) = 0$  for every  $m$ -rectifiable set  $A \subseteq X$ .

**Theorem 2.** *A set  $A \subseteq \mathbb{H}^n$  with  $\mathcal{H}^m(A) < \infty$  is purely  $m$ -unrectifiable if and only if for  $\sigma_{n,m}$ -a.e.  $V \in G(n, m)$ , we have  $\mathcal{H}^m(P_V A) = 0$ .*

The Euclidean version of this result is sometimes also referred to as the Besicovitch–Federer projection theorem; see Theorem 18.1 in [23].

**2. Proofs of Theorems 1 and 2.** First, we recall some preliminaries on hyperbolic geometry and fix the notation used in the sequel. For a more detailed account on hyperbolic geometry as it is used here, we recommend the textbooks [5, 7].

Consider the Poincaré model of the hyperbolic  $n$ -space  $\mathbb{H}^n$ , that is, the metric space  $(D^n, d_{\mathcal{P}})$  where  $D^n := \{x \in \mathbb{R}^n : |x| < 1\}$  is the open unit ball in  $\mathbb{R}^n$  and the Poincaré metric  $d_{\mathcal{P}}$  is given by

$$d_{\mathcal{P}}(x, y) = 2 \operatorname{arctanh} \left( \frac{|x - y|}{(1 - 2\langle x, y \rangle + |x|^2 + |y|^2)^{\frac{1}{2}}} \right).$$

for all  $x, y \in D^n$ .

Let  $\Gamma$  be a circle in  $\mathbb{R}^n$  that intersects  $\partial D^n$  orthogonally. Then  $\Gamma \cap D^n$  is a hyperbolic geodesic in the Poincaré model  $(D^n, d_{\mathcal{P}})$ . The same holds for  $L \cap D^n$  for  $L \in G(n, 1)$ . Conversely, every geodesic of hyperbolic space displayed in the Poincaré model is distance minimizing with respect to  $d_{\mathcal{P}}$  and is either of the type  $\Gamma \cap D^n$  or  $L \cap D^n$ . Moreover, the Poincaré model is known to be a conformal model of hyperbolic space, i.e., the angle in which two curves in the hyperbolic  $n$ -space intersect equals the Euclidean angle in which their representatives in  $(D^n, d_{\mathcal{P}})$  intersect. This makes the Poincaré model a natural choice for studying orthogonal projections of hyperbolic  $n$ -space.

Choose  $0$  to be the representative of the base point  $p \in \mathbb{H}^n$  in the model  $(D^n, d_{\mathcal{P}})$ . This choice is made without loss of generality since the hyperbolic




FIGURE 1. The projection  $P_V^P : D^3 \rightarrow D^3 \cap V$

space is homogeneous with respect to its group of isometries. Then, for all  $V \in G(n, m)$ , the hyperbolic  $m$ -plane  $\exp_p(V)$  corresponds to the  $m$ -dimensional disc  $V \cap D^n$  in the model  $(D^n, d_p)$ . For each  $V \in G(n, m)$ , define

$$P_V^P : D^n \rightarrow V \cap D^n$$

to be the closest-point projection onto  $V \cap D^n$  with respect to the metric  $d_p$ ; see Fig. 1. Then, the family of hyperbolic orthogonal projections  $P_V : \mathbb{H}^n \rightarrow \exp_p(V)$ ,  $V \in G(n, m)$ , can be viewed as  $P_V^P : D^n \rightarrow V \cap D^n$ ,  $V \in G(n, m)$ . Moreover, by conformality of the Poincaré model  $(D^n, d_p)$ , the family the projections  $P_V^P : D^n \rightarrow V \cap D^n$  are orthogonal projections along geodesics in  $(D^n, d_p)$ .

Now, consider the mapping  $\Psi : D^n \rightarrow D^n$ , defined by

$$\Psi(x) := \frac{\tanh(2\operatorname{arctanh}|x|)}{|x|}x,$$

for  $x \in D^n \setminus \{0\}$ , and  $\Psi(0) = 0$ . Notice that  $\Psi$  is a bijection with inverse defined by

$$\Psi^{-1}(y) = \frac{\tanh\left(\frac{1}{2}\operatorname{arctanh}|y|\right)}{|y|}y$$

for  $x \in D^n \setminus \{0\}$ , and  $\Psi^{-1}(0) = 0$ . One can check that  $\Psi$  maps every geodesic  $\Gamma \cap D^n$  (where either  $\Gamma \in G(n, 1)$  or  $\Gamma$  is a circle that intersects  $\partial D^n$  orthogonally) to the Euclidean line segment that connects the endpoints  $p_1, p_2$  of  $\Gamma \cap D^n$ ; see Fig. 2.

The metric space  $(D^n, d_\kappa)$  where  $d_\kappa(x, y) := d_p(\Psi^{-1}(x), \Psi^{-1}(y))$ , for all  $x, y \in D^n$ , is often called the Klein model or the projective model of hyperbolic space; see [5] for details. Note that the Klein model is not a conformal model




FIGURE 2. The mapping  $\Psi : D^3 \rightarrow D^3$  where  $\Gamma$  is a geodesic in  $(D^3, d_{\mathcal{P}})$

of hyperbolic space. However, if  $\Gamma_1$  and  $\Gamma_2$  are representatives of hyperbolic geodesics in  $(D^n, d_{\mathcal{K}})$  and if  $0 \in \Gamma_1$ , then the respective geodesics in hyperbolic space intersect orthogonally if and only if  $\Gamma_1$  and  $\Gamma_2$  intersect orthogonally in the Euclidean sense in  $D^n$ .

The symmetry of  $\Psi$  yields the following relation between the hyperbolic orthogonal projections  $P_V^{\mathcal{P}} : D^n \rightarrow V \cap D^n$ ,  $V \in G(n, m)$ , in the Poincaré model and the Euclidean orthogonal projections  $P_V^{\mathbb{E}} : \mathbb{R}^n \rightarrow V$ ,  $V \in G(n, m)$ , restricted to  $D^n$ .

**Lemma 3.** *For all  $V \in G(n, m)$ , the following holds:  $P_V^{\mathcal{P}} = \Psi^{-1} \circ P_V^{\mathbb{E}} \circ \Psi$ .*

*Proof.* Let  $x \in D^n$  and  $V \in G(n, m)$ . By  $\Gamma$  denote the circular arc in  $D^n$  that is perpendicular to  $V$  and  $\partial D^n$  and contains  $x$ . Then, by definition,  $P_V^{\mathcal{P}}(x)$  is the unique intersection point of  $V$  and  $\Gamma$ . Since  $\Gamma$  intersects  $V$  orthogonally, the set  $\Gamma \cap \partial D^n = \{p_1, p_2\}$  is symmetric under the reflection through  $V$ . Thus, the line segment  $\Psi(\Gamma)$  connecting  $p_1$  and  $p_2$  intersects  $V$  orthogonally; see Fig. 2. By definition,  $\Psi(x)$  is the unique intersection point of  $\Gamma$  with the ray that emerges from the origin and goes through  $x$  within  $D^n$ . Then, since  $\Psi(x) \in \Psi(\Gamma)$ , and  $\Psi(\Gamma)$  intersects  $V$  orthogonally,  $P_V^{\mathbb{E}}(\Psi(x))$  is the point where  $\Psi(\Gamma)$  intersects  $V \cap D^n$ . On the other hand,  $\Psi(P_V^{\mathcal{P}}(x))$  is the intersection point of  $\Psi(\Gamma)$  and the ray that emerges from the origin and passes through  $P_V^{\mathcal{P}}(x)$ . However, this intersection point is exactly  $P_V^{\mathbb{E}}(\Psi(x))$ ; see Fig. 2.  $\square$

*Proof of Theorems 1 and 2.* Note that the restriction of the mapping  $\Psi : D^n \rightarrow D^n$  to  $D^n \setminus \{0\}$  is a  $C^\infty$ -diffeomorphism. Moreover, the metric  $d_{\mathcal{P}}$  is locally bi-Lipschitz to the Euclidean metric on  $D^n$ . Hence, for every set  $A$ , every  $m$ -plane  $V \in G(n, m)$  and every  $s > 0$ ,  $P_V^{\mathbb{E}}(A)$  is an  $\mathcal{H}^s$ -zero

set if and only if  $P_V^{\mathcal{P}}(A)$  is an  $\mathcal{H}^s$ -zero set. In particular, it follows that  $\dim P^{\mathbb{E}}(A) = \dim P_V^{\mathcal{P}}(A)$ . Moreover,  $P_V^{\mathbb{E}}(A)$  has non-empty interior in  $V$  if and only if  $P_V^{\mathcal{P}}(A)$  has non-empty interior in  $V$ . Hence, Theorems 1 and 2 follow from their well-known analogs for orthogonal projection onto  $m$ -planes in  $\mathbb{R}^n$ .  $\square$

**3. Remark on transversality and projection theorems.** In [27] Peres and Schlag establish a very general projection theorem for families of (abstract) projections from compact metric spaces to the Euclidean space. Namely, their result states that if a sufficiently regular family of projections satisfies a certain transversality condition, then this yields bounds for the Sobolev dimension of the push-forward (by the projections) of certain measures. While Peres and Schlag's main applications of this result concern Bernoulli convolutions, all the classical Marstrand-type projection theorems for Euclidean spaces  $\mathbb{R}^n$  can be deduced as corollaries from their result; see Section 6 in [27] and Section 18.3 in [24]. Moreover, Hovila et. al. [16] have proven that if a family of abstract projections satisfies transversality with sufficiently good transversality constants, then a Besicovitch–Federer type characterization of purely unrectifiable sets in terms of this family of projections follows. Therefore, transversality has proven to be a powerful method in establishing Marstrand-type as well as Besicovitch–Federer type projection theorems in various settings. In particular, the works [15] (Heisenberg groups) and [4] (Riemannian surfaces of constant curvature) are based on Peres and Schlag's notion of transversality.

In fact, it is possible to establish transversality for the family of orthogonal projections in the Poincaré model,  $P_V^{\mathcal{P}} : D^n \rightarrow \exp_p(V) \cap D^n$ ,  $V \in G(n, m)$ . This is worked out in detail in the second author's PhD thesis [18, Section 6.2]. The transversality constants obtained (namely  $L = 2$  and  $\delta = 0$  in the notation of [18]) are sufficient to imply both Marstrand-type as well as Besicovitch–Federer-type projection theorems. In particular, Theorem 2 can be deduced as a corollary from this result. However, the upper bounds for the dimension of the exceptional set of planes for Marstrand-type projection theorems in general depend on the transversality constants; see Theorem 7.3 in [27]. In particular, in the cases where  $\dim A > m$ , the bounds obtained by establishing transversality are worse than the bounds in Conclusions (2.b) and (3.b) of Theorem 1. The transversality constants  $L = 2$  and  $\delta = 0$  obtained in [18] could still be improved. In particular, a lengthy but straightforward calculation shows that  $L$  can be improved to 3. However, in order to obtain Theorem 1 as a consequence of transversality, one would need  $L = \infty$ . However, this is not possible due to insufficient regularity of the mapping  $\Psi : D^n \rightarrow D^n$  in the origin.

**Acknowledgements.** The authors would like to thank the referee and the editor for their careful and efficient handling of our paper.

## References

- [1] Balogh, Z.M., Durand-Cartagena, E., Fässler, K., Mattila, P., Tyson, J.T.: The effect of projections on dimension in the Heisenberg group. *Rev. Mat. Iberoam.* **29**(2), 381–432 (2013)
- [2] Balogh, Z.M., Fässler, K., Mattila, P., Tyson, J.T.: Projection and slicing theorems in Heisenberg groups. *Adv. Math.* **231**(2), 569–604 (2012)
- [3] Balogh, Z.M., Iseli, A.: Marstrand type projection theorems for normed spaces. To appear in *J. Fractal Geom.* (2018). [arXiv:1802.10563](https://arxiv.org/abs/1802.10563)
- [4] Balogh, Z.M., Iseli, A.: Dimensions of projections of sets on Riemannian surfaces of constant curvature. *Proc. Am. Math. Soc.* **144**(7), 2939–2951 (2016)
- [5] Benedetti, R., Petronio, C.: *Lectures on Hyperbolic Geometry*. Universitext. Springer, Berlin (1992)
- [6] Besicovitch, A.S.: On the fundamental geometrical properties of linearly measurable plane sets of points (III). *Math. Ann.* **116**(1), 349–357 (1939)
- [7] Bridson, M.R., Haefliger, A.: *Metric spaces of non-positive curvature*. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 319. Springer, Berlin (1999)
- [8] Chen, C.: Restricted families of projections and random subspaces. (2017). [arXiv:1706.03456](https://arxiv.org/abs/1706.03456)
- [9] Falconer, K.J.: Hausdorff dimension and the exceptional set of projections. *Mathematika* **29**(1), 109–115 (1982)
- [10] Falconer, K.J., Howroyd, J.D.: Projection theorems for box and packing dimensions. *Math. Proc. Cambridge Philos. Soc.* **119**(2), 287–295 (1996)
- [11] Falconer, K.J., Howroyd, J.D.: Packing dimensions of projections and dimension profiles. *Math. Proc. Cambridge Philos. Soc.* **121**(2), 269–286 (1997)
- [12] Fässler, K., Orponen, T.: On restricted families of projections in  $\mathbb{R}^3$ . *Proc. Lond. Math. Soc.* (3) **109**(2), 353–381 (2014)
- [13] Federer, H.: The  $(\varphi, k)$  rectifiable subsets of  $n$ -space. *Trans. Am. Math. Soc.* **62**, 114–192 (1947)
- [14] Fraser, J.M., Orponen, T.: The Assouad dimensions of projections of planar sets. *Proc. Lond. Math. Soc.* (3) **114**(2), 374–398 (2017)
- [15] Hovila, R.: Transversality of isotropic projections, unrectifiability, and Heisenberg groups. *Rev. Mat. Iberoam.* **30**(2), 463–476 (2014)
- [16] Hovila, R., Järvenpää, E., Järvenpää, M., Ledrappier, F.: Besicovitch-Federer projection theorem and geodesic flows on Riemann surfaces. *Geom. Dedic.* **161**, 51–61 (2012)
- [17] Iseli, A.: Marstrand-type projection theorems for linear projections and in normed spaces. (2018). [arXiv:1809.00636](https://arxiv.org/abs/1809.00636)
- [18] Iseli, A.: Dimension and projections in normed spaces and Riemannian manifolds. PhD thesis, Universität Bern, Switzerland, 2018. [http://biblio.unibe.ch/download/eldiss/18iseli\\_a.pdf](http://biblio.unibe.ch/download/eldiss/18iseli_a.pdf)

- [19] Kaufman, R.: On Hausdorff dimension of projections. *Mathematika* **15**, 153–155 (1968)
- [20] Marstrand, J.M.: Some fundamental geometrical properties of plane sets of fractional dimensions. *Proc. London Math. Soc.* **3**(4), 257–302 (1954)
- [21] Mattila, P.: Hausdorff dimension, projections, intersections, and Besicovitch sets. (2017). [arXiv:1712.09199](https://arxiv.org/abs/1712.09199)
- [22] Mattila, P.: Hausdorff dimension, orthogonal projections and intersections with planes. *Ann. Acad. Sci. Fenn. Ser. A I Math.* **1**(2), 227–244 (1975)
- [23] Mattila, P.: *Geometry of Sets and Measures in Euclidean Spaces. Fractals and Rectifiability* Cambridge Studies in Advanced Mathematics, vol 44. Cambridge University Press, Cambridge (1995)
- [24] Mattila, P.: *Fourier Analysis and Hausdorff Dimension.* Cambridge Studies in Advanced Mathematics, vol. 150. Cambridge University Press, Cambridge (2015)
- [25] Orponen, T.: On the dimension and smoothness of radial projections. (2017). [arXiv:1710.11053](https://arxiv.org/abs/1710.11053)
- [26] Orponen, T., Venieri, L.: Improved bounds for restricted families of projections to planes in  $\mathbb{R}^3$ . *Int. Math. Res. Not.* (2018). <https://doi.org/10.1093/imrn/rny193>
- [27] Peres, Y., Schlag, W.: Smoothness of projections, Bernoulli convolutions, and the dimension of exceptions. *Duke Math. J.* **102**(2), 193–251 (2000)

ZOLTÁN M. BALOGH AND ANNINA ISELI  
Mathematisches Institut,  
Universität Bern,  
Sidlerstrasse 5,  
3012 Bern,  
Switzerland  
e-mail: [zoltan.balogh@math.unibe.ch](mailto:zoltan.balogh@math.unibe.ch)

ANNINA ISELI  
e-mail: [annina.iseli@math.unibe.ch](mailto:annina.iseli@math.unibe.ch)

Received: 6 August 2018