



Correction to: The impact of different doses of indocyanine green on the sentinel lymph-node mapping in early stage endometrial cancer

Andrea Papadia¹ · Alessandro Buda² · Maria Luisa Gasparri¹ · Giampaolo Di Martino² · Beatrice Bussi² · Debora Verri² · Michael D. Mueller¹

Published online: 8 September 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to:
Journal of Cancer Research and Clinical Oncology
<https://doi.org/10.1007/s00432-018-2716-3>

correct value is 0.084. The corresponding OR value is also corrected in the below table (Table 3).

Unfortunately, the *P* value at multivariate analysis for ICG concentration in Table 3 was incorrectly published. The

Table 3 Univariate and Multivariate analysis factors associated with Bilateral Detection rate (All patients, *n* = 168)

	Bilateral detection (yes/no)			
	Univariate analysis		Multivariate analysis	
	OR (95% CI)	<i>p</i>	OR (95% CI)	<i>p</i>
Age (< 64 vs ≥ 64)	1.37 (0.54, 3.45)	0.49		
BMI (≥ 35 vs < 35)	0.44 (0.15, 1.29)	0.13	0.34 (0.11, 1.07)	0.066
Tumor volume (≥ 2 vs < 2 cm)	1.34 (0.54, 3.33)	0.52		
LVSI (positive vs negative)	0.73 (0.15, 3.45)	0.69		
ICG concentration (5 vs 1.25 mg/ml)	2.2 (0.88, 5.52)	0.09	2.64 (0.81, 6.92)	0.084

Multivariable models were carried out for variables reporting a *p* value ≤ 0.25 in the univariate analysis

The original article can be found online at <https://doi.org/10.1007/s00432-018-2716-3>.

✉ Andrea Papadia
andrea.papadia@insel.ch

¹ Department of Obstetrics and Gynecology, University Hospital of Bern, University of Bern, Effingerstrasse 102, 3010 Bern, Switzerland

² Unit of Gynecologic Oncology Surgery, Department of Obstetrics and Gynecology, San Gerardo Hospital, University of Milano-Bicocca, Monza, Italy