



Analytical Models of Exoplanetary Atmospheres. VI. Full Solutions for Improved Two-stream Radiative Transfer, Including Direct Stellar Beam

Kevin Heng , Matej Malik , and Daniel Kitzmann

University of Bern, Center for Space and Habitability, Gesellschaftsstrasse 6, CH-3012, Bern, Switzerland; kevin.heng@csh.unibe.ch*Received 2018 April 13; revised 2018 June 22; accepted 2018 July 3; published 2018 August 2*

Abstract

Two-stream radiative transfer is used frequently in Earth, planetary, and exoplanetary sciences due to its simplicity and ease of implementation. However, a longstanding limitation of the two-stream approximation is its inaccuracy in the presence of medium-sized or large aerosols. This limitation was lifted by the discovery of the improved two-stream technique, where the accuracy of the scattering greenhouse effect is matched to that of multi-stream calculations by construction. In this study, we derive the full solutions for improved two-stream radiative transfer, following its introduction by Heng & Kitzmann, and include contributions from the direct stellar beam. The generalization of the original two-stream flux solutions comes in the form of a correction factor, traditionally set to unity, which is the ratio of a pair of first Eddington coefficients. We derive an analytical expression for this correction factor and also provide a simple fitting function for its ease of use by other workers. We prove that the direct stellar beam is associated with a second Eddington coefficient that is on the order of unity. Setting this second Eddington coefficient to $2/3$ and $1/\sqrt{3}$ reproduces the Eddington and quadrature closures, respectively, associated with the direct beam. We use our improved two-stream solutions for the fluxes to derive two-stream source function solutions for both the intensity and fluxes.

Key words: methods: analytical – planets and satellites: atmospheres

1. Introduction

A workhorse of radiative transfer is the two-stream method, which approximates the passage of radiation through an atmosphere as a pair of incoming and outgoing fluxes (Schuster 1905). Instead of solving a partial differential equation for the angle-dependent intensity, one solves an ordinary differential equation for the angle-integrated flux. The price to pay is mathematical underdetermination. To mathematically close the system, one has to assume that ratios of moments of the intensity are constants known as Eddington coefficients.

There is a rich literature on two-stream radiative transfer. Joseph et al. (1976) and Wiscombe (1977) introduced ad hoc modifications to the scattering phase function to treat forward-peaked scattering of radiation due to large aerosols, known as the “delta-Eddington approximation.” Meador & Weaver (1980) presented a unified theoretical framework that described different ways of specifying two-stream closures. Toon et al. (1989) presented a specific formulation of the two-stream method that has been influential in the planetary and exoplanetary sciences (e.g., Marley & McKay 1999), partly because it allows for fast computation via the inversion of a tridiagonal matrix. Toon et al. (1977, 1989) introduced the “two-stream source function” method, which reduces the radiative transfer equation from an integro-differential equation to a differential equation for the intensity by utilizing a sleight: the two-stream solution is inserted into the integral involving the scattering phase function and intensity, by assuming that the intensity is related to the two-stream flux by a constant factor. Heng et al. (2014) introduced a

formalism that unified the two-stream solutions with analytical solutions for temperature-pressure profiles, based partly on generalizing the relevant parts of the monograph of Pierrehumbert (2010).

Kitzmann et al. (2013) realized that the original two-stream method performs poorly in the presence of medium-sized or large aerosols, as it underestimates the backscattered radiation by $\sim 10\%$. Kitzmann (2016) demonstrated that this shortcoming of the two-stream method translates into an overestimation of the scattering greenhouse effect for early Mars by about 50 K. Motivated by the work of Kitzmann et al. (2013) and Kitzmann (2016), Heng & Kitzmann (2017) discovered a simple improvement to the two-stream method that allows it to match the accuracy of 32-stream calculations (at the $\sim 0.01\%$ – 1% level or better, depending on the optical depth) in the presence of medium-sized or large aerosols.

The overarching intention of this study is to fully flesh out the improved two-stream method introduced by Heng & Kitzmann (2017) and also use it as input for the two-stream source function method of Toon et al. (1989). The entirety of the paper is devoted to these novel derivations. At the heart of the improved two-stream flux solutions is a correction factor E , shown in Figure 1, which is usually set to unity in the original two-stream method. We provide a convenient fitting function in Equation (31) for the reader to compute E .

2. Basic Theory

The radiative transfer equation may generally be written as (Chandrasekhar 1960; Mihalas 1970, 1978; Meador & Weaver 1980; Toon et al. 1989)

$$\mu \frac{\partial I}{\partial \tau} = I - (1 - \omega_0)B - \omega_0 \int_0^{4\pi} \mathcal{P}(I + I_{\text{beam}}) d\Omega', \quad (1)$$

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.




Figure 1. Ratio of first Eddington coefficients (E) as a function of the single-scattering albedo (ω_0) and scattering asymmetry factor (g_0), which is used to improve the accuracy of two-stream radiative transfer in the presence of medium-sized or large aerosols. The solid, dashed, and dotted-dashed curves correspond to $E = 1.05, 1.1$, and 1.2 , respectively. In the original two-stream formulation, one has $E = 1$. Pure absorption and scattering correspond to $\omega_0 = 0$ and 1 , respectively. Small and large aerosols are represented by $g_0 = 0$ and $g_0 \sim 1$, respectively. Equation (31) provides a fitting function for E that is accurate at the $\sim 0.1\%$ level.

where I is the wavelength-, frequency-, or wavenumber-dependent intensity¹ (which is why we have omitted subscripts on all quantities that commit us to any of these choices, including for the Planck function), τ is the optical depth, ω_0 is the single-scattering albedo, and B is the Planck function. The scattering phase function, $\mathcal{P}(\mu', \phi', \mu, \phi)$, relates the incident polar (θ') and azimuthal (ϕ') angles to the emergent polar (θ) and azimuthal (ϕ) angles. We further define $\mu \equiv \cos \theta$, $\mu' \equiv \cos \theta'$, $d\Omega \equiv d\mu d\phi$, and $d\Omega' \equiv d\mu' d\phi'$. Physically, one has to account for the scattered intensity coming from all incident directions.

In addition to the diffuse radiation field, we assume the presence of a direct beam of radiation due to the star, which impinges on the atmosphere with $\mu' = -\mu_*$ and $\phi' = \phi_*$. Some portion of this incident stellar beam is absorbed, while the rest is scattered into the diffuse radiation field, which is represented by

$$I_{\text{beam}} = F_* \exp\left(\frac{\tau}{\mu'}\right) \delta(\mu' + \mu_*) \delta(\phi' - \phi_*), \quad (2)$$

where F_* is the incident stellar flux at the top of the atmosphere, and the pair of delta functions enforces the directionality of the beam.

¹ The intensity is sometimes referred to as “radiance” by Earth atmospheric scientists.

The radiative transfer equation is multiplied by some arbitrary function $\mathcal{H}(\mu)$ and integrated over all emergent angles (Pierrehumbert 2010; Heng et al. 2014; Heng 2017),

$$\begin{aligned} \frac{\partial}{\partial \tau} \int_0^{2\pi} \left(\int_0^1 \mu \mathcal{H} I d\mu - \int_{-1}^0 \mu \mathcal{H} I d\mu \right) d\phi \\ = \int_0^{4\pi} \mathcal{H} I d\Omega - (1 - \omega_0) \int_0^{4\pi} \mathcal{H} B d\Omega - \mathcal{I}, \end{aligned} \quad (3)$$

where we have

$$\begin{aligned} \mathcal{I} &\equiv \omega_0 \int_0^{4\pi} (I + I_{\text{beam}}) \mathcal{G} d\Omega', \\ \mathcal{G} &\equiv \int_0^{4\pi} \mathcal{H} \mathcal{P} d\Omega. \end{aligned} \quad (4)$$

Equation (3) is the starting point for the suite of derivations we will perform in the current study. The minus sign associated with the gradient terms comes about because we have rescaled the optical depth to be always positive, as has been previously elucidated in Heng et al. (2014) and Chapter 3.5.1 of Heng (2017).

Mathematically, the contribution of the direct beam to \mathcal{I} may be evaluated in two ways: either by evaluating the delta functions first (over $d\Omega'$) or by evaluating \mathcal{G} first. Since these mathematical operations commute, the resulting expressions must agree. The former approach corresponds to the derivation for isotropic scattering, while the latter is for non-isotropic (or anisotropic) scattering.

3. Improved Two-stream with Direct Beam and Isotropic Scattering

In the isotropic limit, Equation (1) becomes

$$\mu \frac{\partial I}{\partial \tau} = I - (1 - \omega_0) B - \frac{\omega_0 J}{4\pi} - \omega_0 \mathcal{P}_* F_* \exp\left(-\frac{\tau}{\mu_*}\right), \quad (5)$$

where we define

$$\mathcal{P}_* \equiv \mathcal{P}(-\mu_*, \phi_*, \mu, \phi). \quad (6)$$

The total intensity is $J \equiv \int_0^{4\pi} I(\mu', \tau) d\Omega'$ and it is related to the total flux by $\epsilon = F_+/J$. The ratio of the flux to the total intensity in one hemisphere only is denoted by ϵ' , where we have assumed that this ratio is the same in both hemispheres.

By integrating over the incoming (\downarrow) and outgoing (\uparrow) hemispheres, we obtain the incoming and outgoing fluxes,²

$$\begin{aligned} \frac{\partial F_{\uparrow}}{\partial \tau} &= \gamma_a F_{\uparrow} - \gamma_s F_{\downarrow} - \gamma_B B - \omega_0 F_* \chi_{\uparrow} \exp\left(-\frac{\tau}{\mu_*}\right), \\ \frac{\partial F_{\downarrow}}{\partial \tau} &= -\gamma_a F_{\downarrow} + \gamma_s F_{\uparrow} + \gamma_B B + \omega_0 F_* \chi_{\downarrow} \exp\left(-\frac{\tau}{\mu_*}\right). \end{aligned} \quad (7)$$

² The flux is sometimes referred to as “irradiance” by Earth atmospheric scientists.

The coefficients of the various terms are given by

$$\begin{aligned}\gamma_a &\equiv \frac{1}{\epsilon'} - \frac{\omega_0}{2\epsilon} = 2E - \omega_0, \\ \gamma_s &\equiv \frac{\omega_0}{2\epsilon} = \omega_0, \\ \gamma_B &\equiv 2\pi(1 - \omega_0).\end{aligned}\quad (8)$$

The derivation technique was established in Heng et al. (2014) and also Chapter 3 of Heng (2017), and we will not repeat it here. Instead, we will point out key, novel aspects of the current derivation that involve both the improvement to the two-stream method and the inclusion of the direct beam.

Previously, Heng (2017) designated both ϵ and ϵ' as the ‘‘first Eddington coefficients.’’ By demanding that the limiting case of an opaque, purely absorbing ($\omega_0 = 0$) atmosphere emits πB of flux in each hemisphere, one obtains $\epsilon = 1/2$ (Toon et al. 1989; Pierrehumbert 2010). By demanding that a purely scattering ($\omega_0 = 1$) atmosphere remains in radiative equilibrium, one sets $\epsilon' = \epsilon$ (Toon et al. 1989). However, since the two-stream solutions are formally invalid in the limit of $\omega_0 = 1$, the second demand is academic, as has already been realized by Heng & Kitzmann (2017). Following Heng & Kitzmann (2017), we set

$$E \equiv \frac{\epsilon}{\epsilon'}, \quad (9)$$

and proceed to obtain an analytical expression for E in terms of ω_0 and the scattering asymmetry factor (g_0).

We have written

$$\begin{aligned}\chi_{\uparrow} &\equiv \int_0^{2\pi} \int_0^1 \mathcal{P}_* d\mu d\phi, \\ \chi_{\downarrow} &\equiv \int_0^{2\pi} \int_{-1}^0 \mathcal{P}_* d\mu d\phi.\end{aligned}\quad (10)$$

Since the integration involves only μ and ϕ , we expect that \mathcal{P} has the same mathematical behavior as \mathcal{P}_* with respect to this procedure. Physically, for isotropic scattering we expect $\chi_{\uparrow} = \chi_{\downarrow} = 1/2$. We keep these quantities distinct, because we will explicitly prove in Section 4 that $\chi_{\uparrow} = \chi_{\downarrow}$ is a natural outcome of setting $g_0 = 0$ for the more general expressions involving the gradients of the fluxes. When scattering is non-isotropic, we generally expect that $\chi_{\uparrow} \neq \chi_{\downarrow}$, but because the scattering phase function needs to be properly normalized, we have

$$\chi_{\uparrow} + \chi_{\downarrow} = 1. \quad (11)$$

In both Meador & Weaver (1980) and Toon et al. (1989), these authors have written $\gamma_3 = \chi_{\uparrow}$, $\gamma_4 = \chi_{\downarrow}$ and $\gamma_3 + \gamma_4 = 1$.

The equations governing the net ($F_- \equiv F_{\uparrow} - F_{\downarrow}$) and total ($F_+ \equiv F_{\uparrow} + F_{\downarrow}$) fluxes are

$$\begin{aligned}\frac{\partial F_+}{\partial \tau} &= (\gamma_a + \gamma_s)F_- + \omega_0 F_* (\chi_{\downarrow} - \chi_{\uparrow}) \exp\left(-\frac{\tau}{\mu_*}\right), \\ \frac{\partial F_-}{\partial \tau} &= (\gamma_a - \gamma_s)F_+ - 2\gamma_B B - \omega_0 F_* \exp\left(-\frac{\tau}{\mu_*}\right).\end{aligned}\quad (12)$$

We will defer seeking the solutions to F_+ and F_- , and hence F_{\uparrow} and F_{\downarrow} , until we deal with the non-isotropic treatment, because the mathematical forms of these governing equations are almost identical.

4. Improved Two-stream with Direct Beam and Anisotropic Scattering

4.1. Governing Equation for Net Flux

For non-isotropic scattering, we return to using Equation (3) as our starting point. If we set $\mathcal{H} = 1$, then we obtain $\mathcal{G} = 1$ (Pierrehumbert 2010; Heng et al. 2014; Heng 2017). The equation governing the net flux becomes

$$\frac{\partial F_-}{\partial \tau} = (1 - \omega_0)(J - 4\pi B) - \omega_0 F_* \exp\left(-\frac{\tau}{\mu_*}\right). \quad (13)$$

There is an unsatisfactory ambiguity if $\epsilon \neq \epsilon'$, because one could convert J to the total flux using $J = F_+/\epsilon$ or $J = J_{\uparrow} + J_{\downarrow} = F_+/\epsilon'$. We resolve this ambiguity by demanding that the equation reduces to its counterpart in the isotropic limit, which yields

$$\frac{\partial F_-}{\partial \tau} = 2F_+(E - \omega_0) - 4\pi B(1 - \omega_0) - \omega_0 F_* \exp\left(-\frac{\tau}{\mu_*}\right). \quad (14)$$

4.2. Governing Equation for Total Flux

To obtain the equation governing the total flux, we set $\mathcal{H} = \mu$, which yields $\mathcal{G} = g_0 \mu'$ (Pierrehumbert 2010; Heng et al. 2014; Heng 2017), from which it follows that

$$\frac{\partial K_-}{\partial \tau} = F_+(1 - \omega_0 g_0) + \omega_0 g_0 \mu_* F_* \exp\left(-\frac{\tau}{\mu_*}\right), \quad (15)$$

where $K_- \equiv K_{\uparrow} - K_{\downarrow}$ and K_{\uparrow} and K_{\downarrow} are the second moments of the intensity. Following Heng et al. (2014) and Heng (2017), we define a second Eddington coefficient,

$$\epsilon_2 \equiv \frac{K_-}{F_+}. \quad (16)$$

For the diffuse emission (which is usually dominant in the infrared range of wavelengths), we demand that $\epsilon_2 = F_+/2EF_-$, in order for the governing equation for the total flux to reduce to its counterpart in the limit of isotropic scattering. The governing equation becomes

$$\frac{\partial F_+}{\partial \tau} = 2EF_-(1 - \omega_0 g_0) + \gamma_* \exp\left(-\frac{\tau}{\mu_*}\right), \quad (17)$$

where we have defined

$$\gamma_* \equiv \frac{\omega_0 g_0 \mu_* F_*}{\epsilon_2}. \quad (18)$$

4.3. Coefficients in Governing Equations

For the direct beam (which is usually dominant in the optical/visible range of wavelengths), we intuitively expect that $\epsilon_2 \sim 1$. We will now demonstrate this claim.

By comparing Equations (14) and (17) with the pair of equations in (12), we may infer that

$$\begin{aligned}\gamma_a - \gamma_s &= 2(E - \omega_0), \\ \gamma_a + \gamma_s &= 2E(1 - \omega_0 g_0),\end{aligned}\quad (19)$$

which yields

$$\begin{aligned}\gamma_a &= 2E - \omega_0(1 + E g_0), \\ \gamma_s &= \omega_0(1 - E g_0).\end{aligned}\quad (20)$$

In the limit of $g_0 = 0$ (isotropic scattering), the beam term in Equation (17) vanishes. By comparing to the first equation in (12), we obtain $\chi_l = \chi_f$.

Another useful comparison is between Equations (14) and (17) and the pair of equations in (7), which yields

$$\begin{aligned}\chi_f &= \frac{1}{2} \left(1 - \frac{\mu_* g_0}{\epsilon_2} \right), \\ \chi_l &= \frac{1}{2} \left(1 + \frac{\mu_* g_0}{\epsilon_2} \right).\end{aligned}\quad (21)$$

When we compare χ_f with the γ_3 values listed in Table 1 of both Meador & Weaver (1980) and Toon et al. (1989), we find that $\epsilon_2 = 2/3$ reproduces the Eddington closure, while $\epsilon_2 = 1/\sqrt{3} \approx 0.58$ reproduces the quadrature closure. Regardless of the choice of value for the second Eddington coefficient, we always have $\chi_f + \chi_l = 1$, which ensures that the scattering phase function is normalized properly.

The physical interpretation of the second Eddington coefficient is that, in the limit of $\epsilon_2/\mu_* = 1$, an extra fraction $g_0/2$ of the stellar direct beam gets scattered into the forward/incoming direction, such that when $g_0 = 1$, all of the stellar beam is deposited downward. Generally, this fraction is $\mu_* g_0/2\epsilon_2$, which allows for backscattered stellar radiation, even when $g_0 = 1$, by setting $\mu_*/\epsilon_2 \neq 1$.

4.4. Solutions for the Two-stream Fluxes

The recipe to solve for the fluxes has previously been established by both Heng et al. (2014) and Chapter 3 of Heng (2017). We will only highlight the key points and novel generalizations. One first merges Equations (14) and (17) into a single second-order ordinary differential equation for F_+ . The solution is

$$\begin{aligned}F_+ &= A_1 \exp(\alpha\tau) + A_2 \exp(-\alpha\tau) \\ &+ 2\pi \left(\frac{1 - \omega_0}{E - \omega_0} \right) [B_i + B'(\tau - \tau_i)] \\ &+ C_* \exp\left(-\frac{\tau}{\mu_*}\right),\end{aligned}\quad (22)$$

where we have $\alpha \equiv \sqrt{(\gamma_a + \gamma_s)(\gamma_a - \gamma_s)}$ and the coefficients A_1 and A_2 are determined by applying boundary conditions to the solution. A linear expansion in the Planck function is performed to allow for the two-stream solutions to treat non-isothermal atmospheric layers described by a gradient $B' \equiv \partial B/\partial\tau$, which has been shown to significantly increase computational efficiency (Malik et al. 2017). The index $i = 1, 2$ will be explained shortly. Note the correction factor in front of the Planck function associated with $E \neq 1$. The coefficient C_* is derived using the method of undetermined coefficients,

$$C_* = \frac{\omega_0 F_* [2E(1 - \omega_0 g_0) + g_0/\epsilon_2]}{4E(E - \omega_0)(1 - \omega_0 g_0) - 1/\mu_*^2}.\quad (23)$$

From the expressions for F_f and F_l , one realizes that a more compact way of expressing the algebra is to define the coupling

coefficients,

$$\zeta_{\pm} \equiv \frac{1}{2} \left[1 \pm \sqrt{\frac{E - \omega_0}{E(1 - \omega_0 g_0)}} \right].\quad (24)$$

These are termed coupling coefficients, because they couple the outgoing and incoming fluxes in the presence of scattering (see Section 3.2 of Heng 2017). As already described in Toon et al. (1989), C_* becomes indeterminate when the denominator in Equation (23) becomes zero, which occurs when $\mu_* = 1/2\sqrt{E(E - \omega_0)(1 - \omega_0 g_0)}$. Toon et al. (1989) suggest that, ‘‘In practice, if the equality happens to occur, this problem can be eliminated by simply choosing a slightly different value of μ_* .’’

The final two-stream solutions for the incoming and outgoing fluxes are expressed in the convention of a pair of atmospheric layers, subscripted by ‘‘1’’ and ‘‘2’’ with the former residing above the latter. In this convention, F_{l2} and F_{l1} are the boundary conditions, which are used to eliminate A_1 and A_2 . This procedure is documented in Section 3.8.2 of Heng (2017). When the dust has settled on the algebra, the two-stream solutions are

$$\begin{aligned}F_{l1} &= \frac{1}{(\zeta_- \mathcal{T})^2 - \zeta_+^2} \{ (\zeta_-^2 - \zeta_+^2) \mathcal{T} F_{l2} - \zeta_- \zeta_+ (1 - \mathcal{T}^2) F_{l1} \\ &+ \Pi [\mathcal{B}_{1+} (\zeta_-^2 \mathcal{T}^2 - \zeta_+^2) + \mathcal{B}_{2+} \mathcal{T} (\zeta_+^2 - \zeta_-^2) \\ &+ \mathcal{B}_{1-} \zeta_- \zeta_+ (1 - \mathcal{T}^2)] + C_* \mathcal{T} \mathcal{T}_{\text{above}} (\zeta_+^2 - \zeta_-^2) \\ &+ \mathcal{T}_{\text{above}} [\zeta_+ (C_+ \zeta_- - C_- \zeta_+) + \zeta_- \mathcal{T}^2 (C_+ \zeta_- - C_- \zeta_+)] \},\end{aligned}\quad (25)$$

and

$$\begin{aligned}F_{l2} &= \frac{1}{(\zeta_- \mathcal{T})^2 - \zeta_+^2} \{ (\zeta_-^2 - \zeta_+^2) \mathcal{T} F_{l1} - \zeta_- \zeta_+ (1 - \mathcal{T}^2) F_{l2} \\ &+ \Pi [\mathcal{B}_{2-} (\zeta_-^2 \mathcal{T}^2 - \zeta_+^2) + \mathcal{B}_{1-} \mathcal{T} (\zeta_+^2 - \zeta_-^2) \\ &+ \mathcal{B}_{2+} \zeta_- \zeta_+ (1 - \mathcal{T}^2)] + C_* \mathcal{T} \mathcal{T}_{\text{above}} (\zeta_+^2 - \zeta_-^2) \\ &\times \mathcal{T}_* \mathcal{T}_{\text{above}} [\zeta_+ (C_- \zeta_- - C_+ \zeta_+) + \zeta_- \mathcal{T}^2 (C_+ \zeta_- - C_- \zeta_+)] \},\end{aligned}\quad (26)$$

where, for compactness of notation, we have defined

$$\begin{aligned}\Pi &\equiv \pi \left(\frac{1 - \omega_0}{E - \omega_0} \right), \\ C_{\pm} &\equiv \frac{1}{2} \left[C_* \pm \frac{(C_*/\mu_* + \gamma_*)}{2E(1 - \omega_0 g_0)} \right], \\ \Delta &\equiv \tau_2 - \tau_1, \\ \mathcal{T} &\equiv \exp(-\alpha\Delta), \\ \mathcal{T}_* &\equiv \exp\left(-\frac{\Delta}{\mu_*}\right), \\ \mathcal{T}_{\text{above}} &\equiv \exp\left(-\frac{\tau_1}{\mu_*}\right), \\ \mathcal{B}_i &\equiv B_i \pm \frac{B'}{2E(1 - \omega_0 g_0)}.\end{aligned}\quad (27)$$

The optical depth of the atmospheric layer is given by Δ . The transmissivity of the layer is represented by \mathcal{T} . Extinction of

the direct stellar beam above and through the layer are given by $\mathcal{T}_{\text{above}}$ and \mathcal{T}_* , respectively. The gradient of the Planck function is $B' = (B_2 - B_1)/\Delta$.

4.5. A Useful Limit: $g_0 = E = 1$

It is instructive to examine the limit of $g_0 = 1$ and $E = 1$, because in the two-stream approximation it behaves as if one is in the limit of pure absorption. In this limit, the coupling coefficients become $\zeta_- = 0$ and $\zeta_+ = 1$, implying that several of the direct beam terms in Equations (25) and (26) vanish. For the purpose of developing physical intuition, we further let $\mu_* = \epsilon_2$, which yields $\gamma_* = \omega_0 F_*$ and

$$C_* = \frac{\omega_0 F_*}{2(1 - \omega_0) - 1/\epsilon_2}. \quad (28)$$

Furthermore, we get $C_- = 0$ and $C_+ = C_*$, which implies that the beam terms in the outgoing flux solution (F_{\uparrow}) vanish. The only non-vanishing beam terms in the incoming flux solution (F_{\downarrow}) are $C_* \mathcal{T}_{\text{above}}(\mathcal{T}_* - \mathcal{T})$. One may verify that regardless of whether $2(1 - \omega_0) > 1/\epsilon_2$ or $2(1 - \omega_0) < 1/\epsilon_2$, $C_* \mathcal{T}_{\text{above}}(\mathcal{T}_* - \mathcal{T})$ is always positive.

5. The E -factor

The ratio of the first Eddington coefficients, E , lies at the heart of our improved two-stream method. In the optically thick limit, the fraction of flux reflected by an atmospheric layer is ζ_-/ζ_+ , a fact that may be verified by setting $\mathcal{T} = 0$ in Equations (25) and (26) and ignoring the blackbody and direct beam terms. The expression for $R_\infty = \zeta_-/\zeta_+$ may be manipulated to obtain an analytical expression for E . In other words, E is *constructed* to reproduce the reflectivity in the optically thick limit (R_∞), as long as one has a way to numerically evaluate R_∞ .

Previously, Heng & Kitzmann (2017) performed a visual inspection of $\gamma_a = 2 - \omega_0(1 + g_0)$ and reasoned that it generalized to $\gamma_a = 2E - \omega_0(1 + g_0)$, while leaving γ_s unchanged. This led to $\zeta_\pm = \frac{1}{2}[1 \pm \sqrt{(E - \omega_0)/(E - \omega_0 g_0)}]$ and $E = \omega_0(1 - g_0 r^2)/(1 - r^2)$. With the formal, complete derivation performed in the present study, we obtain a different expression for E ,

$$E = \frac{\omega_0}{1 - r^2(1 - \omega_0 g_0)}, \quad (29)$$

where $r \equiv (1 - R_\infty)/(1 + R_\infty)$.

The two-stream fluxes depend on ω_0 and g_0 only via the coupling coefficients ζ_- and ζ_+ . When one inserts either expression for E into its corresponding expression for ζ_\pm , one obtains an identical result,

$$\zeta_\pm = \frac{1 \pm r}{2}. \quad (30)$$

In other words, E is *constructed* such that the coupling coefficients only depend on R_∞ . Similarly, because of the ambiguity with relating J , J_\uparrow , and J_\downarrow to the fluxes via ϵ and ϵ' , there are multiple ways of expressing the coupling coefficients in terms of E . Nevertheless, they all reduce to the same

expression for ζ_\pm as stated above. This property implies that the results shown in Figures 2, 3 and 4 of Heng & Kitzmann (2017) are identical³ even when the expression for E in Equation (29) is used, which is why we have not reproduced them in the current study. The conclusions of Heng & Kitzmann (2017) remain qualitatively *and* quantitatively unchanged.

Figure 1 shows the calculations of E using Equation (29). R_∞ is numerically evaluated using a brute-force, 32-stream discrete ordinates method (Stamnes et al. 1988) as implemented by the DISORT code (Hamre et al. 2013). In these numerical calculations, one has to specify an explicit form of the scattering phase function. A Henyey–Greenstein scattering phase function (Henyey & Greenstein 1941) is used when $g_0 \neq 0$; when $g_0 = 0$, we instead use the scattering phase function associated with Rayleigh scattering. To empower the reader in reproducing our calculations of E , we provide a fitting function,

$$E = 1.225 - 0.1582g_0 - 0.1777\omega_0 - 0.07465g_0^2 + 0.2351\omega_0g_0 - 0.05582\omega_0^2. \quad (31)$$

The mean and maximum errors associated with this fitting function are about 0.17% and 1.12%, respectively, for $g_0 \neq 0$. For $g_0 = 0$, the error may be as high as 2%.

6. Improved Two-stream Source Function Method

Starting from Equation (1), it is always possible to rewrite the radiative transfer equation as

$$\frac{\partial}{\partial \tau} \left[I \exp\left(-\frac{\tau}{\mu}\right) \right] = -\frac{(1 - \omega_0)}{\mu} B \exp\left(-\frac{\tau}{\mu}\right) - \frac{\omega_0}{\mu} \exp\left(-\frac{\tau}{\mu}\right) \int_0^{4\pi} \mathcal{P}(I + I_{\text{beam}}) d\Omega'. \quad (32)$$

In the case of pure absorption ($\omega_0 = 0$), there is an exact solution to the preceding equation if one performs a series expansion of B in terms of τ (or assume it to be constant). Generally (when $\omega_0 \neq 0$), the integral is challenging to evaluate.

The trick employed by the two-stream source function method of Toon et al. (1989) is to argue that $I \propto F_\uparrow, F_\downarrow$ in the integral. The scattering phase function is assumed to be $(1 + g_0)/4\pi$ and $(1 - g_0)/4\pi$ in the forward and reverse directions, respectively. Since the two-stream fluxes do not depend on Ω' , the integral becomes trivial to evaluate. The direct beam term is ignored, because the two-stream source function method was originally designed to treat diffuse thermal emission.

If we focus only on the integral term involving the diffuse emission,

$$\omega_0 \int_0^{4\pi} \mathcal{P} I d\Omega' = \frac{\omega_0}{2\pi} [(1 \pm g_0)F_\uparrow + (1 \mp g_0)F_\downarrow], \quad (33)$$

where we need $I = F_{\uparrow\downarrow}/\pi$ in order for the preceding expression to reduce correctly to its isotropic limit of $\omega_0 J/4\pi$ when

³ In a departure from Heng & Kitzmann (2017), we do not modify the expression for \mathcal{T} in an ad hoc way, which will produce small differences at optical depths of order unity and less.

$g_0 = 0$. This is a departure from Toon et al. (1989), who assumed $I = F_{\uparrow\downarrow}/\epsilon$. The minus and plus signs depend on whether one is writing the integral term for the outgoing (\uparrow) or incoming (\downarrow) flux.

We proceed somewhat differently from previous studies. In Toon et al. (1977), who first introduced the two-stream source function approximation, the form of their governing equations and the absence of g_0 in them suggest that their formalism does not generally treat $g_0 \neq 0$ scenarios. In Toon et al. (1989), who generalized their formalism to also consider $g_0 \neq 0$ values, their Equations (53) and (54) imply that their two-stream fluxes may be written as exponentials and linear terms involving the optical depth. The coefficients of these terms do not appear to depend on the optical depth. This is contrary to what we find for our two-stream fluxes in Equations (25) and (26), where the denominator contains the transmission function.

In our approach, we consider the two-stream fluxes to already have undergone the integration from τ_1 to τ_2 , such that

$$\begin{aligned} & \int_{\tau_1}^{\tau_2} \frac{\omega_0}{\mu} \exp\left(-\frac{\tau}{\mu}\right) \int_0^{4\pi} \mathcal{P}I \, d\Omega' \, d\tau \\ &= \frac{\omega_0}{2\pi\mu} [(1 \pm g_0)F_{\uparrow 1} + (1 \mp g_0)F_{\downarrow 2}] \int_{\tau_1}^{\tau_2} \exp\left(-\frac{\tau}{\mu}\right) d\tau. \end{aligned} \quad (34)$$

Note that $F_{\uparrow 1}$ and $F_{\downarrow 2}$ are our two-stream fluxes given by Equations (25) and (26), respectively.

If we again define $\Delta \equiv \tau_2 - \tau_1$ and

$$\mathcal{T}_0 \equiv \exp\left(-\frac{\Delta}{\mu}\right), \quad (35)$$

then the solution to Equation (32) is

$$\begin{aligned} I_i &= I_2 \mathcal{T}_0 + (B_i - B' \tau_i)(1 - \omega_0)(1 - \mathcal{T}_0) \\ &+ B'(1 - \omega_0)[\mu(1 - \mathcal{T}_0) + \tau_1 - \tau_2 \mathcal{T}_0] \\ &+ \frac{\omega_0}{2\pi} [(1 \pm g_0)F_{\uparrow 1} + (1 \mp g_0)F_{\downarrow 2}](1 - \mathcal{T}_0), \end{aligned} \quad (36)$$

where we have $i = 1$ and 2 for the incoming and outgoing directions, respectively. In the preceding expression, μ may be either positive or negative.

From the two-stream source function solution for the intensity in Equation (36), one may integrate over μ to obtain the two-stream-source function fluxes,⁴

$$\begin{aligned} \hat{F}_{\uparrow 1} &= F_{\uparrow 2} \mathcal{T}' + \pi B_2(1 - \omega_0)(1 - \mathcal{T}') \\ &+ \pi B'(1 - \omega_0) \left\{ \frac{2}{3} [1 - \exp(-\Delta)] - \Delta \left(1 - \frac{\mathcal{T}'}{3}\right) \right\} \\ &+ \frac{\omega_0}{2} [(1 + g_0)F_{\uparrow 1} + (1 - g_0)F_{\downarrow 2}](1 - \mathcal{T}'), \\ \times \hat{F}_{\downarrow 2} &= F_{\downarrow 1} \mathcal{T}' + \pi B_1(1 - \omega_0)(1 - \mathcal{T}') \\ &- \pi B'(1 - \omega_0) \left\{ \frac{2}{3} [1 - \exp(-\Delta)] - \Delta \left(1 - \frac{\mathcal{T}'}{3}\right) \right\} \\ &+ \frac{\omega_0}{2} [(1 - g_0)F_{\uparrow 1} + (1 + g_0)F_{\downarrow 2}](1 - \mathcal{T}'), \end{aligned} \quad (37)$$

⁴ Here, ‘‘two-stream-source function flux’’ refers to the fluxes derived using the two-stream source function approximation.

where we have used the symbols $\hat{F}_{\uparrow 1}$ and $\hat{F}_{\downarrow 2}$ to distinguish the two-stream-source function fluxes from the two-stream fluxes. We have defined

$$\mathcal{T}' \equiv 2E_3(\Delta) \equiv 2 \int_1^\infty x^{-3} \exp(-x \Delta) dx, \quad (38)$$

where E_3 is the exponential integral of the third order (Abramowitz & Stegun 1970; Arfken & Weber 1995). The reversal in the minus and plus signs for the two-stream-source function fluxes accounts for the change in the forward versus reverse directions for the incoming versus outgoing fluxes. For example, when $g_0 = 1$, there should be no correction due to backscattering and all of the scattered flux should be in the forward direction.

7. Summary and User’s Manual

The computational implementation of the two-stream solutions of radiative transfer has previously been described elsewhere (e.g., Malik et al. 2017). Instead, we focus on summarizing the different variants of the two-stream method from a practical standpoint.

1. If the reader is interested in modeling only diffuse infrared emission in the absence of medium-sized or large aerosols, the original two-stream formulation may suffice. For a concise statement of the two-stream fluxes with the hemispheric closure, see Equation (3.58) of Heng (2017). Alternatively, use Equations (25) and (26) of the present study with $E = 1$ and discard/ignore the direct beam terms.
2. If the reader is interested in including the direct stellar beam in the original two-stream solutions, see Meador & Weaver (1980) and Toon et al. (1989).
3. If the reader is interested in accurately modeling diffuse infrared emission in the presence of medium-sized or large aerosols, then our improved two-stream formulation in Equations (25) and (26) should be used. Consult Table 1 of Meador & Weaver (1980) or Toon et al. (1989) for the various choices of closure associated with the direct stellar beam (i.e., value of ϵ_2). The E -factor in Equation (29) should be used in tandem with the coupling coefficients in Equation (24). For convenience, we provide a simple fitting function for E in Equation (31).
4. Our improved two-stream solutions may be used to derive the two-stream source function solutions, which we present in Equations (36) and (37) for the intensity and fluxes, respectively.

For completeness, the summary of key symbols used and their correspondence to the symbols used in Toon et al. (1989) are listed in Table 1. Appendix lists some typographical errors we found in Meador & Weaver (1980) and Toon et al. (1989).

We acknowledge financial support from the Swiss National Science Foundation, the European Research Council via Consolidator Grant ERC-2017-CoG-771620-EXOKLEIN (awarded to K.H.), the PlanetS National Center of Competence in Research (NCCR), the Center for Space and Habitability (CSH), and the Swiss-based MERAC Foundation.

Table 1
Summary of Symbols Used, Including Correspondence to Toon et al. (1989)

Symbol	Name	Symbol Used By Toon et al. (1989)
ω_0	single-scattering albedo	ω_0
g_0	scattering asymmetry factor	g
μ_*	cosine of stellar beam angle	μ_0
F_*	stellar flux	πF_s
$\epsilon = 1/2$	first Eddington coefficient	μ_1
ϵ'	first Eddington coefficient	...
$\epsilon_2 \sim 1$	second Eddington coefficient	...
$E \equiv \epsilon/\epsilon'$	ratio of first Eddington coefficients	1
ζ_{\pm}	coupling coefficients	...
ζ_-/ζ_+	asymptotic value of reflectivity	Γ
$A_1 \zeta_+$...	k_1
$A_2 \zeta_-$...	$k_2 \Gamma$
$\Delta \equiv \tau_2 - \tau_1$	optical depth of atmospheric layer	τ
$T \equiv \exp(-\alpha \Delta)$	transmissivity or transmission function	$\exp(-\lambda \tau)$
$T_* \equiv \exp(-\Delta/\mu_*)$	extinction of direct beam through atmospheric layer	$\exp(-\tau/\mu_0)$
T_{above}	extinction of direct beam above atmospheric layer	$\exp(-\tau_c/\mu_0)$
F_{\uparrow}	outgoing flux	F^+
F_{\downarrow}	incoming flux	F^-
$F \equiv F_{\uparrow} - F_{\downarrow}$	net flux	F_{net}
$F_+ \equiv F_{\uparrow} + F_{\downarrow}$	total flux	...
γ_a	coefficient in equation for $\frac{\partial F_{\uparrow\downarrow}}{\partial \tau}$	γ_1
γ_s	coefficient in equation for $\frac{\partial F_{\uparrow\downarrow}}{\partial \tau}$	γ_2
γ_B	coefficient in equation for $\frac{\partial F_{\uparrow\downarrow}}{\partial \tau}$	$2\pi(1 - \omega_0)$
$\gamma_* \equiv \omega_0 g_0 \mu_* F_*/\epsilon_2$	coefficient of direct beam term in equation for $\frac{\partial F_{\uparrow\downarrow}}{\partial \tau}$...
$\alpha \equiv \sqrt{(\gamma_a + \gamma_s)(\gamma_a - \gamma_s)}$	diffusivity factor	λ
χ_{\uparrow}	...	$\gamma_3 = \beta_{0\nu}$
χ_{\downarrow}	...	$\gamma_4 = 1 - \beta_{0\nu}$

Appendix

Typographical Errors in Meador & Weaver (1980) and Toon et al. (1989)

A.1. Toon et al. (1989)

We use the same notation as Toon et al. (1989): (μ', ϕ') and (μ, ϕ) , for the incident and emergent directions, respectively. Equation (1) of Toon et al. (1989) concisely states the functional dependence of their scattering phase function: $P_{\nu}(\mu, \mu', \phi, \phi')$. The scattering phase function associated with the direct beam is stated in their Equation (3): $P_{\nu}(\mu, -\mu_0, \phi, \phi_0)$. Based on their stated convention and also their Equation (10), we may conclude that Equation (9) of Toon et al. (1989) is in error, as it should instead be

$$\beta_{0\nu} = \frac{1}{2} \int_0^1 P_{\nu}(\mu, -\mu_0) d\mu. \quad (39)$$

Specifically, the integration should be over μ and not μ' .

In Equation (12), there is a typographical error: S_n^+ should instead be S_n^- . The logical flow of the text into Equations (13) and (14) of Toon et al. (1989) makes this clear.

A.2. Meador & Weaver (1980)

Our notations for μ, μ', ϕ and ϕ' are the same as those of Meador & Weaver (1980). Equation (1) of Meador &

Weaver (1980) concisely states the functional dependence of their scattering phase function: $P_{\nu}(\mu, \phi; \mu', \phi')$. It also states the functional dependence of the scattering phase function associated with the direct beam: $P_{\nu}(\mu, \phi; -\mu_0, \phi_0)$. Equation (7) of Meador & Weaver (1980) follows logically from their Equation (3). However, their Equation (9) suffers from the same logical inconsistency as what we pointed out for Toon et al. (1989), which is that the integration should be over μ and not μ' . This is clearly seen in the transition from Equations (7) to (9). It is not an issue of a mere change of notation for the integration variable, because it needs to correspond to the correct variable in the scattering phase function. Equation (9) of Meador & Weaver (1980) should instead be


$$\beta_i = \frac{1}{2\omega_0} \int_0^1 P_{\nu}(\mu, -\mu_i) d\mu. \quad (40)$$

It is likely that this error in Meador & Weaver (1980) propagated into Toon et al. (1989).

ORCID iDs

Kevin Heng  <https://orcid.org/0000-0003-1907-5910>

Matej Malik  <https://orcid.org/0000-0002-2110-6694>

Daniel Kitzmann  <https://orcid.org/0000-0003-4269-3311>

References

- Abramowitz, M., & Stegun, I. A. 1970, *Handbook of Mathematical Functions* (New York: Dover) 9th Printing
- Arfken, G. B., & Weber, H. J. 1995, *Mathematical Methods for Physicists* (4th ed.; San Diego: Academic)
- Chandrasekhar, S. 1960, *Radiative Transfer* (New York: Dover)
- Hamre, B., Starnes, S., Starnes, K., & Starnes, J. J. 2013, in *AIP Conf. Proc.* 1531 (Melville, NY: AIP), [923](#)
- Heng, K. 2017, *Exoplanetary Atmospheres: Theoretical Concepts and Foundations* (Oxford: Princeton Univ. Press)
- Heng, K., & Kitzmann, D. 2017, [ApJS](#), **232**, [20](#)
- Heng, K., Mendonça, J. M., & Lee, J.-M. 2014, [ApJS](#), **215**, [4](#)
- Heney, L. G., & Greenstein, J. L. 1941, [ApJ](#), **93**, [70](#)
- Joseph, J. H., Wiscombe, W. J., & Weinman, J. A. 1976, [JAtS](#), **33**, [2452](#)
- Kitzmann, D., Patzer, A. B. C., & Rauer, H. 2013, [A&A](#), **557**, [A6](#)
- Kitzmann, K. 2016, [ApJL](#), **817**, [L18](#)
- Malik, M., Grosheintz, L., Mendonça, J. M., et al. 2017, [AJ](#), **153**, [56](#)
- Marley, M. S., & McKay, C. P. 1999, [Icar](#), **138**, [268](#)
- Meador, W. E., & Weaver, W. R. 1980, [JAtS](#), **37**, [630](#)
- Mihalas, D. 1970, *Stellar Atmospheres* (1st ed.; San Francisco: Freeman)
- Mihalas, D. 1978, *Stellar Atmospheres* (2nd ed.; San Francisco: Freeman)
- Pierrehumbert, R. T. 2010, *Principles of Planetary Climate* (New York: Cambridge Univ. Press)
- Schuster, A. 1905, [ApJ](#), **21**, [1](#)
- Starnes, K., Tsay, S.-C., Wiscombe, W., & Jayaweera, K. 1988, [ApOpt](#), **27**, [2502](#)
- Toon, O. B., McKay, C. P., & Ackerman, T. P. 1989, [JGR](#), **94**, [16287](#)
- Toon, O. B., Pollack, J. B., & Sagan, C. 1977, [Icar](#), **30**, [663](#)
- Wiscombe, W. J. 1977, [JAtS](#), **34**, [1408](#)