


Accepted Manuscript

Revisiting systemic treatment of bacterial endophthalmitis: a review of intravitreal penetration of systemic antibiotics

Lisa Brockhaus, David Goldblum, Laura Eggenschwiler, Stefan Zimmerli, Catia Marzolini



PII: S1198-743X(19)30039-4

DOI: <https://doi.org/10.1016/j.cmi.2019.01.017>

Reference: CMI 1563

To appear in: *Clinical Microbiology and Infection*

Received Date: 14 December 2018

Accepted Date: 28 January 2019

Please cite this article as: Brockhaus L, Goldblum D, Eggenschwiler L, Zimmerli S, Marzolini C, Revisiting systemic treatment of bacterial endophthalmitis: a review of intravitreal penetration of systemic antibiotics, *Clinical Microbiology and Infection*, <https://doi.org/10.1016/j.cmi.2019.01.017>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Revisiting systemic treatment of bacterial endophthalmitis: a review of intravitreal penetration of systemic antibiotics

Lisa Brockhaus¹, David Goldblum², Laura Eggenschwiler², Stefan Zimmerli³, Catia Marzolini¹

¹ Division of Infectious Diseases and Hospital Epidemiology, Departments of Medicine and Clinical Research, University Hospital of Basel and University of Basel, Basel, Switzerland.

² Department of Ophthalmology, University Hospital of Basel and University of Basel, Switzerland.

³ Department of Infectious Diseases, University Hospital of Bern and University of Bern, Switzerland

Narrative review

Word count: Abstract: 221

Text: 2497

Tables: 2

Corresponding author:

Lisa Brockhaus, MD

Division of Infectious Diseases

University Hospital Basel

Switzerland

Phone: +41 61 265 50 53

E-mail: lisa.brockhaus@unibas.ch

Keywords

Endophthalmitis, systemic antibiotic, vitreous concentration, intravitreal penetration, blood-retinal barrier

Abstract

Background: Adjunctive systemic antibiotic therapy for treatment of bacterial endophthalmitis is controversial but common practice due to the severity of the disease. In absence of guidance documents, several antibiotic regimens are being used without applying evidence-based prescribing, thus leading to inappropriate treatment of this serious eye condition.

Objectives: To summarize available data on intravitreal penetration of systemically administered antibiotics and to discuss their usefulness from a microbiological and pharmacological point of view.

Sources: We performed a systematic PubMed search of studies investigating antibiotic concentrations in the vitreous after systemic administration in humans, and selected animal models.

Content: The best-documented agents achieving therapeutic levels in the vitreous are meropenem, linezolid and moxifloxacin. Vancomycin, cefazoline, ceftriaxone, ceftazidime, imipenem and trimethoprim-sulfamethoxazole reach levels justifying their use in specific situations. Available data do not support the use of ciprofloxacin, levofloxacin, aminoglycosides, aminopenicillins, piperacillin, cefepime, and clarithromycin. With very limited but available promising data, the use of daptomycin and rifampicin deserves further investigation.

Implications: The choice of the adjunctive systemic antibiotic agent – in situations where considered relevant for treatment - must to date be made on an individual base, considering microbiological aspects as well as operative status and inflammation of the eye. This review

gives a systematic overview of antibiotic options and provides guidance to the clinician striving for optimal systemic antibiotic treatment of bacterial endophthalmitis.

ACCEPTED MANUSCRIPT

Introduction

Vitreotomy and intravitreal antibiotics are nowadays considered as the gold standard treatment of bacterial endophthalmitis [1]. Adjunctive use of systemic antibiotics is controversial but common practice justified by the severity of the disease. Data on intravitreal antibiotic levels reached by systemic administration are sparse and comprehensive recommendations for systemic use have not been established [1–3]. The availability of new antibiotic agents, changing resistance patterns, and new surgical techniques using implants such as keratoprosthesis justify revisiting their systemic use for endophthalmitis.

This review summarizes available data on intravitreal penetration of systemically administered antibiotics and discusses preferred regimens for the treatment of bacterial endophthalmitis from a microbiological and pharmacological perspective.

Definition and commonly isolated microorganisms

Endophthalmitis refers to the inflammation of the internal eye affecting the vitreous cavity and the anterior chamber, resulting from exogenous (mostly surgery related) or, more rarely (5-10%) hematogenous insertion of microorganisms [4].

The most commonly isolated microorganisms are coagulase-negative staphylococci (40-70%), *Staphylococcus aureus* (10-17%), streptococci (5-15%), other Gram-positive cocci including enterococci (5%), as well as Gram-negative bacilli (5-10%) including *Haemophilus influenzae* and *Pseudomonas aeruginosa* [5–8]. The microbiologic spectrum of less common forms like post-traumatic or endogenous endophthalmitis is more varied. For instance, bacillus sp. is regularly found after open-globe injuries [9]. The bacterial spectrum encountered may further vary according to local epidemiology and peri-operative prophylactic regimens [10].

Current treatment recommendations

The current mainstays of bacterial endophthalmitis treatment are vitrectomy and intravitreal antibiotics [1]. Vitrectomy aims to reduce the bacterial load with the intention of a local “source control”. Although complete vitrectomy would ensure maximal eradication of the infected tissue, partial vitrectomy is often preferred in clinical practice, because of the lower associated risk of iatrogenic retinal detachment.

Intravitreal antibiotics are injected immediately after vitrectomy and their administration is usually repeated after 48 hours if the clinical course is not favourable. The most commonly used regimens are vancomycin combined with ceftazidime or amikacin. The downside of repeated injections is an increased risk of retinal toxicity.

Intravitreal dexamethasone is often added to reduce intraocular inflammation despite conflicting evidence [1]. Corticosteroids accelerate blood-retinal barrier restitution and thus influence antibiotic penetration.

The role of systemic antibiotics

The single large clinical trial evaluating the role of systemic antibiotics for the treatment of endophthalmitis is the Endophthalmitis-Vitrectomy-Study conducted in the 1990s [11]. In this randomized study, no significant difference in visual acuity was found in patients receiving intravitreal antibiotics followed by intravenous antibiotic therapy compared to patients receiving only intravitreal treatment. However, this finding has been questioned because of the exclusion of patients with severe endophthalmitis and the choice of adjunctive antibiotics: ceftazidime has poor activity against the dominant Gram-positive organisms and amikacin has very limited intraocular penetration. Since then, only small studies have evaluated the efficacy of systemic antibiotic therapy in endophthalmitis with varying methodologies and results [12-13].

Some recommendations advocate the adjunctive use of systemic antibiotics in severe acute purulent postoperative endophthalmitis [1]. Recommended regimens include vancomycin

combined with ceftazidime [14] or imipenem with ciprofloxacin [15]. For the treatment of endogenous endophthalmitis the use of systemic antibiotics is undisputed [2, 4].

The pharmacokinetic rationale for adjunctive systemic antibiotics is the rapid elimination of intravitreally applied antibiotics, with almost complete removal after 24h [16], whereas systemic administration favors intraocular antibiotic accumulation over time.

Discussing the controversial benefit of adjunctive systemic antibiotics in terms of visual outcome is beyond the scope of this review. The imminent poor outcome constitutes a strong argument for clinicians to use systemic therapy. We strongly believe that adjunctive systemic therapy should not be denied on an individual base, provided that all efforts are made to isolate the causative pathogen and to apply evidence-based prescribing of antibiotic agents.

Intravitreal penetration of systemic antibiotics

Factors determining the penetration of antibiotics into the eye

The penetration of antibiotics into the posterior segment of the eye after systemic administration is limited by two blood-retinal barrier mechanisms (BRB): The retinal pigment endothelial cells located within the retinal cell layers (outer BRB) and the retinal capillary endothelial cells (inner BRB) [17]. Of note, entry into the anterior segment of the eye is limited by the blood-aqueous-barrier characterized by less restrictive properties, thereby resulting in different aqueous and vitreous drug concentrations.

Drug permeability across the blood-retinal barriers depends on drug characteristics such as the molecular size, lipophilicity, ionization and protein binding. Of interest, BRB was shown to be more permeable than the blood-brain-barrier (BBB) owing to morphological differences [17]. As in the BBB, ocular inflammation increases drug permeability across the BRB.

Elimination of antibiotics from the vitreous occurs via two routes: passive diffusion to the anterior chamber and through the Schlemm's channel (anterior route), and retrograde

transport through the blood-retinal barrier (posterior route). Clearance pathways from the vitreous depends not only on physico-chemical drug properties and ocular inflammation, but also on the surgical status [18]. Post-operative aphakia – after removal of an artificial intra-ocular lens – as well as vitrectomy influence elimination of antibiotics [19].

Pharmacokinetic studies

The knowledge of antibiotics pharmacokinetics in the eye is derived from two types of studies: single concentration measurements in human eyes performed at the time of surgery, and rabbit models allowing for repetitive drug measurements. Several limitations should be considered: although single drug measurements are of high value for antibiotics characterized by a concentration-dependent killing effect (aminoglycosides, daptomycin), this approach is less informative for antibiotics with a killing profile that depends on time-above-MIC (beta-lactams) or AUC/MIC (fluoroquinolones, vancomycin, linezolid).

- 1) Common pharmacodynamic index values (such as C_{max}/MIC used in most of the assessed studies) do not necessarily reflect efficacy in the complex microenvironment of the eye.
- 2) Comprehensive MIC-studies of endophthalmitis isolates are missing. In this review we use EUCAST breakpoints and wild-type distributions (ECOFF) [20] (table 2) as a reference to estimate potential efficacies of antibiotics.
- 3) Animal studies must be interpreted with caution, as they do not fully reflect the pharmacokinetics in humans.

Suitable publications were identified by a PubMed search using the terms [antibiotic name] and [vitreous] and [systemic/oral/intravenous]. If no publications were found, the search was complemented by the terms [eye] and [penetration]. Rabbit model studies were included for

antibiotics with limited human data or if they provided additional relevant information. Studies are summarized in table 1.

Review of available literature

Vancomycin

Vancomycin did not show any accumulation in phakic (including inflamed) eyes in a rabbit study [21]. Aphakic-vitreotomized eyes showed vitreous levels just above breakpoints of commonly involved Gram-positive organisms after one dose. In aphakic eyes, comparable levels were only reached after prolonged therapy. The poor intravitreal penetration of vancomycin is consistent with the limited CSF penetration [22] and explained by its high molecular weight and hydrophilicity.

The rationale for its continued empiric use [14] despite limited data, is its microbiological spectrum covering almost 100% of Gram-positive organisms causing endophthalmitis [7]. Available data nevertheless suggest that systemic administration should only be considered in aphakic eyes. Given the delay in achieving sufficient concentrations, systemic administration should follow immediately intravitreal injection of the same agent.

Penicillins

Poor vitreal penetration of ampicillin and amoxicillin were demonstrated in rabbit models [23-24]. Similarly, insufficient vitreous concentrations of piperacillin were demonstrated in human eyes with diverse operative status [25]. The vitreal penetration of penicillin G has not been studied but based on CSF penetration data [22] and drug properties, penetration into the vitreous is anticipated to be minimal. Nevertheless, in analogy to CSF infections, high systemic doses might provide effective vitreous levels for streptococcal endophthalmitis.

Available data suggest that systemic administration of most penicillins is not appropriate to treat endophthalmitis. Penicillin G vitreous levels remain to be investigated.

Cephalosporins

Cefazoline vitreous concentrations are issued from a large rabbit study [26] showing levels well above streptococcal breakpoints in aphakic-vitreotomized inflamed eyes but undetectable levels in phakic non-inflamed eyes.

Ceftriaxone was detectable in human phakic non-inflamed eyes after multiple dosing [27] at levels well above streptococci and enterobacteriaceae breakpoints, but below the ECOFF of *S.aureus*. Whether the observed levels result from accumulation after repetitive dosing, or rapid penetration as previously shown in CSF studies [28], is not known.

Vitreous levels of ceftazidime are based on two rabbit studies [29-30]. Levels were above enterobacteriaceae breakpoints in aphakic-vitreotomized inflamed eyes. Only delayed penetration was observed in non-vitreotomized inflamed eyes and undetectable levels were found in phakic non-inflamed eyes [29].

Cefepime showed low vitreous levels in human phakic non-inflamed eyes [31], nonetheless penetration in inflamed eyes have not been studied.

In summary, cefazoline, ceftriaxone and ceftazidime can be considered as targeted therapy when the pathogen is identified and the MIC of the isolate has been determined, yet on a thin evidence base. Ceftazidime is likely to be effective against most enterobacteriaceae in aphakic-vitreotomized eyes. Since it exhibits very limited anti-streptococcal and no anti-staphylococcal activity, it should not be used to cover Gram-positive organisms. Cefepime cannot be recommended as observed levels are clearly below those of other cephalosporins, yet investigation in inflamed eyes is warranted. Neither ceftazidime nor cefepime can be recommended to treat pseudomonas endophthalmitis based on available data.

Carbapenems

Two studies in human phakic non-inflamed eyes showed imipenem vitreous concentrations around 2.0 ug/ml [32-33]. Meropenem was shown to rapidly achieve four-fold higher

vitreous concentrations in a comparable study [34]. This finding is consistent with observed high meropenem levels in CSF [22] and explained by favorable physicochemical properties. Unlike imipenem, the observed levels of meropenem clearly exceed breakpoints for relevant Gram-positive and Gram-negative organisms.

The potential value of carbapenems in empirical treatment – as single agents - is limited by the high prevalence of oxacillin-resistant coagulase negative staphylococci also in ophthalmic isolates [8]. For empiric combination therapy, we would advocate the use of meropenem rather than imipenem based on the above-discussed data. Furthermore, meropenem appears to be the preferred option for targeted *Pseudomonas* treatment, particularly when considering the insufficient concentrations of ciprofloxacin, piperacillin, ceftazidime and cefepime as discussed above.

Rifampicin

High rifampicin vitreous levels were observed in phakic non-inflamed rabbit eyes [35], however with doses that, corrected for weight, would largely exceed tolerated doses in humans.

Human studies are limited to aqueous levels, observed to be 0.2-1.3 ug/ml [36]. As rifampicin vitreous levels were consistently shown to be half of aqueous levels [31], human vitreous levels could be expected to exceed 0.1 ug/ml, which is well above breakpoints for staphylococci.

A role of rifampicin in endophthalmitis after foreign-body implantation, incomplete vitrectomy, and aggressive *S.aureus*-infection deserves further consideration due to its bactericidal and biofilm-active-properties. Importantly, rifampicin has to be combined with an effective second anti-staphylococcal agent to prevent resistance.

Linezolid

Mean vitreous levels of linezolid in phakic non-inflamed human eyes range from 1.2 to 3.7 ug/ml after one dose [37–40], with two studies showing further accumulation after two doses (4.5 and 5.7 ug/ml) [37, 39]. The good penetration into the vitreous is consistent with CSF penetration [22].

Considering breakpoints of Gram-positive organisms, sufficient vitreous levels are likely to be attained after two doses. The drug, however, is bacteriostatic.

The well documented ocular penetration and comprehensive coverage of Gram-positive germs make of linezolid a potential alternative to vancomycin. Caution is needed due to its toxicity (myelosuppression, peripheral neuropathy, optic neuropathy), although relatively infrequent in short-term administration [41].

Daptomycin

Knowledge of daptomycin penetration is limited to one case report of a patient treated for MRSA-endophthalmitis in a strongly inflamed, phakic, non-vitreotomized eye [42]. Single dose administration (10 mg/kg) resulted in a vitreous concentration of 12.4 ug/ml 42h post administration (patient had renal insufficiency). This finding appears promising considering low staphylococcal breakpoints and the drug's bactericidal properties, but its use remains experimental to date.

Aminoglycosides

Vitreous concentrations far below breakpoints of relevant pathogens were shown after intravenous administration of amikacin and gentamycin in a rabbit study [43], despite study conditions expected to enhance vitreous levels (inflammation, aphakia and vitrectomy).

Given the availability of better alternatives, there is no role for systemic administration of aminoglycosides in endophthalmitis.

Fluoroquinolones

Despite relatively good vitreous/serum (V/S) ratios owing to favourable physicochemical properties, observed concentrations in vitreous proved insufficient for ciprofloxacin [44–50] and levofloxacin [38, 51-52].

Moxifloxacin demonstrated concentrations well above breakpoints of relevant organisms after two doses [53-54], whereas concentrations were significantly lower after a single dose [55-56]. The low concentrations of the Vedantham study can be explained by an inadequate short sampling time of 90 minutes after oral administration [56]. Moxifloxacin maximal CSF levels were shown to occur 2-3 hours after maximal systemic levels [57].

In summary, there are sufficient data to oppose the use of ciprofloxacin and levofloxacin for the treatment of endophthalmitis. Conversely, several studies consistently demonstrating satisfactory moxifloxacin levels are available. Given the observed higher levels with moxifloxacin 800mg, this increased dosage can be considered with a careful monitoring for side effects. Safety data on this dosage are not comprehensive to date [58].

Moxifloxacin lacks activity against most oxacillin-resistant coagulase-negative staphylococci [8]), and streptococci rapidly develop resistance particularly when drug concentrations are low. Therefore, the use of moxifloxacin for empirical treatment of endophthalmitis is limited.

Trimethoprim-Sulfamethoxazole

Sulfonamides and trimethoprim have demonstrated a moderate penetration in the vitreous in one human study [59]. The doses used, however, were below those for the treatment of meningitis [60]. The observed concentrations are not exceeding breakpoints of all relevant organisms although higher concentrations might be reached with increased doses. Concentrations are far below the breakpoint of *Stenotrophomonas maltophilia* (4 ug/ml) suggesting difficulty in treating this pathogen.

Due to limited data and better alternatives for Gram-positive organisms, there is no current role for TMP/SMX in systemic endophthalmitis treatment.

Clarithromycin

Based on one human study of phakic non-inflamed eyes showing insufficient vitreous levels [61], there is no argument to recommend clarithromycin for endophthalmitis treatment. Similarly insufficient levels have been reported in CSF, where the use of clarithromycin is limited to case reports of successful treatment of atypic organisms [22].

Conclusion

Data on the intravitreal concentrations of systemic antibiotics are generally scarce and are based on a single study for many agents. Relatively good evidence exists for therapeutic vitreous levels of meropenem, linezolid and high-dose moxifloxacin. None covers the required bacterial spectrum when used empirically as single agents, but the combination of linezolid with meropenem in empirical treatment of endophthalmitis may offer broad activity against the majority of pathogens. Vancomycin, cefazoline, ceftriaxone, ceftazidime, imipenem, daptomycin and TMP-SMX exhibit levels supporting their use in specific situations for targeted therapy. The operative status of the infected eye needs also to be considered. Available data do not support the use of ciprofloxacin, levofloxacin, aminoglycosides, aminopenicillins, piperacillin, cefepime or clarithromycin. Rifampicin may be considered for combination therapy in complicated staphylococcal infections. Further data on daptomycin vitreous penetration would be valuable.

The choice of the adjunctive systemic antibiotic agent – in situations where considered relevant for treatment - must to date be made on an individual base, taking into account suspected or detected organisms, operative status, intraocular inflammatory activity and drug side effect profiles. Future research should assess the clinical outcome after use of systemic antibiotics with documented good intraocular penetration (e.g. meropenem, linezolid and moxifloxacin), and assess the role of rifampicin for staphylococcal infections.

Transparency declaration

The authors declare no competing interests regarding this manuscript. This work did not receive any funding.

Access to data

All studies included in this review are publicly available.

Author contributions

LB has done the literature search, analyzed and compiled the data. DG, LE have provided input in relation to ophthalmological and ophthalmological-surgical aspects. CM has contributed to the pharmacological content and analysis of data. SZ has participated in the interpretation of data and revised the manuscript. LB and CM have drafted the manuscript. All authors have revised the manuscript for intellectual content and approved the final submitted version.

Acknowledgement

We thank Prof. Manuel Battegay for helpful comments on an earlier draft of this paper.

References

- [1] Barry P, Cordovés L, Gardner S. ESCRS Guidelines for Prevention and Treatment of Endophthalmitis Following Cataract Surgery: Data, Dilemmas and Conclusions. 2013. <http://www.es CRS.org/endophthalmitis>
- [2] Birnbaum F, Gupta G. Endogenous Endophthalmitis: Diagnosis and Treatment. *EyeNet Mag* 2016;6:33–35.
- [3] Wakely L, Sheard R, Hospital RH. Recent advances in Endophthalmitis Management. *Focus* 2014;2:5-6.
- [4] Kernt M, Kampik A. Endophthalmitis: Pathogenesis, clinical presentation, management, and perspectives. *Clin Ophthalmol* 2010;4:121–135.
- [5] Ong A, Te NA, Zagora S, Symes R, Yates W, Chang A et al. Post-surgical versus post-intravitreal injection endophthalmitis: changing patterns in causative flora. *Clin Exp Ophthalmol* 2018;jul:1–6. (Epub ahead of print)
- [6] Han DP, Wisniewski SR, Wilson LA, Barza M, Vine AK, Doft BH et al. Spectrum and susceptibilities of microbiologic isolates in the endophthalmitis vitrectomy study. *Am J Ophthalmol* 1996;122(1),1–17.
- [7] Gentile RC, Shukla S, Shah M, Ritterband DC, Engelbert M, Davis A et al. Microbiological spectrum and antibiotic sensitivity in endophthalmitis: A 25-year review. *Ophthalmology* 2014;121(8):1634–1642.
- [8] Asbell PA, Mah FS, Sanfilippo CM, DeCory HH. Antibiotic susceptibility of bacterial pathogens isolated from the aqueous and vitreous humor in the Antibiotic Resistance Monitoring in Ocular Microorganisms (ARMOR) surveillance study. *J Cataract Refract Surg* 2016;42(12):1841–1843.
- [9] Miller JJ, Scott IU, Flynn HW jr, Smiddy WE, Murray TG, Berrocal A et al. Endophthalmitis Caused by Bacillus Species. *Am J Ophthalmol* 2008;145(5):883–888.
- [10] Friling E, Lundström M, Stenevi U, Montan P. Six-year incidence of endophthalmitis after cataract surgery: Swedish national study. *J Cataract Refract Surg* 2013;39(1):15–21.
- [11] Endophthalmitis Vitrectomy Study Group. Results of the Endophthalmitis Vitrectomy Study. A randomized trial of immediate vitrectomy and of intravenous antibiotics for the treatment of postoperative bacterial endophthalmitis. *Arch Ophthalmol* 1995;113(12):1479–1496.
- [12] Tappeiner C, Schuerch K, Goldblum D, Zimmerli S, Fleischhauer JC, Frueh BE. Combined meropenem and linezolid as a systemic treatment for postoperative endophthalmitis. *Klin Monatsbl Augenheilkd* 2010;227(4):257–261.
- [13] Hooper CY, Lightman SL, Pacheco P, Tam PMK, Khan A, Taylor SRJ. Adjunctive antibiotics in the treatment of acute bacterial endophthalmitis following cataract surgery. *Acta Ophthalmol* 2012;90(7):572–573.
- [14] Behrens-Baumann W. Current Therapy for Postoperative Endophthalmitis. *Klin Monatsbl für Augenheilk* 2008;225(11):919–923.
- [15] German Society for Cataract & Refractive Surgeons. Leitlinie zur Prophylaxe und Therapie von Endophthalmitiden. 2005. <http://www.dgii.org/de/empfehlungen-101.html>

- [16] Ficker L, Meredith T, Gardner S, Wilson LA. Cefazolin levels after intravitreal injection. Effects of inflammation and surgery. *Invest Ophthalmol Vis Sci* 1990;31(3):502–505.
- [17] Del Amo E, Rimpelä AK, Heikkinen E, Kari OK, Ramsay E, Lajunen T, et al. Progress in Retinal and Eye Research Pharmacokinetic aspects of retinal drug delivery. *Prog Retin Eye Res* 2017;57:134–185.
- [18] Luaces-Rodríguez A, González-Barcia M, Blanco-Teijeiro M, Gil-Martinez M, Gonzalez F, Gomez-Ulla F et al. Review of Introcular Pharmacokinetics of Anti-Infectives Commonly Used in the Treatment of Infectious Endophthalmitis. *Pharmaceutics* 2018;10(66):1–15.
- [19] Radhika M, Mithal K, Bawdekar A, Dave V, Jindal A, Relhan N, et al. Pharmacokinetics of intravitreal antibiotics in endophthalmitis. *J Ophthalmic Inflamm Infect* 2014;4(22):1–9.
- [20] European Comitee for Susceptibility Testing.
http://www.eucast.org/clinical_breakpoints. Version 8.1 (accessed in May 2018)
- [21] Meredith TA, Aguilar HE, Shaarawy A, Kincaid M, Dick J, Niesman MR. Vancomycin levels in the vitreous cavity after intravenous administration. *Am J Ophthalmol* 1995;119(6):774–778.
- [22] Nau R, Sorgel F, Eiffert H. Penetration of drugs through the blood-cerebrospinal fluid/blood-brain barrier for treatment of central nervous system infections. *Clin Microbiol Rev* 2010;23(4):858–883.
- [23] Salminen L. Ampicillin penetration into the rabbit eye. *Acta Ophthalmol* 1978;56:977–938.
- [24] Faigenbaum S, Boyle G, Prywes A, Abel R, Leopold I. Intraocular Penetration of Amoxicillin. *Am J Ophthalmol* 1976;82(4):598–603.
- [25] Robinet A, Le Bot M, Colin J, Riche C. Penetration of Piperacillin into the vitreous after Intravenous Administration. *Retina* 1998;18(6):526–530.
- [26] Martin DF, Ficker LA, Gardner SK, Louis A, Doses RA. Vitreous cefazolin levels after intravenous injection. *Arch Ophthalmol* 1990;108:411–414.
- [27] Sharir M, Triesrer G, Kneer J, Rubinstein E. The Intravitreal Penetration of Ceftriaxone in Man following Systemic Administration. *Invest Ophthalmol Vis Sci* 1989;30(10):2179–2183.
- [28] Del Rio M, McCracken G, Nelson J, Chrane D, Shelton S. Pharmacokinetics and cerebrospinal fluid bactericidal activity of ceftriaxone in the treatment of pediatric patients with bacterial meningitis. *Antimicrob Agents Chemother* 1982;22(4):622–627.
- [29] Aguilar H, Meredith T, Shaarawy A, Kincaid M, Dick J. Vitreous cavity penetration of ceftazidime after intravenous administration. *Retina* 1995;15:154–159.
- [30] Walstad R, Blika S. Penetration of Ceftazidime into the Normal Rabbit and Human Eye. *Scand J Infect Dis* 1985;44:63–67.
- [31] Aras C, Ozdamar A, Ozturk R, Karacorla M, Ozkan S. Intravitreal penetration of cefepime after systemic administration to humans. *Ophthalmologica* 2002;216:261–264.
- [32] Adenis J, Mounier M, Salomon J, Denis F. Human vitreous penetration of imipenem.

- Eur J Ophthalmol 1994;4(2):115–117.
- [33] Axelrod J, Newton J, Klein R, Bergen R, Sheikh M. Penetration of Imipenem Into Human Aqueous and Vitreous Humor. *Am J Ophthalmol* 1987;104:649–653.
- [34] Schauersberger J, Amon M, Wedrich A, Nepp J, El Menyawi I, Derbolav A, et al. Penetration and decay of meropenem into the human aqueous humor and vitreous. *J Ocul Pharmacol Ther* 1999;15(5):439–445.
- [35] Wong K, D’Amico D, Oum B, Baker P, Kenyon K. Intraocular penetration of rifampin after oral administration. *Graefe’s Arch Exp Ophthalmol* 1990;228:40–43.
- [36] Outman WR, Levitz RE, Hill DA, Nightingale CH. Intraocular penetration of rifampin in humans. *Antimicrob. Agents Chemother* 1992;36(7):1575–1576.
- [37] Fiscella RG, Lai W, Buerk B, Khan M, Rodvold KA, Pulido JS, et al. Aqueous and vitreous penetration of linezolid after oral administration. *Ophthalmology* 2004;111(6):1191–1195.
- [38] George J, Fiscella R, Blair M, Rodvold K, Ulanski L, Stokes J, et al. Aqueous and vitreous penetration of linezolid and levofloxacin after oral administration. *J Ocul Pharmacol Ther* 2010;26(6):579–586.
- [39] Horcajada JP, Atienza R, Sarasa M, Soy D, Adán A, Mensa J. Pharmacokinetics of linezolid in human non-inflamed vitreous after systemic administration. *J Antimicrob Chemother*, vol. 63, no. 3, pp. 550–552, 2009.
- [40] Ciulla T, Comer G, Peloquin C, Wheeler J. Human vitreous distribution of linezolid after a single oral dose. *Retina* 2005;25:619–624.
- [41] Vazquez J, Arnold AC, Swanson RN, Biswas P, Bassetti M. Safety of long-term use of linezolid : results of an open-label study. *Ther Clin Risk Manag* 2016;12:1347–1354.
- [42] Sheridan K, Potoski B, Shields R, Nau G. Presence of Adequate Intravitreal Concentration of Daptomycin After Systemic Intravenous Administration in an Patient with Endogenous Endophthalmitis. *Pharmacotherapy* 2010;30:1247–1251.
- [43] El-Massry A, Meredith TA, Aguilar HE; Shaarawy A, Kincaid M, Dick J, et al. Aminoglycoside levels in the rabbit vitreous cavity after intravenous administration. *Am J Ophthalmol* 1996;122(5):684–689.
- [44] Morlet N, Graham GG, Gatus B, McLachlan AJ, Salonikas C, Naydoo D, et al. Pharmacokinetics of Ciprofloxacin in the Human Eye: a Clinical Study and Population Pharmacokinetic Analysis. *Antimicrob Agents Chemother* 2000;44(6):1674–1679.
- [45] Cekic O, Batman C, Yasar Ü, Basci N, Bozkurt A, Kayaalp S. Human aqueous and vitreous humour levels of ciprofloxacin following oral and topical administration. *Eye*, 1999;13:555–558.
- [46] El Baba F, Trousdale M, Gauderman W, Wagner D, Liggett P. Intravitreal penetration of oral ciprofloxacin in humans. *Ophthalmology* 1992;99(4):483–486.
- [47] Keren G, Alhalel A, Bartov E, Kitzes-Cohen R, Rubinstein E, Segev S, et al. The intravitreal penetration of orally administered ciprofloxacin in humans. *Invest Ophthalmol Vis Sci* 1991;32(8):2388–2392.
- [48] Sweeney G, Fern A, Lindsay G, Doig M. Penetration of ciprofloxacin into the aqueous humour of the uninflamed human eye after oral administration. *J Antimicrob Chemother* 1990;26(1):99–105.

- [49] Mounier M, Adenis J, Denis F. Intraocular penetration of ciprofloxacin after infusion and oral administration. *Pathol Biol* 1988;36(5):724–727.
- [50] Madu AA, Mayers M, Perkins R, Liu W, Drusano GL, Aswani R, et al. Aqueous and vitreous penetration of ciprofloxacin following different modes of systemic administration. *Exp Eye Res* 1996;63(2):129–136.
- [51] Fiscella R, Nguyen TKP, Cwik MJ, Philpotts BA, Friedlander D, Alter M et al. Aqueous and vitreous penetration of levofloxacin after oral administration. *Ophthalmology* 1999;106:2286–2290.
- [52] Herbert EN, Pearce IA, McGalliard J, Wong D, Groenewald C. Vitreous penetration of levofloxacin in the uninflamed phakic human eye. *Br J Ophthalmol* 2002;86(4):387–389.
- [53] Hariprasad S, Shah GK, Mieler WF, Feiner L, Blinder KJ, Holekamp NM, et al. Vitreous and Aqueous Penetration of Orally Administered Moxifloxacin in Humans. *Arch Ophthalmol* 2006;124:178–182.
- [54] Fuller J, Lott MN, Henson N, Bhatti A, Singh H, McGwin G, et al. Vitreal penetration of oral and topical moxifloxacin in humans. *Am J Ophthalmol* 2007;143(2):338–340.
- [55] Lott MN, Fuller J, Hancock H, Singh J, Singh H, McGwin G, et al. Vitreal penetration of oral moxifloxacin in humans. *Retina* 2008;28(3):473–476.
- [56] Vedantham V, Lalitha P, Velpandian T, Ghose S, Mahalaksmi R, Ramasamy K. Vitreous and aqueous penetration of orally administered moxifloxacin in humans. *Eye* 2006;20:1273–1278.
- [57] Alffenaar JWC, van Altena R, Bökkerink HJ, Luijckx GJ, van Soolingen D, Aarnoutse RE, et al. Pharmacokinetics of Moxifloxacin in Cerebrospinal Fluid and Plasma in Patients with Tuberculous Meningitis. *Clin Infect Dis* 2009;49(7):1080–1082.
- [58] Ruslami R, Rizal Ganiem A, Dian S, Apriani L, Achmad TH, van der Ven AJ, et al. Intensified regimen containing rifampicin and moxifloxacin for tuberculous meningitis: An open-label, randomised controlled phase 2 trial. *Lancet Infect Dis* 2013;13(1):27–35.
- [59] Feiz V, Nijm L, Glickman R, Morse L, Telander D, Park S, et al. Vitreous and aqueous penetration of orally administered trimethoprim-sulfamethoxazole combination in humans. *Cornea* 2013;2(10):1315–1320.
- [60] Goodwin C, Bucens M, Davis R, Norcott T. High-dose co-trimoxazole and its penetration through uninflamed meninges. *Med J Aust* 1981;2(1):24–27.
- [61] Al-Sibai MB, Al-Kaff AS, Raines D, El-Yazigi A. Ocular penetration of oral clarithromycin in humans. *J Ocul Pharmacol Ther* 1998;14(6):575–583.

table 1. Studies evaluating vitreous concentrations after systemic administration of antibiotics

antibiotic	model	operative status ^a	inflammation status ^b	subject number	dose	mean Cmax after 1 dose (ug/ml)	V/S	vitreous antibiotic level > MIC ^c	references
Vancomycin	rabbit	ph/aph/a-v	inf+non-inf	58	15 mg/kg iv	5.4 (a-v inf), 1.1 (aph inf), 0.0 (ph inf)		yes in a-v inf	Meredith 1994 [21]
Ampicillin	rabbit	ph	non-inf	49	50 mg/kg iv	0.1		no	Salminen 1978 [23]
Amoxicillin	rabbit	ph	non-inf	52	50/500 mg/kg iv (!)	0.2/1.6	0.02		Faigenbaum [24]
Piperacillin	human	ph/ps/aph	inf+non-inf	45	4 g iv	undetectable		no	Robinet 1998 [25]
Cefazoline	rabbit	ph/a-v	inf+non-inf	40	50 mg/kg iv	6.7 (a-v inf), 3.0 (ph inf)		yes in a-v inf	Martin 1990 [26]
Ceftriaxon	human	ph	non-inf	17	2g im bid	5.1 after 6 doses	0.04	partially	Sharir 1998 [27]
Ceftazidime	rabbit	ph/aph/a-v	inf+non-inf	46	50 mg/kg iv	35.4 (a-v inf), 0 (aph inf, ph inf)	0.3	partially	Aguilar 1995 [29]
Ceftazidime	rabbit	ph	non-inf	15	50 mg/kg iv	<1		no	Walstad 1985 [30]
Cefepime	human	ph	non-inf	30	1g/2g iv	1.9/2.9	0.08	no	Aras 2002 [31]
Imipenem	human	ph/ps	non-inf	10	0.5g/1g iv	0.2/1.1	0.08	partially	Adenis 1994 [32]
Imipenem	human	ph	non-inf	10	1g iv	2.5	0.1	partially	Axelrod 1987 [33]
Meropenem	human	ph	non-inf	14	2g iv	8.9	0.3	yes	Schauersberger 1999 [34]
Rifampicin	rabbit	ph	non-inf	?	150/300/600mg po (!)	2.2/2.6/15.2		yes	Wong 1990 [35]
Linezolid	human	ph	non-inf	29	600mg po	2.3	0.3	partially	Fiscella 2004 [37]
Linezolid	human	ph	non-inf	16	600mg po	3.7	0.8	yes	George 2010 [38]
Linezolid	human	ph	non-inf	24	600mg iv/po	3.7	0.6	yes	Horcajada 2008 [39]
Linezolid	human	ph	non-inf	12	600mg po	1.2	0.1	partially	Ciulla 2005 [40]
Daptomycin	human	ph	inf	1	10 mg/kg iv	12.4	0.3	yes	Sheridan 2010 [42]
Amikacin	rabbit	a-v	inf	7	6 mg/kg iv	8.5		no	El-Massry 1996 [43]
Gentamycin	rabbit	a-v	inf	7	1.6 mg/kg iv	1.8		no	El-Massry 1996 [43]
Ciprofloxacin	human	ph	non-inf		various	0.1-0.5		no	Morlet 2000 [44], et al. [45-50]
Levofloxacin	human	ph	non-inf	45	500mg po od/bid	0.6; 2.5 after 2 doses	0.3	no	Fiscella 1999 [51]
Levofloxacin	human	ph	non-inf	10	500 mg po	0.8	0.3	no	Herbert 2002 [52]
Levofloxacin	human	ph	non-inf	16	750 mg po	2.8	0.5	partially	George 2010 [38]
Moxifloxacin	human	ph	non-inf	13	2x400 mg po bid	1.3 after 2 doses	0.4	yes	Hariprasad 2006 [53]
Moxifloxacin	human	ph/ps	non-inf	8	2x400 mg po bid	1.5 after 2 doses		yes	Fuller 2007 [54]
Moxifloxacin	human	ph/ps	non-inf	21	1x400 mg po	0.6		yes	Lott 2008 [55]
Moxifloxacin	human	ph/ps	non-inf	27	1x400 mg po	0.1	0.1	no	Vedantham 2006 [56]
Trimethoprim	human	ph	non-inf	10	160 mg po bid	1.8 after 3 doses	0.40	partially	Feiz 2013 [60]
Sulfamethoxazole	human	ph	non-inf	10	800 mg po bid	5.9 after 3 doses	0.15		Feiz 2013 [60]
Clarithromycin	human	ph	non-inf	21	500 mg po bid	0.3 after 6 doses	0.2	no	Al-Sibai 1998 [61]

a) ph= phakic, ps = pseudophakic, aph = aphakic, a-v = aphakic-vitreotomized

b) inf = inflamed, non-inf = non-inflamed

c) refers to MICs of wild-type pathogens covered by the specific antibiotic agent

d) coverage dependent on species (see text for details)

(!) exceeding human doses

V/S = vitreous/serum antibiotic concentration ratio

ACCEPTED MANUSCRIPT

table 2. EUCAST MIC breakpoints (susceptible \leq) of highly prevalent or otherwise significant organisms in bacterial endophthalmitis (ug/ml). If breakpoints are not defined, ECOFFs (epidemiological cut-off values) are given if reasonable (*).

antibiotic	Str. group A/B/C/G	Str. <i>pneumoniae</i>	Str. viridans group	coagulase-neg. staphylococci	<i>S.aureus</i>	Enterococci	<i>Haemophilus influenzae</i>	Entero- bacteriaceae	<i>Pseudomonas aeruginosa</i>	observed maximum levels (ug/ml)
Vancomycin	2	2	2	4	2	4	NA	NA	NA	5.4
Benzylpenicillin	0.25	0.06	0.25	[0.125]	[0.125]	NA	NA	NA	NA	
Ampicillin, Amoxicillin	a)	0.5	0.5	a)	a)	4	1	[8]	NA	0.1 (- 1.6)
Piperacillin-Tazobactam	a)	a)	a)	b)	b)	c)	c)	8	16	0
Cefazoline ¹	a)	NA	0.5	b)	b)	NA	NA	NA	NA	3.0-6.7
Ceftriaxone	a)	0.5	0.5	b)	8*	NA	0.125	1	NA	5.1
Ceftazidime	NA	NA	NA	NA	NA	NA	NA	1	8	0-35.4
Cefepime	a)	1	0.5	b)	8*	NA	0.25	1	8	2.9
Imipenem	a)	2	2	0.125*	0.125*	4	2	2	4	1.1-2.5
Meropenem	a)	2	2	0.5*	0.5*	NA	2	2	2	8.9
Rifampicin	0.06		NA	0.06	0.06	NA	NA	NA	NA	15.2
Linezolid	2	2	NA	4	4	4	NA	NA	NA	1.2-3.7
Daptomycin	1	NA	NA	1	1	4 ^{d)}	NA	NA	NA	12.4
Amikacin	NA	NA	NA	8	8	NA	NA	8	8	8.5
Gentamycin	NA	NA	NA	1	1	NA	NA	2	4	1.8
Ciprofloxacin	NA	NA	NA	1	1	[4]	0.06	0.25	0.5	0.1-0.5
Levofloxacin	2	2	NA	1	1	[4]	0.06	0.5	[1]	0.6-2.8
Moxifloxacin	0.5	0.5	NA	0.25	0.25	NA	0.125	0.25	NA	0.1-1.5
TMP-SMX ²	1	1	NA	2	2	[0.03]	0.5	2	NA	1.8
Clarithromycin	0.25	0.25	NA	1	1	NA	NA	NA	NA	0.3

a) inferred from benzylpenicillin susceptibility

b) inferred from oxacillin/cefoxitin susceptibility

c) inferred from ampicillin susceptibility

d) CLSI breakpoint

¹ cefazoline non-species-related breakpoint 1ug/ml

² breakpoints are expressed as TMP concentration with a 1:19 TMP-SMX ratio

* ECOFF given; no breakpoints defined

[] square brackets: unfavourable therapy due to resistance rate and/or insufficient evidence for efficacy

NA not applicable