Do clinical investigations predict long-term wheeze? A follow-up of paediatric respiratory outpatients

Carmen CM. de Jong¹, Eva SL. Pedersen¹, Myrofora Goutaki¹, Daniel Trachsel³, Juerg Barben⁴, Claudia E. Kuehni^{1,2}

¹ Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland

² Children's University Hospital of Bern, University of Bern, Bern, Switzerland

³ Paediatric Respiratory Medicine, Children's University Hospital of Basel, Basel, Switzerland

⁴ Paediatric Respiratory Medicine, Children's Hospital of Eastern Switzerland, St. Gallen, Switzerland

Correspondence to

Claudia E. Kuehni

Institute of Social and Preventive Medicine

Mittelstrasse 43, CH-3012 Bern, Switzerland

Tel.: +41 (0)31 631 35 07

E-mail: claudia.kuehni@ispm.unibe.ch

Key words: Asthma, cohort, epidemiology, prognosis, respiratory, wheeze

Number of figures and tables: 4

Material in the electronic repository: 4 tables, 1 figure, 1 supplementary text and 4 supplementary questionnaires

Running title: Predicting wheeze: a follow-up study (max. 50 characters)

Do clinical investigations predict long-term outcome?

A follow-up of paediatric respiratory outpatients

de Jong CCM, Pedersen ESL, Goutaki M, Trachsel M, Barben J, Kuehni CE

Abstract (250/250)

Introduction: The contribution of clinical investigations to prediction of long-term outcomes of children investigated for asthma is unclear.

Aim: We performed a broad range of clinical tests and investigated whether they helped to predict long-term wheeze among children referred for evaluation of possible asthma.

Methods: We studied children aged 6-16 years referred to two Swiss pulmonary outpatient clinics with a history of wheeze, dyspnoea, or cough in 2007. The initial assessment included spirometry, fractional exhaled nitric oxide, skin prick tests, and bronchial provocation tests (BPT) by exercise, methacholine, and mannitol. Respiratory symptoms were assessed with questionnaires at baseline and at follow-up seven years later. Associations between baseline factors and wheeze at follow-up were investigated by logistic regression.

Results: At baseline, 111 children were examined in 2007. Seven years after baseline, 85 (77%) completed the follow-up questionnaire, among whom 61 (72%) had wheeze at baseline, while at follow-up 39 (46%) reported wheeze. Adjusting for age and sex, the following characteristics predicted wheeze at adolescence: wheeze triggered by pets (odds ratio 4.2, 95% CI 1.2-14.8), pollen (2.8, 1.1-7.0), and exercise (3.1, 1.2-8.0). Of the clinical tests, only a positive exercise test (3.2, 1.1-9.7) predicted wheeze at adolescence.

Conclusion: Reported exercise-induced wheeze and wheeze triggered by pets or pollen were important predictors of wheeze persistence into adolescence. None of the clinical tests predicted wheeze more strongly than reported symptoms. Clinical tests might be important for asthma diagnosis but medical history is more helpful in predicting prognosis in children referred for asthma.

Introduction

Asthma is the most prevalent chronic respiratory disease in childhood and adolescence, which leads to many health care visits¹⁻³. Its key symptoms are wheeze, cough, and difficulty breathing, but symptoms vary substantially between individuals and across ages^{1,2}. Some children who present with asthma symptoms continue to have problems later in life, while others do not. Better knowledge of their individual prognoses might affect their follow-up and answer questions of parents in the clinics⁴⁻⁶. Assessing prognosis of asthma symptoms from school age into adulthood and identifying children at high-risk of symptom persistence is challenging⁴.

Studies investigating prognosis of asthma or wheeze in school-aged children are conducted with either clinical asthma cohorts or symptomatic children of a population-based cohort⁷.

Studies in clinical asthma cohorts have found that lower FEV₁, asthma severity, sensitisation to indoor allergens, eczema, hay fever, skin test reactivity, and bronchial hyper-responsiveness were associated with asthma persistence⁸⁻¹⁰. Studies in population-based cohorts have found that wheeze persistence was predicted by frequent attacks of wheeze, female sex, sensitization to furred animals or house dust mites, rhinitis, and bronchial hyper-responsiveness¹¹⁻¹⁶.

For clinical practice, two knowledge gaps remain. First, few studies have examined the prediction of long-term prognosis, but none have done this for school-aged children seen in outpatient clinics for possible asthma. Second, many tests are performed in clinics to diagnose these children, but it is unclear whether these tests predict prognosis more accurately than reported symptoms alone. We determined whether clinical tests in addition to reported symptoms help predict wheeze in adolescence in school-aged children referred for possible asthma.

Methods

Study population and study design

Of the 124 children invited, 111 were recruited from the respiratory outpatient clinics of two paediatric hospitals in Switzerland, 84 from St. Gallen and 27 from Basel, who were eligible if they had been referred for evaluation of current wheeze, dyspnoea, or cough. Children with a known chronic respiratory disease such as cystic fibrosis or primary ciliary dyskinesia, or a respiratory tract infection during four weeks prior to the visit were excluded. At baseline in 2007-2008, parents completed a questionnaire and children underwent a set of standardised clinical tests during two different visits within one week as part of the study protocol^{17,18}. At follow-up, seven years after baseline, in 2014 to 2015, we sent a questionnaire to the 12-23 year-old adolescents or young adults (from now on referred to as adolescents) (E-figure 1).

Ethical approval was obtained from the local Ethics committee and all parents gave informed consent during the first visit at baseline and by sending back the questionnaire at follow-up (EKSG 07/001).

Baseline assessment

The parental questionnaire included ISAAC key questions¹⁹ plus additional questions on type and triggers of respiratory symptoms, atopic symptoms, previous treatments and environmental exposures (Supplementary questionnaire 1 (German, original) and 2 (English, translation)). The study physician reported clinical test results, final diagnosis, and prescribed medication in a uniform way. Physicians diagnosed the children after all clinical tests were done, taking into consideration medical history, clinical examination, and all test results. Vocal cord dysfunction was diagnosed based on medical history, physical examination and normal expiratory curves in spirometry.

The baseline assessment consisted of two visits. At the initial baseline visit, children performed spirometry, fractional exhaled nitric oxide (FeNO) measurement, a skin prick test (SPT), bronchial provocation test (BPT) by exercise and, by methacholine. At the second

baseline visit, children did a BPT by mannitol. All clinical tests were performed according to published guidelines²⁰⁻²⁴. A detailed description of the test procedures has been published elsewhere^{17,18,25} and is included in the online supplementary material (E-text). Lung function measurements were compared to reference values from Zapletal et al²⁶. We considered the exercise test as positive in the event of a $\geq 15\%$ decrease in the FEV₁ after the exercise challenge test, and the methacholine test as positive when the minimal dose causing a $\geq 20\%$ decrease of FEV₁ was $< 1\text{mg}$ (the provocation dose, PD 20). The mannitol dry powder challenge test was considered as positive when a 15% fall in FEV₁ was measured before a cumulative dose of 635 mg was reached, or when a 10% fall in FEV₁ between two doses was reached. FeNO was measured using the portable NIOX MINO® device, and was considered as positive when FeNO was higher than 26ppb¹⁸. We performed skin prick tests for birch, grass, mugwort, alternaria, cat, house dust mites (*D. pteronyssinus*), and positive and negative controls¹⁸. These allergens cover 95% of inhaled allergens in Switzerland²⁷. The test was considered to be positive if any mean wheal diameter was $\geq 3\text{mm}$.

Assessment at follow-up

The follow-up questionnaire was very similar to the baseline questionnaire, but the questions were addressed directly to the adolescents instead of their parents (supplementary questionnaire 3 (German, original) and 4 (English, translation)).

Definitions of wheeze and frequent wheeze

We assessed wheeze at follow-up with the question, "Have you had a whistling sound in the chest in the last 12 months?" If a child had had more than three attacks of wheeze in the last 12 months, we considered the child to have had frequent wheeze.

Statistical analysis

We compared the participants with information at baseline and follow-up to those without follow-up information to test for selection bias, using chi-square test. The participants with

information at baseline and follow-up were included in the analysis.

We investigated the association between exercise-induced wheeze and a positive exercise test at baseline using the Fisher's exact test, and the Mann-Whitney-U test when looking at the association of reported exercise-induced wheeze and the fall of FEV₁% predicted during the exercise test.

We investigated the association between symptoms (table 1) and clinical test results (table 2) at baseline with any wheeze and frequent wheeze at follow-up using logistic regression, adjusting for sex and age. For comparison, we repeated the analysis including only children diagnosed with asthma (N=62). We did not consider interactions or a multivariable model because of the sample size. We used STATA software (version 14; College Station, Texas) to analyse the data.

Results

Characteristics of the study population at baseline and at follow-up

Eighty-five (77%) of the 111 children who participated in the baseline study completed the follow-up questionnaire. The median age was 12 years at baseline (range 6-16) and 18 at follow-up (12-23); 60% (51/85) were male. Wheeze was reported by 61 (72%) at baseline, and 7 reported cough without wheeze, 12 (14%) reported exercise-related breathing problems and 5 (6%) reported allergic rhinitis. Among those with wheeze, 27 (44%) had more than three attacks during 12 months prior to the baseline visit (Table 1). Symptoms at baseline were very similar in children who did not take part in the follow-up (E-table 1 online supplementary material). Asthma medication was prescribed at the baseline visit for 71 (85%) children, of whom 47 (55%) received inhaled short-acting β 2-agonists (SABA) alone, 6 received SABA and inhaled corticosteroids (ICS), and 18 received long-acting β 2-agonists (LABA) and ICS. At follow-up, 39 (46%) participants reported wheeze of whom 30 had more than 3 attacks during the last year. At follow-up, 44 adolescents (52%) reported using

inhalers, including 21 using SABA alone, 2 using SABA and ICS, and 21 using LABA and ICS (Table 1).

Table 2 shows the clinical test results and diagnoses at baseline. All tests were completed in at least 90% of the children. The main reason for not completing a BPT was exhaustion^{17,18}. For the 78 children who completed the BPT by methacholine at baseline, the test was positive in 76% and the median provocation dose was 0.14mg. Eighty-two completed the BPT by mannitol, of whom 28% tested positive. The median provocation dose was 635 mg. Of the 76 children who completed the BPT by exercise, the median fall of FEV₁ was 8% predicted. The test was positive ($\geq 15\%$ decrease in the FEV₁) in 18 (24%) children. SPT was positive in 33 (39%) children and FeNO was positive in 35 (41%). Doctors diagnosed 62 (73%) children with asthma or episodic viral wheeze. The other children were mostly diagnosed with cough not due to asthma or vocal cord dysfunction. At baseline, self-reported exercise-induced wheeze was associated with a positive exercise test ($p=0.022$, E-table 2).

Baseline factors associated with wheeze and frequent wheeze at follow-up

Four respiratory symptoms and one clinical test at baseline were associated with *any* wheeze at follow-up. Of the reported symptoms, frequent wheeze (>3 attacks) (OR 2.86, 95% CI 1.10-7.43), exercise-induced wheeze (3.07, 1.19-7.96), wheeze triggered by pets (4.22, 1.21-14.76), and wheeze triggered by pollen (2.78, 1.11-6.98) were associated with wheeze at follow-up. For the clinical tests, only a positive exercise test was significantly associated with wheeze seven years later (3.20, 1.05-9.70). Results remained very similar after adjusting for age and sex (Table 3). When we repeated the analysis for children diagnosed with asthma (N=62), we found mostly comparable results (E-table 3). However, associations tended to be less strong (lower odds ratios) in particular for exercise induced wheeze (1.79, 0.58-5.48) and positive exercise test (2.00, 0.63-6.39)

Two respiratory symptoms were associated with *frequent wheeze* at follow-up. These were exercise-induced wheeze (OR 3.05, 95% CI 1.07-8.67) and wheeze triggered by pets (3.79, 1.15-12.48; E-table 4). None of the clinical test results were associated with frequent wheeze at follow-up.

Discussion

Among school-aged children referred to a respiratory outpatient clinic for evaluation of wheeze, cough, or dyspnoea, 46% reported wheeze seven years later. Reported exercise-induced wheeze and wheeze triggered by pets or pollen at baseline predicted wheeze at follow-up. Of the clinical tests, only a positive exercise challenge test predicted wheeze at follow-up, but no more strongly than reported exercise-induced wheeze. When we repeated the analysis based on children with asthma only, associations were weaker, probably because the same characteristics that predicted persistence (exercise induced wheeze, positive exercise test) had already been used by the clinicians to decide on a diagnosis of asthma.

A few studies have examined the prediction of prognosis by clinical testing, but ours is the only study to have done this for so many clinical tests in school-aged children referred to a respiratory outpatient clinic. We did not find an association between FEV₁ or bronchial provocation test by methacholine at baseline and wheeze seven years later; previous studies have reported contradictory findings. Both the CAMP cohort of 909 children aged 5-12 years with diagnosed asthma and another Dutch clinical cohort study of 5-14 year-old children diagnosed with asthma found that asthma persistence at ages 15-20 and 32-42, respectively, was associated with decreased FEV₁ at school-age^{8,9}. In contrast, the population-based Tasmanian cohort did not find an association between FEV₁ at age 7 and wheeze persistence at age 29-32¹⁵. The CAMP study also found that a lower methacholine provocation concentration was associated with asthma persistence from age 5-12 until age 15-20⁹. In contrast, the population based Dunedin cohort of 613 children reporting wheeze at age 9 and a Norwegian cohort of 62 children reporting asthma at age 10, found that

233 bronchial provocation test by methacholine was not associated with persistence at age 26
234 and 16 respectively, which is in line with our findings^{12,28}. The Norwegian cohort did not find
235 bronchial provocation test by exercise to be associated with asthma persistence from age 10
236 to age 16, which is in contrast to the association we found²⁸. This heterogeneity between
237 studies could be because children with wheeze from population-based cohorts might have
238 milder disease than those in clinical studies. FeNO was not associated with wheeze
239 persistence in children aged 6-16 years old suspected for asthma in our study. In contrast,
240 FeNO was reported to predict asthma in pre-school children with wheeze²⁹⁻³². To our
241 knowledge, no studies assess the predictive value of FeNO on wheeze persistence at school
242 age. Available publications assessed the predictive value of FeNO on asthma control,
243 relapse or exacerbations in asthmatic children, but with a short follow-up³³⁻³⁶. .
244 Our observation that frequent attacks of wheeze at school age predicted wheeze persistence
245 seven years later is in line with findings from the Melbourne and Tasmanian cohorts^{10,11}. In
246 contrast to their findings, we found no significant association between either eczema or hay
247 fever at baseline and wheeze persistence. This could be because those cohorts used
248 different outcomes—severe wheeze and atopic asthma, respectively—or simply because we
249 had low numbers and limited power.

250 A possible limitation of our study was that the bronchial provocation tests were done within a
251 short period of time. This could have influenced the methacholine test result, which was
252 performed after the exercise test on the same day and was positive in 76% of the children.
253 Most likely the bronchial provocation test by mannitol was not influenced by the short time
254 interval. We assured an appropriate interval of at least 24 hours without a change in
255 respiratory health or medication in this time interval. A second limitation was the small
256 sample size, which limited statistical power and did not allow us to perform a multivariable
257 analysis including all symptoms and test results simultaneously. Adolescents might have
258 underreported respiratory symptoms, which might have led to an underestimation of the
259 proportion of adolescents with wheeze. However, since this underreporting is not likely to be

associated with symptoms or positive test results at baseline, this should not have influenced the results relating to risk factors.

The main strength of our study is its clinical design, which reflects the typical mix of patients in a paediatric outpatient clinic. All children were first-time referrals to the paediatric respiratory clinic for evaluation of possible asthma. Therefore, the study population is representative of daily clinical work, in contrast to many clinical studies that selectively include well-defined moderate to severe asthmatics and leave out patients with unclear degrees of airway reactivity. Our study also profited from a very detailed baseline examination. Children in the study had an extensive array of examinations for lung function, BPT and allergy, which allowed us to assess the contribution of clinical tests in predicting long-term wheeze in addition to reported symptoms among those referred for evaluation of possible asthma.

Conclusion

This study is an initial step towards finding out whether clinical tests can predict wheeze later in life. Though clinical tests might be important for asthma diagnosis, our results suggest that they do not strongly predict prognosis of wheeze. In contrast, our data underline the importance of a detailed history, as school-age children reporting exercise-related wheeze and wheeze triggered by allergens were at higher risk and thus might profit from more frequent follow-up.

General acknowledgements

We thank all participants and lab technicians of the pulmonology department in the children's hospitals in Basel and St. Gallen for their assistance in our study, Marie-Pierre Strippoli (ISPM, Bern) for her work on the study at baseline, Bettina Meier (ISPM, Bern) for entering the follow-up questionnaires into the database and sending the participation reminders. We thank Christopher Ritter (ISPM, Bern) for his editorial assistance and Niels Hagenbuch (ISPM, Bern) for his statistical support.

288

289 **Funding**

290 This study was funded by the Swiss National Science Foundation: 32003B_162820 and by
291 AstraZeneca (Switzerland), the Lung League St. Gallen, and the Schmidheiny Foundation
292 (Heerbrugg, St. Gallen).

293

294 **Author contributions**

295 Claudia Kuehni and Jürg Barben conceptualised and designed the study. Daniel Trachsel
296 and Jürg Barben supervised data collection. Carmen de Jong analysed the data and drafted
297 the manuscript. Eva Pedersen and Myrona Goutaki supported the statistical analysis and
298 gave input for interpretation of the data. All authors critically revised the manuscript and
299 approved the final manuscript as submitted.

300

301

References

1. Braig S, Brandt S, Wabitsch M, Florath I, Brenner H, Rothenbacher D, Genuneit J. Age-specific influence of wheezing phenotypes on pre-adolescent and adolescent health-related quality of life. *Pediatr Allergy Immunol.* 2014;25(8):781-787.
2. Jurca M, Pescatore AM, Goutaki M, Spycher BD, Beardsmore CS, Kuehni CE. Age-related changes in childhood wheezing characteristics: A whole population study. *Pediatr Pulmonol.* 2017;52(10):1250-1259.
3. Stevens CA, Turner D, Kuehni CE, Couriel JM, Silverman M. The economic impact of preschool asthma and wheeze. *Eur Respir J.* 2003;21(6):1000-1006.
4. Fuchs O, Bahmer T, Rabe KF, von Mutius E. Asthma transition from childhood into adulthood. *Lancet Respir Med.* 2017;5(3):224-234.
5. Spycher BD, Cochrane C, Granell R, Sterne JAC, Pedersen E, Gaillard EA, Henderson J, Kuehni CE. Temporal stability of multiple trigger and episodic viral wheeze in early childhood. *Eur Respir J.* 2017;50(5):1700014.
6. van Wonderen KE, Geskus RB, van Aalderen WM, Mohrs J, Bindels PJ, van der Mark LB, Ter Riet G. Stability and predictiveness of multiple trigger and episodic viral wheeze in preschoolers. *Clin Exp Allergy.* 2016;46(6):837-847.
7. Sears MR. Predicting asthma outcomes. *J Allergy Clin Immunol.* 2015;136(4):829-836.
8. Vonk JM, Postma DS, Boezen HM, Grol MH, Schouten JP, Koeter GH, Gerritsen J. Childhood factors associated with asthma remission after 30 year follow up. *Thorax.* 2004;59(11):925-929.
9. Covar RA, Strunk R, Zeiger RS, Wilson LA, Liu AH, Weiss S, Tonascia J, Spahn JD, Szeftler SJ. Predictors of remitting, periodic, and persistent childhood asthma. *J Allergy Clin Immunol.* 2010;125(2):359-366.
10. Wolfe R, Carlin JB, Oswald H, Olinsky A, Phelan PD, Robertson CF. Association between allergy and asthma from childhood to middle adulthood in an Australian cohort study. *Am J Respir Crit Care Med.* 2000;162(6):2177-2181.
11. Martin PE, Matheson MC, Gurrin L, Burgess JA, Osborne N, Lowe AJ, Morrison S, Mészáros D, Giles GG, Abramson MJ, et al. Childhood eczema and rhinitis predict atopic but not nonatopic adult asthma: a prospective cohort study over 4 decades. *J Allergy Clin Immunol.* 2011;127(6):1473-1479.
12. Sears MR, Greene JM, Willan AR, Wiecek EM, Taylor DR, Flannery EM, Cowan JO, Herbison GP, Silva PA, Poulton R. A longitudinal, population-based, cohort study of childhood asthma followed to adulthood. *N Engl J Med.* 2003;349(15):1414-1422.
13. Andersson M, Hedman L, Bjerg A, Forsberg B, Lundback B, Ronmark E. Remission and persistence of asthma followed from 7 to 19 years of age. *Pediatrics.* 2013;132(2):e435-442.
14. Burgess JA, Walters EH, Byrnes GB, Matheson MC, Jenkins MA, Wharton CL, Johns DP, Abramson, Hopper JL, Dharmage SC. Childhood allergic rhinitis predicts asthma incidence and persistence to middle age: a longitudinal study. *J Allergy Clin Immunol.* 2007;120(4):863-869.
15. Jenkins MA, Hopper JL, Bowes G, Carlin JB, Flander LB, Giles GG. Factors in childhood as predictors of asthma in adult life. *BMJ.* 1994;309(6947):90-93.
16. Oswald H, Phelan PD, Lanigan A, Hibbert M, Bowes G, Olinsky A. Outcome of childhood asthma in mid-adult life. *BMJ.* 1994;309(6947):95-96.
17. Barben J, Kuehni CE, Strippoli MP, Schiller B, Hammer J, Trachsel D, Swiss Paediatric Respiratory Research Group. Mannitol dry powder challenge in comparison with exercise testing in children. *Pediatr Pulmonol.* 2011;46:842-848.
18. Barben J, Strippoli MP, Trachsel D, Schiller B, Hammer J, Kuehni CE. Effect of Mannitol dry powder challenge on exhaled nitric oxide in children. *PLoS One.* 2013;8(1):e54521.

19. Asher MI, Keil U, Anderson HR, Beasley R, Crane J, Martinez F, Mitchell EA, Pearce N, Sibbald B, Stewart AW, et al. International study of asthma and allergies in childhood (ISAAC): rationale and methods. *Eur Respir J*. 1995;8:483-491.
20. Crapo RO, Casaburi R, Coates AL, Enright PL, Hankinson JL, Irvin CG, MacIntyre NR, McKay RT, Wanger JS, Anderson SD, et al. Guidelines for methacholine and exercise challenge testing-1999. *Am J Respir Crit Care Med*. 2000;161(1):309-329.
21. Anderson SD, Brannan J, Spring J, Spalding N, Rodwell LT, Chan K, Gonda I, Walsh A, Clark AR. A new method for bronchial-provocation testing in asthmatic subjects using a dry powder of mannitol. *Am J Respir Crit Care Med*. 1997;156(3.1):758-765.
22. Standardization of Spirometry, 1994 Update. American Thoracic Society. *Am J Respir Crit Care Med*. 1995;152(3):1107-1136.
23. Recommendations for standardized procedures for the on-line and off-line measurement of exhaled lower respiratory nitric oxide and nasal nitric oxide in adults and children-1999. This official statement of the American Thoracic Society was adopted by the ATS Board of Directors, July 1999. *Am J Respir Crit Care Med*. 1999;160(6):2104-2117.
24. Barben J RJ. Measurement of bronchial responsiveness in children. In: Hammer J, Eber E, editors. Paediatric pulmonary function testing. *Prog Respir Res*. Basel: Karger;2005;33:125-136.
25. Barben J, Roberts M, Chew N, Carlin JB, Robertson CF. Repeatability of bronchial responsiveness to mannitol dry powder in children with asthma. *Pediatr Pulmonol*. 2003;36(6):490-494.
26. Zapletal A, Paul T. Lung function in children and adolescents. Methods, reference values. *Prog Respir Res*. 1987;22:1-220.
27. Braun-Fahrlander C, Wüthrich B, Gassner M, Gritze I, Neu U, Varonier H. Prävalenz und Risikofaktoren einer allergischen Sensibilisierung bei Schulkindern in der Schweiz. *Allergologie*. 1999;22:54-64.
28. Riiser A, Hovland V, Carlsen KH, Mowinckel P, Lodrup Carlsen KC. Does bronchial hyperresponsiveness in childhood predict active asthma in adolescence? *Am J Respir Crit Care Med*. 2012;186(6):493-500.
29. Pijnenburg MW. The Role of FeNO in Predicting Asthma. *Front Pediatr*. 2019;7:41.
30. Singer F, Luchsinger I, Inci D, Knauer N, Latzin P, Wildhaber JH, Moeller A. Exhaled nitric oxide in symptomatic children at preschool age predicts later asthma. *Allergy*. 2013;68(4):531-538.
31. Elliott M, Heltshe SL, Stamey DC, Cochrane ES, Redding GJ, Debley JS. Exhaled nitric oxide predicts persistence of wheezing, exacerbations, and decline in lung function in wheezy infants and toddlers. *Clin Exp Allergy*. 2013;43(12):1351-1361.
32. Caudri D, Wijga AH, Hoekstra MO, Kerkhof M, Koppelman GH, Brunekreef B, Smit HA, de Jongste JC. Prediction of asthma in symptomatic preschool children using exhaled nitric oxide, Rint and specific IgE. *Thorax*. 2010;65(9):801-807.
33. Lehtimäki L, Csonka P, Mäkinen E, Isojärvi J, Hovi SL, Ahovuo-Saloranta A. Predictive value of exhaled nitric oxide in the management of asthma: a systematic review. *Eur Respir J*. 2016;48(3):706-714.
34. Pijnenburg MW, Hofhuis W, Hop WC, De Jongste JC. Exhaled nitric oxide predicts asthma relapse in children with clinical asthma remission. *Thorax*. 2005;60(3):215-218.
35. Visitsunthorn N, Mahawichit N, Maneechotesuwan K. Association between levels of fractional exhaled nitric oxide and asthma exacerbations in Thai children. *Respirology*. 2017;22(1):71-77.
36. Yang S, Park J, Lee YK, Kim H, Hahn YS. Association of longitudinal fractional exhaled nitric oxide measurements with asthma control in atopic children. *Respir Med*. 2015;109(5):572-579.

402 **Supplementary data**

403 **E-table 1** Comparison of characteristics of the children included in the follow-up study and
404 the children that did not take part in the follow-up study
405

406 **E-table 2** Association between reported exercise-induced wheeze and exercise test result at
407 baseline N=76
408

409 **E-table 3** Association between baseline factors and wheeze at follow-up in children with
410 diagnosed asthma
411

412 **E-table 4** Association between baseline factors and frequent wheeze at follow up
413

414 **E-figure 1** Flowchart of patient recruitment at baseline and follow-up
415

416 **E-text:** methods of clinical tests

417 **Supplementary questionnaire 1:** Baseline questionnaire in German (original)

418 **Supplementary questionnaire 2:** Baseline questionnaire in English

419 **Supplementary questionnaire 3:** Follow-up questionnaire in German (original)

420 **Supplementary questionnaire 4:** Follow-up questionnaire in English