

Target journal: *International Journal of Cancer*

Title: Cervical cancer risk in women living with HIV across four continents: a multicohort study

Running head: Regional cervical cancer risk in HIV-positive women

Eliane Rohner¹, Lukas Bütikofer², Kurt Schmidlin¹, Mazvita Sengayi³, Mhairi Maskew⁴, Janet Giddy⁵, Katayoun Taghavi¹, Richard D. Moore⁶, James J. Goedert⁷, M. John Gill⁸, Michael J. Silverberg⁹, Gypsyamber D'Souza¹⁰, Pragna Patel¹¹, Jessica L. Castilho¹², Jeremy Ross¹³, Annette Sohn¹³, Firouze Bani-Sadr¹⁴, Ninon Taylor¹⁵, Vassilios Paparizos¹⁶, Fabrice Bonnet^{17,18}, Annelies Verbon¹⁹, Jörg Janne Vehreschild^{20,21}, Frank A. Post²², Caroline Sabin²³, Amanda Mocroft²³, Fernando Drona²⁴, Niels Obel²⁵, Sophie Grabar²⁶⁻²⁸, Vincenzo Spagnuolo²⁹, Eugenia Quiros-Roldan³⁰, Cristina Mussini³¹, José M. Miro³², Laurence Meyer^{33,34}, Barbara Hasse³⁵, Deborah Konopnicki³⁶, Bernardino Roca³⁷, Diana Barger¹⁸, Gary M. Clifford³⁸, Silvia Franceschi³⁹, Matthias Egger^{1,40}, and Julia Bohlius¹ for the AIDS-defining Cancer Project Working Group of IeDEA and COHERE in EuroCoord

Affiliations: ¹Institute of Social and Preventive Medicine, University of Bern, Switzerland; ²CTU Bern, University of Bern, Switzerland; ³National Cancer Registry, National Health Laboratory Service, Johannesburg, South Africa; ⁴Health Economics and Epidemiology Research Office, Department of Internal Medicine, School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa; ⁵Department of Medicine, McCord Hospital, Durban, South Africa; ⁶Johns Hopkins University, School of Medicine, Baltimore, Maryland; ⁷Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, Maryland; ⁸University of Calgary, Alberta, Canada; ⁹Division of Research, Kaiser Permanente Northern California, Oakland, USA; ¹⁰Johns Hopkins Bloomberg School of

Public Health, Baltimore, Maryland; ¹¹Division of Global HIV and TB, Centers for Disease Control and Prevention, Atlanta, Georgia; ¹²Division of Infectious Diseases, Vanderbilt University Medical Center, Nashville, USA; ¹³TREAT Asia/amfAR - The Foundation for AIDS Research, Bangkok, Thailand; ¹⁴Reims Champagne-Ardenne University, Faculté de médecine, CHU Reims, Hôpital Robert Debré, Tropical and Infectious Diseases, Reims, France; ¹⁵IIIrd Medical Department with Haematology, Medical Oncology, Haemostaseology, Infectious Diseases and Rheumatology, Oncologic Center, Paracelsus Medical University, Salzburg, Austria, Present address: Department of Dermatology, University Hospital Salzburg, Paracelsus Medical University, Salzburg, Austria; ¹⁶AIDS Unit, Clinic of Venereologic and Dermatologic Diseases, Athens Medical School, "Syngros" Hospital, Athens, Greece; ¹⁷CHU de Bordeaux, Service de Médecine Interne et Maladies Infectieuses, Hôpital Saint-André, Bordeaux, France; ¹⁸ Univ. Bordeaux, ISPED, Centre INSERM U1219-Bordeaux Population Health, F-33000 Bordeaux, France; ¹⁹Department Medical Microbiology and Infectious Diseases, Erasmus Medical Center, Rotterdam, The Netherlands; ²⁰Department I of Internal Medicine, University Hospital of Cologne, Cologne, Germany; ²¹German Centre for Infection Research, partner site Bonn-Cologne, Cologne, Germany; ²²King's College Hospital NHS Foundation Trust, London, UK; ²³Institute for Global Health, UCL, London, United Kingdom; ²⁴Department of Infectious Diseases, Hospital Ramón y Cajal, Madrid, Spain; ²⁵Department of Infectious Diseases, Copenhagen University Hospital, Copenhagen, Denmark; ²⁶Sorbonne Universités, UPMC Univ Paris 06, UMR_S 1136, Institut Pierre Louis d'Epidémiologie et de Santé Publique, F-75013, Paris, France; ²⁷INSERM, UMR_S 1136, Institut Pierre Louis d'Epidémiologie et de Santé Publique, F-75013, Paris, France; ²⁸Université Paris Descartes et Assistance Publique-Hôpitaux de Paris, Groupe hospitalier Cochin Hôtel-Dieu, Paris, France; ²⁹Vita-Salute San Raffaele University, Faculty of Medicine and Surgery, Milan, Italy;

³⁰Infectious and Tropical Diseases Institute, University of Brescia, Brescia, Italy; ³¹Infectious Diseases Clinics, University Hospital, Modena, Italy; ³²Infectious Diseases Service, Hospital Clinic – IDIBAPS, University of Barcelona, Barcelona, Spain; ³³INSERM, U1018, Epidemiology of HIV, Reproduction, Paediatrics, CESP, University Paris-Sud, Paris, France; ³⁴Department of Public Health and Epidemiology, Bicêtre Hospital, AP-HP, Le Kremlin Bicêtre, Paris, France; ³⁵Division of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, University of Zurich, Switzerland; ³⁶Department of Infectious Diseases, St Pierre University Hospital, Université Libre de Bruxelles, Brussels, Belgium; ³⁷Hospital General Universitario, Castellón, Spain; ³⁸International Agency for Research on Cancer, Lyon, France; ³⁹Centro di Riferimento Oncologico di Aviano (CRO) IRCCS, Aviano, Italy; ⁴⁰Centre for Infectious Disease Epidemiology and Research, University of Cape Town, Cape Town, South Africa.

Corresponding author: Eliane Rohner, MD, Institute of Social and Preventive Medicine, University of Bern, Mittelstrasse 43, 3012 Bern, Switzerland; e-mail: eliane.rohner@ispm.unibe.ch

Key words: Cervical cancer; HIV; incidence rate; cohort study

Novelty and impact: Our study compares invasive cervical cancer (ICC) rates among women living with HIV who initiated antiretroviral therapy (ART) across different geographic regions. It shows that ICC incidence rates are particularly high in women living with HIV in South Africa or Latin America. ICC prevention through early ART initiation, equitable access to effective cervical cancer screening, and promotion of human papillomavirus vaccination should be key elements of global efforts to reduce cancer-related health inequities.

Abbreviations: aHR, adjusted hazard ratio; ART, antiretroviral therapy; CI, confidence interval; COHERE, Collaboration of Observational HIV Epidemiological Research in Europe; HIV, human immunodeficiency virus; HPV, human papillomavirus; ICC, invasive cervical cancer; IeDEA, International epidemiology Databases to Evaluate AIDS; IQR, interquartile range; pys, person-years; NNRTI, non-nucleoside reverse transcriptase inhibitor; PI, protease inhibitor.

Abstract

We compared invasive cervical cancer (ICC) incidence rates in Europe, South Africa, Latin and North America among women living with HIV who initiated antiretroviral therapy (ART) between 1996 and 2014. We analyzed cohort data from the International epidemiology Databases to Evaluate AIDS (IeDEA) and the Collaboration of Observational HIV Epidemiological Research in Europe (COHERE) in EuroCoord. We used flexible parametric survival models to determine regional ICC rates and risk factors for incident ICC. We included 64,231 women from 45 countries. During 320,141 person-years (pys), 356 incident ICC cases were diagnosed (Europe 164, South Africa 156, North America 19, Latin America 17). Raw ICC incidence rates per 100,000 pys were 447 in South Africa (95% confidence interval [CI] 382-523), 136 in Latin America (95% CI 85-219), 76 in North America (95% CI 48-119), and 66 in Europe (95% CI 57-77). Compared with European women ICC rates at 5 years after ART initiation were more than double in Latin America (adjusted hazard ratio [aHR] 2.43, 95% CI 1.27-4.68) and 11-times higher in South Africa (aHR 10.66, 95% CI 6.73-16.88), but similar in North America (aHR 0.79, 95% CI 0.37-1.71). Overall, ICC rates increased with age (>50 years versus 16-30 years, aHR 1.57, 95% CI 1.03-2.40) and lower CD4 cell counts at ART initiation (per 100 cell/ μ l decrease, aHR 1.25, 95% CI 1.15-1.36). Improving access to early ART initiation and effective cervical cancer screening in women living with HIV should be key parts of global efforts to reduce cancer-related health inequities.

Introduction

Vast global inequities in the burden of invasive cervical cancer (ICC) exist.^{1,2} While access to effective screening and treatment of pre-cancerous cervical lesions has substantially reduced the risk of developing ICC in high-income countries, ICC remains a common cause of premature mortality and morbidity in women in low- and middle-income countries.^{1,2} ICC disproportionately affects women living with human immunodeficiency virus (HIV), who are more likely to have persistent co-infection with high-risk human papillomavirus (HPV) types,³ to develop pre-cancerous cervical lesions,⁴ and to progress to ICC than HIV-negative women.⁴ The advent and scale-up of combination antiretroviral therapy (ART) has led to a dramatic decline in morbidity and mortality from many HIV-associated diseases,⁵ but these decreases have not occurred for ICC.^{6,7} Indeed, as life expectancy after starting ART increases, there is more time for pre-cancerous cervical lesions to develop into ICC, but early initiation of ART seems to lower HPV co-infection rates and improve control of pre-cancerous cervical lesions.⁸

Global inequities in ICC incidence rates among women living with HIV have not been assessed previously. Our aim was to assess such inequities by comparing ICC incidence rates across different geographic regions among women who had initiated ART. Additionally, we examined risk factors for developing ICC in these women.

Methods

Databases

We analyzed routinely collected clinical, demographic, laboratory, and treatment data of women enrolled in observational HIV cohorts that participate in the International epidemiology Databases to Evaluate AIDS (IeDEA) or the Collaboration of Observational HIV

Epidemiological Research in Europe (COHERE) in EuroCoord. IeDEA has regional data centers in the Asia-Pacific, Australia, North America, Latin America and four African regions. Cohorts from the following IeDEA regions initially contributed data to this study: the North American AIDS Cohort Collaboration on Research and Design (NA-ACCORD),⁹ the Caribbean, Central and South America network for HIV epidemiology (CCASAnet),¹⁰ IeDEA Southern Africa,¹¹ and IeDEA Asia-Pacific.¹² For the IeDEA Southern Africa region, we restricted the analysis to two cohorts from South Africa that reduced under-reporting of cancer cases in the HIV cohorts through record linkages with the National Cancer Registry.¹³ COHERE is a collaboration of observational HIV cohorts across Europe.¹⁴ It contributed data from 24 cohorts, covering 36 countries. All cohorts obtained ethical approval from local ethics committees or institutional review boards, and the Cantonal Ethics Committee of Bern (number 028/2015) also granted ethical approval for this study.

Inclusion criteria and definitions

We restricted the analysis to cohorts that systematically collected cancer data or had enhanced their data through record linkages with cancer registries. We included women living with HIV who started ART after 1995 at 16 years or older. We excluded women who started ART before enrolment into cohort, women without follow-up after ART initiation, and women without any CD4 cell count measurements at ART initiation or during follow-up. We also excluded cohorts with less than 100 eligible women and the Asia-Pacific region because of small sample size (post-hoc decision). We analyzed ICC cases diagnosed any time after ART initiation as incident cases and excluded women diagnosed with ICC before or at ART initiation (prevalent ICC cases) from the analysis. We defined ART as a combination of at least three antiretroviral drugs from any class, including protease inhibitors (PIs), nucleoside reverse transcriptase inhibitors, and non-nucleoside reverse transcriptase inhibitors (NNRTIs). We

assumed that women remained on ART and did not consider treatment interruptions and terminations. CD4 cell count at ART initiation was defined as the cell count closest to ART initiation, during the period within 180 days before to seven days after initiation.

Statistical analyses

We calculated raw ICC incidence rates by dividing the number of incident ICC cases by person-years (pys) at risk. Time at risk was measured from ART initiation to ICC diagnosis, last follow-up visit, death, or database closure, whichever happened first. We used proportional hazard flexible parametric survival models¹⁵ to estimate regional ICC incidence rates and to identify risk factors for developing ICC. We compared ICC rates at 2 years and 5 years after ART initiation across geographic regions. We used restricted cubic splines with 4 degrees of freedom and allowed for time-dependent region-effects with 2 degrees of freedom to model the baseline hazard. We performed likelihood ratio tests to test interactions between risk factors and regions. We assessed the following potential risk factors in the analysis: age at ART initiation (16-30, 31-50, >50 years); first-line ART regimen (NNRTI-based, PI-based, other); calendar period of ART initiation (1996-1998, 1999-2003, 2004-2007, 2008-2014); CD4 cell count at ART initiation; and current (time-updated) CD4 cell count. We treated CD4 cell count at ART initiation and current CD4 cell count as continuous variables. Analyses including CD4 cell count at ART initiation were restricted to women with available data on this variable. HIV RNA load at ART initiation was assessed in descriptive analyses.

We fit a crude model that included only the time-dependent region-effects, resulting in region-specific baseline hazards, and no other risk factors. The main adjusted model included region, CD4 cell count at ART initiation, age at ART initiation, first-line ART regimen, and calendar period of ART initiation. From the main adjusted model, we predicted ICC

incidence rates for women with a specific set of risk factors, i.e. for women who initiated an NNRTI-based regimen between 2008-2014 at age 31-50 years with a CD4 cell count of 200 cells/ μ l. In a sensitivity analysis, the adjusted model included current (time-updated) CD4 cell count instead of CD4 cell count at ART initiation. In a second sensitivity analysis, we excluded ICC cases diagnosed within the first three months after ART initiation as prevalent cases and women with less than three months of follow-up. Results are presented as medians with interquartile ranges (IQR), number and percentages of women, incidence rates per 100,000 pys and hazard ratios (HRs) with 95% confidence intervals (CIs). We used Stata 14 (Stata Corporation, College Station, Texas, USA) and R (R Foundation, Vienna, Austria) for our analyses.

Results

Descriptive analyses

The merged dataset included information on 126,063 women living with HIV. We excluded 44,419 women because they did not receive ART and another 14,413 women for reasons detailed in Supplementary Figures S1-S5. We made a post-hoc decision to exclude the Asia-Pacific region because too few eligible women remained after applying our exclusion criteria.

We included data on 64,231 women living with HIV, drawn from 36 cohorts and 45 countries across Europe, North America, Latin America, and South Africa (Figure 1). Overall, median age at ART initiation was 34.9 years (IQR 29.3-41.9), and was higher in North America (38.6 years) than in other regions (Table 1). Median CD4 cell count at ART initiation was 115 cells/ μ l (IQR 50-182) in South Africa, 178 cells/ μ l (IQR 74-281) in Latin America, and 241

cells/ μ l in both North America and Europe (Table 1). In South Africa, less than 1% of women started ART before 2004, but 26% of women in Latin America, 40% in Europe and 70% in North America initiated ART between 1996 and 2003. Most women in South Africa (93%) and Latin America (70%) received an NNRTI-based first-line regimen, but the majority of women in the European (55%) and North American (60%) cohorts received a PI-based first-line regimen. Median follow-up after ART initiation was around 5 years in Europe, North, and Latin America, but shorter in South Africa (2.1 years).

Over 320,141 pys of follow-up, 356 incident ICC cases were diagnosed (164 in Europe, 156 in South Africa, 19 in North America, and 17 in Latin America). In women who developed ICC, median time from ART initiation to ICC diagnosis was 1.9 years (IQR 0.7-4.2), and it ranged from 1.7 years in South Africa and North America to 2.6 years in Latin America (Supplementary Table S1). Median age at ICC diagnosis was 33 years in Latin America, and 38-40 years in South Africa, North America, and Europe. Median CD4 cell count at ICC diagnosis ranged from 275 cells/ μ l in Latin America to 370 cells/ μ l in North America.

Comparing ICC risk across regions

The raw ICC incidence rate was highest in South Africa, with 447/100,000 pys (95% CI 382-523), followed by Latin America (136/100,000 pys; 95% CI 85-219), North America (76/100,000 pys; 95% CI 48-119), and Europe (66/100,000 pys; 95% CI 57-77). In Europe, North America, and Latin America, there was some evidence for a decrease in crude and adjusted ICC incidence rates after more than one year on ART, except in South Africa (Figure 2). In crude analyses, ICC rates at 5 years after ART initiation were 11-times higher in women living with HIV in South Africa than in their European counterparts (HR 11.06, 95% CI 7.80-15.68). The much higher ICC rate in South African women was not explained by differences in CD4 cell

count at ART initiation, age at ART initiation, first-line ART regimen, or calendar period of ART initiation (adjusted HR [aHR] 10.66, 95% CI 6.73-16.88; see Table 2). In crude (HR 2.32, 95% CI 1.24-4.31) and adjusted analyses (aHR 2.43, 95% CI 1.27-4.68), ICC rates at 5 years after ART initiation were more than twice as high in Latin American as in European women. In North American and European women, ICC rates after ART initiation were comparable in crude (HR 0.98, 95% CI 0.48-1.99) and adjusted analyses (aHR 0.79, 95% CI 0.37-1.71). The regional comparisons of ICC rates were similar at 2 years after ART initiation (Table 2). Also at 2 years after ART initiation, ICC rates were much higher in South Africa than in Europe (aHR 6.23, 95% CI 4.29-9.05). When we excluded ICC cases diagnosed within the first three months after ART initiation in a sensitivity analysis, results did not meaningfully change (Supplementary Table S2).

Risk factors for incident ICC

We did not find evidence of regional variation in the effect of CD4 cell count at ART initiation, age at ART initiation, first-line ART regimen, or calendar period of ART initiation on the risk of developing ICC (all p-values for interaction ≥ 0.13 , see Table 3). Across all regions combined, the risk of developing ICC increased among women who initiated ART at lower CD4 cell counts (per 100 cell/ μl decrease, aHR 1.25, 95% CI 1.15-1.36), and with older age at ART initiation (>50 years versus 16-30 years, aHR 1.57, 95% CI 1.03-2.40). There was no association between type of first-line ART regimen and the risk of developing incident ICC (PI-based versus NNRTI-based, aHR 1.05, 95% CI 0.79-1.41), and we did not observe a relevant decline in ICC rates by calendar period of ART initiation. The effects of the risk factors assessed in the main adjusted model remained similar when we excluded ICC cases diagnosed within the first three months after ART initiation from the analysis (Supplementary Table S3).

In a sensitivity analysis, we assessed the effect of current CD4 cell count on the risk of developing ICC and found that it varied across regions (p-value for interaction = 0.017). In analyses adjusted for age, first-line ART regimen, and calendar period of ART initiation, we did not find an association between current CD4 cell count and risk of developing ICC in South Africa (per 100 cells/ μ l decrease, aHR 1.00, 95% CI 0.92-1.10) or North America (aHR 1.08, 95% CI 0.90-1.30). However, a decrease of 100 cells/ μ l in current CD4 cell count increased the risk of developing ICC by 18% in European women (aHR 1.18, 95% CI 1.10-1.27) and 41% in Latin American women (aHR 1.41, 95% CI 1.07-1.86; see Supplementary Table S4 and Supplementary Figure S6).

Discussion

Across geographic regions, we found large inequities for cervical cancer incidence in women living with HIV. ICC incidence rates were high in women living with HIV in all regions studied, but the risk of developing ICC was much higher in women who had initiated ART in South Africa or Latin America than in women who had initiated ART in Europe or North America. Across all regions combined, the risk of developing ICC increased with older age and lower CD4 cell counts at ART initiation.

We believe this is the first study to provide a comparison of ICC incidence rates among women living with HIV across several geographic regions. To improve comparability of results across regions, we applied the same inclusion criteria and statistical methods across the whole dataset. With more than 60,000 women and 356 ICC cases included, this is also the largest study of ICC incidence in women living with HIV. However, several limitations of our study need to be acknowledged. Less than 20 ICC cases each were recorded in Latin America and

North America. Thus, our comparison of ICC rates between those regions and Europe are of limited precision. ICC case identification and validation are likely to vary across regions and may have affected observed regional differences in ICC rates. Our results for South Africa may not be generalizable to Southern Africa as a region, given that we restricted our analyses to two urban cohorts in South Africa, which had been linked with the National Cancer Registry to reduce under-reporting of ICC cases. Because we included all women who started ART, irrespective of whether they remained in treatment, our results may not be representative of women who stayed continuously on ART. HIV RNA measurements at ART initiation were missing for one-third of women included in Latin America, and almost 80% of women from South Africa. Therefore, we could not use HIV RNA load to evaluate treatment response over time. Information on duration of HIV infection, HPV co-infection status, cervical cancer screening history, and smoking status was generally not available. Thus, we could not explore their effects on the risk of being diagnosed with ICC or adjust the regional comparisons for these potential confounders. Furthermore, as data on history of hysterectomy were not available, we could not exclude women who were no longer at risk of developing ICC. It would also have been interesting to assess ICC-related inequities in more depth, but we did not have data on ICC stage at diagnosis, for example.

We found that across all regions women living with HIV were at high risk of developing ICC after ART initiation. Most previous studies did not restrict their analyses to women who had initiated ART, but rather report ICC incidence estimates for women living with HIV irrespective of ART use.^{6,16-18} The raw ICC incidence rates in women living with HIV who had initiated ART, ranging from 66/100,000 pys in Europe to 447/100,000 pys in South Africa, were substantially higher than the ICC incidence rates reported for women from the general population in the included regions ($\leq 30/100,000$ pys).² In our study, ICC rates after ART

initiation were by far highest in South Africa, followed by Latin America, and they were lower in women who had started ART in North America or Europe. These findings corroborate the regional ICC incidence rate pattern in the general population,² but the difference between South Africa and other regions is even more pronounced among women living with HIV.

The high ICC incidence rates we found in women from South Africa are similar to ICC incidence rates in women living with HIV in the United States in the early 1990s.⁷ In the United States, ICC incidence rates in women living with HIV had already dropped in the mid-1990s, before ART became available, and this drop has partly been attributed to better screening and more effective treatment of pre-cancerous cervical lesions.⁷ The extent to which ART protects women living with HIV from developing ICC is still being explored. Although ART reduces the prevalence of high-risk HPV in women living with HIV and promotes regression of cervical lesions,⁸ many women in our analyses, notably in South Africa, may have started ART too late, when potentially irreversible pre-cancerous cervical lesions were already present. Furthermore, not all women in our study would have achieved sustained suppression of HIV RNA, and high HIV RNA loads have been associated with an increased risk of HPV infection and cervical pre-cancerous lesions.¹⁹ Low CD4 cell counts have also been associated with a higher risk of HPV infection³ and development of severe cervical lesions.²⁰ Accordingly, several studies showed an increased ICC risk in women with low nadir,²¹ baseline,¹⁸ or current CD4 cell counts.¹⁷ It remains a matter of debate at what stage of cervical carcinogenesis the effect of HIV-related immunodeficiency is largest. Across all regions combined, we found that the risk of developing ICC increased in women who initiated ART at low CD4 cell counts. High current CD4 cell counts had a protective effect in Latin America and Europe, but not in North America and South Africa.

Our analyses revealed massive regional differences in ICC rates in women living with HIV. Several factors could account for this finding. HPV prevalence in women living with HIV in sub-Saharan Africa or Latin America is higher than in North America or Europe,²² and this may contribute to the increased ICC burden in South African and Latin American women living with HIV. Women in South Africa and Latin America also tended to initiate ART at lower CD4 cell counts than women in Europe, and low CD4 cell counts at ART initiation increased the risk of developing ICC. Nevertheless, in our analyses large regional differences in ICC rates persisted after adjusting for CD4 cell counts. Therefore, inequities in access to effective cervical cancer screening and treatment of pre-cancerous cervical lesions are likely to be the main driver of regional variation in ICC rates in women living with HIV. Substantial global efforts are needed to improve cervical cancer screening and treatment for women living with HIV, and to promote national HPV vaccination programmes. Unfortunately, most Southern African countries and some regions of Latin America lack the resources to treat ICC.^{23,24}

The availability of HPV vaccination and the long natural history from HPV infection through cervical intraepithelial neoplasia to invasive cancer make ICC particularly amenable to primary and secondary prevention.²⁵ However, it has been estimated that in 2014 less than one-third of female adolescents aged 10-20 years in high-income countries and only 1% in low-income countries had received the full course of HPV vaccine.²⁶ At present, data on HPV vaccination coverage among women living with HIV are lacking.²⁷ Access to screening services with early detection and treatment of pre-cancerous cervical lesions remains key for ICC prevention in women living with HIV. However, there are extensive regional differences in access to effective cervical cancer screening. Less than 10% of women living in low-income countries have access to effective cervical cancer screening as compared to more than 60% in high-income countries.²⁸ Integrating cervical cancer screening services into established HIV

care programmes may facilitate screening access for women living with HIV and improve sustainability of screening programmes.²⁹ Yet, it remains unclear how many women living with HIV actually receive regular screening for pre-cancerous cervical lesions. HIV cohorts and integrated cervical cancer screening services often do not systematically collect patient-level data on screening and treatment of pre-cancerous cervical lesions.³⁰ Rigorous patient-level monitoring of cervical cancer screening and treatment programmes is essential to identify coverage gaps and target interventions.³⁰

Conclusion

Our finding that women living with HIV who initiated ART in South Africa or Latin America were at much higher risk of developing ICC than women in North America or Europe reveals drastic global health inequities. ICC prevention through early ART initiation and scale up of effective cervical cancer screening services for women living with HIV, alongside the promotion of global access to HPV vaccination should be key parts of international efforts to reduce cancer-related health inequities.

Funding

Research reported in this publication was supported by the National Institute of Allergy and Infectious Diseases (NIAID), the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD), the National Cancer Institute (NCI), the National Institute of Mental Health, and the National Institute on Drug Abuse of the U.S. National Institutes of Health (NIH) under Award Number U01AI069924 (Southern Africa), U01AI069907 (Asia-Pacific), U01AI069923 (Caribbean, Central, and South America), and U01-AI069918 (North America). The North American AIDS Cohort Collaboration on Research and Design (NA-ACCORD) was also supported by NIH grants U01AI069918, F31AI124794, F31DA037788, G12MD007583, K01AI093197, K01AI131895, K23EY013707, K24AI065298, K24AI118591, K24DA000432, KL2TR000421, M01RR000052, N01CP01004, N02CP055504, N02CP91027, P30AI027757, P30AI027763, P30AI027767, P30AI036219, P30AI050410, P30AI094189, P30AI110527, P30MH62246, R01AA016893, R01CA165937, R01DA011602, R01DA012568, R01 AG053100, R24AI067039, U01AA013566, U01AA020790, U01AI031834, U01AI034989, U01AI034993, U01AI034994, U01AI035004, U01AI035039, U01AI035040, U01AI035041, U01AI035042, U01AI037613, U01AI037984, U01AI038855, U01AI038858, U01AI042590, U01AI068634, U01AI068636, U01AI069432, U01AI069434, U01AI103390, U01AI103397, U01AI103401, U01AI103408, U01DA03629, U01DA036935, U01HD032632, U10EY008057, U10EY008052, U10EY008067, U24AA020794, U54MD007587, UL1RR024131, UL1TR000004, UL1TR000083, UL1TR000454, UM1AI035043, Z01CP010214 and Z01CP010176; contracts CDC-200-2006-18797 and CDC-200-2015-63931 from the Centers for Disease Control and Prevention, USA; contract 90047713 from the Agency for Healthcare Research and Quality, USA; contract 90051652 from the Health Resources and Services Administration, USA; grants CBR-86906, CBR-94036, HCP-97105 and TGF-96118 from the Canadian Institutes of Health

Research, Canada; Ontario Ministry of Health and Long Term Care; and the Government of Alberta, Canada. Additional support was provided by the National Cancer Institute, National Institute for Mental Health and National Institute on Drug Abuse. The COHERE study group has received unrestricted funding from: Agence Nationale de Recherches sur le SIDA et les Hépatites Virales (ANRS), France; HIV Monitoring Foundation, The Netherlands; and the Augustinus Foundation, Denmark. The research leading to these results has received funding from the European Union Seventh Framework Programme (FP7/2007-2013) under EuroCoord grant agreement no. 260694. A list of the funders of the participating cohorts can be found at www.COHERE.org. JMM received a personal 80:20 research grant from the Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain during 2017-19. JLC was supported by NIH grant K23AI120875. This study was also made possible by the generous support of the American people through the United States Agency for International Development (INROADS USAID-674-A-12-00029), and by a grant from the Swiss National Science Foundation (Ambizione-PROSPER PZ00P3_160407 to JB, special project funding grant 174281 to ME). The content is solely the responsibility of the authors and does not necessarily represent the official views of the funders.

Acknowledgements

We thank all patients, care providers and data managers in the different IeDEA regions and COHERE in EuroCoord. We would also like to acknowledge Kali Tal for her editorial suggestions. More detailed acknowledgements concerning the participating consortia can be found in the supplementary material.

Conflicts of interest

FB received fees from ViiV Healthcare, Janssen, BMS, Gilead, and MSD for educational presentations and research grants from Gilead and Janssen. MJG has served as ad hoc member on Advisory HIV Boards to Merck, ViiV and Gilead in the past three years. AM has received honoraria, lecture fees, consultancy or travel support from Gilead and ViiV. MS has received research grants to his institution from Merck and Gilead. AS has received grants and travel support to her institution from ViiV Healthcare. JJV has personal fees from Merck/MSD, Gilead, Pfizer, Astellas Pharma, Basilea, German Centre for Infection Research (DZIF), University Hospital Freiburg/ Congress and Communication, Academy for Infectious Medicine, University Manchester, German Society for Infectious Diseases (DGI), Ärztekammer Nordrhein, University Hospital Aachen, Back Bay Strategies, German Society for Internal Medicine (DGIM) and grants from Merck/MSD, Gilead, Pfizer, Astellas Pharma, Basilea, German Centre for Infection Research (DZIF), German Federal Ministry of Education and Research (BMBF).

References

- 1 Ginsburg O, Bray F, Coleman MP, et al. The global burden of women's cancers: a grand challenge in global health. *Lancet* 2017; **389**: 847–60.
- 2 Ferlay J, Soerjomataram I, Ervik M, et al. GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11 [Internet]. Lyon, France: International Agency for Research on Cancer; 2013. Available from: <http://globocan.iarc.fr>, accessed on 18/04/2018.
- 3 Palefsky JM, Minkoff H, Kalish LA, et al. Cervicovaginal Human Papillomavirus Infection in Human Immunodeficiency Virus-1 (HIV)-Positive and High-Risk HIV-Negative Women. *J Natl Cancer Inst* 1999; **91**: 226–36.
- 4 Denslow SA, Rositch AF, Firnhaber C, Ting J, Smith JS. Incidence and progression of cervical lesions in women with HIV: a systematic global review. *Int J STD AIDS* 2014; **25**: 163–77.
- 5 Palella FJ, Delaney KM, Moorman AC, et al. Declining Morbidity and Mortality among Patients with Advanced Human Immunodeficiency Virus Infection. *N Engl J Med* 1998; **338**: 853–60.
- 6 Biggar RJ, Chaturvedi AK, Goedert JJ, Engels EA, HIV/AIDS Cancer Match Study. AIDS-related cancer and severity of immunosuppression in persons with AIDS. *J Natl Cancer Inst* 2007; **99**: 962–72.
- 7 Blattner WA, Nowak RG. Epidemiology of AIDS-Defining Malignancies. In: *Cancers in People with HIV and AIDS*. New York, NY: Springer New York, 2014: 17–30.
- 8 Kelly H, Weiss HA, Benavente Y, et al. Association of antiretroviral therapy with high-risk human papillomavirus, cervical intraepithelial neoplasia, and invasive cervical cancer in women living with HIV: a systematic review and meta-analysis. *Lancet HIV*

- 2018; **5**: e45–58.
- 9 Gange SJ, Kitahata MM, Saag MS, et al. Cohort profile: the North American AIDS Cohort Collaboration on Research and Design (NA-ACCORD). *Int J Epidemiol* 2007; **36**: 294–301.
 - 10 McGowan CC, Cahn P, Gotuzzo E, et al. Cohort Profile: Caribbean, Central and South America Network for HIV research (CCASAnet) collaboration within the International Epidemiologic Databases to Evaluate AIDS (IeDEA) programme. *Int J Epidemiol* 2007; **36**: 969–76.
 - 11 Egger M, Ekouevi DK, Williams C, et al. Cohort Profile: the international epidemiological databases to evaluate AIDS (IeDEA) in sub-Saharan Africa. *Int J Epidemiol* 2012; **41**: 1256–64.
 - 12 Asia-Pacific | IeDEA. Available at: <http://www.iedea.org/regions/asia-pacific>; accessed on 18/04/2018.
 - 13 Sengayi M, Spoerri A, Egger M, et al. Record linkage to correct under-ascertainment of cancers in HIV cohorts: The Sinikithemba HIV clinic linkage project. *Int J Cancer* 2016; **139**: 1209–16.
 - 14 Chêne G, Phillips A, Costagliola D, et al. Cohort Profile: Collaboration of Observational HIV Epidemiological Research Europe (COHERE) in EuroCoord. *Int J Epidemiol* 2017; **46**: 797–797n.
 - 15 Lambert PC, Royston P. Further development of flexible parametric models for survival analysis. *Stata J* 2009; **9**: 265–90.
 - 16 Polesel J, Franceschi S, Suligo B, et al. Cancer incidence in people with AIDS in Italy. *Int J Cancer* 2010; **127**: 1437-1445.
 - 17 Guiguet M, Boué F, Cadranel J, Lang J-M, Rosenthal E, Costagliola D. Effect of

- immunodeficiency, HIV viral load, and antiretroviral therapy on the risk of individual malignancies (FHDH-ANRS CO4): a prospective cohort study. *Lancet Oncol* 2009; **10**: 1152–9.
- 18 Abraham AG, D’Souza G, Jing Y, et al. Invasive cervical cancer risk among HIV-infected women: a North American multicohort collaboration prospective study. *J Acquir Immune Defic Syndr* 2013; **62**: 405–13.
- 19 Minkoff H, Zhong Y, Burk RD, et al. Influence of adherent and effective antiretroviral therapy use on human papillomavirus infection and squamous intraepithelial lesions in human immunodeficiency virus-positive women. *J Infect Dis* 2010; **201**: 681–90.
- 20 Silverberg MJ, Leyden WA, Chi A, et al. Human Immunodeficiency Virus (HIV)- and Non-HIV-Associated Immunosuppression and Risk of Cervical Neoplasia. *Obstet Gynecol* 2018; **131**: 47–55.
- 21 Patel P, Hanson DL, Sullivan PS, et al. Incidence of types of cancer among HIV-infected persons compared with the general population in the United States, 1992-2003. *Ann Intern Med* 2008; **148**: 728–36.
- 22 Clifford GM, Tully S, Franceschi S. Carcinogenicity of Human Papillomavirus (HPV) Types in HIV-Positive Women: A Meta-Analysis From HPV Infection to Cervical Cancer. *Clin Infect Dis* 2017; **64**: 1228–35.
- 23 Goss PE, Lee BL, Badovinac-Crnjevic T, et al. The Lancet Oncology Commission Planning cancer control in Latin America and the Caribbean. *Lancet Oncol* 2013; **14**: 391–436.
- 24 Kingham TP, Alatise OI, Vanderpuye V, et al. Treatment of cancer in sub-Saharan Africa. *Lancet Oncol* 2013; **14**: e158–67.
- 25 World Health Organization. Guidelines for screening and treatment of precancerous

- lesions for cervical cancer prevention. Geneva: World Health Organization, 2013.
- 26 Bruni L, Diaz M, Barrionuevo-Rosas L, et al. Global estimates of human papillomavirus vaccination coverage by region and income level: a pooled analysis. *Lancet Glob Health* 2016; **4**: e453–63.
- 27 Kojic EM, Rana AI, Cu-Uvin S. Human Papillomavirus Vaccination in HIV-infected Women: Need for Increased Coverage. *Expert Rev Vaccines* 2016; **15**: 105–17.
- 28 Gakidou E, Nordhagen S, Obermeyer Z. Coverage of Cervical Cancer Screening in 57 Countries: Low Average Levels and Large Inequalities. *PLoS Med* 2008; **5**: e132.
- 29 Sigfrid L, Murphy G, Haldane V, et al. Integrating cervical cancer with HIV healthcare services: A systematic review. *PLoS One* 2017; **12**: e0181156.
- 30 Drummond JL, Were MC, Arrossi S, Wools-Kaloustian K. Cervical cancer data and data systems in limited-resource settings: Challenges and opportunities. *Int J Gynaecol Obstet* 2017; **138 Suppl 1**: 33–40.

Table 1: Characteristics of included women at ART initiation.

	South Africa	Latin America	North America	Europe
	N (%)	N (%)	N (%)	N (%)
All women	13,569 (100%)	2,261 (100%)	4,120 (100%)	44,281 (100%)
Median follow-up time (IQR) [years]	2.1 (0.8-4.1)	5.0 (2.1-8.5)	5.4 (2.3-9.9)	4.8 (2.0-8.5)
Median age at ART initiation (IQR) [years]	35.1 (29.7-41.7)	35.4 (29.3-43.6)	38.6 (32.4-44.8)	34.6 (29.0-41.5)
Age at ART initiation [years]				
16-30	3,584 (26%)	627 (28%)	693 (17%)	12,934 (29%)
31-50	8,969 (66%)	1,363 (60%)	2,947 (72%)	26,944 (61%)
>50	1,016 (7%)	271 (12%)	480 (12%)	4,403 (10%)
First-line ART regimen				
NNRTI-based	12,594 (93%)	1,579 (70%)	1,261 (31%)	16,724 (38%)
PI-based	939 (7%)	623 (28%)	2,476 (60%)	24,325 (55%)
Other ART	36 (<1%)	59 (3%)	383 (9%)	3,232 (7%)
Year of ART initiation				
1996-1998	0 (0%)	26 (1%)	1,600 (39%)	4,260 (10%)
1999-2003	51 (<1%)	560 (25%)	1,276 (31%)	13,236 (30%)
2004-2007	6,791 (50%)	801 (35%)	864 (21%)	12,821 (29%)
2008-2014	6,727 (50%)	874 (39%)	380 (9%)	13,964 (32%)
Median CD4 cell count at ART initiation (IQR) [cells/μl]	115 (50-182)	178 (74-281)	241 (107-385)	241 (129-363)
CD4 cell count at ART initiation [cells/μl]				
< 50	3,113 (23%)	340 (15%)	577 (14%)	4,676 (11%)
50-99	2,402 (18%)	274 (12%)	303 (7%)	3,271 (7%)
100-199	4,733 (35%)	465 (21%)	663 (16%)	7,890 (18%)
200-349	1,821 (13%)	601 (27%)	1,047 (25%)	13,122 (30%)
350-499	276 (2%)	154 (7%)	587 (14%)	6,146 (14%)
500-699	139 (1%)	74 (3%)	349 (8%)	3,069 (7%)
\geq 700	66 (<1%)	30 (1%)	170 (4%)	1,638 (4%)
Missing	1,019 (8%)	323 (14%)	424 (10%)	4,469 (10%)

Median HIV RNA at ART initiation (IQR) [log₁₀	4.4 (2.6-5.2)	4.8 (4.0-5.3)	4.3 (3.4-5.0)	4.5 (3.7-5.1)
HIV RNA at ART initiation [log₁₀ copies/ml]				
< 2.7	788 (6%)	91 (4%)	608 (15%)	4,704 (11%)
2.7-3.9	404 (3%)	260 (11%)	806 (20%)	7,319 (17%)
4.0-4.9	870 (6%)	585 (26%)	1,218 (30%)	14,051 (32%)
≥ 5.0	984 (7%)	568 (25%)	946 (23%)	11,764 (27%)
Missing	10,523 (78%)	757 (33%)	542 (13%)	6,443 (15%)

ART, antiretroviral therapy; HIV, human immunodeficiency virus; IQR, interquartile range; NNRTI, non-nucleoside reverse-transcriptase inhibitor; PI, protease-inhibitor; RNA, ribonucleic acid.

Table 2: Comparison of ICC rates between different regions and Europe: Crude and adjusted HRs for ICC at 2 years and 5 years after ART initiation in women living with HIV.

	At 2 years		At 5 years	
	Crude HR (95% CI)	Adjusted HR* (95% CI)	Crude HR (95% CI)	Adjusted HR* (95% CI)
Region				
Europe	1.00	1.00	1.00	1.00
North America	1.81 (0.94 - 3.48)	1.51 (0.73 – 3.12)	0.98 (0.48 - 1.99)	0.79 (0.37 - 1.71)
Latin America	1.93 (1.09 - 3.42)	1.83 (0.99 - 3.37)	2.32 (1.24 - 4.31)	2.43 (1.27 - 4.68)
South Africa	6.84 (5.20 - 9.00)	6.23 (4.29 - 9.05)	11.06 (7.80 - 15.68)	10.66 (6.73 - 16.88)

* Adjusted for CD4 cell count at ART initiation, age at ART initiation, first-line ART regimen, and calendar period of ART initiation.

ART, antiretroviral therapy; CI, confidence interval; HR, hazard ratio; ICC, invasive cervical cancer.

Table 3: Adjusted hazard ratios for the effect of different factors on the risk of developing incident ICC in women who have initiated ART.

	Hazard ratio* (95% CI)	p-value for interaction**
CD4 cell count at ART initiation		0.76
per 100 cells/ μ l decrease	1.25 (1.15 - 1.36)	
Age at ART initiation [years]		0.34
16-30	1.00	
31-50	1.38 (1.05 - 1.81)	
>50	1.57 (1.03 - 2.40)	
First-line ART regimen		0.21
NNRTI-based	1.00	
PI-based	1.05 (0.79 - 1.41)	
Other ART	0.57 (0.27 - 1.18)	
Year of ART initiation		0.13
1996-1998	1.49 (0.92 - 2.42)	
1999-2003	1.19 (0.80 - 1.77)	
2004-2007	0.83 (0.61 - 1.14)	
2008-2014	1.00	

* Adjusted for region, CD4 cell count at ART initiation, age at ART initiation, calendar year of ART initiation, and first-line ART regimen.

** Derived from likelihood ratio test comparing the adjusted model with and without the interaction of a specific variable with region.

ART, antiretroviral therapy; CI, confidence interval; ICC, invasive cervical cancer; NNRTI, non-nucleoside reverse-transcriptase inhibitor; PI, protease-inhibitor

Figure 1: Map of countries that contributed cohort data; number of included women, by country.





Figure 2: Regional ICC incidence rates with 95% confidence intervals by time since ART initiation, predicted from the crude model (red) and the main adjusted model (blue) for women who initiated an NNRTI-based regimen between 2008-2014, at age 31-50 years, with a CD4 cell count of 200 cells/ μ l.



ART, antiretroviral therapy; ICC, invasive cervical cancer; NNRTI, non-nucleoside reverse-transcriptase inhibitor; pys, person-years.

Cervical cancer risk in women living with HIV across four continents: a multicohort study

The AIDS-defining Cancer Project Working Group of IeDEA and COHERE in EuroCoord

Supplementary material

Figures

- **Supplementary Figure S1:** Identification of study population for analysis. The flow diagram shows the number of included and excluded patients.
- **Supplementary Figure S2:** Identification of study population for the European region.
- **Supplementary Figure S3:** Identification of study population for the North American region.
- **Supplementary Figure S4:** Identification of study population for the Latin American region.
- **Supplementary Figure S5:** Identification of study population for the South African region.
- **Supplementary Figure S6:** ICC incidence rates by time since ART initiation and current CD4 cell count (cells/ μ l) predicted from the second adjusted model (sensitivity analysis) for women living with HIV who initiated an NNRTI-based first-line ART regimen between 2008-2014 at age 31-50 years.


Tables

- **Supplementary Table S1:** Characteristics of women with incident ICC.
- **Supplementary Table S2:** Comparison of ICC rates between different regions and Europe: Crude and adjusted HRs for ICC at 2 years and 5 years after ART initiation in women living with HIV; sensitivity analysis excluding ICC cases diagnosed within the first three months after ART initiation.
- **Supplementary Table S3:** Sensitivity analysis of regional risk factors for incident ICC in women who have initiated ART, excluding ICC cases diagnosed within the first three months after ART initiation.
- **Supplementary Table S4:** Sensitivity analysis of regional risk factors for incident ICC in women who have initiated ART, including current CD4 cell count instead of CD4 cell count at ART initiation.

Acknowledgements


- **Detailed acknowledgements of consortia contributing data to this analysis.**

Supplementary Figure S1: Identification of study population for analysis. The flow diagram shows the number of included and excluded women.




ART, antiretroviral therapy; ICC, invasive cervical cancer.


Supplementary Figure S2: Identification of study population for the European region.




Supplementary Figure S3: Identification of study population for the North American region.




Supplementary Figure S4: Identification of study population for the Latin American region.



Supplementary Figure S5: Identification of study population for the South African region.



Supplementary Figure S6: ICC incidence rates by time since ART initiation and current CD4 cell count (cells/ μ l) predicted from the second adjusted model (sensitivity analysis) for women living with HIV who initiated an NNRTI-based first-line ART regimen between 2008-2014 at age 31-50 years.



ART, antiretroviral therapy; ICC, invasive cervical cancer; NNRTI, non-nucleoside reverse-transcriptase inhibitor.

Supplementary Table S1: Characteristics of women with incident ICC.

	South Africa	Latin America	North America	Europe
	N (%)	N (%)	N (%)	N (%)
All ICC cases	156 (100%)	17 (100%)	19 (100%)	164 (100%)
Median time to ICC diagnosis (IQR) [years]	1.7 (0.9-3.2)	2.6 (0.7-5.6)	1.7 (1.2-6.0)	2.2 (0.7-5.3)
Median age at ICC diagnosis (IQR) [years]	38.3 (33.5; 42.7)	33.0 (28.8; 43.4)	39.0 (29.6; 43.1)	39.7 (33.8; 45.4)
Age at ICC diagnosis [years]				
16-30	13 (8%)	5 (29%)	5 (26%)	16 (10%)
31-50	131 (84%)	10 (59%)	13 (68%)	124 (76%)
>50	12 (8%)	2 (12%)	1 (5%)	24 (15%)
First-line ART regimen				
NNRTI-based	150 (96%)	14 (82%)	3 (16%)	55 (34%)
PI-based	5 (3%)	3 (18%)	15 (79%)	103 (63%)
Other ART	1 (1%)	0 (0%)	1 (5%)	6 (4%)
Year of ART initiation				
1996-1998	0 (0%)	1 (6%)	10 (53%)	36 (22%)
1999-2003	0 (0%)	6 (35%)	6 (32%)	71 (43%)
2004-2007	114 (73%)	3 (18%)	3 (16%)	31 (19%)
2008-2014	42 (27%)	7 (41%)	0 (0%)	26 (16%)
Median CD4 cell count at ICC diagnosis (IQR) [cells/μl]	292 (170-450)	275 (190-406)	370 (200-586)	350 (183-523)
Women with missing CD4 cell count at ICC diagnosis	37 (24%)	6 (35%)	1 (5%)	23 (14%)
Median HIV RNA at ICC diagnosis (IQR) [log₁₀ copies/ml]	1.7 (1.7-1.7)	1.7 (1.6-3.3)	2.3 (2.3-2.8)	2.0 (1.7-3.7)
Women with missing HIV RNA at ICC diagnosis	51 (33%)	4 (24%)	1 (5%)	31 (19%)

ART, antiretroviral therapy; HIV, human immunodeficiency virus; IQR, interquartile range; NNRTI, non-nucleoside reverse-transcriptase inhibitors; PI, protease-inhibitors; RNA, ribonucleic acid.

Supplementary Table S2: Comparison of ICC rates between different regions and Europe: Crude and adjusted HRs for ICC at 2 years and 5 years after ART initiation in women living with HIV; sensitivity analysis excluding ICC cases diagnosed within the first three months after ART initiation.

Region	At 2 years		At 5 years	
	Crude HR (95% CI)	Adjusted HR* (95% CI)	Crude HR (95% CI)	Adjusted HR* (95% CI)
Europe	1.00	1.00	1.00	1.00
North America	1.31 (0.71 - 2.41)	1.09 (0.55 - 2.15)	0.99 (0.49 - 2.01)	0.80 (0.37 - 1.74)
Latin America	1.81 (1.02 - 3.24)	1.94 (0.98 - 3.82)	2.32 (1.23 - 4.39)	2.42 (1.22 - 4.78)
South Africa	7.36 (5.52 - 9.83)	6.85 (4.60 - 10.19)	10.53 (7.08 - 15.67)	9.87 (5.89 - 16.53)

* Adjusted for CD4 cell count at ART initiation, age at ART initiation, first-line ART regimen, and calendar period of ART initiation.

ART, antiretroviral therapy; CI, confidence interval; HR, hazard ratio; ICC, invasive cervical cancer.

Supplementary Table S3: Sensitivity analysis of risk factors for incident ICC in women who have initiated ART, excluding ICC cases diagnosed within the first three months after ART initiation.

	Hazard ratio* (95% CI)
CD4 cell count at ART initiation	
per 100 cells/ μ l decrease	1.23 (1.12 - 1.34)
Age at ART initiation [years]	
16-30	1.00
31-50	1.32 (1.00 - 1.76)
>50	1.48 (0.94 - 2.33)
First-line ART regimen	
NNRTI-based	1.00
PI-based	1.07 (0.79 - 1.45)
Other ART	0.39 (0.16 - 0.97)
Year of ART initiation	
1996-1998	1.71 (1.01 - 2.90)
1999-2003	1.37 (0.88 - 2.13)
2004-2007	0.98 (0.69 - 1.39)
2008-2014	1.00

* Adjusted for region, CD4 cell count at ART initiation, age, calendar year of ART start, and first-line ART regimen.

** Derived from likelihood ratio test comparing the adjusted model with and without the interaction of a specific variable with region.

ART, antiretroviral therapy; CI, confidence interval; ICC, invasive cervical cancer; NNRTI, non-nucleoside reverse-transcriptase inhibitor; PI, protease-inhibitor

Supplementary Table S4: Sensitivity analysis of risk factors for incident ICC in women who have initiated ART, including current CD4 cell count instead of CD4 cell count at ART initiation.

	Hazard ratio* (95% CI)	p-value for interaction**
Current CD4 cell count per 100 cells/μl decrease		0.017
South Africa	1.00 (0.92 - 1.10)	
Latin America	1.41 (1.07 - 1.86)	
North America	1.08 (0.90 - 1.30)	
Europe	1.18 (1.10 - 1.27)	
Age at ART initiation [years]		0.055
16-30	1.00	
31-50	1.39 (1.07 - 1.81)	
>50	1.54 (1.02 - 2.32)	
First-line ART regimen		0.103
NNRTI-based	1.00	
PI-based	1.01 (0.77 - 1.33)	
Other ART	0.48 (0.23 - 1.00)	
Year of ART initiation		0.129
1996-1998	1.46 (0.92 - 2.32)	
1999-2003	1.14 (0.77 - 1.66)	
2004-2007	0.84 (0.62 - 1.13)	
2008-2014	1.00	

* Adjusted for current CD4 cell count and its interaction with region, age, calendar year of ART start, and first-line ART regimen.

** Derived from likelihood ratio test comparing the adjusted models with and without the interaction of a specific variable with region.

ART, antiretroviral therapy; CI, confidence interval; ICC, invasive cervical cancer; NNRTI, non-nucleoside reverse-transcriptase inhibitor; PI, protease-inhibitor

Acknowledgements

COHERE Steering Committee - Contributing Cohorts: Ali Judd (AALPHI), Robert Zangerle (AHIVCOS), Giota Touloumi (AMACS), Josiane Warszawski (ANRS CO1 EPF/ANRS CO11 OBSERVATOIRE EPF), Laurence Meyer (ANRS CO2 SEROCO), François Dabis (ANRS CO3 AQUITAINE), Murielle Mary Krause (ANRS CO4 FHDH), Jade Ghosn (ANRS CO6 PRIMO), Catherine Leport (ANRS CO8 COPILOTE), Linda Wittkop (ANRS CO13 HEPAVIH), Peter Reiss (ATHENA), Ferdinand Wit (ATHENA), Maria Prins (CASCADE), Heiner Bucher (CASCADE), Diana Gibb (CHIPS), Gerd Fätkenheuer (Cologne-Bonn), Julia Del Amo (CoRIS), Niels Obel (Danish HIV Cohort), Claire Thorne (ECS), Amanda Mocroft (EuroSIDA), Ole Kirk (EuroSIDA), Christoph Stephan (Frankfurt), Santiago Pérez-Hoyos (GEMES-Haemo), Osamah Hamouda (German ClinSurv), Barbara Bartmeyer (German ClinSurv), Nikoloz Chkhartishvili (Georgian National HIV/AIDS), Antoni Noguera-Julian (CORISPE-cat), Andrea Antinori (ICC), Antonella d'Arminio Monforte (ICONA), Norbert Brockmeyer (KOMPNET), Luis Prieto (Madrid PMTCT Cohort), Pablo Rojo Conejo (CORISPE-Madrid), Antoni Soriano-Arandes (NENEXP), Manuel Battegay (SHCS), Roger Kouyos (SHCS), Cristina Mussini (Modena Cohort), Pat Tookey (NSHPC), Jordi Casabona (PISCIS), Jose M. Miró (PISCIS), Antonella Castagna (San Raffaele), Deborah Konopnick (St. Pierre Cohort), Tessa Goetghebuer (St Pierre Paediatric Cohort), Anders Sönnnerborg (Swedish InfCare), Eugenia Quiros-Roldan (The Italian Master Cohort), Caroline Sabin (UK CHIC), Ramon Teira (VACH), Myriam Garrido (VACH). David Haerry (European AIDS Treatment Group)

COHERE Executive Committee: Stéphane de Wit (Chair, St. Pierre University Hospital), Jose M^a Miró (PISCIS), Dominique Costagliola (FHDH), Antonella d'Arminio-Monforte (ICONA), Antonella Castagna (San Raffaele), Julia del Amo (CoRIS), Amanda Mocroft (EuroSida), Dorthe Raben (Head, Copenhagen Regional Coordinating Centre), Geneviève Chêne (Head, Bordeaux Regional Coordinating Centre). Paediatric Cohort Representatives: Ali Judd, Pablo Rojo Conejo.

COHERE Regional Coordinating Centres: Bordeaux RCC: Diana Barger, Christine Schwimmer, Monique Termote, Linda Wittkop; Copenhagen RCC: Maria Campbell, Casper M. Frederiksen, Nina Friis-Møller, Jesper Kjaer, Dorthe Raben, Rikke Salbøl Brandt.

COHERE Project Leads and Statisticians: Juan Berenguer, Julia Bohlius, Vincent Bouteloup, Heiner Bucher, Alessandro Cozzi-Lepri, François Dabis, Antonella d'Arminio Monforte, Mary-Anne Davies, Julia del Amo, Maria Dorrucchi, David Dunn, Matthias Egger, Hansjakob Furrer, Marguerite Guiguet, Sophie Grabar, Ali Judd, Ole Kirk, Olivier Lambotte, Valérie Leroy, Sara Lodi, Sophie Matheron, Laurence Meyer, Jose M. Miro, Amanda Mocroft, Susana Monge, Fumiyo Nakagawa, Roger Paredes, Andrew Phillips, Massimo Puoti, Eliane Rohner, Michael Schomaker, Colette Smit, Jonathan Sterne, Rodolphe Thiebaut, Claire Thorne, Carlo Torti, Marc van der Valk, Linda Wittkop.

leDEA-SA site investigators and cohorts: Gary Maartens, Aid for AIDS, South Africa; Michael Vinikoor, Centre for Infectious Disease Research in Zambia (CIDRZ), Zambia; Monique van Lettow, Dignitas, Malawi; Robin Wood, Gugulethu ART Programme, South Africa; Nosisa Sipambo, Harriet Shezi Clinic, South Africa; Frank Tanser, Africa Centre for Health & Population Studies (Hlabisa), South Africa; Andrew Boulle, Khayelitsha ART Programme, South Africa; Geoffrey Fatti, Kheth'Impilo, South Africa; Sam Phiri, Lighthouse Clinic, Malawi; Cleophas Chimbetete, Newlands Clinic, Zimbabwe; Karl Technau, Rahima Moosa Mother and Child Hospital, South Africa; Brian Eley, Red Cross Children's Hospital, South Africa; Josephine Muhairwe, SolidarMed Lesotho; Anna Jores, SolidarMed Mozambique; Cordelia Kunzekwenyika, SolidarMed Zimbabwe, Matthew P Fox, Themba Lethu Clinic, South Africa; Hans Prozesky, Tygerberg Academic Hospital, South Africa.

We also would like to acknowledge colleagues at the National Health Laboratory Service and the National Cancer Registry in South Africa.

leDEA-SA data centers: Nina Anderegg, Marie Ballif, Lina Bartels, Julia Bohlius, Frédérique Chammartin, Benedikt Christ, Cam Ha Dao Ostinelli, Matthias Egger, Lukas Fenner, Per von Groote, Andreas Haas, Taghavi Katayoun, Eliane Rohner, Lilian Smith, Adrian Spörri, Gilles Wandeler, Elizabeth Zaniewski, Kathrin Zürcher, Institute of Social and Preventive Medicine, University of Bern, Switzerland; Andrew Boulle, Morna Cornell, Mary-Ann Davies, Victoria Iyun, Leigh Johnson, Mmamapudi Kubjane, Nicola Maxwell, Tshabakwane Nembandona, Patience Nyakato, Ernest Mokotoane, Gem Patten, Michael Schomaker, Priscilla Tsondai, Renee de Waal, School of Public Health and Family Medicine, University of Cape Town, South Africa.

NA-ACCORD Collaborating Cohorts and Representatives: AIDS Clinical Trials Group Longitudinal Linked Randomized Trials: Constance A. Benson and Ronald J. Bosch; AIDS Link to the IntraVenous Experience: Gregory D. Kirk; Fenway Health HIV Cohort: Stephen Boswell, Kenneth H. Mayer and Chris Grasso; HAART Observational Medical Evaluation and Research: Robert S. Hogg, P. Richard Harrigan, Julio SG Montaner, Benita Yip, Julia Zhu, Kate Salters and Karyn Gabler; HIV Outpatient Study: Kate Buchacz and John T. Brooks; HIV Research Network: Kelly A. Gebo and Richard D. Moore; Johns Hopkins HIV Clinical Cohort: Richard D. Moore; John T. Carey Special Immunology Unit Patient Care and Research Database, Case Western Reserve University: Benigno Rodriguez; Kaiser Permanente Mid-Atlantic States: Michael A. Horberg; Kaiser Permanente Northern California: Michael J. Silverberg; Longitudinal Study of Ocular Complications of AIDS: Jennifer E. Thorne; Multicenter Hemophilia Cohort Study–II: Charles Rabkin; Multicenter AIDS Cohort Study: Joseph B. Margolick, Lisa P. Jacobson and Gypsyamber D'Souza; Montreal Chest Institute Immunodeficiency Service Cohort: Marina B. Klein; Ontario HIV Treatment Network Cohort Study: Sean B. Rourke, Anita R. Rachlis, and Patrick Cupido;

Retrovirus Research Center, Bayamon Puerto Rico: Robert F. Hunter-Mellado and Angel M. Mayor; Southern Alberta Clinic Cohort: M. John Gill; Study of the Consequences of the Protease Inhibitor Era: Steven G. Deeks and Jeffrey N. Martin; Study to Understand the Natural History of HIV/AIDS in the Era of Effective Therapy: Pragna Patel and John T. Brooks; University of Alabama at Birmingham 1917 Clinic Cohort: Michael S. Saag, Michael J. Mugavero and James Willig; University of North Carolina at Chapel Hill HIV Clinic Cohort: Joseph J. Eron and Sonia Napravnik; University of Washington HIV Cohort: Mari M. Kitahata, Heidi M. Crane and Daniel R. Drozd; Vanderbilt Comprehensive Care Clinic HIV Cohort: Timothy R. Sterling, David Haas, Peter Rebeiro, Megan Turner, Sally Bebawy and Ben Rogers; Veterans Aging Cohort Study: Amy C. Justice, Robert Dubrow, and David Fiellin; Women's Interagency HIV Study: Stephen J. Gange and Kathryn Anastos.

NA-ACCORD Study Administration: Executive Committee: Richard D. Moore, Michael S. Saag, Stephen J. Gange, Mari M. Kitahata, Keri N. Althoff, Michael A. Horberg, Marina B. Klein, Rosemary G. McKaig and Aimee M. Freeman; Administrative Core: Richard D. Moore, Aimee M. Freeman and Carol Lent; Data Management Core: Mari M. Kitahata, Stephen E. Van Rompaey, Heidi M. Crane, Daniel R. Drozd, Liz Morton, Justin McReynolds and William B. Lober; Epidemiology and Biostatistics Core: Stephen J. Gange, Keri N. Althoff, Alison G. Abraham, Bryan Lau, Jinbing Zhang, Jerry Jing, Sharada Modur, Cherise Wong, Brenna Hogan, Fidel Desir, Bin Liu and Bin You.

leDEA Caribbean, Central, and South America (CCASAnet): Fundación Huésped, Argentina: Pedro Cahn, Carina Cesar, Valeria Fink, Omar Sued, Emanuel Dell'Isola, Cleyton Yamamoto; Instituto Nacional de Infectologia-Fiocruz, Brazil: Beatriz Grinsztejn, Valdilea Veloso, Paula Luz, Raquel de Boni, Sandra Cardoso Wagner, Ruth Friedman, Ronaldo Moreira; Universidade Federal de Minas Gerais, Brazil: Jorge Pinto, Flavia Ferreira, Marcelle Maia; Universidade Federal de São Paulo, Brazil: Regina Célia de Menezes Succi, Daisy Maria Machado, Aida de Fátima Barbosa Gouvêa; Fundación Arriarán, Chile: Marcelo Wolff, Claudia Cortes, Maria Fernanda Rodriguez, Gladys Allendes; Les Centres GHESKIO, Haiti: Jean William Pape, Vanessa Rouzier, Adias Marcelin, Christian Perodin; Hospital Escuela Universitario, Honduras: Marco Tulio Luque; Instituto Hondureño de Seguridad Social, Honduras: Denis Padgett; Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico: Juan Sierra Madero, Brenda Crabtree Ramirez, Paco Belaunzaran, Yanink Caro Vega; Instituto de Medicina Tropical Alexander von Humboldt, Peru: Eduardo Gotuzzo, Fernando Mejia, Gabriela Carriquiry; Vanderbilt University Medical Center, USA: Catherine C McGowan, Bryan E Shepherd, Timothy Sterling, Karu Jayathilake, Anna K Person, Peter F Rebeiro, Mark Giganti, Jessica Castilho, Stephany N Duda, Fernanda Maruri, Hilary Vansell.

leDEA Asia-Pacific TAHOD study members: PS Ly* and V Khol, National Center for HIV/AIDS, Dermatology & STDs, Phnom Penh, Cambodia; FJ Zhang* ‡, HX Zhao and N Han, Beijing Ditan Hospital, Capital Medical University, Beijing, China; MP Lee*, PCK Li, W Lam and YT Chan, Queen Elizabeth Hospital, Hong Kong, China; N Kumarasamy*, S Saghayam and C Ezhilarasi, Chennai Antiviral Research and Treatment Clinical Research Site (CARTCRS), YRGCARE Medical Centre, VHS, Chennai, India; S Pujari*, K Joshi, S Gaikwad and A Chitalikar, Institute of Infectious Diseases, Pune, India; TP Merati*, DN Wirawan and F Yuliana, Faculty of Medicine Udayana University & Sanglah Hospital, Bali, Indonesia; E Yuniastuti*, D Imran and A Widhani, Faculty of Medicine Universitas Indonesia - Dr. Cipto Mangunkusumo General Hospital, Jakarta, Indonesia; J Tanuma*, S Oka and T Nishijima, National Center for Global Health and Medicine, Tokyo, Japan; JY Choi*, Na S and JM Kim, Division of Infectious Diseases, Department of Internal Medicine, Yonsei University College of Medicine, Seoul, South Korea; BLH Sim*, YM Gani, and R David, Hospital Sungai Buloh, Sungai Buloh, Malaysia; A Kamarulzaman*, SF Syed Omar, S Ponnampalavanar and I Azwa, University Malaya Medical Centre, Kuala Lumpur, Malaysia; R Ditangco*, E Uy and R Bantique, Research Institute for Tropical Medicine, Manila, Philippines; WW Wong* †, WW Ku and PC Wu, Taipei Veterans General Hospital, Taipei, Taiwan; OT Ng*, PL Lim, LS Lee and PS Ohnmar, Tan Tock Seng Hospital, Singapore; A Avihingsanon*, S Gatechompol, P Phanuphak and C Phadungphon, HIV-NAT/Thai Red Cross AIDS Research Centre, Bangkok, Thailand; S Kiertiburanakul*, S Sungkanuparph, L Chumla and N Sanmeema, Faculty of Medicine Ramathibodi Hospital, Mahidol University, Bangkok, Thailand; R Chaiwarith*, T Sirisanthana, W Kotarathitum and J Preparattanapan, Research Institute for Health Sciences, Chiang Mai, Thailand; P Kantipong* and P Kambua, Chiangrai Prachanukroh Hospital, Chiang Rai, Thailand; KV Nguyen*, HV Bui, DTH Nguyen and DT Nguyen, National Hospital for Tropical Diseases, Hanoi, Vietnam; DD Cuong*, NV An and NT Luan, Bach Mai Hospital, Hanoi, Vietnam; AH Sohn*, JL Ross* and B Petersen, TREAT Asia, amfAR - The Foundation for AIDS Research, Bangkok, Thailand; MG Law*, A Jiamsakul* and DC Boettiger, The Kirby Institute, UNSW Australia, Sydney, Australia.

* TAHOD Steering Committee member; † Steering Committee Chair; ‡ co-Chair.

leDEA Asia-Pacific AHOD study members: New South Wales: D Ellis, Coffs Harbour Medical Centre, Coffs Harbour; M Bloch, S Agrawal, T Vincent, Holdsworth House Medical Practice, Sydney; D Allen, Holden Street Clinic, Gosford; D Smith, A Rankin, Lismore Sexual Health & AIDS Services, Lismore; D Baker*, East Sydney Doctors, Surry Hills; DJ Templeton*, CC O'Connor, O Thackeray, RPA Sexual Health, Camperdown; E Jackson, K McCallum, Blue Mountains Sexual Health and HIV Clinic, Katoomba; N Ryder, G Sweeney, Clinic 468, HNE Sexual Health, Tamworth; D Cooper, A Carr, K Macrae, K Hesse, St Vincent's Hospital, Darlinghurst; R Finlayson, S Gupta, Taylor Square Private Clinic, Darlinghurst; J Langton-Lockton, J Shakeshaft, Nepean Sexual Health and HIV Clinic, Penrith; K

Brown, S Idle, N Arvela, Illawarra Sexual Health Service, Warrawong; R Varma, H Lu, Sydney Sexual Health Centre, Sydney; D Couldwell, S Eswarappa, Western Sydney Sexual Health Clinic; DE Smith*, V Furner, D Smith, G Cabrera, Albion Street Centre; S Fernando, Clinic 16 – Royal North Shore Hospital; A Cogle*, National Association of People living with HIV/AIDS; C Lawrence*, National Aboriginal Community Controlled Health Organisation; B Mulhall*, Department of Public Health and Community Medicine, University of Sydney; M Boyd*, University of Adelaide; M Law*, K Petoumenos*, R Puhr*, R Huang*, A Han*, The Kirby Institute, University of NSW. Northern Territory: M Gunathilake, R Payne, Communicable Disease Centre, Darwin. Queensland: M O’Sullivan, A Croydon, Gold Coast Sexual Health Clinic, Miami; D Russell, C Cashman, C Roberts, Cairns Sexual Health Service, Cairns; D Sowden, K Taing, P Marshall, Clinic 87, Sunshine Coast-Wide Bay Health Service District, Nambour; D Orth, D Youds, Gladstone Road Medical Centre, Highgate Hill; D Rowling, N Latch, E Warzywoda, Sexual Health and HIV Service in Metro North, Brisbane; B Dickson*, CaraData. South Australia: W Donohue, O’Brien Street General Practice, Adelaide. Victoria: R Moore, S Edwards, S Boyd, Northside Clinic, North Fitzroy; NJ Roth*, H Lau, Prahran Market Clinic, South Yarra; T Read, J Silvers*, W Zeng, Melbourne Sexual Health Centre, Melbourne; J Hoy*, K Watson*, M Bryant, S Price, The Alfred Hospital, Melbourne; I Woolley, M Giles*, T Korman, J Williams*, Monash Medical Centre, Clayton. Western Australia: D Nolan, A Allen, G Guelfi. Department of Clinical Immunology, Royal Perth Hospital, Perth. New Zealand: G Mills, C Wharry, Waikato District Hospital Hamilton; N Raymond, K Bargh, Wellington Hospital, Wellington. CoDe reviewers: D Templeton, M Giles, K Brown and J Hoy.

* AHOD Steering Committee member