



Draft Genome Sequence of Methicillin-Resistant *Staphylococcus aureus* Strain AW7, Isolated from a Patient with Bacteremia

David R. Cameron,^a  Alban Ramette,^b Josef Prazak,^a José Entenza,^c Matthias Haenggi,^a Yok-Ai Que,^a Grégory Resch^c

^aDepartment of Intensive Care Medicine, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland

^bInstitute for Infectious Diseases, University of Bern, Bern, Switzerland

^cDepartment of Fundamental Microbiology, University of Lausanne, Lausanne, Switzerland

ABSTRACT Methicillin-resistant *Staphylococcus aureus* (MRSA) strain AW7 is a commonly used challenge strain in experimental models of MRSA infection. Here, we report its draft genome sequence.

Methicillin-resistant *Staphylococcus aureus* (MRSA) strain AW7 (complete taxonomy is *Bacteria*, *Firmicutes*, *Bacilli*, *Bacillales*, *Staphylococcaceae*, *Staphylococcus aureus* strain AW7) was isolated from a patient with bacteremia in Switzerland in the 1980s. The strain has been used for many decades in experimental endocarditis models aimed at testing antibiotic efficacy (1, 2). More recently, we have used AW7 in a rodent model of ventilator-associated pneumonia to assess phage therapy (3).

AW7 was grown overnight at 37°C in tryptic soy broth. Genomic DNA was prepared using a Qiagen blood and tissue kit per the manufacturer's instructions and sequenced using the Illumina MiSeq platform (Nextera DNA Flex; assay V2 2 × 150 bp, 300 cycles), generating 6,274,783 reads. Sequencing data were processed using the shovill pipeline with the `-trim` option and default parameters (v0.9.0). The average read length was 148 bp with an avgQscore of 37.6. Contigs were assembled using SPAdes (v3.11.1; `-only-assembler`; k-mer sizes of 31, 51, 71, 91, and 111) (4) and polished using Pilon (v1.22) (5). The sequence comprised 63 contigs (>200 bp long; N_{50} , 140,137 bp) with a combined length of 2,910,678 bp (G+C content of 32.7%) and an average coverage of ~62×. The draft genome was annotated using the NCBI Prokaryotic Genome Annotation Pipeline (v4.8) (6), resulting in 2,918 predicted protein-coding regions.

AW7 belongs to multilocus sequence type 247 (ST247) (7) and harbors a staphylococcal cassette chromosome *mec* type I (1B) [SCC*mec* I (1B)] element (as determined by SCC*mec* finder v2.1). In addition to SCC*mec*, which facilitates methicillin resistance, AW7 harbors genes predicted to code for additional resistance determinants for β -lactams (*blaZ*), aminoglycosides [*aph(3')*-III, *ant(6)-Ia*, *ant(9)-Ia*, and *aac(6')*-*aph(2'')*], tetracycline (*tetM*), macrolide-lincosamide-streptogramin B (*ermA*) (2), and chloramphenicol (*cat*) (8). The genome contains three prophages, as determined by PHAST (9), including a β -hemolysin-converting phage that harbors virulence factors enterotoxin A (*sea*), staphylokinase (*sak*), and complement inhibitor (*scn*). Additional virulence factors include toxin genes α -hemolysin (*hla*), γ -hemolysin (*hlgABC*), and leukocidin (*lukDE*), as well as exoenzymes aureolysin (*aur*), serine protease A (*spIA*), serine protease B (*spIB*), and serine protease E (*spIE*) (10).

The genome sequence of AW7 provides a useful resource for future studies of MRSA infection in experimental animal models.

Data availability. This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [SRL00000000](https://doi.org/10.7892/boris.133752), with SRA accession number [PRJNA530527](https://doi.org/10.7892/boris.133752).

Citation Cameron DR, Ramette A, Prazak J, Entenza J, Haenggi M, Que Y-A, Resch G. 2019. Draft genome sequence of methicillin-resistant *Staphylococcus aureus* strain AW7, isolated from a patient with bacteremia. *Microbiol Resour Announc* 8:e00806-19. <https://doi.org/10.1128/MRA.00806-19>.

Editor Steven R. Gill, University of Rochester School of Medicine and Dentistry

Copyright © 2019 Cameron et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to David R. Cameron, davidrobert.cameron@insel.ch.

Received 15 July 2019

Accepted 10 September 2019

Published 3 October 2019

ACKNOWLEDGMENT

This work was supported by Swiss National Foundation grants 320030_176216 and CR3113_166124.

REFERENCES

1. Vouillamoz J, Entenza JM, Feger C, Glauser MP, Moreillon P. 2000. Quinupristin-dalfopristin combined with beta-lactams for treatment of experimental endocarditis due to *Staphylococcus aureus* constitutively resistant to macrolide-lincosamide-streptogramin B antibiotics. *Antimicrob Agents Chemother* 44:1789–1795. <https://doi.org/10.1128/aac.44.7.1789-1795.2000>.
2. Entenza JM, Drugeon H, Glauser MP, Moreillon P. 1995. Treatment of experimental endocarditis due to erythromycin-susceptible or -resistant methicillin-resistant *Staphylococcus aureus* with RP 59500. *Antimicrob Agents Chemother* 39:1419–1424. <https://doi.org/10.1128/aac.39.7.1419>.
3. Prazak J, Iten M, Cameron DR, Save J, Grandgirard D, Resch G, Goepfert C, Leib SL, Takala J, Jakob SM, Que YA, Haenggi M. 2019. Bacteriophages improve outcome in experimental *Staphylococcus aureus* ventilator associated pneumonia. *Am J Respir Crit Care Med*. <https://doi.org/10.1164/rccm.201812-2372OC>.
4. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
5. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
6. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569>.
7. Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG. 2000. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J Clin Microbiol* 38:1008–1015.
8. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.
9. Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. 2011. PHAST: a fast phage search tool. *Nucleic Acids Res* 39:W347–W352. <https://doi.org/10.1093/nar/gkr485>.
10. Joensen KG, Scheutz F, Lund O, Hasman H, Kaas RS, Nielsen EM, Aarestrup FM. 2014. Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *J Clin Microbiol* 52:1501–1510. <https://doi.org/10.1128/JCM.03617-13>.