

- 1 NON-MEDICAL FACTORS SIGNIFICANTLY INFLUENCE THE LENGTH OF
- 2 HOSPITAL STAY AFTER SURGERY FOR DEGENERATIVE SPINE DISORDERS.
- 3

4 **Background.** Unnecessarily long hospital stays are costly and inefficient. Studies have
5 shown that the length of hospital stay (LOS) for spine surgical procedures is influenced
6 by various disease-related or medical factors but few have examined the role of
7 sociodemographic/socioeconomic (SDE) factors.

8 **Methods.** This was a retrospective analysis of data from 10,770 patients (5,056 men,
9 5,714 women; 62±15y) with degenerative spinal disorders, collected prospectively in an
10 in-house database within the framework of EUROSPINE's Spine Tango Registry.
11 Surgeons completed the Tango surgery form (clinical history, demographics, surgical
12 measures, complications), and patients, a baseline Core Outcome Measures Index.
13 Stepwise linear regression analyses examined SDE predictors of LOS, controlling for
14 potential medical/biological factors.

15 **Results.** The mean LOS was 7.9 ± 5.2 days. The final model accounted for 42% of
16 variance in LOS, with SDE variables explaining 13% variance and medical/surgical
17 predictors, 29%. In the final model, the SDE factors age and being female were
18 significant independent predictors of LOS, whereas others were either non-significant
19 (insurance status, being of Swiss nationality, being a smoker) or reached only
20 borderline significance ($p < 0.1$) (BMI). Controlling for all other SDE and medical/surgical
21 confounders, being female was associated with 1.11-day longer LOS (95%CI, 0.96-
22 1.27; $p < 0.0001$).

23 **Conclusions.** Patients of advanced age and female gender are at increased risk of
24 longer hospital stay after surgery for degenerative spinal disorders. Further studies
25 should seek to understand the reasoning behind the gender disparity, in order to
26 minimize potentially unnecessary costs of prolonged LOS. Targeted pre-operative
27 discharge planning may improve the utilisation of hospital resources.

28 **Keywords:** length of stay; spine surgery; degenerative disorders of the cervical and
29 thoracic/lumbar spine; non-medical predictors

30

31 **Introduction**

32 Inpatient stays after orthopaedic surgery can be very costly. The use of hospital
33 resources such as equipment, staffing and bed capacity are all related to the length of
34 stay (LOS). The LOS is a metric that is commonly used to plan resource utilisation and
35 to monitor the quality of care. Generally, LOSs have decreased steadily over the years
36 as a result of surgical innovations and improvements in perioperative management
37 [1][2]. Shorter LOSs are also being promoted in systematic developments such as the
38 introduction of prospective payment systems and diagnosis-related groups for hospital
39 reimbursements [3], which ensure payment of a given amount for a procedure
40 regardless of the hospitalisation episode. However, hospitals are constantly under
41 clinical and financial pressure to become more cost efficient and LOS is a common
42 target for cost-containing initiatives. Intuitively, the decision to discharge a patient after
43 surgery should be a medical one. The type and extent of surgery, as well as various
44 intraoperative variables, will play a role in how a patient feels after the operation and
45 how long their hospital stay will be. A longer, more complex surgery, with a greater level
46 of surgical invasiveness [4] can be expected to result in a longer stay than would a
47 shorter, more straightforward surgery. However, surgical variables are likely not the only
48 factors governing the LOS; for any given procedure, with its given degree of complexity,
49 various non-surgical and even non-medical factors can also be expected to play a role
50 and partially account for inter-individual differences in LOS.

51 Only few data are available on non-medical or sociodemographic/socioeconomic (SDE)
52 predictors of LOS in relation to spine surgery. In addition, some of the existing data are
53 inconsistent. Previous studies retrospectively reviewing factors related to prolonged
54 LOS have identified increased age as an important factor, explained by the fact that
55 aged tissue takes physiologically longer to heal and older patients may be less able to
56 function independently at home upon discharge. Other non-medical factors such as
57 insurance status, race/ethnicity or geographic location were found to be significant
58 predictors of prolonged LOS in anterior cervical spine procedures [5]. Female gender
59 has been found to be a significant predictor of LOS in some studies [6, 7] while other
60 reports have found no relationship between gender and LOS [8]. Often, the identification
61 of a given variable as a risk factor (or not) is influenced by the inclusion, in the model, of

62 other closely related or "proxy" measures of that variable. Furthermore, some of the
63 prior LOS studies in the spine literature have substantial limitations such as small
64 sample sizes with inadequate statistical power, examination of just one specific
65 pathology, or a study design in which important medical confounders of LOS were not
66 included in the predictor models [7, 9].

67 Against this background, the present study sought to determine the influence of non-
68 medical factors on LOS after surgery for degenerative spine disorders. It involved the
69 data from a large consecutive series of patients, collected over 13 years, and made use
70 of multivariable regression models to account for a high quality set of potential
71 confounders such as the invasiveness of surgery, surgical complications and other
72 intraoperative factors.

73 **Patients and methods**

74 The study was a retrospective analysis of data that had been collected prospectively
75 from consecutive patients operated in our Spine Centre, part of a tertiary care
76 orthopaedic hospital in Switzerland. The data were collected in our local spine
77 outcomes database between 2005 and 2017 using the framework of the EUROSPINE
78 Spine Tango Registry (<https://www.eurospine.org/spine-tango.htm>) and were
79 supplemented with additional data (insurance status, nationality, etc.) systematically
80 imported into the database from the clinic information system. The use of this routinely
81 collected data, given with the patients' informed consent, was approved by the Ethics
82 Committee (KEK-ZH-Nr 2014-0418).

83

84 We identified 13,368 consecutive patients who were documented as having undergone
85 spine surgery for degenerative diseases of the spine, with either a primary intervention
86 or revision surgery. The cervical interventions primarily included anterior or posterior
87 fusion (\pm instrumentation) procedures. Thoracic/lumbar interventions included posterior
88 decompression and/or spondylodesis with rigid stabilization either with posterior and/or
89 interbody fusion (open or mini-open/MISS), as well as complex deformity corrections.
90 Overall, 10,770 of 13,368 patients were retained in complete case analyses (19% lost

91 due to missing data; i.e. “listwise deletion”; Fig 1). Their characteristics are described in
92 Table 1.

93 Of the variables documented in the local database, predictors of LOS (i.e. number of
94 days from surgery to discharge) were chosen based on the available literature, and
95 were categorised into blocks representing particular stages of the clinical pathway:

96 1) Non-medical patient characteristics; 2) Preoperative medical patient characteristics;
97 3) Surgical details; 4) Perioperative outcomes. The patient demographics and variables
98 considered in the analysis are illustrated in Table 1. Non-medical patient characteristics
99 comprised age, sex, BMI, smoking status, nationality and insurance status.

100 Preoperative medical characteristics included the Core Outcome Measures Index
101 (COMI)[10] pre-operative score (measure of symptoms/disease burden/reduced quality
102 of life before surgery), region operated (cervical vs thoracic/lumbar spine), morbidity
103 state (American Society of Anesthesiologists (ASA) class [11]), number of affected
104 segments, and previous surgery involving the same spine level. It must be noted that
105 there was an unknown degree of uncertainty around the spinal level of the repeat
106 surgical procedures included in the analysis because this item is not specifically
107 measured in Spine Tango for each recorded instance of repeat surgery (only “number of
108 previous surgeries” and whether “any previous surgery done on the same level”). An
109 assumption was therefore made that cases of multiple repeat procedures were likely to
110 involve the same level repeatedly and the predictor was constructed accordingly. It is
111 shown in the models with the caveat “at least one previous surgery involving same
112 level”. As the predictor derived this way exhibited qualities suggesting a degree of
113 ordinality in the initial statistical models, it was retained despite the limitation. Surgical
114 procedures were characterised by whether instrumented fusion was used or not and
115 whether the surgery was conventional or minimally/less invasive (MISS/LISS). In a
116 subset of data collected using a later iteration of the Spine Tango form, introduced in
117 2012 (Fig 1), more detailed surgical information was available, allowing calculation of a
118 surgical invasiveness index score for each case (based on the formulae of Mirza et al.
119 [4]). This represented a composite score, given by the number of levels operated and
120 whether anterior and/or posterior decompression, fusion and/or stabilisation procedures
121 had been carried out. The rich set of potential predictors was completed by the

122 perioperative outcome variables blood loss, duration of surgery, general medical
123 complications and surgical complications arising during surgery and before discharge,
124 as documented on the Spine Tango form.

125 The local registry data represented a nearly complete record (99% coverage) of all
126 relevant surgical procedures in the target period and the missing 1% were not expected
127 to bias the results in either direction. The 19% cases excluded from the analysis by
128 listwise deletion of those without a full complement of predictor variables appeared to be
129 a random sub-sample of all potential participants, based on observable characteristics.

130 >>>>Insert Table 1 here<<<<<

131 The dataset was analysed through a series of stepwise linear regression models. We
132 entered the aforementioned blocks of variables sequentially, retaining or dropping
133 variables after each step. Predictors were retained in models if they showed at least
134 borderline statistical significance ($p < 0.1$) and if they proved generally robust in the
135 context of a growing number of covariates. The distributional properties of the outcome
136 variable made it suitable for standard linear regression (Gaussian distributional
137 assumption / identity link function), resulting in model residuals following an
138 approximately normal distribution. Alternative GLM specifications were initially
139 considered but deemed unnecessary. Continuous predictors were entered as centred
140 variables where this was deemed useful in order to facilitate model interpretation. This
141 allowed interpretation of the constant as the predicted length of stay for an “average”
142 patient with reference values of the categorical predictors. All analyses were done in
143 Stata 15.1.

144 **Results**

145 The mean (\pm SD) LOS was 7.9 ± 5.2 days. The main regression models are shown in
146 Table 2. Model 1 includes the non-medical patient characteristics age, sex and BMI.
147 The latter accounted for 13% of variance. We also tested for effects of smoking status,
148 nationality (Swiss vs. other nationality) and type of health insurance, each of which was
149 statistically insignificant. Preoperative medical patient characteristics were added in
150 model 2, and showed that LOS was associated strongly and positively with previous

151 surgeries, COMI-score, morbidity and the number of affected spinal segments. The
152 model's predictive power improved to explain nearly 23% of variance. Key surgical
153 details were added in model 3. The number of cases in which MISS/LISS procedures
154 were used was very small (see Table 1), but minimally invasive procedures were
155 nonetheless associated with reduced LOS (-1.5 days), while the use of rigid stabilisation
156 was associated with a pronounced increase of LOS (+3.8 days). The final model
157 included a set of perioperative outcomes: in addition to the amount of blood loss and the
158 duration of surgery, we observed particularly strong effects for the presence of a
159 general medical complication (+2.2 days) and of a surgical complication (+4.3 days).
160 Predictive power peaked at nearly 43% of variance and the predictors included in
161 previous steps generally retained their effect directions, although they naturally revealed
162 gradually reducing effect sizes. Notably, female patients, even after controlling for a
163 large set of covariates that were strong determinants of LOS, had a systematically
164 longer LOS than male patients (+1.1 days). Further interaction analysis of model 4
165 (details not shown) revealed that the influence of sex on LOS varied somewhat by age
166 group. Table 2 shows the average effect over the entire age distribution, and, whilst an
167 effect was present in all age groups, it was shown to increase with age: in the age group
168 under 45 y, women stayed 0.86 days longer than men; in the age group 45-64 y, 1.02
169 days longer; and in the age group 65+ y, 1.29 days longer.

170 The "average patient" in model 4 (a male recipient of back surgery of average age and
171 BMI with a low risk profile, e.g. no complications or previous surgery, low blood loss and
172 an uncomplicated standard surgical approach) was predicted to spend 4.8 days in
173 hospital (model constant).

174 An alternative model specification relying on the more parsimonious invasiveness index
175 (replacing the individual variables MISS, rigid stabilisation, number of affected levels,
176 blood loss and duration of surgery)¹ showed the same significant predictors as the full
177 model, but previous surgery had a higher degree of stability in the model, the effect of

¹ Calculation of this score was only possible for the data collected since 2012 (see earlier) and therefore the model is based on a smaller, more recent subsample of patients (n=5,409) (Fig 1).

178 BMI achieved statistical significance, and gender still showed approximately the same
179 effect size, with females having +1.2 days longer LOS than males (Table 3).

180 Finally, given the long observation period (2005-2017), additional tests were carried out
181 for possible effects of time. However, whilst mean LOS varied from year to year there
182 was no clear trend towards either lower or higher LOS and no effect could be detected
183 in the statistical models (controlling for individual years or pooled time periods).

184

185 >>>>insert Table 2 here<<<<<<

186 >>>>insert Table 3 here<<<<<<

187

188 **Discussion**

189 This study was designed to determine the influence of non-medical factors on LOS after
190 surgery for degenerative spinal disorders, whilst controlling for known medical and
191 surgical determinants of LOS. Female gender and age, in particular, and (depending on
192 the model in question) BMI were each found to be significantly and independently
193 associated with a longer LOS, whilst insurance status, being of Swiss nationality, and
194 being a smoker were non-significant. The non-medical factors accounted for almost
195 one-third of the total variance in LOS explained by the model.

196 We intentionally included all relevant medical/surgical factors in our predictor models in
197 order to ascertain the unique influence of the sociodemographic/economic factors. The
198 effect of age was not simply a reflection of greater comorbidity or a higher incidence of
199 complications, since these factors were controlled for in the models by inclusion of the
200 ASA grade [12] and the presence of medical/surgical complications as documented on
201 the Spine Tango form. Instead, it is likely that the finding is explained by older people
202 recovering more slowly from surgery and anaesthesia (as well as from any associated
203 complications), with older tissue taking longer to heal. The effect of age may also reflect
204 a sociological component, in that older patients may be considered less able to function

205 independently at home upon discharge. Our findings regarding age as a predictor of
206 LOS are in line with almost all previous studies [7, 8, 13–15]. Age, of course, is not a
207 modifiable risk factor and hence there are few practical implications that can emerge
208 from this finding; nonetheless, a knowledge of the independent influence of age on LOS
209 could allow for better discharge planning and the consideration of less costly
210 alternatives for postoperative care.

211 A significant finding from our study was that gender seemed to play a relevant role in
212 governing LOS. The effect was consistent across different age groups, although more
213 marked in older patients. All else being equal, LOS was just over one day longer for
214 women undergoing spine surgery than for men. A similar gender difference in LOS has
215 been reported before, in two small studies in the spine literature [7, 9] and also in
216 studies of patients undergoing shoulder [16], knee or hip surgery [17], although it has
217 not been a consistent finding in all [5, 18] and few studies have demonstrated an effect
218 beyond that mediated by an increase in complications in women. A very recent analysis
219 of the NSQIP database by Heyer et al. 2019 [19], also reported that female patients
220 stayed significantly longer in hospital. The authors hypothesised that this may have
221 been due to the greater need for blood transfusion, higher incidence of urinary tract
222 infections (UTIs), or more dependent preoperative status of women necessitating
223 subsequent discharge to nursing or rehabilitation facilities; however, despite having
224 measured these explanatory variables, they did not go on to include them in a
225 multivariable model to test their hypothesis with respect to LOS [19]. In our study,
226 complications as well as preoperative status were included in the multivariable model in
227 which female gender still exerted an independent effect on LOS, and hence these
228 factors are unlikely to account for our findings. We were unable to account for the
229 severity of complications, and, if women had suffered more severe complications than
230 men, this could possibly have explained their longer LOS; however, at least based on
231 the types of complications recorded, there was no evidence to suggest that this was the
232 case. Another possible explanation for why women stay longer in hospital after surgery
233 is that more elderly women than men tend to live alone, since men have a shorter life
234 expectancy. A lack of family support has been shown to be associated with a prolonged
235 LOS [20]. A further possible explanation is that women typically experience slightly

236 higher pain levels than men after surgery and for some degenerative disorders may be
237 less satisfied with the surgical result [21], leading the medical team to err on the side of
238 caution, delaying discharge to allow longer for recovery. In the past, women were
239 considered to be the weaker sex and this idea might still prevail in the minds of either
240 patients or doctors, biasing their appraisal of readiness for discharge. Interestingly, a
241 previous study showed that spine surgeons tend to over-rate the 12-month outcome in
242 men more so than in women, in comparison to the patient's own ratings [22]. Whether or
243 not this is also the case in the early stages after surgery, remains to be known. A
244 previous study showed that a lack of paid employment was significantly associated with
245 a longer hospital stay after spine surgery [23]; although the findings were not specifically
246 related to gender, women are still less likely to be in paid employment than men. It is
247 also conceivable that the "working man" wishes to get back to his comfortable home
248 environment more readily than does the "home-maker" wife, for whom a return to home
249 effectively constitutes a return to work. Policy makers may wish to clarify the possible
250 explanations for the gender difference and evaluate whether less costly social (rather
251 than medical) care could be used to assist with earlier discharge in women. Discharge
252 planning prior to admission with a focus on a clear endpoint in the patient's care could
253 ensure that all the support processes are in place to help the discharge happen in a
254 timely fashion [24].

255 In the present study, BMI had a slight (but in most models statistically significant)
256 influence on LOS. The impact of a higher BMI or obesity on LOS is controversial, and
257 may depend on the other variables included in the given predictor models. Obesity may
258 increase the risk of postoperative complications, exerting its effect through greater
259 comorbidity or by simply making access to the surgical site more difficult. In the present
260 study, these covariates were controlled for in our model, perhaps minimising any effect
261 of BMI *per se*. Kalanithi et al showed that spinal fusion patients who were morbidly
262 obese were at significantly increased risk of a longer hospital stay, generating greater
263 hospital charges [25]. Possibly, in this respect, preoperative weight loss programmes or
264 lifestyle modification counselling may prove cost effective and at the same time allow
265 patients to benefit from improved postoperative outcomes. However, in the very obese,
266 the necessary weight loss may be difficult to achieve, and if it requires closely

267 supervised programmes and/or other medical interventions, then this may offset any
268 potential savings in terms of LOS.

269 Factors such as smoking status, nationality and health insurance were not significantly
270 associated with LOS in our study. The literature is inconsistent regarding the influence
271 of smoking status on LOS. Using adjusted analysis, Seican et al. [26] reported that prior
272 smokers undergoing elective spine surgery were significantly more likely to have
273 prolonged hospitalisation and major complications compared with never smokers. It is
274 unclear why smoking showed no significant effect in the present study. We documented
275 current smoking habit rather than a history of smoking and it was based on self-
276 declaration at the time of surgery; possibly, social desirability bias or the adoption of a
277 period of smoking cessation just prior to surgery may have influenced the accuracy of
278 the responses. That nationality and health insurance had no significant effect on LOS
279 was encouraging, and suggested equitable hospital policy was in place with regard to
280 this aspect of care.

281 Despite including a large sample of data, collected systematically on a prospective
282 basis over many years, our study has certain limitations that require mention. First, it is
283 based on a secondary analysis of registry data supplemented with other routinely
284 collected clinical data and, as such, data items were not specifically provided and
285 designed to answer the research question. Additional important confounders may have
286 been omitted by virtue of the study design. Also, it involved the data from just one
287 orthopaedic hospital in Switzerland. The results should ideally be confirmed in a
288 multicentre study. This could be done using data collected with the same documentation
289 forms from other reliable centres in the EUROSPINE Spine Tango registry, and
290 employing the exact same approach to the statistical models. Whilst the extent of
291 surgery and predominant surgical technique (instrumented fusion or not; use of MISS)
292 were controlled for in the model, the degenerative disorders were only grouped as either
293 cervical or thoracic/lumbar, without further differentiation or comparison between
294 diagnoses or specific operative procedures. This will be addressed in our future studies,
295 to create more detailed surgical models. Although standardised forms were used for
296 data collection, there may have been inconsistencies in documentation between the

297 different surgeons. And finally, a variable that was not included in the present study but
298 that has been shown to be relevant in relation to LOS is the discharge destination [27].
299 In terms of achieving a more complete understanding of how non-medical factors affect
300 LOS, the post-operative circumstances the patient has to face, such as going into a
301 rehabilitation program or going home to live alone, should be known.

References

1. Pumberger M, Chiu Y-L, Ma Y, et al (2012) National in-hospital morbidity and mortality trends after lumbar fusion surgery between 1998 and 2008. *J Bone Joint Surg Br* 94-B:359–364. <https://doi.org/10.1302/0301-620x.94b3.27825>
2. Sivasubramaniam V, Patel HC, Ozdemir BA, Papadopoulos MC (2015) Trends in hospital admissions and surgical procedures for degenerative lumbar spine disease in England: A 15-year time-series study. *BMJ Open* 5:1–7. <https://doi.org/10.1136/bmjopen-2015-009011>
3. MedlinePlus Medical Encyclopedia (2013) Health Care Financing Trends. *Health Care Financ Rev* 9:127–132
4. Mirza SK, Deyo RA, Heagerty PJ, et al (2008) Development of an Index to Characterize the “Invasiveness” of Spine Surgery. *Spine (Phila Pa 1976)* 33:2651–2661. <https://doi.org/10.1097/brs.0b013e31818dad07>
5. Alesh H, Li D, Riley LH, Skolasky RL (2015) Health Care Burden of Anterior Cervical Spine Surgery. *J Spinal Disord Tech* 28:5–11. <https://doi.org/10.1097/bsd.0000000000000001>
6. Saltzman BM, Basques B, Leroux T, et al (2018) The influence of gender on early adverse events , hospital charges and length of stay after shoulder arthroplasty. 149–155. <https://doi.org/10.1007/s00264-017-3547-6>
7. Arnold M, Rice LR, Anderson KK, et al (2011) Factors affecting hospital length of stay following anterior cervical discectomy and fusion. 2:11–18
8. Basques BA, Fu MC, Buerba RA, et al (2014) Using the ACS-NSQIP to identify factors affecting hospital length of stay after elective posterior lumbar fusion. *Spine (Phila Pa 1976)* 39:497–502. <https://doi.org/10.1097/BRS.0000000000000184>
9. Sultan AA, Berger RJ, Cantrell WA, et al (2019) Predictors of Extended Length of Hospital Stay in Adolescent Idiopathic Scoliosis Patients Undergoing Posterior Segmental Instrumented Fusion. *Spine (Phila Pa 1976)* 44:715–722. <https://doi.org/10.1097/BRS.0000000000002919>

10. Mannion AF, Porchet F, Kleinstück FS, et al (2009) The quality of spine surgery from the patient's perspective. Part 1: The Core Outcome Measures Index in clinical practice. *Eur Spine J* 18:367–373. <https://doi.org/10.1007/s00586-009-0942-8>
11. De Cassai A, Boscolo A, Tonetti T, et al (2019) Assignment of ASA-physical status relates to anesthesiologists' experience: A survey-based national-study. *Korean J Anesthesiol* 72:53–59. <https://doi.org/10.4097/kja.d.18.00224>
12. Bjorgul K, Novicoff WM, Saleh KJ (2010) American society of anesthesiologist physical status score may be used as a comorbidity index in hip fracture surgery. *J Arthroplasty* 25:134–137. <https://doi.org/10.1016/j.arth.2010.04.010>
13. Nie H, Hao J, Peng C, Ou Y (2013) Clinical Outcomes of Discectomy in Octogenarian Patients With Lumbar Disc Herniation. 26:74–78
14. Zheng F, Cammisa FP, Sandhu HS, et al (2002) Factors predicting hospital stay, operative time, blood loss, and transfusion in patients undergoing revision posterior lumbar spine decompression, fusion, and segmental instrumentation. *Spine (Phila Pa 1976)* 27:818–24
15. Gruskay JA, Fu M, Basques BA, et al (2016) Factors Affecting Length of Stay and Complications After Elective Anterior Cervical Discectomy and Fusion National Surgical Quality Improvement Project Database (ACS NSQIP). 29:34–42
16. Dunn JC, Lanzi J, Kusnezov N, et al (2015) Predictors of length of stay after elective total shoulder arthroplasty in the United States. *J Shoulder Elb Surg* 24:754–759. <https://doi.org/10.1016/j.jse.2014.11.042>
17. Husted H, Holm G, Jacobsen S (2008) Predictors of length of stay and patient satisfaction after hip and knee replacement surgery: Fast-track experience in 712 patients. *Acta Orthop* 79:168–173. <https://doi.org/10.1080/17453670710014941>
18. Herren C, Aghayev E, Kaulhausen T, et al (2014) Einflussfaktoren auf die Verweildauer in der WirbelsäulenchirurgieInfluencing factors on the length of stay in lumbar spine surgery. *Orthopade* 43:1043–1051. <https://doi.org/10.1007/s00132-014-3033-4>

19. Heyer JH, Cao N, Amdur RL, Rao RR (2019) Postoperative Complications Following Orthopedic Spine Surgery: Is There a Difference Between Men and Women? *Int J Spine Surg* 13:6017. <https://doi.org/10.14444/6017>
20. Shelton W, Moore CD, Socaris S, et al (2010) The effect of a family support intervention on family satisfaction, length-of-stay, and cost of care in the intensive care unit. 38:1315–1320. <https://doi.org/10.1097/CCM.0b013e3181d9d9fe>
21. Pochon L, Kleinstück FS, Porchet F, Mannion AF (2016) Influence of gender on patient-oriented outcomes in spine surgery. *Eur Spine J* 25:235–246. <https://doi.org/10.1007/s00586-015-4062-3>
22. Porchet F, Lattig F, Grob D, et al (2010) Comparison of patient and surgeon ratings of outcome 12 months after spine surgery. *J Neurosurg Spine* 12:447–455. <https://doi.org/10.3171/2009.11.spine09526>
23. Walid MS, Robinson ECM, Robinson JS (2011) Higher comorbidity rates in unemployed patients may significantly impact the cost of spine surgery. *J Clin Neurosci* 18:640–644. <https://doi.org/10.1016/j.jocn.2010.08.029>
24. Gonçalves-Bradley DC, Lannin NA, Clemson LM, et al (2016) Discharge planning from hospital (Review) SUMMARY OF FINDINGS FOR THE MAIN COMPARISON. *Cochrane Database Syst Rev* 74. <https://doi.org/10.1002/14651858.CD000313.pub5.www.cochranelibrary.com>
25. Kalanithi PA, Arrigo R, Boakye M (2012) Morbid obesity increases cost and complication rates in spinal arthrodesis. *Spine (Phila Pa 1976)* 37:982–988. <https://doi.org/10.1097/BRS.0b013e31823bbef>
26. Seicean A, Seicean S, Alan N, et al (2013) Effect of smoking on the perioperative outcomes of patients who undergo elective spine surgery. *Spine (Phila Pa 1976)* 38:1294–1302. <https://doi.org/10.1097/BRS.0b013e31828e2747>
27. Kulinskaya E, Kornbrot D, Gao H (2005) Length of stay as a performance indicator: Robust statistical methodology. *IMA J Manag Math* 16:369–381. <https://doi.org/10.1093/imaman/dpi015>

Table 1. Predictor variables as used in the statistical models (n=10770)

Continuous	Mean (SD)	Median (Min-Max)
Age at OP	62.3 (14.5)	64 (6-94)
BMI score	26.4 (4.7)	26 (13-56)
COMI pre-score (0 (best) to 10 (worst) status) ¹	7.5 (1.9)	7.9 (0-10) ²
Duration of surgery (hours)	2.7 (1.7)	2.5 (0.5-10)
Invasiveness Index based on using Spine Tango formula (relevant data available from 2011; n=5,409)	7.9 (8.9)	6.0 (0-84)
Categorical	n	%
Sex		
Male	5,056	47.0
Female	5,714	53.0
Smoker ²		
No	7,540	74.7
Yes	2,550	25.3
Nationality ²		
Swiss	9,429	87.5
Other	1,341	12.5
Insurance status ²		
Privately insured	3,184	29.6
Semi-privately insured	2,890	26.8
General public insurance scheme	4,696	43.6
Previous surgery involving same level		
None	8,034	74.6
1 involving same level	1,495	13.9
2 (at least one involving same level)	678	6.3
3 or more (at least one involving same level)	563	5.2
Spine location		
Cervical	1,663	15.4
Thoracic or lumbar	9,107	84.6

¹ 0.19% of patients reported a baseline COMI score of 0, and 0.56% of ≤ 1 . These scores may seem unlikely, but we have no reason to assume that the responses are invalid. COMI is a self-report measure and thus contains subjective elements. Moreover, most of these patients had conditions that are not typically very symptomatic in terms of pain and disability but are at risk of serious progression if not treated e.g. early cervical myelopathy.

² These covariates were not retained in the final models due to statistical insignificance and instability likely caused by multi-collinearity (smoking status and insurance status correlate with age)

Morbidity state (ASA)		
No disturbance	2,142	19.9
Mild/moderate	5,490	51.0
Severe or worse	3,138	29.1
Type of surgery		
Conventional techniques	10,515	97.6
MISS or LISS	255	2.4
Extent of lesion (number of affected levels)		
1 segment	5,202	48.3
2-3 segments	4,705	43.7
4-5 segments	640	5.9
6 or more segments	223	2.1
Rigid stabilisation (anterior or posterior)		
No	5,323	49.4
Yes	5,447	50.6
Blood loss		
<100ml	2,593	24.1
100-500ml	6,090	56.6
500-1000ml	1,295	12.0
More than 1000ml	792	7.4
Any general medical complication ³		
No	9,956	92.4
Yes	814	7.6
Any surgical complication ⁴		
No	9,885	91.8
Yes	885	8.2

³ General medical complications documented on the Spine Tango form included: anaesthesiological, cardiovascular, pulmonary, cerebral, kidney/urinary, liver/GI, death, other.

⁴ Surgical complications included: wrong level, vascular, neurological, implant malposition, dural tear, wound infection, implant failure, other. An alternative statistical model in which we specifically included the detailed complication information (i.e. the different complication types) did not change the overall findings regarding the significant predictors.

Table 2. Main regression models (full dataset)

VARIABLES	MODEL (1) Coefficient [95% CI]	MODEL (2) Coefficient [95% CI]	MODEL (3) Coefficient [95% CI]	MODEL (4) Coefficient [95% CI]
Age at OP (centred)	0.10*** [0.098 - 0.11]	0.058*** [0.050 - 0.065]	0.064*** [0.057 - 0.071]	0.060*** [0.053 - 0.066]
Sex: female	1.91*** [1.73 - 2.10]	1.70*** [1.52 - 1.88]	1.12*** [0.95 - 1.28]	1.11*** [0.96 - 1.27]
BMI score (centred)	0.096*** [0.077 - 0.12]	0.036*** [0.017 - 0.055]	0.038*** [0.020 - 0.056]	0.015* [-0.0011 - 0.032]
Previous surgery involving same level None = reference				
1 involving same level		0.67*** [0.41 - 0.92]	-0.017 [-0.26 - 0.22]	-0.28** [-0.51 - -0.061]
2 (at least one involving same level)		1.68*** [1.32 - 2.05]	0.55*** [0.21 - 0.89]	0.17 [-0.15 - 0.49]
3 or more (at least one involving same level)		1.96*** [1.56 - 2.35]	0.83*** [0.46 - 1.21]	0.46*** [0.12 - 0.81]
COMI pre-score (centred)		0.18*** [0.13 - 0.22]	0.15*** [0.11 - 0.20]	0.12*** [0.075 - 0.16]
Spine location: Thoracic/lumbar = reference Spine location: cervical		-1.49*** [-1.74 - -1.24]	-3.01*** [-3.26 - -2.77]	-2.19*** [-2.43 - -1.95]
Morbidity state (ASA): no disturbance = reference				
Morbidity state (ASA): mild/moderate		0.78*** [0.52 - 1.05]	0.017 [-0.23 - 0.26]	-0.11 [-0.34 - 0.12]
Morbidity state (ASA): severe or worse		2.24*** [1.92 - 2.57]	1.41*** [1.11 - 1.72]	0.89*** [0.61 - 1.18]
Extent of lesion: 1 segment = reference				
Extent of lesion: 2-3 segments		1.19*** [1.00 - 1.38]	0.88*** [0.70 - 1.05]	0.39*** [0.22 - 0.56]
Extent of lesion: 4-5 segments		2.71*** [2.33 - 3.09]	2.36*** [2.01 - 2.72]	0.93*** [0.58 - 1.27]
Extent of lesion: 6 or more segments		4.53*** [3.90 - 5.15]	3.49*** [2.91 - 4.07]	1.55*** [0.99 - 2.11]

Type of surgery: conventional technologies = reference Type of surgery: MISS or LISS			-1.46*** [-1.99 - -0.93]	-1.17*** [-1.67 - -0.67]
Rigid stabilisation (anterior or posterior)			3.79*** [3.61 - 3.97]	2.34*** [2.11 - 2.56]
Blood loss: <100ml = reference Blood loss: 100-500ml Blood loss: 500-1000ml Blood loss: more than 1000ml				0.57*** [0.36 - 0.77] 1.49*** [1.16 - 1.83] 1.61*** [1.19 - 2.03]
Duration of surgery (centred)				0.42*** [0.34 - 0.50]
Any general medical complication ¹				2.23*** [1.94 - 2.52]
Any surgical complication ¹				4.25*** [3.97 - 4.53]
Constant	6.81*** [6.67 - 6.94]	5.03*** [4.77 - 5.29]	4.72*** [4.48 - 4.97]	4.83*** [4.54 - 5.12]
Observations	10,770	10,770	10,770	10,770
Adjusted R-squared	0.131	0.226	0.333	0.424

*** p<0.01, ** p<0.05, * p<0.1

¹An alternative statistical model in which we specifically included the more detailed complication information (i.e. the different complication types) did not change the overall findings regarding the significance of the other variables explaining LOS.

Table 3. Alternative regression model using invasiveness index (calculable in data collected after 2012)

VARIABLES	Coefficient [95% CI]	
Age at OP (centred)	0.055***	[0.044 - 0.065]
Sex: female	1.15***	[0.91 - 1.40]
BMI score (centred)	0.059***	[0.034 - 0.085]
Previous surgery involving same level None = reference		
1 involving same level	-0.0064	[-0.35 - 0.34]
2 (at least one involving same level)	0.86***	[0.38 - 1.35]
3 or more (at least one involving same level)	0.86***	[0.34 - 1.39]
COMI pre-score (centred)	0.12***	[0.052 - 0.18]
Spine location: thoracic/lumbar = reference		
Spine location: cervical	-1.34***	[-1.68 - -1.01]
Morbidity state (ASA): no disturbance = reference		
Morbidity state (ASA): mild/moderate	0.34	[-0.072 - 0.75]
Morbidity state (ASA): severe or worse	1.18***	[0.69 - 1.67]
Any general medical complication	2.46***	[1.96 - 2.95]
Any surgical complication	5.83***	[5.42 - 6.25]
Invasiveness index¹	0.22***	[0.21 - 0.24]
Constant	4.27***	[3.87 - 4.66]
Observations	5,409	
Adjusted R-squared	0.393	

*** p<0.01, ** p<0.05, * p<0.1

¹ based on the formulae of Mirza et al [4] but calculated on the basis of the fields completed on the Spine Tango Surgery form

Figure 1 Data flow diagram

