

***Haloferax volcanii* for biotechnology applications: challenges, current state and perspectives**

Haque, R. U.^{1,2,3}, Paradisi, F.^{2,4} & Allers, T^{1*}

1. School of Life Sciences, Queens Medical Centre, University of Nottingham, Nottingham NG7 2UH, UK

2. School of Chemistry, University Park, University of Nottingham, Nottingham NG7 2RD, UK

3. Warwick Integrative Synthetic biology Centre, School of Life Sciences, Gibbet Hill Campus, University of Warwick, Coventry CV4 7AL, UK

4. Department of Chemistry and Biochemistry, University of Bern, Freiestrasse 3, 3012 Bern, Switzerland

*Corresponding author thorsten.allers@nottingham.ac.uk Tel. +44 115 823 0304

ORCID ID: Haque, R. U. (0000-0003-1513-459X), Paradisi, F. (0000-0003-1704-0642), Allers, T. (0000-0002-7308-3332)

Abstract

Haloferax volcanii is an obligate halophilic archaeon with its origin in the Dead Sea. Simple laboratory culture conditions and a wide range of genetic tools have made it a model organism for studying haloarchaeal cell biology. Halophilic enzymes of potential interest to biotechnology have opened up the application of this organism in biocatalysis, bioremediation, nanobiotechnology, bioplastics and the biofuel industry. Functionally active halophilic proteins can be easily expressed in a halophilic environment, and an extensive genetic toolkit with options for regulated protein overexpression has allowed the purification of biotechnologically important enzymes from different halophiles in *H. volcanii*. However, corrosion mediated damage caused to stainless-steel bioreactors by high salt concentrations and a tendency to form biofilms when cultured in high volume are some of the challenges of applying *H. volcanii* in biotechnology. The ability to employ expressed active proteins in immobilised cells within a porous biocompatible matrix offers new avenues for exploiting *H. volcanii* in biotechnology. This review critically evaluates the various application potentials, challenges and toolkits available for using this extreme halophilic organism in biotechnology.

Keywords: *Haloferax volcanii*, Biotechnology, Protein Expression, Immobilisation, Model Organism

Acknowledgements

Funding

This work was supported by the Biotechnology and Biological Sciences Research Council (BBSRC) [grant number BB/P002536/1]. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author contributions

All authors contributed equally to writing the manuscript.

Introduction

Halophiles are extremophilic microbes that are widespread in all three domains of life –Eukarya, Bacteria and in particular, Archaea (Kamekura 1998; Oren 2013). They are found in hypersaline environments around the world where the salt concentrations is higher than sea water (around 3.5% dissolved salt), in various geographical niches such as salt lakes, deep sea, salt pans, saline soils, salt mines or salt marshes (Oren 2015). Based on their salt preference, halophiles can be split into three different categories: mild halophiles growing optimally at 1%-3% (0.2 – 0.5 M) NaCl, moderate halophiles with optimum growth at 3%-15% (0.5 M – 2.5 M), and extreme halophiles with optimal growth at 15%-20% (2.5 – 5.2 M) NaCl (Amoozegar et al. 2017; De Lourdes Moreno et al. 2013). Below 0.2 M NaCl (<1%), microbes are classified as non-halophiles and a key difference is that the growth of mesophilic organisms is impaired at higher salinities, where archaeal species dominate. All halophilic archaea belong to the Euryarchaeota kingdom (Oren et al. 2009).

Halophiles have been important in natural salt production since they facilitate salt crystallization through the absorption of solar radiation using their carotenoid pigments, which confers distinct pink-reddish hues to the saltern crystallizer ponds they inhabit. More recently, halophiles have been used in biotechnology as producers of antimicrobials, bioplastics, biofuel, extremozymes, extracellular polysaccharides, retinal proteins, coloured pigments, and compatible solutes (Amoozegar et al. 2019; Atanasova et al. 2013; Koller 2019; Patel and Saraf 2015; Torregrosa-Crespo et al. 2017). This has led to their applications in astrobiology, agriculture, food and nutrition industry, biosensor development and photochemical industry, bioremediation, textile and tanning industry, aquaculture, nanotechnology, sustainable chemical production, and medical application (Le Borgne et al. 2008; Lee 2013; Margesin and Schinner 2001; Rodrigo-Baños et al. 2015). Several reviews on the biotechnologically valuable products generated by halophiles and their potential applications have been recently published (Antunes et al. 2017; DasSarma et al. 2010a; Mohammadipanah et al. 2015; Oren 2010; Yin et al. 2015).

Halophiles are under-represented in biotechnology compared with other extremophiles such as thermophiles or alkaliphiles, which have been used extensively for valuable product generation. This is surprising considering the biotechnological potential of halophiles and their great diversity across all three domains of life. While some halophiles are mechanically fragile, prone to lysis in low salt environment, and their culturing can lead to possible damage of the stainless equipment (due to the required molar concentrations of salt), some examples of their successful application in industrial biotechnology are represented by the application of purple membrane protein bacteriorhodopsin (from *Halobacterium salinarum*) in photochemical application, ectoine, an enzyme immobiliser used in the cosmetic industry (from *Halomonas elongata*), and β -carotene production (from green algae *Dunaliella salina*) (Margesin and Schinner 2001; Oren 2010).

H. volcanii is an extremely halophilic archaeon belonging to the family Haloferacaceae. Easy growth conditions, a short generation time and facile genetics have made it a model organism for haloarchaeal biology (Allers and Mevarech 2005; Leigh et al. 2011). It encodes numerous biotechnologically attractive halophilic enzymes with application potentials in sustainable fine chemical synthesis, bioprocessing, bioremediation, biofuel and bioplastic production (Amoozegar et al. 2019; Koller 2019; Lobasso et al. 2015; Timpson et al. 2013; Uthandi et al. 2009).

Developments of tools such as bioreactor-scale overexpression of functionally active proteins and immobilisation of cells within a porous matrix have opened up new possibilities in exploiting *H. volcanii* for biotechnological applications (Haque et al. 2019; Strillinger et al. 2016). This review summarises the biotechnologically relevant features of this organism, the challenges involved and the tools available for realising its true biotechnological potential.

Genetics and biochemistry of *H. volcanii*

H. volcanii was first isolated from the sediment of the Dead Sea (Mullakhanbhai and Larsen 1975). The organism originally described as *Halobacterium volcanii* was named after the microbiologist Benjamin Elazari-Volcani who reported the presence of indigenous microbial life in the salt rich Dead Sea (Elazari-Volcani, 1943). It grows optimally at 45°C with a generation time of 2 hour (Robinson et al. 2005). It is simple to culture in the laboratory since it grows aerobically in complex, minimal and synthetic media (Mevarech and Werczberger 1985). Compared to other extreme halophiles, *H. volcanii* tolerates a wide range of salt concentrations (1.8 – 3.5 M NaCl) (Allers 2010). It is comparatively resistant to contamination since few other organisms can survive the molar concentration of salt present in the growth media. It can grow up to a temperature of 50°C (Dyall-Smith 2009). It lacks a rigid cell wall and instead presents a single layer of glycoprotein known as surface layer or S-layer (Rodrigues-Oliveira et al. 2017). Protein subunits in S-layers are held together by divalent cation such as Mg²⁺ (Cohen et al. 1991). Hence, the S-layer can be completely removed by treatment of *H. volcanii* with chelating agents such as EDTA.

The *H. volcanii* genome consists of a main chromosome (2.848 Mb), three smaller mini-chromosomes (85.1 kb pHV1, 438 kb pHV3 and 636 kb pHV4) along with a 6.4 kb pHV2 plasmid (Hartman et al. 2010). It has a stable genome when compared with other extreme halophiles such as *Halobacterium salinarum*, which undergoes frequent IS-mediated rearrangement (López-García et al. 1995; Sapienza et al. 1982). One of the major incentives to exploit *H. volcanii* in biotechnology is it has an extensive genetic toolbox to perform a plethora of genetic manipulations. Transformation of DNA into *H. volcanii* is achieved using the PEG protocol (Cline et al. 1989). The method was further improved by generating strains deleted for the *mrr* restriction endonuclease gene, which reduces the transformation efficiency by cleaving the incoming foreign DNA at methylated sites (Allers et al. 2010). The *mrr* deficient *H. volcanii* strain allows direct and efficient transformation, eliminating the need to passage the foreign DNA in an *Escherichia coli dam* mutant. There are multiple shuttle vectors available for use, which are based on pHK2, pHV2 and pHV1 origins (Allers and Mevarech 2005; Allers et al. 2004). The ability of *H. volcanii* to grow in minimal and synthetic media has led to the development of multiple successful auxotrophic selection markers, where genes involved in amino acid or nucleotide biosynthetic pathways are used to complement chromosomal mutations. Uracil (*pyrE2*), leucine (*leuB*), thymidine (*hdrB*) and tryptophan (*trpA*) based selection markers have been used extensively for performing gene deletion/complementation in *H. volcanii* (Allers et al. 2004; Bitan-Banin et al. 2003; Leigh et al. 2011). A CRISPRi toolkit is also available for gene interference based on the clustered regularly interspaced short palindromic repeats (CRISPR)-Cas system (Maier et al. 2019). Gene expression reporters based on β-galactosidase and green fluorescent protein (GFP) are available (Holmes and Dyall-Smith 2000; Reuter and Maupin-Furlow 2005). For protein production, both inducible and constitutive promoter systems have been developed (Haque et al. 2019; Large et al. 2007), and metal based protein

purification is commonly used owing to purification tags in the expression vectors (Allers et al. 2010). Given these genetic tools and a wide range of salt tolerance, it is not surprising that *H. volcanii* has been used as a heterologous host to study genes from other halophiles. For example, gas vesicles produced by the closely related species *Haloferax mediterranei* have been characterised using *H. volcanii* as a heterologous host, which is naturally devoid of gas vesicle genes (Pfeifer et al. 2002).

Biotechnological potential of *H. volcanii*

Due to their remarkable properties such as antioxidants, colouring and light absorption, carotenoids are of great interest in biotechnology and medicine (Eggersdorfer and Wyss 2018; Fiedor and Burda 2014; Zhang et al. 2014a). Recently, there has been growing interest in naturally derived carotenoids owing to the increasing demand for bioactive compounds of natural origin with higher bioaccessibility and environmentally benign bioprocessing methods (Giani et al. 2019). The carotenoid biosynthesis pathway in *H. volcanii* begins with a condensation of two C₂₀ molecules into C₄₀ phytoene, a reaction catalysed by phytoene synthase, encoded by the *crtB* (HVO_2524) gene. Subsequently, phytoene is converted to lycopene by the phytoene desaturase enzyme encoded by *crtI* (HVO_2528). In *H. volcanii*, the *lye* (HVO_2527) gene encoded lycopene elongase subsequently converts lycopene to bacterioruberin. *H. volcanii* predominantly produces the C₅₀ carotenoid bacterioruberin (82%) and its derivatives such as monoanhydrobacterioruberin (7%), bisanhydrobacterioruberin (2%), dihydromonoanhydrobacterioruberin (2%), along with small amount of lycopene (0.3%) (Rønnekleiv, 1995). A recent study has shown that bacterioruberin produced by *H. volcanii* is three times more effective as an antioxidant compared to β-carotene, and improves the cell viability, motility and velocities of ram sperm cells (Zalazar et al. 2019). *HvLON3* strain, a conditional *H. volcanii* mutant for Lon protease produces 10-15 times more bacterioruberin than the parental strain given that Lon protease degrades the phytoene synthase, the rate limiting enzyme in carotenoid biosynthesis (Cerletti et al. 2018). *H. volcanii* is incapable of naturally producing β-carotene since the lycopene cyclase (encoded by *crtY*) is absent, whereas other halophiles such as *Haloarcula marismortui*, *Halobacterium salinarum* possess *crtY* (Baliga et al. 2004; Peck et al. 2002). However, the carotenoid biosynthetic pathway in *H. volcanii* can be genetically manipulated to produce β-carotene through the heterologous expression of *crtY* from *Haloarcula marismortui* (Smith 2018). Compared to other halophilic archaea, *H. volcanii* is better suited for heterologous pathway engineering due to its faster growth, readily available genetic toolkit and stable genome (Leigh et al. 2011). Given the absence of a rigid cell wall, *H. volcanii* affords a cheap and simple protocol for carotenoid extraction compared to the lengthy and complicated extraction procedure from pink shrimp or carrot, which involves the use of expensive equipment (Mezzomo et al. 2011; Rawson et al. 2011; Zalazar et al. 2019). Hence, by remodelling the carotenoid biosynthetic pathway through genetic engineering, production and extraction of β-carotene using *H. volcanii* may provide a more effective, efficient, and affordable method of natural carotenoid synthesis than those currently offered by chemical synthesis route or by extraction route from other microbes, plants and algae.

Enzyme mediated biocatalysis offers a more cost-effective and environmentally sustainable way of generating biotechnologically valuable chemicals when compared with traditional chemical syntheses (Hughes and Lewis 2018). *H. volcanii* possesses alcohol dehydrogenase (ADH) enzymes, which catalyse the production of important building blocks in fine chemical, flavour and fragrance industries such as chiral alcohols (Chapuis and Jacoby

2001; Schmid et al. 2001). It encodes two haloalkaliphilic and thermoactive ADHs: *HvADH1* (HVO_2428) and *HvADH2* (HVO_B0071), which catalyse the reversible oxidation of primary and secondary alcohols into aldehydes and ketones, and the reduction of aldehydes, ketones into their corresponding alcohols in the presence of NADPH/NADH cofactors (Timpson et al. 2013). *HvADH2* is industrially more attractive compared to *HvADH1* due to greater enzyme activity, dual cofactor and broad substrate scope, stability in organic solvents, halophilic and thermophilic conditions (Alsafadi and Paradisi 2013; Timpson et al. 2013). Given its activity in high temperature (> 80°C), it is described as the most thermoactive haloarchaeal ADH to date (Cao et al. 2008; Timpson et al. 2012; Timpson et al. 2013). It accepts a broad range of substrates ranging from medium chain alcohols to branched and aromatic alcohols for the oxidative reaction. For the reduction reaction, it accepts a range of aromatic ketone substrates such as alkyl, methyl, alpha-halo aryl ketones (Alsafadi et al. 2017). Immobilisation within epoxy resin and metal organic framework material further improves the enzyme activity over a broader pH range, temperature and enhances enzyme stability in organic solvents (Alsafadi and Paradisi 2014; Carucci et al. 2018). It is also possible to improve the substrate scope of *HvADH2* for biocatalysis through rational *in silico* enzyme design (Cassidy et al. 2017). *H. volcanii* expression systems can be used to overexpress and characterise industrially attractive enzymes from other halophiles (e.g. *HmADH12* from extreme halophile from the Dead Sea *Haloarcula marismortui* and *HsADH2* from the halophilic archaeon *Halobacterium salinarum* NRC-1) using the tryptophan inducible expression system (Liliensiek et al. 2013; Timpson et al. 2012). The same expression system has also been successfully used to produce haloarchaeal transaminase capable of producing chiral amines (Kelly et al. 2019). In this work, the authors overexpressed a haloarchaeal transaminase from a Triassic period halite deposit that showed halophilic activity, thermo and organic solvent tolerance profile.

Lignin and its derivatives are considered as serious contaminants in paper and pulp industries, tanneries, textile mills due to their poor biodegradability and intense colour (Costa et al. 2017; Raghukumar et al. 2008). Given that waste streams in these industries contain high concentrations of recycled salt, halophilic enzymes capable of lignin degradation could play a pivotal role in biopulping and biobleaching processes (Amoozegar et al. 2019). Laccase is a multicopper oxidase enzyme that couples oxidation of phenolic compounds to the four-electron reduction of molecular oxygen to water (Solomon et al. 1996). It has received attention in the biofuel industry due to its catalytic properties, and helps to break down lignocellulose materials, prominent feedstocks for advanced biofuel production (Kudanga and Le Roes-Hill 2014). It has also been implicated in the detoxification of water and soil, in bilirubin level determination in serum. and for use in biosensors and bioreactors (Sakurai and Kataoka 2007). *H. volcanii* encodes *LccA* (HVO_B0205), a rare archaeal laccase, which is secreted into the culture medium as a highly stable glycoprotein that is active at high salt concentrations (0.1 M to 1.4 M) and temperature (55°C). It accepts a broad range of organic substrates such as bilirubin, syringaldazine (SGZ), 2,2-azino-bis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) and dimethoxyphenol (DMP), and retains its activity in organic solvents such as dimethyl sulfoxide (DMSO) and dimethylformamide (DMF) (Uthandi et al. 2009). Hence, the *LccA* enzyme could serve as valuable biotechnological tool for the removal of phenolic contaminants in phenol-laden saline wastewater derived from various industrial settings (Moussavi et al. 2010). Despite the potential of *LccA* in biofuel industry, its direct *in vivo* lignolytic activity on lignin has still to be experimentally verified (Brink et al. 2019).

As can be seen from the above examples, enzymes from *H. volcanii* show polyextremophilicity, meaning that they are not only active in high salt environments but also are functional at increased temperature and in organic solvents. It has been established that enzymes from halophiles show greater activity and stability in organic solvents compared to their non-halophilic counterparts (Margesin and Schinner 2001). This unusual ability to be active in environments with low water activity (high salt, solvent, and/or desiccation) has made *H. volcanii* and its enzymes ideal candidates for biotechnology advances.

Halocins are antimicrobial compounds naturally produced by halophilic archaea and bacteria living in hypersaline environments (Shand and Leyva 2007). These compounds are believed to confer selective advantages to the organism for survival in a competitive environment, including against other halophiles (Atanasova et al. 2013). Due to the emergence of antimicrobial resistant pathogens, there is a dire need to screen and characterise novel antimicrobials (Baker 2015). This has included the search for halocins that possess a broad range antimicrobial spectrum. *H. volcanii* KPS1 strain produces halocin that is active at high temperatures (up to 80°C), and over a broad range of pH (3.0-9.0) and salinity (Kavitha et al. 2011). Most importantly, it shows antimicrobial activity against both Gram-positive and Gram-negative bacteria. Large amounts of salts are used in tanning processes, which provides a favourable environment for the growth of halophiles. Given its anti-halophilic activity, halocins can be exploited in the textile industry to inhibit the unwanted growth of halophiles that could be damaging for the product (Birbir et al. 2004). The ability of *H. volcanii* to degrade aromatic hydrocarbons such as anthracene, naphthalene, phenanthrene, pyrene and benzoanthracene could also be beneficial for bioremediation in hypersaline environments (Bonfa et al. 2011).

Polyhydroxyalkanoates (PHA) are polyesters composed of hydroxy fatty acids, which serve as intracellular storage material of carbon source and energy (Rehm 2007). They are found in a variety of microbes including haloarcheal species belonging to the genera *Haloarcula*, *Haloferax*, *Halococcus*, *Haloquadratum*, *Halobacterium*, *Haloterrigena*, *Natronobacterium* (Koller 2019). These water-insoluble nanosized cytoplasmic compounds crystallize after solvent extraction and, most importantly, exhibit thermo-plastic and elastomeric properties (Poli et al. 2011). Therefore, PHA compounds have received global attention due to their importance as natural, bio-based, biocompatible and biodegradable alternatives to plastics of petrochemical origin. Poly 3-hydroxybutyrate (PHB) is the best known polyhydroxyalkanoate (PHA). Polymerisation of 3-hydroxybutyryl-CoA results in PHA formation through PHA synthase activity (Rehm 2003). *H. volcanii* produces PHB up to 7% of its cell dry weight when grown on media containing 250 g/L salts, 10 g/L glucose and 1 g/L yeast extract (Fernandez-Castillo et al. 1986). An additional report for *H. volcanii* mediated PHA accumulation from sugarcane bagasse substrate, a fibrous leftover by-product of the sugarcane industry, was recently published (Salgaonkar and Bragança 2017). In contrast, another study reported no detection of PHB from *H. volcanii* (Legat et al. 2010). *H. volcanii* has been successfully used as a heterologous host to generate PHA through the expression of PHA synthase genes from the halophile *Haloarcula hispanica* (Han et al. 2009).

Alongside facile genetics, at least three other traits make *H. volcanii* an attractive host for bioprocessing. Firstly, industrially desirable quasi-sterile conditions are readily achieved, since few other microbes cannot grow in the molar concentrations of salt present in growth media for *H. volcanii*. Secondly, the cost involved in product extraction is significantly reduced since it is easy to lyse *H. volcanii* cells through addition of water and/or the use

of simple chelating agents to disrupt the outer S-layer. Finally, the potential for LPS (lipopolysaccharide) mediated toxicity caused by bacterial hosts is eliminated by the use of *H. volcanii*. These properties would substantially facilitate downstream processing in economic and environmental terms during bioprocessing.

The S-layer of *H. volcanii* is a two-dimensional array of protein or glycoprotein subunits, which form part of the cell envelope components (Tamir and Eichler 2017). The ability of isolated archaeal S-layer proteins to assemble into regular arrays in liquid or on surfaces such as metals, polymers or interfaces such as lipid films, liposomes makes them attractive for potential applications in nanobiotechnology. These arrays show reproducible physiochemical properties down to the nanometre scale (Debabov 2004). The pores in S-layers are identical in size and morphology, and functional groups such as carboxylic acid, amine and hydroxyl groups on the protein lattices and pore areas are aligned in defined positions and orientations. These combined features have led to their applications in the production of isoporous ultrafiltration membranes, bioanalytical sensors, biomimetics, affinity membranes, immunotherapy, drug delivery, vaccine development, immobilisation matrices for binding of functional molecules such as enzymes, antibodies (Ilk et al. 2008; Sleytr et al. 2007). Mammalian olfactory receptor from rat has been heterologously expressed in the lipid bilayer of the *H. volcanii* (Lobasso et al. 2015). Therefore, the lipid matrix of *H. volcanii* could serve as an ideal biomaterial for the future development of nano-vesicle based hybrid biosensor devices.

Challenges of applying *H. volcanii* in biotechnology

Halophiles adapt to high salt environment using two alternative strategies (Oren 2008). The first strategy, termed 'salt-out', is mainly adopted by halotolerant bacteria and eukaryotes. Excess salt is excluded from halotolerant bacteria and eukaryotes, with an increase in compatible organic solutes such as glycerol or glycine betaine in the cytoplasm to maintain an osmotic balance. These compatible solutes do not interfere with intracellular enzymatic activity. The second strategy, termed 'salt-in', is adopted mainly by haloarchaea including *H. volcanii*. Equimolar concentrations of salt, predominantly KCl, are accumulated in the cytoplasm to maintain the osmotic balance (Oren 2013). Since biosynthesis of organic osmotic solutes in the 'salt-out' strategy is energetically costly compared to the 'salt-in' strategy, it is haloarchaea such as *H. volcanii* that thrive in hypersaline conditions (Oren 1999). Halophilic proteins from *H. volcanii* have evolved to function in the presence of molar concentration of salt, they have an overall increase in the number of highly positively charged acidic residues on surface to make the proteins soluble in high salt environment, along with a decrease in overall hydrophobicity through the substitution of large hydrophobic residues on the surface with small hydrophilic ones (Danson and Hough 1997; Mevarech et al. 2000; Wright et al. 2002).

This peculiarity poses a challenge for the expression of active halophilic proteins in industrially attractive heterologous hosts such as *E. coli*, since these proteins often misfold and aggregate in low ionic strength environment. However, there have been a few reports of heterologous expression of halophilic proteins in *E. coli* (Cao et al. 2008; Cendrin et al. 1993; Domenech and Ferrer 2006; Johnsen and Schönheit 2004), which rely on solubilization of inclusion bodies followed by laborious solubilisation and refolding of insoluble proteins in a hypersaline environment (Connaris et al. 1999; Singh and Panda 2005). An additional caveat is the requirement for prior knowledge of the target protein characteristics. In any case, this approach is not

universally effective, as many enzymes remain inactive. For example, enzymatically active protein could not be recovered from the heterologously overexpressed alcohol dehydrogenase (*HmADH12*) of the extreme halophile *Haloarcula marismortui*, despite repeated attempts to solubilise the inclusion bodies in varying salt and pH conditions (Timpson et al. 2012). In another study, no protein expression was observed following attempts of overexpression in *E. coli* of two halophilic alcohol dehydrogenases (ADH/D1 and ADH/A1) from the deep Red Sea brine pools (Strillinger et al. 2016). These drawbacks of heterologous expression have led to the development of halophilic systems to generate biotechnologically important proteins that are functionally active.

One of the major obstacles in exploiting extremozymes is the low yield or activity of enzymes when cultivated on a large bioreactor scale, as required for industrial applications (Elleuche et al. 2014; Sarmiento et al. 2015). For example, it is essential to produce active alcohol dehydrogenase (*HvADH*), laccase (*LccA*) enzymes, and bacterioruberin, and halocins proteins (Table 1) from *H. volcanii* on large scale to realise their true biotechnological potential. It is challenging to ensure expression and purification of bulk quantities of highly active proteins in a halophilic environment. Co-purification of unwanted non-specific proteins with the target protein is often encountered by the researchers following affinity tagged protein purification (Kimple et al. 2013). A bottleneck of large-scale protein overexpression in *H. volcanii* host is their intrinsic ability to form biofilms (Chimileski et al. 2014; Frols et al. 2012), which can interfere with expensive sensors present in the bioreactor during the fermentation process and may also alter the characteristics of expressed proteins. Molar concentrations of salt required to culture *H. volcanii* can rapidly corrode stainless-steel bioreactors. This necessitates the use of reactors made of alternative materials rather than the readily available stainless-steel.

Tools for *H. volcanii* applications in biotechnology

For protein overexpression, a modified strain *H. volcanii* host strain H1424 (Δ *pyrE2* Δ *hdrB* Δ *mrr* *Nph-pitA cdc48d-Ct*) is available along with its corresponding plasmid pTA1228 (Fig. 1a) (Brendel et al 2014; Stroud et al. 2012). Regulated gene expression systems enable the tailoring of expression of biotechnologically important enzymes/proteins according to the industrial need. The pTA1228 plasmid (Fig. 1a) contains the tryptophan inducible promoter *p.tnaA* for protein expression (Large et al. 2007), which is active in the presence of >1 mM tryptophan. The plasmid pTA1228 also encodes an N-terminal hexahistidine (6xHis) tag to facilitate protein purification using immobilised metal-based affinity chromatography (IMAC), since this affinity tag is compatible with high salt media. The host strain H1424 is deleted for the histidine-rich PitA and Cdc48d proteins, major contaminants of the His-tagged proteins purified from *H. volcanii* using IMAC (Bab-Dinitz et al. 2006). Given the essential nature of the *pitA* and *cdc48d*, it was not possible to delete these genes completely, therefore *pitA* was replaced with an orthologue (*Nph-pitA*) from the haloalkaliphile *Natronomonas pharaonis* that does not contain a high number of histidine residues, and a truncated version of *cdc48d* coding for a protein (Cdc48d-Ct) lacking the histidine rich C-terminus was engineered (Allers et al. 2010; Stroud et al 2012). The expression cassette in pTA1228 is flanked by two transcriptional terminators (t.L11e and t.Syn) to prevent read through transcription (Fig. 1a), and the pHV2 replication origin maintains the plasmid at a copy number of 6 per genome (Allers et al. 2004). Incorporation of *pyrE2* and *hdrB* marker genes in pTA1228 enables selection on media lacking

uracil and thymidine, respectively; the latter is lacking in rich media (*Hv*-YPC), facilitating maximal cell growth and protein overexpression (Allers et al. 2004; Bitan-Banin et al. 2003).

To give researchers more options for protein purification, another plasmid pTA1392 (Fig. 1b) is available that encodes a C-terminal Streptavidin II tag (Strep II) for protein purification, alongside the established N-terminal His-tag (Braun et al. 2019); both tags are compatible with high salt concentrations. For protein expression at lower levels, for example if the protein is toxic to the host cells, the pTA1926 plasmid (Fig. 1c) contains the *p.tnaM3* promoter that shows 50% reduced activity compared to *p.tnaA* promoter present in pTA1228 and pTA1392 (Braun et al. 2019). For constitutive overexpression of proteins that are not harmful to the host, the pTA1992 plasmid (Fig. 1d) contains the highly active *p.syn* promoter (Haque et al. 2019; Large et al. 2007). This plasmid has the potential to make biotechnological processes cost effective since its usage alleviates the need for repeated tryptophan supplementation into *H. volcanii* cultures, which is an expensive amino acid.

A simple tryptophan-inducible protein overexpression system was reported that allows halophilic protein overexpression using a stirred tank bioreactor system (Strillinger et al. 2016). This study overexpressed two thermohalophilic alcohol dehydrogenases (ADH/D1 and ADH/A1), which showed about 28 fold more enzyme activity when compared with the uncontrolled shaker flask cultivations. The use of bioreactors has inherent advantages such as control of pH, temperature, supply of oxygens, nutrient media during the fermentation process. To prevent biofilm formation inside bioreactors, a *H. volcanii* strain (H1895) that is deleted for the two essential genes for biofilm formation: an integral membrane protein (HVO_1033) and ATPase subunit (HVO_1034) of the archaeal pilus biosynthesis machinery could be exploited in biotechnology given its inability to form biofilm during protein overexpression (Strillinger et al. 2016; Tripepi et al. 2013; Tripepi et al. 2012).

Alternative bioreactors other than corrosion prone stainless-steel are available for large scale halophilic protein production. A stirred and aerated bioreactor made of corrosion resistant materials such as polyetherether ketone (PEEK), borosilicate glass, or silicon nitride ceramics has been successfully used to generate poly-L-glutamic acid (PGA) or poly- β -hydroxy butyric acid (PHB) from the cultures of extreme halophilic archaeon *Natrialba* spp. (Hezayen et al. 2000). Another extreme halophile *H. mediterranei* has been cultured using a bioreactor with parts made of PEEK and borosilicate glass (Lorantfy et al. 2014). However, these materials are costly and more work is necessary in designing more energy efficient, corrosion-resistant and cost-effective alternative bioreactors for culturing *H. volcanii*.

Recent work has shown *H. volcanii* can be immobilised in inexpensive, non-toxic and highly porous calcium alginate beads following a simple method and used as whole cell biocatalyst (Fig. 2) (Haque et al. 2019). It offers halophilic enzyme expression with high product yield and reusability in a hypersaline environment. Furthermore, it provides data that calcium alginate beads with once immobilised *H. volcanii* can be used repeatedly for 12 successive biocatalytic cycles without compromising the bead morphology. Product yield from a batch of immobilised beads with *H. volcanii* that had been stored at room temperature for a month was very similar to freshly prepared beads. This system is particularly cost-effective with enzymatic processes that require expensive NADPH/NADH cofactor supplementation, since *in situ* cofactors generated by *H. volcanii* are used for the

catalytic reaction, thus making the process self-sufficient. In this reported system, 1.1 g dry weight (from 120 ml broth culture) of immobilised *H. volcanii* whole cell biocatalyst showed high product yield range of 96.4% - 100% in the presence of 5 mM acetophenone substrate after 24 hours. In contrast, a yield range of 48% - 94% was found from 500 ml of *Yarrowia lipolytica* in presence of 3.3 mM of the same substrate after 48 hours (Janeczko et al. 2015). In another work, a product yield range of 53% - 63% was found with 5 g of the fungus, *Aspergillus niger* whole cell biocatalyst in the presence of 1 mM acetophenone substrate after 24 hours (Kurbanoglu et al. 2007). Therefore, the performance of this immobilised *H. volcanii* whole cell biocatalyst is superior compared to the reported above two studies.

Immobilised *H. volcanii* within the calcium alginate beads offer multiple advantages for biotechnological applications (Fig. 2). Firstly, it stabilises the whole cells within the calcium alginate bead matrix leading to higher cell density and specific productivity (Strand et al. 2004). Secondly, the porous alginate bead matrix eases the product separation and flow of material to the *H. volcanii* cells. Thirdly, this type of highly defined and reproducible encapsulation within the natural alginate biopolymer ensures biocompatible microenvironment for the maintenance of viable *H. volcanii* cells (Melvik and Dornish 2004). Finally, immobilised nature of the whole cells means it is ideal for their applications in bioreactors without the problem of cell wash out (Obradovic et al. 2004). Such simple system can easily be scaled up by packing the beads with immobilised *H. volcanii* inside a stirred tank fermenter (Strillinger et al. 2016). Organism such as *Alcaligenes faecalis* immobilised within the calcium alginate beads has been successfully used in packed bed bioreactor as whole cell biocatalyst (Xue et al. 2013; Zhang et al. 2014b). Intermittent or continuous substrate feeding approach using flow biocatalysis can be combined with *in situ* product removal strategy to further increase product yield and alleviate toxicity mediated by the presence of excessive substrate/product in the immobilised system (Freeman et al. 1993; Tamborini et al. 2018)). Therefore, if the biotechnological process involves supplementation or generation of a toxic compound, use of immobilised *H. volcanii* whole cell approach would be preferable over the bioreactor scale fermentation since it would offer more protection to the cells.

Interest in halophiles such as *H. volcanii* has led to the development of curated bioinformatic databases such as HaloWeb, HProtDB, HaloDom. These tools greatly accelerate bioprospecting through genome mining. HaloWeb (<http://halo4.umbi.umd.edu>) provides the complete genome sequences for all haloarchaeal genomes, along with BLAST options against various genomes and genomic maps (DasSarma et al. 2010b). HProtDB (<http://webapp.cabgrid.res.in/protein/>) contains information about physical and biochemical properties of halophilic proteins for 21 strains of Archaea and Bacteria (Sharma et al. 2014). HaloDom catalogues all the halophilic species studied to date across all three domains of life, Archaea, Bacteria and Eukarya (<http://halodom.bio.auth.gr/>). It also provides a 'Halopredictor' function to predict the signatures of halophilic adaptation in the FASTA sequence of a novel protein (Loukas et al. 2018). The 'Halohandbook' is a manual containing detailed laboratory protocols for studying halophiles including *H. volcanii* (Dyall-Smith 2009). It has been compiled with input from leading halophile researchers.

Conclusions

Characteristics such as facile genetics, adaptation to low water environment and high salt tolerance have great potential in making *H. volcanii* a valuable resource for biotechnology. Of particular interest is the use of *H. volcanii* as a heterologous host to study important biotechnological enzymes from other halophiles, ranging from moderate to extreme. However, challenges still remain in applying *H. volcanii* in a broad range of biotechnological applications. With the discovery of novel biotechnologically important proteins, expansion of genetic toolkits and rapidly advanced technologies, the potential for exploiting *H. volcanii* in biotechnology can only grow.

Compliance with Ethical Standards

Competing interests

The authors declare that there are no competing interests.

Funding

This work was supported by the Biotechnology and Biological Sciences Research Council (BBSRC) [grant number BB/P002536/1]. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author contributions

All authors contributed equally to writing the manuscript.

Ethical approval

Not applicable since the work does not involve any study with human participants or animals.

References

- Allers T (2010) Overexpression and purification of halophilic proteins in *Haloferax volcanii*. *Bioeng Bugs* 1(4):288-290 doi:10.4161/bbug.1.4.11794
- Allers T, Barak S, Liddell S, Wardell K, Mevarech M (2010) Improved strains and plasmid vectors for conditional overexpression of His-tagged proteins in *Haloferax volcanii*. *Appl Environ Microbiol* 76(6):1759-1769 doi:10.1128/aem.02670-09
- Allers T, Mevarech M (2005) Archaeal genetics — the third way. *Nat Rev Genet* 6(1):58-73 doi:10.1038/nrg1504
- Allers T, Ngo H-P, Mevarech M, Lloyd RG (2004) Development of additional selectable markers for the halophilic archaeon *Haloferax volcanii* based on the *leuB* and *trpA* genes. *Appl Environ Microbiol* 70(2):943-953 doi:10.1128/aem.70.2.943-953.2004
- Alsafadi D, Alsalmán S, Paradisi F (2017) Extreme halophilic alcohol dehydrogenase mediated highly efficient syntheses of enantiopure aromatic alcohols. *Org Biomol Chem* 15(43):9169-9175 doi:10.1039/C7OB02299A
- Alsafadi D, Paradisi F (2013) Effect of organic solvents on the activity and stability of halophilic alcohol dehydrogenase (ADH2) from *Haloferax volcanii*. *Extremophiles* 17(1):115-122 doi:10.1007/s00792-012-0498-0
- Alsafadi D, Paradisi F (2014) Covalent immobilization of alcohol dehydrogenase (ADH2) from *Haloferax volcanii*: how to maximize activity and optimize performance of halophilic enzymes. *Mol Biotechnol* 56(3):240-247 doi:10.1007/s12033-013-9701-5
- Amoozegar MA, Safarpour A, Noghabi KA, Bakhtiary T, Ventosa A (2019) Halophiles and their vast potential in biofuel production. *Front Microbiol* 10:1895 doi:10.3389/fmicb.2019.01895
- Amoozegar MA, Siroosi M, Atashgahi S, Smidt H, Ventosa A (2017) Systematics of haloarchaea and biotechnological potential of their hydrolytic enzymes. *Microbiology* 163(5):623-645 doi:10.1099/mic.0.000463
- Antunes A, Simões MF, Grötzinger SW, Eppinger J, Bragança J, Bajic VB (2017) Bioprospecting Archaea: focus on extreme halophiles. In: Paterson R, Lima N (eds) *Bioprospecting: success, potential and constraints*. Springer International Publishing, Cham, pp 81-112
- Atanasova NS, Pietilä MK, Oksanen HM (2013) Diverse antimicrobial interactions of halophilic archaea and bacteria extend over geographical distances and cross the domain barrier. *Microbiologyopen* 2(5):811-825 doi:10.1002/mbo3.115
- Bab-Dinitz E, Shmueli H, Maupin-Furlow J, Eichler J, Shaanan B (2006) *Haloferax volcanii* PitA: an example of functional interaction between the Pfam chlorite dismutase and antibiotic biosynthesis monooxygenase families? *Bioinformatics* 22(6):671-675 doi:10.1093/bioinformatics/btk043
- Baker S (2015) A return to the pre-antimicrobial era? *Science* 347(6226):1064 doi:10.1126/science.aaa2868
- Baliga NS, Bonneau R, Facciotti MT, Pan M, Glusman G, Deutsch EW, Shannon P, Chiu Y, Weng RS, Gan RR, Hung P, Date SV, Marcotte E, Hood L, Ng WV (2004) Genome sequence of *Haloarcula marismortui*: a halophilic archaeon from the Dead Sea. *Genome Res* 14(11):2221-2234 doi:10.1101/gr.2700304
- Birbir M, Eryilmaz S, Ogan A (2004) Prevention of halophilic microbial damage on brine cured hides by extremely halophilic halocin producer strains. *J Soc Leath Tech Ch* 88:99-104

- Bitan-Banin G, Ortenberg R, Mevarech M (2003) Development of a gene knockout system for the halophilic archaeon *Haloferax volcanii* by use of the *pyrE* Gene. *J Bacteriol* 185(3):772-778 doi:10.1128/JB.185.3.772-778.2003
- Bonfa MR, Grossman MJ, Mellado E, Durrant LR (2011) Biodegradation of aromatic hydrocarbons by Haloarchaea and their use for the reduction of the chemical oxygen demand of hypersaline petroleum produced water. *Chemosphere* 84(11):1671-1676 doi:10.1016/j.chemosphere.2011.05.005
- Braun F, Thomalla L, van der Does C, Quax TEF, Allers T, Kaefer V, Albers S-V (2019) Cyclic nucleotides in archaea: cyclic di-AMP in the archaeon *Haloferax volcanii* and its putative role. *Microbiologyopen* 8(9):e00829 doi:10.1002/mbo3.829
- Brendel J, Stoll B, Lange SJ, Sharma K, Lenz C, Stachler AE, Maier LK, Richter H, Nickel L, Schmitz RA, Randau L, Allers T, Urlaub H, Backofen R, Marchfelder A (2014) A complex of Cas proteins 5, 6, and 7 is required for the biogenesis and stability of clustered regularly interspaced short palindromic repeats (crispr)-derived rnas (crnas) in *Haloferax volcanii*. *J Biol Chem* 289(10):7164-7177 doi:10.1074/jbc.M113.508184
- Brink DP, Ravi K, Lidén G, Gorwa-Grauslund MF (2019) Mapping the diversity of microbial lignin catabolism: experiences from the eLignin database. *Appl Microbiol Biotechnol* 103(10):3979-4002 doi:10.1007/s00253-019-09692-4
- Cao Y, Liao L, Xu XW, Oren A, Wang C, Zhu XF, Wu M (2008) Characterization of alcohol dehydrogenase from the haloalkaliphilic archaeon *Natronomonas pharaonis*. *Extremophiles* 12(3):471-476 doi:10.1007/s00792-007-0133-7
- Carucci C, Bruen L, Gascón V, Paradisi F, Magner E (2018) Significant enhancement of structural stability of the hyperhalophilic ADH from *Haloferax volcanii* via entrapment on metal organic framework support. *Langmuir* 34(28):8274-8280 doi:10.1021/acs.langmuir.8b01037
- Cassidy J, Bruen L, Rosini E, Molla G, Pollegioni L, Paradisi F (2017) Engineering substrate promiscuity in halophilic alcohol dehydrogenase (HvADH2) by *in silico* design. *PloS One* 12(11):e0187482 doi:10.1371/journal.pone.0187482
- Cendrin F, Chroboczek J, Zaccai G, Eisenberg H, Mevarech M (1993) Cloning, sequencing, and expression in *Escherichia coli* of the gene coding for malate dehydrogenase of the extremely halophilic archaeobacterium *Haloarcula marismortui*. *Biochemistry* 32(16):4308-4313 doi:10.1021/bi00067a020
- Cerletti M, Paggi R, Troetschel C, Ferrari MC, Guevara CR, Albaum S, Poetsch A, De Castro R (2018) LonB protease is a novel regulator of carotenogenesis controlling degradation of phytoene synthase in *Haloferax volcanii*. *J Proteome Res* 17(3):1158-1171 doi:10.1021/acs.jproteome.7b00809
- Chapuis C, Jacoby D (2001) Catalysis in the preparation of fragrances and flavours. *Appl Catal A* 221:93-117 doi:10.1016/S0926-860X(01)00798-0
- Chimileski S, Franklin MJ, Papke RT (2014) Biofilms formed by the archaeon *Haloferax volcanii* exhibit cellular differentiation and social motility, and facilitate horizontal gene transfer. *BMC Biol* 12:65-65 doi:10.1186/s12915-014-0065-5
- Cline SW, Lam WL, Charlebois RL, Schalkwyk LC, Doolittle WF (1989) Transformation methods for halophilic archaeobacteria. *Can J Microbiol* 35(1):148-152 doi:10.1139/m89-022
- Cohen S, Shilo M, Kessel M (1991) Nature of the salt dependence of the envelope of a Dead Sea archaeobacterium, *Haloferax volcanii*. *Arch Microbiol* 156(3):198-203

- Connaris H, Chaudhuri JB, Danson MJ, Hough DW (1999) Expression, reactivation, and purification of enzymes from *Haloferax volcanii* in *Escherichia coli*. *Biotechnol Bioeng* 64(1):38-45
- Costa S, Dedola D, Pellizzari S, Blo R, Rugiero I, Pedrini P, Tamburini E (2017) Lignin biodegradation in pulp-and-paper mill wastewater by selected white rot fungi. *Water* 9:935 doi:10.3390/w9120935
- Danson MJ, Hough DW (1997) The structural basis of protein halophilicity. *Comp Biochem Physiol Part A Physiol* 117(3):307-312 doi:[https://doi.org/10.1016/S0300-9629\(96\)00268-X](https://doi.org/10.1016/S0300-9629(96)00268-X)
- DasSarma P, Coker J, Huse V, DasSarma S (2010a) Halophiles, industrial applications. In *Encyclopedia of industrial biotechnology: bioprocess, bioseparation and cell technology*. Hoboken, NJ, USA: John Wiley & Sons, Inc.2009
- DasSarma SL, Capes MD, DasSarma P, DasSarma S (2010b) HaloWeb: the haloarchaeal genomes database. *Saline Systems* 6:12-12 doi:10.1186/1746-1448-6-12
- De Lourdes Moreno M, Pérez D, García MT, Mellado E (2013) Halophilic bacteria as a source of novel hydrolytic enzymes. *Life* 3(1):38-51
- Debabov VG (2004) Bacterial and archaeal S-layers as object of bionanotechnology. *Mol Biol (Mosk)* 38(4):578-591
- Domenech J, Ferrer J (2006) A new D-2-hydroxyacid dehydrogenase with dual coenzyme-specificity from *Haloferax mediterranei*, sequence analysis and heterologous overexpression. *Biochim Biophys Acta* 1760(11):1667-1674 doi:10.1016/j.bbagen.2006.08.024
- Dyall-Smith M (2009) The Halohandbook- protocols for haloarchaeal genetics. https://haloarchaea.com/wp-content/uploads/2018/10/Halohandbook_2009_v7.3mds.pdf
- Eggersdorfer M, Wyss A (2018) Carotenoids in human nutrition and health. *Arch Biochem Biophys* 652:18-26 doi:10.1016/j.abb.2018.06.001
- Elazari-Volcani B (Wilkansky) (1943) Bacteria in the bottom sediments of the Dead Sea. *Nature* 152(3853):274-275 doi:10.1038/152274c0
- Elleuche S, Schröder C, Sahm K, Antranikian G (2014) Extremozymes—biocatalysts with unique properties from extremophilic microorganisms. *Curr Opin Biotechnol* 29:116-123 doi:<https://doi.org/10.1016/j.copbio.2014.04.003>
- Fernandez-Castillo R, Rodriguez-Valera F, Gonzalez-Ramos J, Ruiz-Berraquero F (1986) Accumulation of poly(beta-hydroxybutyrate) by halobacteria. *Appl Environ Microbiol* 51(1):214-216
- Fiedor J, Burda K (2014) Potential role of carotenoids as antioxidants in human health and disease. *Nutrients* 6(2):466-488 doi:10.3390/nu6020466
- Freeman A, Woodley JM, Lilly MD (1993) *In situ* product removal as a tool for bioprocessing. *Biotechnology (NY)* 11(9):1007-1012 doi:10.1038/nbt0993-1007
- Frols S, Dyall-Smith M, Pfeifer F (2012) Biofilm formation by haloarchaea. *Environ Microbiol* 14(12):3159-3174 doi:10.1111/j.1462-2920.2012.02895.x
- Giani, M., Garbayo, I., Vílchez, C., & Martínez-Espinosa, R. M. (2019). Haloarchaeal carotenoids: healthy novel compounds from extreme environments. *Mar Drugs* 17(9), 524. doi:10.3390/md17090524
- Han J, Lu Q, Zhou L, Liu H, Xiang H (2009) Identification of the polyhydroxyalkanoate (PHA)-specific acetoacetyl coenzyme A reductase among multiple FabG paralogs in *Haloarcula hispanica* and

- reconstruction of the PHA biosynthetic pathway in *Haloferax volcanii*. *Appl Environ Microbiol* 75(19):6168-6175 doi:10.1128/aem.00938-09
- Haque RU, Paradisi F, Allers T (2019) *Haloferax volcanii* as immobilised whole cell biocatalyst: new applications for halophilic systems. *Appl Microbiol Biotechnol* 103(9):3807-3817 doi:10.1007/s00253-019-09725-y
- Hartman AL, Norais C, Badger JH, Delmas S, Haldenby S, Madupu R, Robinson J, Khouri H, Ren Q, Lowe TM, Maupin-Furlow J, Pohlschroder M, Daniels C, Pfeiffer F, Allers T, Eisen JA (2010) The complete genome sequence of *Haloferax volcanii* DS2, a model archaeon. *PloS One* 5(3):e9605 doi:10.1371/journal.pone.0009605
- Hezayen FF, Rehm BH, Eberhardt R, Steinbuchel A (2000) Polymer production by two newly isolated extremely halophilic archaea: application of a novel corrosion-resistant bioreactor. *Appl Microbiol Biotechnol* 54(3):319-325 doi:10.1007/s002530000394
- Holmes ML, Dyall-Smith ML (2000) Sequence and expression of a halobacterial beta-galactosidase gene. *Mol Microbiol* 36(1):114-122 doi:10.1046/j.1365-2958.2000.01832.x
- Hughes G, Lewis JC (2018) Introduction: biocatalysis in industry. *Chem Rev* 118(1):1-3 doi:10.1021/acs.chemrev.7b00741
- Ilk N, Egelseer EM, Ferner-Ortner J, Küpcü S, Pum D, Schuster B, Sleytr UB (2008) Surfaces functionalized with self-assembling S-layer fusion proteins for nanobiotechnological applications. *Colloid Surf A Physicochem Eng Asp* 321(1):163-167 doi:<https://doi.org/10.1016/j.colsurfa.2007.12.038>
- Janeczko T, Bąkowski W, Walczak E, Robak M, Dmochowska-Gładysz J, Kostrzewa-Susłow E (2015) Biotransformation of acetophenone and its halogen derivatives by *Yarrowia lipolytica* strains. *Ann Microbiol* 65(2):1097-1107. doi:10.1007/s13213-014-0955-3
- Johnsen U, Schönheit P (2004) Novel xylose dehydrogenase in the halophilic archaeon *Haloarcula marismortui*. *J Bacteriol* 186(18):6198-6207 doi:10.1128/JB.186.18.6198-6207.2004
- Kamekura M (1998) Diversity of extremely halophilic bacteria. *Extremophiles* 2(3):289-295
- Kavitha PK, Lipton A, A R S, Aishwarya Ms (2011) Growth characteristics and halocin production by a new isolate, *Haloferax volcanii* KPS1 from Kovalam Solar Saltern (India). *Res J Biol Sci* 6: 257-262 doi:10.3923/rjbsci.2011.257.262
- Kelly SA, Magill DJ, Megaw J, Skvortsov T, Allers T, McGrath JW, Allen CCR, Moody TS, Gilmore BF (2019) Characterisation of a solvent-tolerant haloarchaeal (R)-selective transaminase isolated from a Triassic period salt mine. *Appl Microbiol Biotechnol* 103(14):5727-5737 doi:10.1007/s00253-019-09806-y
- Kimble, M. E., Brill, A. L., Pasker, R. L. (2013) Overview of affinity tags for protein purification. *Curr Protoc Protein Sci* 9.9.1–9.9.23. doi:10.1002/0471140864.ps0909s73
- Koller M (2019) Polyhydroxyalkanoate biosynthesis at the edge of water activity-haloarchaea as biopolyester factories. *Bioengineering (Basel)* 6(2) doi:10.3390/bioengineering6020034
- Kudanga T, Le Roes-Hill M (2014) Laccase applications in biofuels production: current status and future prospects. *Appl Microbiol Biotechnol* 98 (15):6525-6542 doi:10.1007/s00253-014-5810-8
- Kurbanoglu EB, Zilbeyaz K, Kurbanoglu NI, Kilic H (2007) Enantioselective reduction of substituted acetophenones by *Aspergillus niger*. *Tetrahedron: Asymmetry* 18(10):1159-1162 doi:<https://doi.org/10.1016/j.tetasy.2007.05.017>

- Large A, Stamme C, Lange C, Duan Z, Allers T, Soppa J, Lund PA (2007) Characterization of a tightly controlled promoter of the halophilic archaeon *Haloferax volcanii* and its use in the analysis of the essential *cctI* gene. *Mol Microbiol* 66(5):1092-1106 doi:10.1111/j.1365-2958.2007.05980.x
- Le Borgne S, Paniagua D, Vazquez-Duhalt R (2008) Biodegradation of organic pollutants by halophilic bacteria and archaea. *J Mol Microbiol Biotechnol* 15(2-3):74-92 doi:10.1159/000121323
- Lee HS (2013) Diversity of halophilic archaea in fermented foods and human intestines and their application. *J Microbiol Biotechnol* 23(12):1645-1653 doi:10.4014/jmb.1308.08015
- Legat A, Gruber C, Zangger K, Wanner G, Stan-Lotter H (2010) Identification of polyhydroxyalkanoates in *Halococcus* and other haloarchaeal species. *Appl Microbiol Biotechnol* 87(3):1119-1127 doi:10.1007/s00253-010-2611-6
- Leigh JA, Albers SV, Atomi H, Allers T (2011) Model organisms for genetics in the domain Archaea: methanogens, halophiles, *Thermococcales* and *Sulfolobales*. *FEMS Microbiol Rev* 35(4):577-608 doi:10.1111/j.1574-6976.2011.00265.x
- Liliensiek AK, Cassidy J, Gucciardo G, Whitely C, Paradisi F (2013) Heterologous overexpression, purification and characterisation of an alcohol dehydrogenase (ADH2) from *Halobacterium* sp. NRC-1. *Mol Biotechnol* 55(2):143-149 doi:10.1007/s12033-013-9666-4
- Lobasso S, Vitale R, Lopalco P, Corcelli A (2015) *Haloferax volcanii*, as a novel tool for producing mammalian olfactory receptors embedded in archaeal lipid bilayer. *Life (Basel)* 5(1):770-782 doi:10.3390/life5010770
- López-García P, St Jean A, Amils R, Charlebois RL (1995) Genomic stability in the archaeae *Haloferax volcanii* and *Haloferax mediterranei*. *J Bacteriol* 177(5):1405-1408 doi:10.1128/jb.177.5.1405-1408.1995
- Lorantfy B, Seyer B, Herwig C (2014) Stoichiometric and kinetic analysis of extreme halophilic Archaea on various substrates in a corrosion resistant bioreactor. *N Biotechnol* 31(1):80-89 doi:10.1016/j.nbt.2013.08.003
- Loukas A, Kappas I, Abatzopoulos TJ (2018) HaloDom: a new database of halophiles across all life domains. *J Biol Res (Thessalon)* 25:2-2 doi:10.1186/s40709-017-0072-0
- Maier L-K, Stachler A-E, Brendel J, Stoll B, Fischer S, Haas KA, Schwarz TS, Alkhnbashi OS, Sharma K, Urlaub H, Backofen R, Gophna U, Marchfelder A (2019) The nuts and bolts of the *Haloferax* CRISPR-Cas system I-B. *RNA Biol* 16(4):469-480 doi:10.1080/15476286.2018.1460994
- Margesin R, Schinner F (2001) Potential of halotolerant and halophilic microorganisms for biotechnology. *Extremophiles* 5(2):73-83
- Melvik JE, Dornish M (2004) Alginate as a carrier for cell immobilisation. In: Nedović V, Willaert R (eds) *Fundamentals of cell immobilisation biotechnology*. Springer Netherlands, Dordrecht, pp 33-51
- Mevarech M, Frolow F, Gloss LM (2000) Halophilic enzymes: proteins with a grain of salt. *Biophys Chem* 86(2-3):155-164 doi:10.1016/s0301-4622(00)00126-5
- Mevarech M, Werczberger R (1985) Genetic transfer in *Halobacterium volcanii*. *J Bacteriol* 162(1):461-462
- Mezzomo N, Maestri B, dos Santos RL, Maraschin M, Ferreira SRS (2011) Pink shrimp (*P. brasiliensis* and *P. paulensis*) residue: influence of extraction method on carotenoid concentration. *Talanta* 85(3):1383-1391 doi:<https://doi.org/10.1016/j.talanta.2011.06.018>

- Mohammadipanah F, Hamed J, Dehghani M (2015) Halophilic bacteria: potentials and applications in biotechnology. In: Maheshwari DK, Saraf M (eds) Halophiles: biodiversity and sustainable exploitation. Springer International Publishing, Cham, pp 277-321
- Moussavi G, Barikbin B, Mahmoudi M (2010) The removal of high concentrations of phenol from saline wastewater using aerobic granular SBR. Chem Eng J 158(3):498-504 doi:<https://doi.org/10.1016/j.cej.2010.01.038>
- Mullakhanbhai MF, Larsen H (1975) *Halobacterium volcanii* spec. nov., a Dead Sea halobacterium with a moderate salt requirement. Arch Microbiol 104(1):207-214 doi:10.1007/bf00447326
- Obradovic B, Nedović VA, Bugarski B, Willaert RG, Vunjak-Novakovic G (2004) Immobilised cell bioreactors. In: Nedović V, Willaert R (eds) Fundamentals of cell immobilisation biotechnology. Springer Netherlands, Dordrecht, pp 411-436
- Oren A (1999) Bioenergetic aspects of halophilism. Microbiol Mol Biol Rev 63(2):334-348
- Oren A (2008) Microbial life at high salt concentrations: phylogenetic and metabolic diversity. Saline Systems 4:2 doi:10.1186/1746-1448-4-2
- Oren A (2010) Industrial and environmental applications of halophilic microorganisms. Environ Technol 31(8-9):825-834 doi:10.1080/09593330903370026
- Oren A (2013) Life at high salt concentrations, intracellular KCl concentrations, and acidic proteomes. Front Microbiol 4(315) doi:10.3389/fmicb.2013.00315
- Oren A (2015) Halophilic microbial communities and their environments. Curr Opin Biotechnol 33:119-124 doi:<https://doi.org/10.1016/j.copbio.2015.02.005>
- Oren A, Arahall DR, Ventosa A (2009) Emended descriptions of genera of the family *Halobacteriaceae*. Int J Syst Evol Microbiol 59(3):637-642 doi:<https://doi.org/10.1099/ijs.0.008904-0>
- Patel S, Saraf M (2015) Perspectives and application of halophilic enzymes. In: Maheshwari DK, Saraf M (eds) Halophiles: biodiversity and sustainable exploitation. Springer International Publishing, Cham, pp 403-419
- Peck RF, Johnson EA, Krebs MP (2002) Identification of a lycopene beta-cyclase required for bacteriorhodopsin biogenesis in the archaeon *Halobacterium salinarum*. J Bacteriol 184(11):2889-2897 doi:10.1128/jb.184.11.2889-2897.2002
- Pfeifer F, Gregor D, Hofacker A, Plosser P, Zimmermann P (2002) Regulation of gas vesicle formation in halophilic archaea. J Mol Microbiol Biotechnol 4(3):175-181
- Poli A, Di Donato P, Abbamondi GR, Nicolaus B (2011) Synthesis, production, and biotechnological applications of exopolysaccharides and polyhydroxyalkanoates by archaea. Archaea 2011, 1-13 doi:10.1155/2011/693253
- Raghukumar C, D'Souza-Ticlo D, Verma AK (2008) Treatment of colored effluents with lignin-degrading enzymes: an emerging role of marine-derived fungi. Crit Rev Microbiol 34(3-4):189-206 doi:10.1080/10408410802526044
- Rawson A, Tiwari BK, Tuohy MG, O'Donnell CP, Brunton N (2011) Effect of ultrasound and blanching pretreatments on polyacetylene and carotenoid content of hot air and freeze dried carrot discs. Ultrason Sonochem 18(5):1172-1179 doi:<https://doi.org/10.1016/j.ultsonch.2011.03.009>

- Rehm BH (2003) Polyester synthases: natural catalysts for plastics. *Biochem J* 376(Pt 1):15-33 doi:10.1042/bj20031254
- Rehm BH (2007) Biogenesis of microbial polyhydroxyalkanoate granules: a platform technology for the production of tailor-made bioparticles. *Curr Issues Mol Biol* 9(1):41-62
- Reuter C, Maupin-Furlow J (2005) Analysis of proteasome-dependent proteolysis in *Haloferax volcanii* cells, using short-lived green fluorescent proteins. *Appl Environ Microbiol* 70:7530-7538 doi:10.1128/AEM.70.12.7530-7538.2004
- Robinson JL, Pyzyna B, Atrasz RG, Henderson CA, Morrill KL, Burd AM, DeSoucy E, Fogleman RE, Naylor JB, Steele SM, Elliott DR, Leyva KJ, Shand RF (2005) Growth kinetics of extremely halophilic archaea (family halobacteriaceae) as revealed by arrhenius plots. *J Bacteriol* 187(3):923-929 doi:10.1128/JB.187.3.923-929.2005
- Rodrigo-Baños M, Garbayo I, Vílchez C, Bonete MJ, Martínez-Espinosa RM (2015) Carotenoids from haloarchaea and their potential in biotechnology. *Mar Drugs* 13(9):5508-5532 doi:10.3390/md13095508
- Rodrigues-Oliveira T, Belmok A, Vasconcellos D, Schuster B, Kyaw CM (2017) Archaeal S-Layers: overview and current state of the art. *Front Microbiol* 8(2597) doi:10.3389/fmicb.2017.02597
- Rønnekleiv M (1995) Bacterial carotenoids 53* C50-carotenoids 23; carotenoids of *Haloferax volcanii* versus other halophilic bacteria. *Biochem Syst Ecol* 23(6):627-634 doi:[https://doi.org/10.1016/0305-1978\(95\)00047-X](https://doi.org/10.1016/0305-1978(95)00047-X)
- Sakurai T, Kataoka K (2007) Basic and applied features of multicopper oxidases, CueO, bilirubin oxidase, and laccase. *Chem Rec* 7(4):220-229 doi:10.1002/tcr.20125
- Salgaonkar BB, Bragança JM (2017) Utilization of sugarcane bagasse by *Halogeometricum borinquense* strain E3 for biosynthesis of poly(3-hydroxybutyrate-co-3-hydroxyvalerate). *Bioengineering (Basel)* 4(2):50 doi:10.3390/bioengineering4020050
- Sapienza C, Rose MR, Doolittle WF (1982) High-frequency genomic rearrangements involving archaeobacterial repeat sequence elements. *Nature* 299(5879):182-185 doi:10.1038/299182a0
- Sarmiento F, Peralta R, Blamey JM (2015) Cold and hot extremozymes: industrial relevance and current trends. *Front Bioeng Biotechnol* 3:148-148 doi:10.3389/fbioe.2015.00148
- Schmid A, Dordick JS, Hauer B, Kiener A, Wubbolts M, Witholt B (2001) Industrial biocatalysis today and tomorrow. *Nature* 409(6817):258-268 doi:10.1038/35051736
- Shand RF, Leyva KJ (2007) Peptide and protein antibiotics from the domain archaea: halocins and sulfolobocins. In: Riley MA, Chavan MA (eds) *Bacteriocins: ecology and evolution*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 93-109
- Sharma N, Farooqi MS, Chaturvedi KK, Lal SB, Grover M, Rai A, Pandey P (2014) The halophile protein database. *Database (Oxford)* 2014:bau114-bau114 doi:10.1093/database/bau114
- Singh SM, Panda AK (2005) Solubilization and refolding of bacterial inclusion body proteins. *J Biosci Bioeng* 99(4):303-310 doi:10.1263/jbb.99.303
- Sleytr UB, Huber C, Ilk N, Pum D, Schuster B, Egelseer EM (2007) S-layers as a tool kit for nanobiotechnological applications. *FEMS Microbiol Lett* 267(2):131-144 doi:10.1111/j.1574-6968.2006.00573.x
- Smith EaP, Ron (2018) *Haloferax volcanii* for carotenoid production. vol Honours Theses, Paper 899

- Solomon EI, Sundaram UM, Machonkin TE (1996) Multicopper oxidases and oxygenases. *Chem Rev* 96(7):2563-2606 doi:10.1021/cr950046o
- Strand BL, Skjåk-Bræk G, Gåserød O (2004) Microcapsule formulation and formation. In: Nedović V, Willaert R (eds) *Fundamentals of cell immobilisation biotechnology*. Springer Netherlands, Dordrecht, pp 165-183
- Stroud A, Liddell S, Allers T (2012) Genetic and biochemical identification of a novel single-stranded dna-binding complex in *Haloferax volcanii*. *Front Microbiol* 3(224) doi:10.3389/fmicb.2012.00224
- Strillinger E, Grotzinger SW, Allers T, Eppinger J, Weuster-Botz D (2016) Production of halophilic proteins using *Haloferax volcanii* H1895 in a stirred-tank bioreactor. *Appl Microbiol Biotechnol* 100(3):1183-1195 doi:10.1007/s00253-015-7007-1
- Tamborini L, Fernandes P, Paradisi F, Molinari F (2018) Flow bioreactors as complementary tools for biocatalytic process intensification. *Trends Biotechnol* 36(1):73-88 doi:10.1016/j.tibtech.2017.09.005
- Tamir A, Eichler J (2017) N-Glycosylation is important for proper *Haloferax volcanii* S-Layer stability and function. *Appl Environ Microbiol* 83(6):e03152-16 doi:10.1128/AEM.03152-16
- Timpson LM, Alsafadi D, Mac Donnchadha C, Liddell S, Sharkey MA, Paradisi F (2012) Characterization of alcohol dehydrogenase (ADH12) from *Haloarcula marismortui*, an extreme halophile from the Dead Sea. *Extremophiles* 16(1):57-66 doi:10.1007/s00792-011-0405-0
- Timpson LM, Liliensiek AK, Alsafadi D, Cassidy J, Sharkey MA, Liddell S, Allers T, Paradisi F (2013) A comparison of two novel alcohol dehydrogenase enzymes (ADH1 and ADH2) from the extreme halophile *Haloferax volcanii*. *Appl Microbiol Biotechnol* 97(1):195-203 doi:10.1007/s00253-012-4074-4
- Torregrosa-Crespo J, Pire C, Martínez-Espinosa R (2017) Biocompounds from Haloarchaea and their uses in biotechnology. In: Sghaier, Haitham; Najjari, Afef; Ghedira, Kais (Eds.). *Archaea - New Biocatalysts, Novel Pharmaceuticals and Various Biotechnological Applications*. InTech, 2017, pp 63-82
- Tripepi M, Esquivel RN, Wirth R, Pohlschroder M (2013) *Haloferax volcanii* cells lacking the flagellin FlgA2 are hypermotile. *Microbiology* 159(Pt 11):2249-2258 doi:10.1099/mic.0.069617-0
- Tripepi M, You J, Temel S, Önder Ö, Brisson D, Pohlschröder M (2012) N-Glycosylation of *Haloferax volcanii* flagellins requires known Agl proteins and is essential for biosynthesis of stable flagella. *J Bacteriol* 194(18):4876 doi:10.1128/JB.00731-12
- Uthandi S, Saad B, Humbard M, Maupin-Furlow J (2009) LccA, an archaeal laccase secreted as a highly stable glycoprotein into the extracellular medium by *Haloferax volcanii*. *Appl Environ Microbiol* 76:733-743 doi:10.1128/AEM.01757-09
- Wright DB, Banks DD, Lohman JR, Hilsenbeck JL, Gloss LM (2002) The effect of salts on the activity and stability of *Escherichia coli* and *Haloferax volcanii* dihydrofolate reductases. *J Mol Biol* 323(2):327-344 doi:10.1016/s0022-2836(02)00916-6
- Xue Y-P, Xu M, Chen H-S, Liu Z-Q, Wang Y-J, Zheng Y-G (2013) A novel integrated bioprocess for efficient production of (R)-(-)-Mandelic Acid with immobilized *Alcaligenes faecalis* ZJUTB10. *Org Process Res Dev* 17(2):213-220 doi:10.1021/op3001993
- Yin J, Chen JC, Wu Q, Chen GQ (2015) Halophiles, coming stars for industrial biotechnology. *Biotechnol Adv* 33(7):1433-1442 doi:10.1016/j.biotechadv.2014.10.008

- Zalazar L, Pagola P, Miro MV, Churio MS, Cerletti M, Martinez C, Iniesta-Cuerda M, Soler AJ, Cesari A, De Castro R (2019) Bacterioruberin extracts from a genetically modified hyperpigmented *Haloferax volcanii* strain: antioxidant activity and bioactive properties on sperm cells. *J Appl Microbiol* 126(3):796-810 doi:10.1111/jam.14160
- Zhang J, Sun Z, Sun P, Chen T, Chen F (2014a) Microalgal carotenoids: beneficial effects and potential in human health. *Food Funct* 5(3):413-425 doi:10.1039/c3fo60607d
- Zhang J-F, Liu Z-Q, Zhang X-H, Zheng Y-G (2014b) Biotransformation of iminodiacetonitrile to iminodiacetic acid by *Alcaligenes faecalis* cells immobilized in ACA-membrane liquid-core capsules. *Chem Pap* 68(1):53-64 doi:10.2478/s11696-013-0423-8

Fig. 1 Regulated and constitutive protein expression system in *H. volcanii* - a) pTA1228 The *p.tnaA* promoter is induced by L-tryptophan and allows for regulated protein expression. An N-terminal histidine (His) tag is available for immobilised metal-based affinity chromatography (IMAC) protein purification. A multiple cloning site is located after His tag. At the 5' end of the halophilic gene, the ATG start codon is replaced with an *NcoI*, *BspHI*, *PciI* or *SphI* site (depending on the coding sequence), and inserted in-frame with the His tag at the *PciI* or *SphI* site; at the 3' end of the gene, an *EcoRI* or *BamHI* site is used. The cloning cassette is flanked by L11e and t.Syn terminators to prevent read-through transcription. **b) pTA1392** A derivative of pTA1228. A C-terminal streptavidin II (StrepII) tag is available along with the His tag. At the 3' end of the gene, the stop codon is replaced with an *NheI* site and inserted in-frame with the StrepII tag. **c) pTA1926** A 50% reduced strength tryptophan inducible promoter, when compared with *p.tnaA*. Both His and Strep II tags are available. **d) pTA1992** A strong *p.syn* promoter for constitutive protein expression with both His and Strep II tags

Fig. 2 Simple method for immobilisation of *H. volcanii* within the calcium alginate hydrogels - a) Calcium alginate beads with encapsulated *H. volcanii* is formed instantaneously following the dropwise addition of resuspended *H. volcanii* cells in 4% sodium alginate solution into 1.5% CaCl₂ solution, **b)** Formation of pink beads due to the presence of high carotenoid pigment confirms entrapment of *H. volcanii* within the calcium alginate beads

Table 1: Biotechnologically important components encoded by *H. volcanii*

Component	Potential applications	References
Phytoene synthase (HVO_2524), Phytoene desaturase (HVO_2528), Lycopene elongase (HVO_2527)	Carotenoid biosynthesis	(Smith 2018; Zalazar et al. 2019)
Alcohol dehydrogenases - <i>HvADH1</i> (HVO_2428) and <i>HvADH2</i> (HVO_B0071)	Fine chemical production through biocatalysis	(Timpson et al. 2013; Alsafadi et al. 2017; Haque et al. 2019)
Laccase (HVO_B0205)	Biopulping, biobleaching, biofuel production	(Uthandi et al. 2009)
Halocin	Antimicrobials, bioremediation	(Kavitha et al. 2011)
Polyhydroxyalkanoate	Bioplastic production	(Koller 2019)
Outer surface layer (S-layer)	Nanobiotechnology	(Ilk et al. 2008; Sleytr et al. 2007; Lobasso et al. 2015)