

## CORRECTION

# Correction: Inactive matrix gla protein plasma levels are associated with peripheral neuropathy in Type 2 diabetes

Anne-Caroline Jeannin, Joe-Elie Salem, Ziad Massy, Carole Elodie Aubert, Cees Vemeer, Chloé Amouyal, Franck Phan, Marine Halbron, Christian Funck-Brentano, Agnès Hartemann, Olivier Bourron

The fifth author's name is spelled incorrectly. The correct name is: Cees Vermeer. The correct citation is: Jeannin A-C, Salem J-E, Massy Z, Aubert CE, Vermeer C, Amouyal C, et al. (2020) Inactive matrix gla protein plasma levels are associated with peripheral neuropathy in Type 2 diabetes. PLoS ONE 15(2): e0229145. <https://doi.org/10.1371/journal.pone.0229145>

## Reference

1. Jeannin A-C, Salem J-E, Massy Z, Aubert CE, Vemeer C, Amouyal C, et al. (2020) Inactive matrix gla protein plasma levels are associated with peripheral neuropathy in Type 2 diabetes. PLoS ONE 15(2): e0229145. <https://doi.org/10.1371/journal.pone.0229145> PMID: 32092076



## OPEN ACCESS

**Citation:** Jeannin A-C, Salem J-E, Massy Z, Aubert CE, Vemeer C, Amouyal C, et al. (2020) Correction: Inactive matrix gla protein plasma levels are associated with peripheral neuropathy in Type 2 diabetes. PLoS ONE 15(5): e0232996. <https://doi.org/10.1371/journal.pone.0232996>

**Published:** May 5, 2020

**Copyright:** © 2020 Jeannin et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.