

Contaminants in Aquatic and Terrestrial Environments

**TRACING URBAN WASTEWATER CONTAMINANTS INTO
THE ATLANTIC OCEAN BY NONTARGET SCREENING**Pablo Antonio Lara-Martín, Aurea C. Chiaia-Hernández, Miriam
Biel-Maeso, Rosa María Baena-Nogueras, and Juliane Hollender*Environ. Sci. Technol.*, **Just Accepted Manuscript** • DOI: 10.1021/acs.est.9b06114 • Publication Date (Web): 02 Mar 2020Downloaded from pubs.acs.org on March 4, 2020**Just Accepted**

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1 TRACING URBAN WASTEWATER CONTAMINANTS INTO THE ATLANTIC
2 OCEAN BY NONTARGET SCREENING

3

4 Pablo A. Lara-Martín^{1*}, Aurea C. Chiaia-Hernández^{2,3}, Miriam Biel-Maeso¹, Rosa M. Baena-
5 Noguerras¹, Juliane Hollender^{2,4}

6

7 ¹Physical Chemistry Department, Faculty of Marine and Environmental Sciences, University of
8 Cadiz, Campus of International Excellence of the Sea (CEI-MAR), 11510 Puerto Real, Spain

9 ²Eawag, Swiss Federal Institute of Aquatic Science and Technology (Eawag), 8600 Dübendorf,
10 Switzerland

11 ³Institute of Geography and Oeschger Centre for Climate Change Research, University of Bern, Bern,
12 3012 Switzerland

13 ⁴Institute of Biogeochemistry and Pollutant Dynamics, IBP, ETH Zurich, 8092 Zurich, Switzerland

14

15

16

17

18

19

20

21

22

23

24

25

26

27 Abstract


28

29 Oceans are the ultimate sink for many of the over 100 million man-made substances. Until now,
30 monitoring was limited to a reduced number of targeted persistent organic pollutants, reaching open
31 waters mainly via atmospheric deposition. However, the composition and fate of the thousands of
32 pollutants reaching the marine environment through wastewater discharges from coastal sources
33 remain largely unexplored. By combining a newly developed non-target screening (NTS) workflow
34 and high resolution mass spectrometry (HRMS), we have identified over 500 sewage-derived
35 contaminants occurring in the ocean. Samples from the NE Atlantic contained this anthropogenic
36 imprint at distances over 50 km from the coastline and > 500 m depth, beyond the continental margin.
37 The range of identified compounds spans from pharmaceuticals and personal care products to food
38 additives and industrial chemicals, including several that have never been reported in the
39 environment, as they escaped conventional targeted analytical methods. Predicting the effects of the
40 continuous input of this chemical “cocktail” on marine ecosystems is a formidable challenge, since
41 40% of the detected compounds lack information regarding their use and ecotoxicity.

42

43 TOC

44



45

46

47

48

49 **1. Introduction**

50

51 Oceans are the vastest ecosystems on Earth, as well as sources of a wide number of resources such as
52 fisheries, minerals, and energy. Coastal zones are the interface between land and the ocean, and have
53 historically attracted human settlements because of these resources, as well as for logistical purposes
54 (e.g., marine trade and transport). The development and utilization of coastal zones have greatly
55 increased during the recent decades, undergoing tremendous socio-economic and environmental
56 changes. The coastal population is projected to be nearly 1 billion by 2030, and most of the world's
57 megacities are located by the coastline (1). These rapid changes in the coastal zones have had
58 detrimental impacts towards marine species and habitats. Anthropogenic stressors on marine
59 ecosystems, such as overfishing and nutrient input, are increasing globally (2). Among them,
60 pollution by man-made chemicals has recently caught the attention of both environmental scientists
61 and public interest, especially after becoming evident to the naked eye in the form of floating plastic
62 debris in vast areas of the ocean (3).

63

64 Plastics are not the only artificial compounds mainly derived from land-based sources that can be
65 detected in seawater. As of 2018, there are over 130 million registered chemical substances in the
66 Chemical Abstracts Service (CAS) Registry File. Approximately 400,000 of them are somehow
67 regulated in the international markets (e.g., U.S. Toxic Substances Control Act, or TSCA) and much
68 less regularly monitored in aquatic systems (4). The vast number of chemical contaminants that
69 consequently may be potentially present, but have not been detected, in the marine environment poses
70 a great threat to the marine diversity and a challenge for both regulators and researchers. For
71 regulators, one of the most common approaches has been to develop lists of substances to focus on
72 their potential hazard to the ecosystems and/or to human health. In this context, there are several
73 national and international legislations such as the US Clean Water Act (CWA), the European Water
74 Framework Directive (WFD, 2000/60/EC1) and the Marine Strategy Framework Directive (MSFD,

75 2008/56/EC2) that established a limited number of substances of wide concern (named priority
76 substances) to be regularly monitored in the coastline. Most of the target substances included in these
77 legislations are either metals (As, Cd, Hg, etc.) or persistent organic pollutants (POPs) such as
78 organochlorine compounds, pesticides, brominated flame retardants, perfluorinated chemicals, and
79 recognized endocrine disruptors (e.g., nonylphenol and phthalates). The most comprehensive list of
80 potential chemical contaminants in the marine environment, based on a compilation of substances
81 from relevant aforementioned legislations, global conventions (e.g., Stockholm Convention), and
82 other international organizations (e.g., US EPA) contains approximately 2700 substances (5). Many
83 of these proposed compounds are considered as contaminants of emerging concern and have not been
84 implemented yet in routine monitoring programs.

85

86 From an analytical point of view, the challenge is even bigger due to the wide range of chemical
87 structures and properties considered and the very low concentrations (< parts per billion, or ppb)
88 occurring in the ocean. To date, most of the effort has been focused on a targeted analysis of selected
89 POPs, susceptible to long-range atmospheric transport and bioaccumulation in marine organisms (6).
90 Analysis of non-regulated substances such as specific sewage-markers (e.g., coprostanol) and, most
91 recently, selected pharmaceuticals has often been limited to enclosed seas such as the Baltic Sea and
92 the Mediterranean Sea (7, 8). The identification of new contaminants is not only a challenge but also
93 one of the most commonly-mentioned bottlenecks in the advance of disciplines such as environmental
94 chemistry and ecotoxicology (9). Recent advances in high resolution mass spectrometry (HRMS),
95 capable of discerning individual chemicals in complex matrices by accurate mass measurement,
96 coupled with chromatographic separation are seen as key tools to address this challenge. By enabling
97 higher throughput analysis of unknown substances at trace levels, the new so-called non-target
98 screening (NTS) strategies aim to capture the universe of anthropogenic organic chemicals occurring
99 in the environment. As we aim to explore the whole chemical composition of samples instead of
100 specific target compounds, complexity in the interpretation of the results increases exponentially. The

101 resulting big data are managed by crafting analytical workflows capable of analyzing HRMS data
102 with openly accessible online chemical compound databases, structure prediction algorithms, and
103 statistical tools (10). To date, much effort has been dedicated to the identification of relevant
104 contaminants and transformation products in wastewater treatment plants (WWTPs) (11, 12) and to
105 characterize components in hydraulic fluids used for fracking (13).

106

107 Here, by using state-of-the-art NTS tools, we present a comprehensive characterization of the oceanic
108 contamination by organic pollutants. As the release of synthetic chemicals into the marine
109 environment may occur through a variety of pathways, including river and atmospheric transport, as
110 well as directly at sea via aquaculture, shipping and fishing activities (5), we have focused our
111 attention on sewage-derived contaminants by performing a directed NTS on these substances.
112 Wastewater is the largest contributing source of emerging contaminants to aquatic environments (14)
113 and a rising issue in coastal areas due to the aforementioned increase of urban settlements and
114 population. The study area selected was the Gulf of Cadiz (NE Atlantic), as well as its coastline
115 (Cadiz Bay, SW Spain) and the main WWTP (Fig. 1). Comparative analyses of wastewater, coastal
116 waters, and offshore waters collected were performed to unravel the occurrence of 537 wastewater
117 contaminants that could be detected from source to the open ocean, at distances further than 50 km
118 from the coast and > 500 m depth.

119

120 **2. Materials and Methods**

121

122 **Water sampling.** All samples were collected in summer 2015 from the inlet and outlet of the WWTP
123 serving Jerez de la Frontera, the main population in the area, and along 6 different transects: 3 within
124 the Cadiz Bay area (Guadalete River estuary: G1-G10, Sancti Petri creek: S1-9, Rio San Pedro creek:
125 R1-8) and 3 offshore (Cadiz Bay coast: CC1-4, Guadalquivir River mouth: GD1-4. and Trafalgar

126 Cape: TF1-5) (Fig. 1). Description of the sampling area and further details on how the sampling was
127 conducted are in ref. 15 and Supporting Information (SI), respectively.

128

129 **Sample extraction and mass spectrometry determination.** Briefly, all water samples were filtered
130 with glass fiber filters and extracted using Oasis HLB cartridges. Extracts were spiked with a mix of
131 internal standards (Table S1) and analyzed by high performance liquid chromatography coupled to a
132 Q Exactive hybrid quadrupole-Orbitrap mass spectrometer equipped with an electrospray ionization
133 source (HPLC-ESI-HRMS) and operating in full-scan mode at a mass resolution of 140,000 FWHM
134 referenced to m/z 400 according to ref. 16. For structural elucidation of prioritized compounds (see
135 next section), targeted MS/MS fragmentation was performed at 17,5000 FWHM. Specific
136 information on SPE conditions and HPLC-HRMS instrumental settings can be found in SI.

137

138 **Selection, prioritization, and identification sewage-derived compounds.** A flowchart of the
139 workflow used and the different nodes involved is presented in Fig. 2. The workflow consisted of a
140 combination of nodes implemented with statistical software R (version 3.4.3) and the vendor software
141 Compound Discoverer 2.1. The resulting dataset was subjected to statistical analyses using principal
142 component analysis (PCA) and agglomerative hierarchical cluster analysis (HCA) to prioritize
143 compounds of interest that were present both in wastewater and surface waters samples (Table S2).
144 Identification of the molecular structure of these substances was carried out through a combination
145 of different nodes (Fig. 2), which were complementary and provided identification confidence levels
146 (17, 18) between 1 and 4. This process involved using reference standards (EAWAG standards, Table
147 S1), library spectrum data search (mzCloud and MassBank), in-silico fragmentation (MetFrag and
148 FISh), online compound databases (ChemSpider and PubChem) and suspect lists (NORMAN
149 network and Environment and Food Safety, or EFS, lists) (Table S3). All the relevant information
150 regarding the identification of sewage-derived compounds found in both coastal and oceanic waters

151 (n = 537) is provided in Table S4. Further details on the different workflow nodes, their optimization,
152 and how the statistical analysis was performed can be found in SI.

153

154 **3. Results and Discussion**

155

156 **Prioritization of sewage-derived compounds.** HRMS analysis of wastewater samples at the influent
157 and effluent from the WWTP serving Jerez, the biggest coastal city (215,000 inhabitants) in the study
158 area (Fig. 1), revealed over 250,000 different compounds occurring in sewage. By compounds, we
159 mean the integration of all the different MS features such as molecular ions, salt adducts, and isotopes.
160 They cover a chemical space containing dissolved polar to semi-polar substances within a mass-to-
161 charge (m/z) window of 100 to 800 Da that are susceptible to electrospray ionization. Key aspects of
162 the methodology developed here for such integration are provided in SI. To isolate and prioritize
163 signals of interest (sewage-derived contaminants) from the background (naturally occurring
164 components making up the dissolved organic matter), the NTS workflow was first calibrated by using
165 a specific subset of samples. More specifically, we conducted a comparative analysis between the
166 aforementioned wastewater samples and sewage-impacted surface waters from Guadalete Estuary
167 (G1-10) (Fig. 1) to screen for common features. This estuary, highly polluted due to continuous
168 sewage inputs from Jerez WWTP and other pollution sources, has been the subject of a series of
169 studies on targeted contaminants, mainly surfactants (active ingredients in detergent formulations)
170 (15, 19, 20).

171

172 We found an average number of approximately 47,000 compounds in the samples analyzed along
173 Guadalete Estuary (G1-10). Principal component analysis (PCA) (Fig. S1A in the Supporting
174 Information Appendix) revealed that locations adjacent to the discharge outlet of the WWTP (G1 and
175 G2) differed substantially from the rest. They featured the highest numbers of compounds in the
176 estuary (> 50,000), and most of these compounds had the highest signal intensities. Therefore, we

177 used the samples from these locations, together with the influent and effluent wastewater samples, to
178 build a preliminary filter for prioritization of sewage-derived substances. The four samples (G1, G2,
179 effluent, and influent) shared 11,256 common compounds. Many of these substances might be
180 organic contaminants in wastewater persistent enough to partially escape sewage treatment and,
181 consequently, be detected in the receiving surface waters. Although the full characterization of the
182 contaminants entering the WWTP is out of the scope of this study, known substances frequently
183 detected in urban wastewater such as caffeine and cocaine (21) were identified using certified
184 standards (Table S1). For a considerable fraction of the detected chemicals (35%), relative signal
185 intensities (previously normalized by considering sample volume and ion suppression using internal
186 standards) were higher in the wastewater effluent than in the influent, suggesting either de-
187 conjugation of parent compounds or transformation into different products at the WWTP. Examples
188 are ranitidine S-oxide and dihydroxycarbamazepine, two pharmaceutical biodegradation products
189 also confirmed with available standards.

190

191 In a second step, the number of possible sewage-derived contaminants was reduced to 6892 by
192 hierarchical cluster analysis (HCA) of the 11,256 pre-selected compounds using all the samples taken
193 along Guadalete Estuary (Fig. 3A). The main criterion was to select those compounds whose
194 concentrations decreased continuously from G1 (adjacent to the WWTP outlet) towards G10 (at the
195 mouth of the estuary), an indication that they are anthropogenically sourced through wastewater
196 discharges and undergo dilution and/or degradation/sorption during their transport (Fig. 3B). The rest
197 of the substances that did not follow this trend were assigned to either natural dissolved organic matter
198 (34%) (Fig. 3C) or different contamination sources (5%) (Fig. 3D). More specifically, the latter group
199 showed a maximum around sample site G8, where a port serving El Puerto de Santa Maria (90,000
200 inhabitants) is located, so its occurrence may be associated with the maritime traffic and naval
201 activities occurring in the estuary (15). The resulting lists of prioritized sewage-derived components
202 (Table S2) were then included into the NTS workflow to limit the number of substances to detect and

203 identify in the rest of samples. These include coastal waters from Cadiz Bay (R1-8 and P1-9, at two
204 tidal creeks) and surface and bottom samples taken offshore at the Gulf of Cadiz (CC1-4, GD1-4, and
205 TF1-5, three transects of approximately 50 km length away from the coastline, covering the
206 continental shelf and the shelf break) (Fig. 1).

207

208 **Identification of wastewater contaminants in the ocean.** The average number of possible organic
209 compounds detected in the coastal ($n = 26$) and oceanic samples ($n = 27$) analyzed was approximately
210 36,500 and 12,000, respectively (Fig. 4). The percentage of those compounds included in the list of
211 sewage-derived contaminants was between 1 and 11%, depending on the sample. Specific examples
212 are plotted in Fig. S2, where red dots are wastewater contaminants and the rest (grey dots) are either
213 contaminants from other sources or naturally occurring organic matter. There were two coastal
214 sampling sites, Guadalete Estuary (G1-10) and Rio San Pedro tidal creek (R1-R8) where more than
215 50,000 substances were measured, indicative of higher concentrations of dissolved organic matter.
216 As discussed in the previous section, this number is partly justified by the continuous input of
217 wastewater from Jerez WWTP into the estuary, which resulted in 6% of total number of compounds,
218 on average, being attributed to sewage-derived chemicals. The tidal creek, however, is not subjected
219 to direct wastewater discharges, as confirmed by the considerably lower percentage ($< 2\%$), number
220 (450-1400), and signal intensities of sewage-derived chemicals detected here. PCA plots (Fig. S1A)
221 also grouped samples from this area differently from the rest. The major contributors to the dissolved
222 organic carbon (DOC) are attributed to effluents from fish farms (22), located in the vicinity of
223 sampling site R1. The characterization of the contaminants derived from aquaculture activities is out
224 of the scope of this work.

225

226 The spatial distribution of sewage-derived compounds showed a general trend, where the numbers
227 and concentrations of contaminants (expressed as normalized signal intensities by considering sample
228 volume and ion suppression using internal standards) progressively declined when moving from the

229 main source of wastewater discharge towards the open ocean (Fig. 4). This was expected due to
230 dilution and co-occurrence of different environmental processes such as biodegradation, photolysis
231 or sorption, aimed to reduce the concentrations of chemicals in aquatic environments (23). The
232 decreasing trend was not only visible when comparing different coastal and ocean transects, but also
233 to a lesser degree, within each transect as the distance increased from the coastline (Fig. S3). From
234 the almost 7000 compounds included in this analysis (Table S2), the number of sewage-derived
235 substances detected offshore was reduced to between 220 and 600. Some sampling sites (e.g.,
236 Trafalgar Cape) were considerably more impacted than the rest (Fig. S1B) possibly due to direct
237 discharges of untreated wastewater from popular touristic spots (e.g., Caños de Meca) not connected
238 to the main sewage network, in agreement to local maximum values for DOC measured there (24).
239 Additionally, there were also differences in the vertical distribution of sewage components along the
240 water column. Higher relative signal intensities and numbers were always detected at the surface (Fig.
241 S3), on average $22.8 \pm 1.5\%$ higher. These differences have been previously reported for target
242 contaminants such as pharmaceuticals and surfactants in urbanized estuaries (25) and can be
243 attributed to positive buoyancy of less dense wastewater plumes entering the ocean via estuaries and
244 marine outfalls.

245

246 As the main goal of this study was to characterize substances entering and occurring in the Atlantic
247 Ocean, identification of compounds by assigning specific compound formulas, structures, and names
248 was performed uniquely for those detected in offshore samples. This resulted in 537 different sewage-
249 related chemicals (Table S4). Confidence identification levels from 1 (the highest confidence,
250 confirmed with reference standards) to 4 (the lowest, tentative chemical compound from a chemical
251 database) were assigned by using a combination of available chemical pure standards, mass spectra
252 libraries, online chemical database searches, and in-silico mass spectral fragmentation algorithms (see
253 Materials and Methods section). Some examples to illustrate different aspects used for identification
254 are shown in Fig. S4. Here, the artificial sweetener sucralose, the food additive sulfurol, and the

255 industrial chemical 17-amino-3,6,9,12,15-pentaoxaheptadecan-1-ol, were identified at levels 1, 2, and
256 3, respectively. Identification confidence levels and elemental composition of all the sewage-derived
257 contaminants detected in oceanic waters from the Gulf of Cadiz is summarized in Fig. 5A, where
258 information on their inclusion in chemical databases and known applications is presented in Fig. 5B.
259 The vast majority of contaminants (85%) were exclusively formed of C, H, N, O, and S atoms,
260 covering a wide range of functional groups including alcohols, carboxylates, alkylamines, sulfonated
261 and sulfated compounds. The rest were either halogenated chemicals (mainly chlorinated substances,
262 see blue bar in Fig. 5A) or had P and/or Si atoms in their composition (see red bar in Fig. 5A). Other
263 chemicals such as hydrocarbons (composed of C and H only) were not detected due to the limitations
264 of the analytical approach used, based on HPLC-ESI-HRMS.

265

266 Only about 20% of the prioritized sewage contaminants could be identified with high degree of
267 confidence (levels 1 and 2, see light green and blue sectors, respectively, in Fig. 5A) due to the limited
268 number of standards available in our laboratory (Table S1) and the still relatively low (although
269 growing) number of HRMS spectra recorded from commercially available standards. For instance,
270 as of 2018, there were approximately 16 thousand substances in one of the mass spectral libraries
271 used here (MassBank); this number represents a tiny fraction of the more than 130 million registered
272 chemical substances in the Chemical Abstracts Service (CAS). Our identification effort still resulted
273 in over 100 chemicals positively identified in the ocean, many of them never reported before. The
274 bulk of the prioritized list (60% of the substances), however, was composed by chemicals identified
275 at level 3 by combining database searches (suspect lists in Table S3, ChemSpider, and PubChem) and
276 in-silico fragmentation tools (MetFrag and FISH). Higher scores in Table S4 (columns P, R, and S)
277 are indicative of higher probabilities of achieving successful identification. For the remaining
278 chemicals (17%), HRMS information was not enough to elucidate their chemical structures. Tentative
279 names were assigned from top-ranked compounds in ChemSpider and PubChem databases (in terms
280 of number of references) that matched their molecular formula.

281

282 **Environmental relevance of the chemicals identified.** We could find 95% of the different
283 substances identified in the ocean in online databases (ChemSpider and PubChem, light blue and light
284 green sectors combined in Fig. 5B), and almost 50% of the possible candidates were already compiled
285 in lists of suspected environmental contaminants (EFS and NORMAN network lists, see Table S3)
286 (light green sector in Fig. 5B). Relevant environmental information such as their uses and applications
287 was found for over 60% of them. This allowed their classification into six different groups of
288 chemicals: pharmaceuticals and related bioactive substances, pesticides, personal care products,
289 surfactants, industrial chemicals, and food additives. The top-ten compounds of each group, in terms
290 of signal intensity and frequency of detection, are presented in Fig. 6. Their relative abundance is also
291 depicted as maximum signal intensity ranges in coastal (G1-10, R1-8, and P1-9, red bars) and oceanic
292 (CC1-4, GD1-4, and TF1-5, blue bars) samples, the latter often being several orders of magnitude
293 lower. The rest of the contaminants were not considered for the analysis shown in Fig. 6 as no data
294 on their possible uses and/or ecotoxicological relevance could be retrieved (red bar in Fig. 5B).

295

296 Pharmaceutically active compounds were, in terms of number of identified chemicals ($n = 144$), the
297 most relevant group of contaminants, comprising up to 25% of the prioritized wastewater components
298 detected in the NE Atlantic. Those compounds showing highest intensities were analgesic (e.g.,
299 antipyrine, also known as phenazone) and psychiatric drugs such as venlafaxine (an antidepressant),
300 carbamazepine, and olanzapine. The former group were reported worldwide at the highest
301 concentrations in coastal WWTPs due to their higher consumption (non-prescription drugs in many
302 cases), whereas the latter are very recalcitrant when undergoing conventional sewage treatment (25).
303 The presence of metabolites (4-formylaminoantipyrine, 4-acetamidoantipyrine, and valsartan acid,
304 among others) and lidocaine (a local anesthetic) was also confirmed (level 1), whereas antivirals such
305 as telvibudine and nevirapine were identified at level 3. Possible effects on marine biota derived from
306 the occurrence of these bioactive substances have been recently reviewed by Mezzelani and co-

307 workers (26), including alterations in the enzymatic and immunological systems of filter feeders. As
308 NTS workflows are not aimed for accurate quantification of the concentrations, we proceeded to re-
309 analyze our samples using specific methodology for target determination of pharmaceuticals. Total
310 concentrations of these contaminants were between 50 and 200 ng L⁻¹, not currently known to be high
311 enough to pose any foreseen risk towards marine biota (20). This preliminary environmental risk
312 assessment, however, was not without limitations due to poor ecotoxicological data regarding the
313 toxicity of these substances towards marine species, as well as unknown additive or synergistic effects
314 with other chemicals co-existing in the water column. Among these chemicals, pesticides are also
315 bioactive substances on their own, designed to kill specific organisms. Fungicides, such as metalaxyl
316 and carbendazim, and herbicides (terbutryn, 2,4-D, and fluometuron) were identified at different
317 confidence levels. Toxicity towards primary producers such as marine microalgae can be expected
318 even at sub-ppb concentrations (27).

319

320 Surfactants were second in number of identified sewage-derived components, but first in terms of
321 signal intensities. Although their removal in conventional WWTPs is very efficient (25) (> 95%),
322 their current worldwide production is the highest of all synthetic chemicals (after plastics), with over
323 15 million tonnes per year (28). As a consequence, these compounds are reported at the highest
324 concentrations in treated effluents and streams in comparison with other organic contaminants not so
325 efficiently eliminated during wastewater treatment but used in lower amounts (e.g., psychiatric drugs)
326 (29). Linear alkylbenzene sulfonates (LAS), polyethylene glycols (PEG), and polypropylene glycols
327 (PPG) were the main components detected in seawater, together with their respective biodegradation
328 products (e.g., sulfophenyl carboxylic acids, or SPCs, and oxidized PEG) and manufacturing
329 impurities (dialkyltetralin sulfonates, or DATS). Rather than individual compounds, they are complex
330 mixtures comprising different homologous and ethoxymer series (Fig. S5) with changing
331 environmental and ecotoxicological properties (30). Although readily biodegradable according to
332 screening tests, their elimination was shown to be considerably slower under marine conditions (31)

333 in comparison with freshwater environments. This is explained by a combination of marine microbial
334 communities being less active than their freshwater counterparts toward xenobiotic chemicals and
335 complexation with ions in seawater that reduces bioavailability, thereby inhibiting biodegradation
336 (30). Surfactants are used in a myriad of applications including cleaning detergents (LAS), personal
337 care products (PEG and PPG), and various industrial uses (e.g., oil, textiles, polymers, agriculture,
338 and paints). The global surfactant market is forecast to grow at a compound annual rate of 6% over
339 the next years, with rising demand for personal care products (PCPs) being the market driver (30).
340 Other chemicals included in the formulation of PCPs that were detected in our samples were
341 fragrances (galaxolidone, a degradation product from galaxolide), UV filters (sulisobenzone), and
342 insect repellents (DEET and picaridin). Some of these compounds, reported in freshwater resources
343 from all continents, are bioaccumulative and suspected endocrine disruptors, leading to potential
344 alterations on fecundity, growth and development of exposed species (14).

345
346 Lastly, industrial chemicals were the third most relevant group identified by our NTS workflow. Their
347 inclusion in the prioritized list of sewage-derived contaminants can be explained by the presence of
348 industrial activities (e.g., wineries, laundry services, food processing, auto repair shops, etc.) within
349 the city and its surroundings, also served by Jerez WWTP. This is a very diverse group comprising
350 substances used in a wide range of applications, including processing aids (siloxanes), buffering
351 agents (CAPSO), reaction intermediates (benzenesulfonamide), complexing agents (21-Crown-7),
352 and corrosion inhibitors (benzotriazoles). The vast majority of the chemicals included in this group
353 (n = 54) has never been reported in the marine environment and there is no literature on their risks
354 towards marine species. This includes food additives such as the flavoring agents sulfurol, identified
355 at level 2, geranyl acetate, and eugenol benzoate (level 3). Some of them are not only synthetically
356 produced, but also natural products. Such is the case of panthenol, a vitamin (B5) but also a
357 moisturizer widely used in PCPs that was recently identified in WWTP using a NTS approach (12)
358 and also detected in our seawater samples (Table S4). To discern between natural and anthropogenic

359 sources and the relative contributions of these sources is challenging and certainly would not be
360 possible without using additional analytical techniques (e.g., compound-specific stable isotope
361 analysis). Nevertheless, the highest concentrations observed in urban wastewater ensure that cities
362 are one of the main sources of this type of chemicals towards the marine environment. Within this
363 category, only non-volatile cyclosiloxanes (D6 and D7), also used in PCPs and with potential
364 carcinogenicity, and benzotriazoles were previously reported in WWTP effluents (32, 33). Both are
365 resistant to biodegradation and very persistent in aquatic environments, and the latter have been
366 proposed as anthropogenic indicators in groundwater and rivers (34). Similarly, the artificial
367 sweetener sucralose, the food additive detected at the highest intensity in oceanic samples, is
368 considered as an ideal marker of sewage contamination in freshwater (35) and, more recently, coastal
369 systems (36) due to its recalcitrance, source specificity, and high solubility. Concentrations up to 70
370 ng L^{-1} were reported in the Gulf Stream (37), the only available data reported from offshore samples.
371 In the present study, we determined an average 350-fold decrease in the relative abundance of these
372 well-known sewage markers (Fig. S6A) from the most contaminated station (G1) to offshore samples.
373 Due to the low reactivity of these markers, it was mostly attributed to dilution of wastewater in the
374 ocean. Other chemicals more prone to undergo degradation (e.g., antipyrine and its main metabolites)
375 showed a decrease in their concentration of more than 1000-fold from the estuary to the ocean. The
376 transformation of the parent compound into different degradation products during their transport was
377 evident as the relative abundances increased for the latter as we moved away from the contamination
378 source (Fig. S6B).

379

380 **Future perspectives.** This work presents a list with over 500 different contaminants detected in
381 coastal and oceanic samples from NE Atlantic. Their occurrence not only along the coastline but also
382 in the whole continental shelf and beyond in a very dynamic environment, as opposed to previous
383 targeted studies in enclosed seas, confirms their ubiquity. The detection of these chemicals in the
384 open sea can only be explained by a combination of persistence in seawater and continuous input of

385 significant amounts from land-based sources such as WWTPs. Immediate efforts should be focused
386 towards development of specific target methodologies aimed at the accurate determination of the
387 concentrations of those contaminants detected by NTS, as well as ecotoxicity assays and/or modeling
388 to discern whether they may pose risks towards marine biota. This is also a long-term task as the
389 number of synthetic chemicals is continuously increasing (e.g., in 2015, 56 new drugs were launched
390 and more of 7000 compounds were in trial) (38). Their global trends in terms of worldwide production
391 and diversification has already outpaced the rates of change of other agents of global change such as
392 rising CO₂ atmospheric concentrations and habitat destruction (39), but pollution by synthetic
393 chemicals has not yet been included in most analyses of global change. In addition, and in spite of
394 nowadays stricter environmental policies, concentrations of many sewage-derived chemicals in the
395 marine environment are expected to keep rising within the next decades as the human population
396 increases while conventional WWTPs are incapable of efficient removal of many contaminants. This
397 growing trend has already been confirmed for pharmaceuticals from recent targeted analyses of dated
398 marine sediment cores (40). Mitigation measures include improving wastewater treatment through
399 the use of additional technologies (e.g., advanced oxidation processes) in current and planned coastal
400 WWTPs, which have proved to increase water quality in sewage-impacted freshwater settings (41).

401

402 This study also explores the boundaries of the current existing methodologies for environmental high-
403 throughput identification of compounds by HRMS. Although targeted analyses using reference
404 standards remains the most reliable way to determine contaminants, it is not feasible to extend this
405 approach over more than a few hundreds of substances of interest (more than 600 were used here) for
406 routine monitoring. As mass spectral libraries are growing and computational mass spectrometry is
407 advancing at a fast pace, the confidence in the identification of unknown substances using NTS
408 workflows will increase in the near future. Already acquired HRMS data such as ours (available in
409 File S1) will also benefit from upcoming achievements in the field (e.g., more robust in-silico
410 fragmentation algorithms and identification of new contaminants) through the use of retrospective

411 screening, as has been recently tested with aqueous environmental samples from different European
412 countries (42). Use of complementary techniques such as GC-HRMS for screening of hydrophobic
413 and semi-volatile chemicals will allow to expand the analytical horizon towards an even more
414 complete picture of the chemical universe in our environment (43). Lastly, by applying NTS
415 workflows similar to the one described here, different potential contamination sources towards the
416 marine environment such as aquaculture and maritime traffic could be investigated to unravel
417 additional contaminants.

418

419 **Associated content**

420 **Supporting Information.** Materials and methods, Figures S1-S7, Tables S1-S4, and File S1 (MS/MS
421 spectra). This material is available free of charge via the Internet at <http://pubs.acs.org>.

422

423 **Author information**

424 ***Corresponding Author.** Address: Campus Río San Pedro s/n, Campus de Excelencia Internacional
425 del Mar (CEI-MAR), 11510 Puerto Real, Cádiz, Spain. Phone: +34 956 016159. E-mail:
426 pablo.lara@uca.es

427 **Notes.** The authors declare no competing financial interest.

428

429 **Acknowledgments**

430

431 Thanks to J.E Schollé and K. Kiefer for their HPLC-HRMS and MetFragR support, respectively, and
432 to S. Salzmann during the revision process. This research was supported by the Spanish Ministry of
433 Economy, Industry and Competitiveness (MINECO) (CTM2015-70731-R) and by the University of
434 Bern through the Interfaculty Research Cooperation One Health.

435

436 **References**

437

- 438 1. Neumann, B.; Vafeidis, A.T.; Zimmermann, J.; Nicholls, R.J. Future coastal population growth
439 and exposure to sea-level rise and coastal flooding – A global assessment. *PLoS ONE* **2015**, 10, 3.
- 440 2. Halpern, B.S.; Frazier, M.; Potapenko, J.; Casey, K.S.; Koenig, K.; Longo, C.; Lowndes, J.S.;
441 Rockwood, R.C.; Selig, E.R.; Selkoe, K.A.; Walbridge, S. Spatial and temporal changes in
442 cumulative human impacts on the world's ocean. *Nat. Commun.* **2015**, 6, 7615.
- 443 3. Cózar, A.; Echevarría, F.; González-Gordillo, J.I.; Irigoien, X.; Úbeda, B.; Hernández-León, S.;
444 Palma, A.T.; Navarro, S.; García-de-Lomas, J.; Ruiz, A.; Fernández-de-Puelles, M.L.; Duarte, C.M.
445 Plastic debris in the open ocean. *PNAS* **2014**, 111, 10239-10244.
- 446 4. CAS, Database Counter, 2018, <http://www.cas.org/content/counter> (accessed 10.10.18).
- 447 5. Tornero, V; Hanke, G. Potential chemical contaminants in the marine environment: An overview
448 of main contaminant lists. ISBN 978-92-79-77045-6, EUR 28925, **2017**.
- 449 6. Luek, J.L.; Dickhut, R.M.; Cochran, M.A.; Falconer, R.L.; Kylin, H. Persistent organic pollutants
450 in the Atlantic and southern oceans and oceanic atmosphere. *Sci. Total Environ.* **2017**, 583, 64-71.
- 451 7. Maldonado, C; Dachs, J.; Bayona, J.M. Trialkylamines and Coprostanol as Tracers of Urban
452 Pollution in Waters from Enclosed Seas: The Mediterranean and Black Sea. *Environ. Sci. Technol.*
453 **1999**, 33, 3290-3296.
- 454 8. Björlenius, B.; Ripszám, M.; Haglund, P.; Lindberg, R.H.; Tysklind, M.; Fick, J. Pharmaceutical
455 residues are widespread in Baltic Sea coastal and offshore waters - Screening for pharmaceuticals
456 and modelling of environmental concentrations of carbamazepine. *Sci. Total Environ.* **2018**, 633,
457 1496-1509.
- 458 9. Schwarzenbach, R.P.; Escher, B.I.; Fenner, K.; Hofstetter, T.B.; Johnson, C.A.; von Gunten, U.;
459 Wehrli, B. The challenge of micropollutants in aquatic systems. *Science* **2006**, 313, 1072-1077.

- 460 10. Hollender, J.; Schymanski, E.L.; Singer, H.; Ferguson, P.L. Non-target screening with high
461 resolution mass spectrometry in the environment: Ready to go?. *Environ. Sci. Technol.* **2017**, 51,
462 11505-11512.
- 463 11. Schollée, J.E.; Schymanski, E.L.; Avak, S.E.; Loos, M.; Hollender, J. Prioritizing Unknown
464 Transformation Products from Biologically-Treated Wastewater Using High-Resolution Mass
465 Spectrometry, Multivariate Statistics, and Metabolic Logic. *Anal. Chem.* **2015**, 87, 12121–12129.
- 466 12. Gago-Ferrero, P.; Schymanski, E.L.; Bletsou, A.A.; Aalizadeh, R.; Hollender, J.; Thomaidis, N.S.
467 Extended Suspect and Non-Target Strategies to Characterize Emerging Polar Organic Contaminants
468 in Raw Wastewater with LC-HRMS/MS. *Environ. Sci. Technol.* **2015**, 49, 12333-12341.
- 469 13. Hoelzer, K.; Summer, A.J.; Karatum, O.; Nelson, R.K.; Drollette, B.D.; O'Connor, M.P.;
470 D'Ambro, E.L.; Getzinger, G.J.; Lee Ferguson, P.; Reddy, C.M.; Elsner, M.; Plata, D.L. Indications
471 of Transformation Products from Hydraulic Fracturing Additives in Shale-Gas Wastewater. *Environ.*
472 *Sci. Technol.* **2016**, 50, 8036-8048.
- 473 14. Montes-Grajales, D.; Fennix-Agudelo, M.; Miranda-Castro, W. Occurrence of personal care
474 products as emerging chemicals of concern in water resources: A review. *Sci. Total Environ.* **2017**,
475 595, 601-604.
- 476 15. Lara-Martín, P.A.; Gómez-Parra, A.; González-Mazo, E. Sources, transport and reactivity of
477 anionic and non-ionic surfactants in several aquatic ecosystems in SW Spain: A comparative study.
478 *Environ. Poll.* **2008**, 156, 36-45.
- 479 16. Chiaia-Hernández, A.; Günthardt, B.F.; Frey, M.P.; Hollender, J. Unravelling Contaminants in
480 the Anthropocene Using Statistical Analysis of Liquid Chromatography–High-Resolution Mass
481 Spectrometry Nontarget Screening Data Recorded in Lake Sediments. *Environ. Sci. Technol.* **2017**,
482 51, 12547-12556.
- 483 17. Schymanski, E.L.; Jeon, J.; Gulde, R.; Fenner, K.; Ruff, M.; Singer, H.P.; Hollender, J. Identifying
484 Small Molecules via High Resolution Mass Spectrometry: Communicating Confidence. *Environ. Sci.*
485 *Technol.* **2014**, 48, 2097–2098.

- 486 18. Ruttkies, C.; Schymanski, E.L.; Wolf, S.; Hollender, J.; Neumann, S. MetFrag relaunched:
487 incorporating strategies beyond in silico fragmentation. *J. Cheminform.* **2016**, *8*, 3.
- 488 19. Corada-Fernández, C.; Lara-Martín, P.A.; Candela, L.; González-Mazo, E. Tracking sewage
489 derived contamination in riverine settings by analysis of synthetic surfactants. *J. Environ. Monitor.*
490 **2011**, *13*, 2010-2017.
- 491 20. Biel-Maeso, M.; Baena-Nogueras, R.M.; Corada-Fernández, C.; Lara-Martín, P.A. Occurrence,
492 distribution and environmental risk of pharmaceutically active compounds (PhACs) in coastal and
493 ocean waters from the Gulf of Cadiz (SW Spain). *Sci. Total Environ.* **2018**, *612*, 649-659.
- 494 21. Choi, P.M.; Tscharke, B.J.; Donner, E.; O'Brien, J.W.; Grant, S.C.; Kaserzon, S.L.; Mackie, R.;
495 O'Malley, E.; Crosbie, N.D.; Thomas, K.V.; Mueller, J.F. Wastewater-based epidemiology
496 biomarkers: Past, present and future. *TrAC* **2018**, *105*, 453-469.
- 497 22. Burgos, M.; Ortega, T.; Bohórquez, J.; Corzo, A.; Rabouille, C.; Forja, J.M. Seasonal variation
498 of early diagenesis and greenhouse gas production in coastal sediments of Cadiz Bay: Influence of
499 anthropogenic activities. *Estuar. Coast. Shelf. Sci.* **2018**, *200*, 99-115.
- 500 23. Jaffé, R. Fate of hydrophobic organic pollutants in the aquatic environment: A review. *Environ.*
501 *Pollut.* **1991**, *69*, 237-257.
- 502 24. González-Ortegón, E.; Amaral, V.; Sánchez-Leal, R.F.; Bellanco, M.J.; Jiménez, M.P.; Forja, J.;
503 Vilas, C.; Tovar-Sanchez, A. Sources and coastal distribution of dissolved organic matter in the Gulf
504 of Cadiz. *Sci. Total Environ.* **2018**, *630*, 1583-1595.
- 505 25. Lara-Martín, P.A.; González-Mazo, E.; Petrovic, M.; Barceló, D.; Brownawell, B.J. Occurrence,
506 distribution and partitioning of nonionic surfactants and pharmaceuticals in the urbanized Long Island
507 Sound Estuary (NY). *Mar. Pollut. Bull.* **2014**, *85*, 710-719.
- 508 26. Mezzelani, M.; Gorbi, S.; Regoli, F. Pharmaceuticals in the aquatic environments: Evidence of
509 emerged threat and future challenges for marine organisms. *Mar. Environ. Res.* **2018**, *140*, 41-60.

- 510 27. Moro, L.; Pezzotti, G.; Turemis, M.; Sanchis, J.; Farré, M.; Denaro, R.; Giacobbe, M.G.; Crisafi,
511 F.; Giardi, M.T. Fast pesticide pre-screening in marine environment using a green microalgae-based
512 optical bioassay. *Mar. Pollut. Bull.* **2018**, 129, 212-221.
- 513 28. Acmite Market Intelligence, 2016. Market Report: Global Surfactant Market, fourth ed., **2016**.
- 514 29. Kolpin, D.W.; Furlong, E.T.; Meyer, M.T.; Thurman, E.M.; Zaugg, S.D.; Barber, L.B.; Buxton,
515 H.T. Pharmaceuticals, hormones, and other organic wastewater contaminants in U.S. streams,
516 1999–2000: A national reconnaissance. *Environ. Sci. Technol.* **2002**, 36, 1202–11.
- 517 30. Jackson, M.; Eadsforth, C.; Schowanek, D.; Delfosse, T.; Riddle, A.; Budgen, N. Comprehensive
518 review of several surfactants in marine environments: Fate and ecotoxicity. *Environ. Toxicol. Chem.*
519 **2016**, 35, 1077-1086.
- 520 31. Perales, J.A.; Manzano, M.A.; Garrido, M.C.; Sales, D.; Quiroga, J.M. Biodegradation kinetics
521 of linear alkylbenzene sulphonates in sea water. *Biodegradation* **2007**, 18, 63-70.
- 522 32. Giger, W.; Schaffner, C.; Kohler, H.P. Benzotriazole and tolyltriazole as aquatic contaminants.
523 1. Input and occurrence in rivers and lakes. *Environ. Sci. Technol.* **2006**, 40, 7186-7192.
- 524 33. Capela, D.; Ratola, N.; Alves, A.; Homem, V. Volatile methylsiloxanes through wastewater
525 treatment plants – A review of levels and implications. *Environ. Int.* **2017**, 102, 9-29.
- 526 34. Seit, W.; Winzenbacher, R. A survey on trace organic chemicals in a German water protection
527 area and the proposal of relevant indicators for anthropogenic influences. *Environ. Monit. Assess*
528 **2017**, 189, 244.
- 529 35. Lange, F.T.; Scheurer, M.; Brauch, H.J. Artificial sweeteners--a recently recognized class of
530 emerging environmental contaminants: a review. *Anal. Bioanal. Chem.* **2012**, 403, 2503-2518.
- 531 36. Baena-Nogueras, R.M.; Traverso-Soto, J.M.; Biel-Maeso, M.; Villar-Navarro, E.; Lara-Martín,
532 P.A. Sources and trends of artificial sweeteners in coastal waters in the bay of Cadiz (NE Atlantic).
533 *Mar. Pollut. Bull.* **2018**, 13, 607-616.

- 534 37. Mead, R.N.; Morgan, J.B.; Avery, G.B.; Kieber, R.J.; Kirk, A.M.; Skrabal, S.A.; Willey, J.D.
535 Occurrence of the artificial sweetener sucralose in coastal and marine waters of the United States.
536 *Mar. Chem.* **2009**, 116, 13–17.
- 537 38. International Federation of Pharmaceutical Manufacturers, Associations (IFMPA). The
538 Pharmaceutical Industry and Global Health: Facts and Figures 2017. ISBN 978-2-940498-50-5, **2017**.
- 539 39. Bernhard, E.S.; Rosi, E.J.; Gessner, M.O. Synthetic chemicals as agents of global change. *Front.*
540 *Ecol. Environ.* **2017**, 15, 84-90.
- 541 40. Lara-Martín, P.A.; Renfro, A.A.; Cochran, J.K.; Brownawell, B.J. Geochronologies of
542 Pharmaceuticals in a Sewage-Impacted Estuarine Urban Setting (Jamaica Bay, New York). *Environ.*
543 *Sci. Technol.* **2015**, 49, 1202–1211.
- 544 41. Salimi, M.; Esrafil, A.; Gholami, M.; Jonidi Jafari, A.; Rezaei Kalantary, R.; Farzadkia, M.;
545 Kermani, M.; Sobhi, H.R. Contaminants of emerging concern: a review of new approach in AOP
546 technologies. *Environ. Monit. Assess.* **2017**, 189, 414.
- 547 42. Alygizakis, N.A.; Samanipour, S.; Hollender, J.; Ibáñez, M.; Kaserzon, S.; Kokkali, V.; van
548 Leerdam, J.A.; Mueller, J.F.; Pijnappels, M.; Reid, M.J.; Schymanski, E.L.; Slobodnik, J.; Thomaidis,
549 N.S.; Thomas, K.V. Exploring the Potential of a Global Emerging Contaminant Early Warning
550 Network through the Use of Retrospective Suspect Screening with High-Resolution Mass
551 Spectrometry. *Environ. Sci. Technol.* **2018**, 52, 5135-5144.
- 552 43. Hernández, F.; Ibáñez, M.; Portolés, T.; Cervera, M.I.; Sancho, J.V.; López, F.J. Advancing
553 towards universal screening for organic pollutants in waters. *J. Hazard. Mater.* 2015, 282, 86-95.

554
555
556
557
558
559
560

561 **Figure Captions**

562

563 **Fig. 1.** Map showing the location of the Gulf of Cadiz (SW Spain) and sampling sites: GD1-4 =
564 Guadalquivir River mouth transect, CC1-4 = Cadiz Coast transect, and TF1-5 = Trafalgar Cape
565 transect. The inset shows the Cadiz Bay and sampling sites: G1-10 = Guadalete Estuary transect,
566 R1-8 = Rio San Pedro transect, and P1-9 = Sancti Petri transect. The location of the main WWTP is
567 also depicted (Schlitzer, R. Ocean Data View, odv.awi.de, 2020).

568

569 **Fig. 2.** Workflow tree and nodes involved in the non-target screening (NTS) and identification of
570 sewage-derived compounds in coastal and oceanic samples. Maximum confidence identification level
571 provided by different nodes is also depicted: 1 (chemical standards), 2 (spectral libraries), 3 (in-silico
572 fragmentation algorithms), and 4 (online compound databases and suspect lists). A detailed
573 explanation of each node can be found in Supporting Information.

574

575 **Fig. 3.** Hierarchical cluster analysis (HCA) of potential sewage-derived compounds ($n = 11,256$)
576 along the Guadalete Estuary (G1-10) (A) and selected clusters: potential contaminants from Jerez
577 WWTP (61%) (B), naturally occurring organic matter (34%) (C), and other pollution sources (5%)
578 (D). Examples of compounds for each cluster are plotted in red.

579

580 **Fig. 4.** Box plot showing the variability (minimum, first quartile, median, third quartile, and
581 maximum) in the number of all organic compounds and potential sewage-derived contaminants
582 detected in sampling stations at the Gulf of Cadiz (GD1-4 = Guadalquivir River mouth transect, CC1-
583 4 = Cadiz Coast transect, and TF1-5 = Trafalgar Cape transect) and the Cadiz Bay (P1-9 = Sancti
584 Petri transect, R1-8 = Rio San Pedro transect, and G1-10 = Guadalete Estuary transect). Normalized
585 signal intensity ranges of sewage-derived contaminants are also plotted.

586

587 **Fig. 5.** Characterization of the sewage-derived contaminants (n = 537) detected in oceanic waters
588 from the Gulf of Cadiz: identification confidence levels (1-4, sector graph) and elemental composition
589 (bar graph) (A), and percentage of these chemicals found in online databases and suspect lists (sector
590 graph) and known uses and applications (bar graph) (B). For those compounds included in the red
591 N/A sector and bar in Fig. 5B, information was not found on their identity or uses/applications,
592 respectively.

593

594 **Fig. 6.** Maximum normalized signal intensities of the top 10 sewage-derived contaminants in coastal
595 (Cadiz Bay) and oceanic (Gulf of Cadiz) waters sorted by use/application class. Confidence
596 identification levels (1 = green, 2 = blue, 3 = orange) are presented on the right of the compound
597 names.




Fig. 1





Fig. 2

**Fig. 3**




Fig. 4




Fig. 5




Fig. 6