

1 Note

2

3 **Black queen cell virus and drifting of honey bee workers**

4 (*Apis mellifera*)

5

6

7 **Gina Retschnig^{a,b,*}, Liv A. Kellermann^{a,b,c}, Marion M. Mehmann^{a,b}, Orlando**
8 **Yañez^{a,b}, Pius Winiger^{a,b}, Geoffrey R. Williams^{a,b,d}, Peter Neumann^{a,b}**

9

10

11 ^a Institute of Bee Health, Vetsuisse Faculty, University of Bern,

12 Schwarzenburgstrasse 161, 3003 Bern, Switzerland

13 ^b Agroscope, Swiss Bee Research Centre, Schwarzenburgstrasse 161, 3003 Bern,

14 Switzerland

15 ^c Department of Agronomy, Bern University of Applied Sciences HAFL, Länggasse

16 85, 3052 Zollikofen, Switzerland

17 ^d Department of Entomology & Plant Pathology, Auburn University, 301 Funchess

18 Hall, Auburn, AL 36849, USA

19

20

21 *Correspondence: E-mail: gina.retschnig@vetsuisse.unibe.ch,

22 Tel.: +41 (0)31 631 57 68

23 ORCID ID: 0000-0001-6249-9853

24 **Black queen cell virus (BQCV) / *Apis mellifera* / drifting / Virus**

25

26

27

28

29

30 **Summary**

31 Social insects may accidentally drift into foreign nests due to orientation errors. Even
32 though pathogens have been reported to promote drifting, no data currently exist about
33 the potential impact of titers of the widespread black queen cell virus (BQCV) on the
34 orientation abilities of honey bee workers, *Apis mellifera*. Here, we investigated titers
35 of BQCV in naturally infected drifted and non-drifted workers. The data show
36 significantly higher virus titers in the drifted workers (Wilcoxon rank sum test, $P <$
37 0.01). Our results suggest that high BQCV loads may compromise honey bee
38 orientation, possibly by affecting learning performance similar to other viruses. If
39 future work demonstrates that the correlation found here represents a causal
40 relationship between higher viral titers and drifting, this will be the first identification
41 of clinical symptoms of BQCV in adult honey bee hosts.

42

43 Honey bees may not return to their home colonies due to orientation errors, which
44 is known as drifting (Rauschmayer 1928; Neumann et al. 2000). Infections with
45 pathogens might enhance chances of orientation errors, thereby promoting both drifting
46 and pathogen transmission between colonies (Fries and Camazine 2001). While honey
47 bee viruses are common (Chen and Siede 2007), no significant association between the
48 presence of ten common viruses and drifting of honey bee workers has been reported
49 yet (Forfert et al. 2015). However, there may be an important difference with respect
50 to phenotype between the sheer detection of a pathogen vs. the actual pathogen load,
51 i.e. only higher virus loads may affect honey bee behavior as in case of learning and
52 deformed wing virus (Iqbal and Mueller 2006). Here, we compared for the first time
53 natural infection levels with the nearly ubiquitous black queen cell virus (BQCV)
54 between drifted and non-drifted honey bee workers.

55 For this study, freshly emerged honey bee workers from four local colonies were
56 individually marked and randomly introduced into three queenright 2-frame
57 observation hives (N=200 each) that were installed in one row at intervals of ~1.5 m;
58 each was equipped with an optically distinguishable hive entrance. All labelled workers
59 that were present in the hives after 14 days were recaptured (N=178) and assigned to
60 drifting status based on their markings (drifted N=13; non-drifted N=165). Then,
61 BQCV levels were quantified using standard qPCR (Gauthier et al. 2007). Briefly, total
62 RNA of individually homogenized workers was extracted using the Nucleospin RNA
63 II kit (Macherey-Nagel, Düren, Germany) and reverse transcription was conducted with
64 the Thermoscript™ RT system (Invitrogen, Carlsbad, USA) according to
65 manufacturer guidelines. qPCR was then performed using a KAPA SYBR FAST
66 Universal Mastermix kit (KAPA Biosystems, Wilmington, USA) in an Eco™ Real

67 Time PCR System (Illumina, San Diego, USA) (de Miranda et al. 2013) with specific
68 primers for BQCV published in Gauthier et al. 2007.


69 The analyses showed a significant effect of the observation hive on drifting
70 ($\chi^2 = 10.52, P < 0.01$): 15.39 %, 4.62 % and 0 % of the bees drifted from hive 1, 2 and
71 3, respectively. Despite the limited sample size of drifting workers (N=13), they
72 showed significantly higher BQCV titers compared to non-drifting ones (N = 165,
73 Wilcoxon rank sum test, $P < 0.01$) (Figure 1).

74 In a previous study, no significant differences in the prevalence of BQCV and nine
75 other common viruses were found between drifted and non-drifted honey bee workers
76 (Forfert et al. 2015). However, the significantly higher viral titers of drifted workers in
77 our study suggest that only high BQCV loads may compromise orientation abilities of
78 their honey bee hosts, very similar to effects on learning performance by deformed
79 wing virus (Iqbal and Mueller 2006). If that holds true, this compromised orientation
80 ability would constitute the first reported clinical symptom of BQCV in adult honey
81 bees (Chen and Siede 2007).

82

83 **Acknowledgments**

84 We wish to thank Peter Retschnig and Kaspar Roth for technical assistance. Financial
85 support was granted to PN by the EU 7th Framework Program BEE DOC (244956 CP-
86 FP), the Vinetum Foundation, the Swiss National Science Foundation and the Ricola
87 Foundation Nature and Culture.



89

90 **Figure I.** Drifting of adult worker honey bees and BQCV copies/bee. Drifted workers
91 (N = 13) showed significantly higher BQCV titers compared to non-drifted ones
92 (N = 165; Wilcoxon rank sum test, $P < 0.01=*$). Boxplots show the inter-quartile range
93 (box), median (black line within box), data range (vertical lines) and outliers (grey
94 dots). NCSS 10 was used for statistical analyses and the Figure.

95

96 **References**
97
98 Chen Y, Siede R (2007) Honey bee viruses. *Adv Vir Res* 70:33-80.
99 [https://doi.org/10.1016/S0065-3527\(07\)70002-7](https://doi.org/10.1016/S0065-3527(07)70002-7)
100
101 de Miranda J, Bailey L, Ball BV, Blanchard P, Budge GE, Chejanovsky N, Chen Y-
102 P, Gauthier L, Genersch E, de Graaf DC, Ribière M, Ryabov E, De Smet L, van
103 der Steen JJM (2013) Standard methods for virus research in *Apis mellifera*. *J Apic*
104 *Res* 52(4). <https://doi.org/10.3896/IBRA.1.52.4.22>
105
106 Forfert N, Natsopoulou ME, Frey E, Rosenkranz P, Paxton RJ, Moritz RFA (2015)
107 Parasites and pathogens of the honeybee (*Apis mellifera*) and their influence on
108 inter-colonial transmission. *PLoS ONE* 10(10).
109 <https://doi.org/10.1371/journal.pone.0140337>
110
111 Fries I, Camazine S (2001) Implications of horizontal and vertical pathogen
112 transmission for honey bee epidemiology. *Apidologie* 32:199-214.
113 <https://doi.org/10.1051/apido:2001122>
114
115 Gauthier L, Tentcheva D, Tournaire M, Dainat B, Cousserans F, Colin ME, Bergoin M
116 (2007) Viral load estimation in asymptomatic honey bee colonies using the
117 quantitative RT-PCR technique. *Apidologie* 38:426–435.
118 <https://doi.org/10.1051/apido:2007026>
119
120 Iqbal J, Mueller U (2007) Virus infection causes specific learning deficits in honeybee
121 foragers. *Proc R Soc L B* 274:1517-1521. <https://doi.org/10.1098/rspb.2007.0022>
122
123 Neumann P, Moritz R, Mautz D (2000) Colony evaluation is not affected by drifting of
124 drone and worker honeybees (*Apis mellifera* L.) at a performance testing apiary.
125 *Apidologie* 31:67-79. <https://doi.org/10.1051/apido:2000107>
126
127 Rauschmayer F (1928) Das Verfliegen der Bienen und die optische Orientierung am
128 Bienenstand. *Arch. Bienenk.* 9:249-322
129