



Maternal, Fetal, and Placental Selectins in Women With Pre-eclampsia; Association With the Renin-Angiotensin-System

Hiten D. Mistry^{1*}, Melissa V. Hott Ogalde¹, Fiona Broughton Pipkin¹, Geneviève Escher^{2,3} and Lesia O. Kurlak¹

¹ Department of Obstetrics and Gynaecology, University of Nottingham, Nottingham, United Kingdom, ² Department of Nephrology and Hypertension, University of Bern, Bern, Switzerland, ³ Department of Biomedical Research, University of Bern, Bern, Switzerland

OPEN ACCESS

Edited by:

Kirsty Pringle,
The University of Newcastle, Australia

Reviewed by:

Süleyman Cansun Demir,
Çukurova University, Turkey
Tanja Premru-Sršen,
University Medical Centre
Ljubljana, Slovenia

*Correspondence:

Hiten D. Mistry
hiten.mistry@nottingham.ac.uk

Specialty section:

This article was submitted to
Obstetrics and Gynecology,
a section of the journal
Frontiers in Medicine

Received: 28 October 2019

Accepted: 15 May 2020

Published: 12 June 2020

Citation:

Mistry HD, Ogalde MVH,
Broughton Pipkin F, Escher G and
Kurlak LO (2020) Maternal, Fetal, and
Placental Selectins in Women With
Pre-eclampsia; Association With the
Renin-Angiotensin-System.
Front. Med. 7:270.
doi: 10.3389/fmed.2020.00270

Selectins [endothelial (E), platelet (P), and leucocytes (L)] are a class of cell adhesion molecules, stimulated in response to inflammation. Pre-eclampsia is characterized by inflammation, and angiotensin II is pro-inflammatory. We hypothesized that circulating maternal and fetal concentrations and placental expression of selectins would be increased in women with pre-eclampsia and would be associated with the angiotensin receptors (AT1R and AT2R). Maternal and fetal blood and placental tissue was collected at delivery from White European normotensive controls ($n = 17$) and women with pre-eclampsia ($n = 17$). Soluble (s) E-, P- and L-selectin protein concentrations were measured by ELISA and placental protein expression was examined by immunohistochemistry. Maternal sE-selectin concentrations were increased in pre-eclampsia ($P < 0.001$); conversely fetal sE- and sP-selectin levels were lower in pre-eclampsia ($P < 0.05$ for both). Staining was mainly localized to the syncytiotrophoblast for all selectins. E-selectin expression was increased, while P-selectin was decreased in placental from pre-eclampsia ($P < 0.05$ for both); no differences were observed for L-selectin expression. Both E- and L-selectin were positively correlated ($P < 0.008$; $P < 0.02$) with AT2R placental expression, whilst P-selectin was negatively associated with AT1R ($P < 0.005$), all only in the pre-eclampsia group. This novel study reports maternal, fetal and placental expression of selectins in pre-eclampsia. The increased E-selectins reflect the endothelial dysfunction, characteristic of pre-eclampsia. In contrast, the reduced P-selectins and the positive association of placental AT2Rs with both E- and L-selectin in pre-eclampsia could be a protective mechanism to limit the endothelial dysfunction.

Keywords: selectins, angiotensin receptors, endothelial dysfunction, pre-eclampsia, inflammation

INTRODUCTION

Pre-eclampsia is a pregnancy-specific syndrome characterized by *de novo* hypertension and proteinuria after 20 weeks gestation (1). It affects between 2 and 8% of all pregnancies and is a major cause of both maternal and perinatal morbidity and mortality worldwide (2, 3). The consequences of pre-eclampsia are not restricted to pregnancy, with both mothers and their

offspring being at increased risk of cardiovascular, metabolic and renal disease in later life (4, 5). The etiology of pre-eclampsia remains unknown, but endothelial dysfunction due to abnormal placentation, oxidative stress, extensive leucocyte activation and inappropriate activation of the renin-angiotensin system (RAS) appear to be key contributors (6–10). The endothelium is the main regulator of vascular homeostasis. In addition to contributing to vasodilation, the intact endothelium also has anti-aggregatory, anti-inflammatory and anti-proliferative properties (11). Hypertension is associated with endothelial dysfunction, which in turn results from a dysregulation of locally-acting mediators, such as the vasodilators nitric oxide, and prostacyclin and the vasoconstrictor endothelin, and angiotensin II (Ang II)

The RAS is an important regulator of blood pressure and is one of the first hormonal systems to recognize pregnancy (10). Moreover, the RAS also plays a role in various pathological situations involving regulation of cell proliferation, inflammation, oxidative stress, and angiogenesis (12, 13). We have previously demonstrated a disrupted placental RAS in pre-eclampsia with the angiotensin II type 1 receptor (AT1R), activation of which results in vasoconstriction, being more highly expressed in placenta from women with pre-eclampsia (14).

Selectins are type 1 transmembrane proteins and are involved in leucocyte trafficking, mediating the initial tethering and rolling interactions between blood and lymphatic organs via the endothelium (15). Selectins were so named because of their ability to selectively bind carbohydrate moieties via their ligand P-selectin glycoprotein ligand-1 (PSGL-1) (15). There are three types of selectins, endothelial E-, platelets P- and leucocyte L-selectins; CD62E, CD62P and CD62L, respectively. All 3 selectins are composed of common extracellular domains and are distinguished from one another by their variable number of consensus repeats and very different cytoplasmic tails (16). E-selectin is normally absent from resting endothelium, but is transcriptionally induced in venules of multiple organs by inflammatory mediators, such as tumor necrosis factor (TNF) and IL-1 (17). P-selectin is constitutively synthesized in both platelets and endothelial cells and stored in Weibel-Palade bodies. They are rapidly translocated to the surface upon stimulation with inflammatory mediators, such as thrombin, histamine, and platelet-activating factor (17).

The shortest selectin molecule, present in leukocytes, is L-selectin, which, in the microvasculature, facilitates neutrophil rolling and migration during inflammatory responses (18). Elevated sE-selectin levels may be indicative of endothelial cell activation and damage. In addition to the conventional roles described above, the selectins play an essential role in human implantation; L-selectin and its ligands participate in the adhesion of the blastocyst to the endometrium at the maternal-fetal interface (19). Moreover, P- and E-selectins are also involved in immune recognition between maternal decidua and the embedded embryo, as well as trophoblast migration within decidual spiral arterioles (19). Reduced expression of all selectins has been reported in both decidual cells in uterine decidual stroma and in the placenta (cytotrophoblasts and syncytiotrophoblasts) of early miscarriages (6–11 weeks) compared to controls (terminations for social reasons) (20).

Concentrations of both soluble (s) E- and P-selectins have been reported to be increased in patients with chronic hypertension with the most severe endothelial dysfunction (21). Previous studies have reported elevated maternal plasma sE-selectin concentrations in women with pre-eclampsia, both at the time of delivery (22–26) and also prior to clinical diagnosis (22). However, the available data concerning both sP- and sL-selectins (22, 23, 26, 27) are conflicting.

Associations between the RAS and the selectins have previously been reported: Ang II is a potent stimulus to angiogenesis, both directly and by supplying angiogenic factors by platelet aggregation (28) and inducing the expression of P-selectin on platelets, while lessening its expression in endothelial cells (29). In addition, platelets express AT1R on their surface, and AT1R antagonists reduce P-selectin expression in platelets (30) and inhibit P-selectin-mediated platelet adhesion to the microvessels (31). Ang II also acts directly on naive T cells inducing upregulation of L-selectin via the AT1R (32). Finally, Ang II treatment significantly increased E-selectin mRNA expression in vascular endothelial cells collected from lung tissues in a mouse model of pulmonary metastasis (33).

We hypothesized that both soluble and placental expression of selectins would be increased in women with pre-eclampsia reflecting the endothelial dysfunction and inflammation characteristic of the syndrome. These selectins would also be associated with RAS receptors, contributing to the mechanistic changes associated with pre-eclampsia. This study therefore aimed to measure both soluble and placental expression of all selectins in matched maternal, fetal, and placental samples from normotensive control women and women with pre-eclampsia.

MATERIALS AND METHODS

Subjects and Sample Collection

The study population consisted of 17 normotensive control women and 17 women with pre-eclampsia, collected over a period of 18 months (Table 1). Nottingham University Hospital Ethics Committee approved the investigations and written informed consent to participation was obtained from each woman. Cases were defined on admission with a clinical diagnosis of pre-eclampsia, defined as a systolic blood pressure of ≥ 140 mm Hg and diastolic pressure (Korotkoff V) of ≥ 90 mm Hg on 2 occasions after 20 weeks' gestation in a previously normotensive women and proteinuria > 300 mg/L, 500 mg/day or 2+ on dipstick analysis of midstream urine (MSU) if 24-h collection result was not available (1). Although samples were collected under the criteria defined originally (1), they still fit the recently updated definition (34). For subgroup analysis, the pre-eclampsia group was further split by early- (diagnosis ≤ 34 weeks) and late- (diagnosis > 34 weeks) onset pre-eclampsia (35). Medical and obstetric histories, including delivery data, were obtained for each woman. The birthweight centile for each baby was computed, correcting for gestation age, sex, maternal parity, and body mass index (BMI) (36). Venous blood samples were collected before delivery and umbilical venous (fetal) blood were collected immediately after and processed as previously described (6); samples were stored in aliquots at -80°C until

TABLE 1 | Clinical and obstetric data of subject groups*.

Parameter	NC (n = 17)	PE (n = 17)
Age (yrs)	28.2 ± 7.2	31 ± 6.5
Booking body mass index (Kg/m ²)	26.4 ± 5.5	24.8 ± 3.5
Smoking status		
Non-smoker	9 (53)	11 (65)
Smoker	8 (47)	6 (35)
Parity		
Nulliparous	11 (65)	10 (59)
Multiparous	6 (35)	7 (41)
Max. systolic blood pressure (mmHg)	114 ± 3.8	155 ± 3.8 [†]
Max. diastolic blood pressure (mmHg)	75 ± 2.1	97 ± 5.0 [†]
Proteinuria (g/L) Median [min, max]	-	1.0 [0.3, 9.4]
Gestational age at delivery (Wks)	40.0 ± 1.0	37.7 ± 1.8
Mean birthweight (g)	3439 ± 498	2993 ± 735 [†]
Corrected birthweight centile	27.9 [17.5, 66.8]	35.1 [8.6, 76.3]
Caesarean section	4 (24)	13 (76)
Early-onset PE	-	6 (35)

*Data represented as means ± SD or median [IQR] as appropriate, except for smoking status, parity and Caesarean sections and early-onset PE, which are shown as number (percentage). NC, normotensive control; PE, pre-eclampsia; BMI, body mass index. [†] $P < 0.05$ between normotensive controls and women with pre-eclampsia.

analysis. In addition, placental tissue samples were collected from a standardized location midway between cord insertion and placental border and processed for immunohistochemistry as previously described (6).

Soluble Selectin Assays

Soluble (s) E-, P-, and L-selectin were measured in maternal and fetal EDTA plasma using ELISAs (KA0116, KA0548, and KA0117, respectively; Abnova, Taiwan) following the manufacturers' instructions. Plasma samples were diluted 1:100, run in duplicate and blinded to outcome group. Intra- and inter-assay variations were: sE-selectin: 5.4 and 6%; sP-selectin: 5.5 and 6.1%; sL-selectin: 3.7 and 4.2%.

Placental Selectin Protein Expression/Localization

Immunohistochemical analysis was performed as previously described (37), using antibodies to E-selectin (mouse monoclonal, BBA16; 5 µg/mL; R&D Systems), P-selectin (mouse monoclonal, BBA30; 10 µg/mL; R&D Systems) and L-selectin (rabbit polyclonal, GTX59778; 0.1 µg/mL; GeneTex Inc.). All slides were assessed by the same observer, blinded to pregnancy outcome. Quantification was performed as described previously (12), using the Positive Pixel Algorithm of Aperio ImageScope software; a visual check was also performed to establish localization of staining. The AT1R and AT2R protein expression analysis was performed as previously reported (14).

Statistical Analysis

All tests were performed using SPSS version 24. Summary data are presented as means ± standard deviation (SD) or median and interquartile range (IQR) as appropriate. The Kolmogorov-Smirnov test indicated the experimental data were not normally-distributed. The Kruskal-Wallis test followed by Mann-Whitney *U*-test was used for multiple group analysis; the Wilcoxon paired tests were used for maternal and fetal selectin concentrations. Potential associations between continuous data were tested using Spearman's Rho correlation tests. The null hypothesis was rejected where $P < 0.05$.

RESULTS

Subjects

Demographics and clinical characteristics of the participants are summarized in **Table 1**. By definition, blood pressures were significantly increased ($P < 0.05$) and significant proteinuria was present in the pre-eclampsia group. Birthweights were also lower in the women who suffered from pre-eclampsia. The groups were matched for maternal age, BMI, smoking status, parity, and gestational age at delivery.

Soluble Selectin Concentrations in Maternal and Fetal Plasma

Concentrations of all soluble selectins are summarised in **Figure 1**. Maternal sE-selectin was increased in pre-eclampsia compared to controls ($P < 0.05$; **Figure 1A**). No significant differences were observed for sP- or sL-selectins ($P > 0.05$ for both; **Figures 1B,C**). In contrast, in the fetus, both sE- and sP-selectin concentrations were lower in the pre-eclampsia group ($P < 0.05$ for both; **Figures 1A,B**), but did not differ for sL-selectin ($P > 0.05$; **Figure 1C**).


When comparing paired maternal-fetal samples, both sE- and sP-selectin concentrations were increased in fetal compared to maternal plasma only in the normotensive group ($P < 0.05$ for both; **Figures 1A,B**); fetal plasma from women who had pre-eclampsia remaining similar to their paired maternal plasma ($P > 0.05$). Conversely, fetal sL-selectin concentrations were reduced compared to their paired maternal samples in both groups ($P < 0.05$; **Figure 1C**).

When sub-grouped by early-/late-onset pre-eclampsia, only maternal sE-selectin concentrations were raised in plasma from early-onset ($n = 6$) pre-eclampsia ($P = 0.003$), compared to both late-onset ($n = 11$) pre-eclampsia ($P = 0.01$) and normotensive controls ($P = 0.002$; **Figure 1D**). No differences were seen for either sP- or sL-selectin concentrations ($P > 0.05$ for both; data not shown).

Ratio between maternal and fetal sE-, sP- and sL-selectin was calculated (**Figure 2**). Significantly increased ratios were observed for both sE- ($P = 0.001$) and sP- ($P = 0.02$) selectin in pre-eclampsia, compared to normotensive controls. No differences were observed for sL-selectin ($P > 0.05$).

Placental Selectin Protein Expression

Placental expression of all 3 selectins was determined and localization was mainly in the syncytiotrophoblast layer,



stromal areas and some fetal vessel staining (Figure 3). Placental E-selectin expression was increased in pre-eclampsia (median [IQR], positivity; normotensive controls: 0.56 [0.51, 0.65]; pre-eclampsia: 0.71 [0.60, 0.80]; $P = 0.008$; Figure 3A), whereas P-selectin was reduced in pre-eclampsia (normotensive controls 0.47 [0.34, 0.59]; pre-eclampsia 0.35 [0.22, 0.41]; $P = 0.008$; Figure 3B). E-selectin expression levels were raised ($P = 0.002$), while P-selectin expression was reduced in both the early- and late-onset pre-eclampsia groups, compared to normotensive control samples ($P < 0.05$; Figures 3D,E).

Although no differences were observed between controls and pre-eclamptic women for L-selectin ($P > 0.05$; Figure 3C), when sub-grouped by onset of pre-eclampsia, a significant difference again became apparent the expression being increased in early-onset pre-eclampsia group expression compared to both normotensive controls ($P = 0.008$) and late-onset pre-eclampsia ($P = 0.002$; Figure 3F).

Association of Placental Expression of Selectins With AT1R and AT2R



We compared placental expression of selectins with previously measured AT1R and AT2R placental protein expression. In the pre-eclampsia samples, but not those from normotensive women, placental expression of both E-selectin ($r = 0.55$; $P = 0.008$; Figure 4A) and L-selectin ($r = 0.61$; $P = 0.02$; Figure 4B) positively correlated with AT2R expression. In contrast, again only in the pre-eclampsia group, P-selectin displayed a negative association with AT1R expression ($r = 0.68$; $P = 0.005$; Figure 4C).

DISCUSSION

This is the first study to present data on all selectins in paired maternal/fetal plasma together with their placental expression in pregnancy and pre-eclampsia. It helps clarify the conflicting data found previously in relation to P- and L-selectin in pregnancy and pre-eclampsia.

The increased maternal sE-selectin in pre-eclampsia is in accordance with previous reports (22, 23, 25, 26). We have now shown that there is also increased placental E-selectin, reflecting the overall endothelial dysfunction associated with pre-eclampsia. Furthermore, the markedly increased concentrations of maternal sE-selectin, found only in the early-onset pre-eclampsia group (Figure 1D), presumably reflect the severity of the syndrome, as has been suggested by others (24). This is further supported by the finding that sE-selectin is increased prior to clinical diagnosis of pre-eclampsia (22, 38). Moreover, Chen et al., have shown that maternal sE-selectin was not associated with all cases of preterm birth, but specifically only when complicated by pre-eclampsia (39). The increased placental E-selectin expression we have shown corresponds with increased maternal sE-selectin and suggests that the placental dysfunction contributes to the endothelial damage in pre-eclampsia. Shaw et al., reported increased E-selectin mRNA expression in endothelial cell HUVECs and HECCs after incubation with maternal perfusates obtained from placenta of women with pre-eclampsia (40), which might amplify the underlying difference.

The interesting finding of reduced fetal sE-selectin in pre-eclampsia could either be a protective adaptation to limit the endothelial damage to the fetus, or reflects an underlying inability to synthesize E-selectin. Since E-selectin can be induced on fetal endothelial cells to the same extent as adult vessels by 32 weeks' gestation (17, 41) and the gestational age of the early-onset PE




group was nearer 34 weeks, this supports the view that the difference is attributable to the pre-eclampsia pathology and not an immaturity of development. sE-selectin also contributes to homing in the endothelial progenitor cells and promotes tube formation in microvasculature as a repair mechanism (42). Given that there is reduced tubule formation in HUVECS isolated from pre-eclamptic pregnancies (43), this further suggests that the functional capacity of fetal endothelial cells is impaired in pre-eclampsia.

Nevertheless, the positive correlation with placental AT2R expression only in the pre-eclamptic samples supports the hypothesis of a protective mechanism; AT2R forms part of the

“protective” arm of the RAS and is strongly upregulated following tissue damage such a vascular and neuronal injury (44). AT2R confers this role through its anti-inflammatory, anti-fibrotic and anti-apoptotic properties. Moreover, in lung tissues, which have many parallels with the placenta, Ang II treatment significantly increased E-selectin mRNA expression in vascular endothelial cells, further supporting the association (33).

We did not identify any differences in maternal sP-selectin between groups, which is in contrast to a previous study, which reported increased levels in pre-eclampsia (23, 26). A potential explanation for this difference lies in the later gestation age at sampling in our study when compared to the previous study. The



lack of differences in sP-selectins between groups could have been due to our not measuring platelet sP-selectin levels. Nevertheless, by using whole plasma, we were able to measure all 3 selectins.

As with sE-selectins, the lower fetal P-selectin concentrations in the pre-eclamptic group could be a protective mechanism to limited endothelial damage to the fetus. The reduced placental P-selectin expression in pre-eclampsia could also contribute to lower fetal sP-selectin. In addition, the decreased P-selectin could also be explained by the known reduction in platelet numbers in pre-eclampsia (34). Furthermore, the inverse relationship with AT1R, suggests this is independent of AT1R expressed on platelets. Ang II can increase P-selectin expression within the vascular wall (45) and thus increase atherosclerotic plaque progression. Therefore, the reduced placental P-selectin expression in pre-eclampsia could be a feedback mechanism to limit this damage and is supported by this negative correlation with AT1R.

Inspection of **Figures 2A,B** led us to suggest that the ratio between maternal and fetal sE- or sP-selectin might differ between normotensive and pre-eclamptic women, and in early and late-onset pre-eclampsia. As **Figure 3** shows, this is indeed the case, again emphasizing a possible protective mechanism.

The lack of differences observed between groups in sL-selectin in plasma may be because differences are only seen in specific cell types, for example the decreased levels in pre-eclampsia reported in neutrophils, T-cells, and monocytes (46). However, although not significant, there did appear to be a trend toward reduced maternal sL-selectin concentrations in the early-onset pre-eclamptic group. This is in line with previous studies where the majority of the pre-eclamptic samples measured were from early-onset pre-eclampsia (23, 47). To investigate this fully, future studies will examine whether any differences are observed when leukocytes and platelets are isolated and analyzed to confirm if the differences in this study are confirmed, or were masked by measuring in whole plasma. In contrast, there is increased placental expression in placenta from the early-onset pre-eclampsia group, which is not reflected in parallel increases in the maternal or fetal circulation. This again suggests a possible protective mechanism, further supported by the positive

association seen with placental AT2R expression in the pre-eclamptic group. It has been suggested that these lower levels result in increased leukocyte activation as seen in some chronic inflammatory states (e.g., chronic artery disease (48)). We can thus speculate that the increased placental L-selectin in early-onset pre-eclampsia could lead to increased leukocyte activation and therefore atherosclerosis in these placenta.

To conclude, this study is the first to report matched maternal, fetal and placental expression of all 3 selectins in pre-eclampsia. The increased E-selectins reflect the endothelial dysfunction, characteristic of pre-eclampsia. In contrast, the reduced P-selectins and the positive association with the placental AT2R with both E- and L-selectin in pre-eclampsia could be a protective mechanism to limit the endothelial dysfunction. Due to the roles of the selectins, this observational study further supports the importance of endothelial cells, platelets, and leukocytes in the pathogenesis of pre-eclampsia.

DATA AVAILABILITY STATEMENT

The datasets generated for this study are available on request to the corresponding author.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Nottingham University Hospital Ethics Committee. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

HM, LK, MO, and GE developed and performed measurement techniques, analyzed data, and wrote the manuscript. LK and HM collected samples and clinical information. FB directed the project, analyzed data, and assisted with writing the manuscript. All authors provided critical reviews of the manuscript.

ACKNOWLEDGMENTS

We thanks Heidi Jamin and Beatrice Rohrbach for running the ELISA assays and all the women who participated in the study.

REFERENCES

- Brown MA, Lindheimer MD, de Swiet M, Van Assche A, Moutquin JM. The classification and diagnosis of the hypertensive disorders of pregnancy: statement from the International Society for the Study of Hypertension in Pregnancy (ISSHP). *Hypertens Pregnancy*. (2001) 20:IX-XIV. doi: 10.3109/10641950109152635
- Stegers EA, von Dadelszen P, Duvekot JJ, Pijnenborg R. Pre-eclampsia. *Lancet*. (2010) 376:631–44. doi: 10.1016/S0140-6736(10)60279-6
- Duley L. The global impact of pre-eclampsia and eclampsia. *Semin Perinatol*. (2009) 33:130–7. doi: 10.1053/j.semperi.2009.02.010
- Bellamy L, Casas JP, Hingorani AD, Williams DJ. Pre-eclampsia and risk of cardiovascular disease and cancer in later life: systematic review and meta-analysis. *BMJ*. (2007) 335:974. doi: 10.1136/bmj.39335.385301.BE
- Mistry HD, Kurlak LO, Mansour YT, Zurkinden L, Mohaupt MG, Escher G. Increased maternal and fetal cholesterol efflux capacity and placental CYP27A1 expression in preeclampsia. *J Lipid Res*. (2017) 58:1186–95. doi: 10.1194/jlr.M071985
- Mistry HD, Wilson V, Ramsay MM, Symonds ME, Broughton Pipkin F. Reduced selenium concentrations and glutathione peroxidase activity in pre-eclamptic pregnancies. *Hypertension*. (2008) 52:881–8. doi: 10.1161/HYPERTENSIONAHA.108.116103
- Poston L. Endothelial dysfunction in pre-eclampsia. *Pharmacol Rep*. (2006) 58 Suppl:69–74.
- Poston L, Igosheva N, Mistry HD, Seed PT, Shennan AH, Rana S, et al. Role of oxidative stress and antioxidant supplementation in pregnancy disorders. *Am J Clin Nutr*. (2011) 94(6 Suppl):1980S–5S. doi: 10.3945/ajcn.110.001156
- Campello E, Spiezia L, Radu CM, Dhima S, Visentin S, Valle FD, et al. Circulating microparticles in umbilical cord blood in normal pregnancy and pregnancy with preeclampsia. *Thromb Res*. (2015) 136:427–31. doi: 10.1016/j.thromres.2015.05.029
- Broughton Pipkin F. The renin-angiotensin system in pre-eclampsia. In: Lyall F, Belfort M, editors. *Pre-Eclampsia. Etiology Clinical Practice*. Cambridge: Cambridge University Press (2007) 78–91.
- Ross R. Atherosclerosis—an inflammatory disease. *N Engl J Med*. (1999) 340:115–26. doi: 10.1056/NEJM199901143400207
- Williams PJ, Mistry HD, Innes BA, Bulmer JN, Broughton Pipkin F. Expression of AT1R, AT2R and AT4R and their roles in extravillous trophoblast invasion in the human. *Placenta*. (2010) 31:448–55. doi: 10.1016/j.placenta.2010.02.014
- Farag E, Maheshwari K, Morgan J, Sakr Esa WA, Doyle DJ. An update of the role of renin angiotensin in cardiovascular homeostasis. *Anesth Analg*. (2015) 120:275–92. doi: 10.1213/ANE.0000000000000528
- Mistry HD, Kurlak LO, Broughton Pipkin F. The placental renin-angiotensin system and oxidative stress in pre-eclampsia. *Placenta*. (2013) 34:182–6. doi: 10.1016/j.placenta.2012.11.027
- Zarbock A, Ley K, McEver RP, Hidalgo A. Leukocyte ligands for endothelial selectins: specialized glycoconjugates that mediate rolling and signaling under flow. *Blood*. (2011) 118:6743–51. doi: 10.1182/blood-2011-07-343566
- Kappelmayer J, Nagy B Jr. The interaction of selectins and PSGL-1 as a key component in thrombus formation and cancer progression. *Biomed Res Int*. (2017) 2017:6138145. doi: 10.1155/2017/6138145
- Auvinen K, Jalkanen S, Salmi M. Expression and function of endothelial selectins during human development. *Immunology*. (2014) 143:406–15. doi: 10.1111/imm.12318
- Walcheck B, Moore KL, McEver RP, Kishimoto TK. Neutrophil-neutrophil interactions under hydrodynamic shear stress involve L-selectin and PSGL-1. A mechanism that amplifies initial leukocyte accumulation of P-selectin *in vitro*. *J Clin Invest*. (1996) 98:1081–7. doi: 10.1172/JCI118888
- Feng Y, Ma X, Deng L, Yao B, Xiong Y, Wu Y, et al. Role of selectins and their ligands in human implantation stage. *Glycobiology*. (2017) 27:385–91. doi: 10.1093/glycob/cwx009
- Eskicioglu F, Lacin S, Ozbilgin K, Kose C. The role of selectins in the first trimester pregnancy loss. *Ginekol Pol*. (2014) 85:287–93. doi: 10.17772/gp/1725
- de la Sierra A, Larrousse M. Endothelial dysfunction is associated with increased levels of biomarkers in essential hypertension. *J Hum Hypertens*. (2010) 24:373–9. doi: 10.1038/jhh.2009.91
- Carty DM, Anderson LA, Freeman DJ, Welsh PI, Brennan JE, Dominiczak AF, et al. Early pregnancy soluble E-selectin concentrations and risk of preeclampsia. *J Hypertens*. (2012) 30:954–9. doi: 10.1097/HJH.0b013e328352573b
- Chaiworapongsa T, Romero R, Yoshimatsu J, Espinoza J, Kim YM, Park K, et al. Soluble adhesion molecule profile in normal pregnancy and pre-eclampsia. *J Matern Fetal Neonatal Med*. (2002) 12:19–27. doi: 10.1080/jmf.12.1.19.27
- Veas CJ, Aguilera VC, Munoz JJ, Gallardo VI, Miguel PL, Gonzalez MA, et al. Fetal endothelium dysfunction is associated with circulating maternal levels of sE-selectin, sVCAM1, and sFlt-1 during pre-eclampsia. *J Mat -Fetal Neon Med*. (2011) 24:1371–7. doi: 10.3109/14767058.2011.556204
- Krauss T, Azab H, Dietrich M, Augustin HG. Fetal plasma levels of circulating endothelial cell adhesion molecules in normal and preeclamptic pregnancies. *Eur J Obstet Gynecol Reprod Biol*. (1998) 78:41–5. doi: 10.1016/S0301-2115(98)00010-4
- Docheva N, Romero R, Chaemsaitong P, Tarca AL, Bhatti G, Pacora P, et al. The profiles of soluble adhesion molecules in the “great obstetrical syndromes”(). *J Matern Fetal Neonatal Med*. (2019) 32:2113–36. doi: 10.1080/14767058.2018.1427058
- Nasrollahi S, Hoseini Panah SM, Tavilani H, Tavasoli S, Naderan M, Shoar S. Antioxidant status and serum levels of selectins in pre-eclampsia. *J Obstet Gynaecol*. (2015) 35:16–8. doi: 10.3109/01443615.2014.935710
- Senchenkova EY, Russell J, Almeida-Paula LD, Harding JW, Granger DN. Angiotensin II-mediated microvascular thrombosis. *Hypertension*. (2010) 56:1089–95. doi: 10.1161/HYPERTENSIONAHA.110.158220
- Amano H, Ito Y, Ogawa F, Eshima K, Suzuki T, Oba K, et al. Angiotensin II type 1A receptor signaling facilitates tumor metastasis formation through P-selectin-mediated interaction of tumor cells with platelets and endothelial cells. *Am J Pathol*. (2013) 182:553–64. doi: 10.1016/j.ajpath.2012.10.026
- Alexandru N, Popov D, Georgescu A. Platelet dysfunction in vascular pathologies and how can it be treated. *Thromb Res*. (2012) 129:116–26. doi: 10.1016/j.thromres.2011.09.026
- Ishikawa M, Sekizuka E, Yamaguchi N, Nakadate H, Terao S, Granger DN, et al. Angiotensin II type 1 receptor signaling contributes to platelet-leukocyte-endothelial cell interactions in the cerebral microvasculature. *Am J Physiol Heart Circ Physiol*. (2007) 292:H2306–15. doi: 10.1152/ajpheart.00601.2006
- Silva-Filho JL, Souza MC, Henriques MG, Morrot A, Savino W, Caruso-Neves C, et al. Renin-angiotensin system contributes to naive T-cell migration *in vivo*. *Arch Biochem Biophys*. (2015) 573:1–13. doi: 10.1016/j.abb.2015.02.035
- Ishikane S, Hosoda H, Nojiri T, Tokudome T, Mizutani T, Miura K, et al. Angiotensin II promotes pulmonary metastasis of melanoma through the activation of adhesion molecules in vascular endothelial cells. *Biochem Pharmacol*. (2018) 154:136–47. doi: 10.1016/j.bcp.2018.04.012
- Brown MA, Magee LA, Kenny LC, Karumanchi SA, McCarthy FP, Saito S, et al. The hypertensive disorders of pregnancy: ISSHP classification, diagnosis and management recommendations for international practice. *Pregnancy Hypertens*. (2018) 13:291–310. doi: 10.1016/j.preghy.2018.05.004
- Tranquilli AL, Brown MA, Zeeman GG, Dekker G, Sibai BM. The definition of severe and early-onset preeclampsia. Statements from the International Society for the Study of Hypertension in Pregnancy (ISSHP). *Pregnancy Hypertens*. (2013) 3:44–7. doi: 10.1016/j.preghy.2012.11.001

36. Gardosi J, Chang A, Kalyan B, Sahota D, Symonds EM. Customised antenatal growth charts. *Lancet*. (1992) 339:283–7. doi: 10.1016/0140-6736(92)91342-6
37. Mistry HD, McCallum LA, Kurlak LO, Greenwood IA, Broughton Pipkin F, Tribe RM. Novel expression and regulation of voltage-dependent potassium channels in placentas from women with preeclampsia. *Hypertension*. (2011) 58:497–504. doi: 10.1161/HYPERTENSIONAHA.111.173740
38. Chavarria ME, Lara-Gonzalez L, Garcia-Paleta Y, Vital-Reyes VS, Reyes A. Adhesion molecules changes at 20 gestation weeks in pregnancies complicated by preeclampsia. *Eur J Obstet Gynecol Reprod Biol*. (2008) 137:157–64. doi: 10.1016/j.ejogrb.2007.06.014
39. Chen X, Scholl TO. Maternal biomarkers of endothelial dysfunction and preterm delivery. *PLoS ONE*. (2014) 9:e85716. doi: 10.1371/journal.pone.0085716
40. Shaw J, Tang Z, Schneider H, Salje K, Hansson SR, Guller S. Inflammatory processes are specifically enhanced in endothelial cells by placental-derived TNF-alpha: Implications in preeclampsia (PE). *Placenta*. (2016) 43:1–8. doi: 10.1016/j.placenta.2016.04.015
41. Nussbaum C, Gloning A, Pruenster M, Frommhold D, Bierschenk S, Genzel-Boroviczeny O, et al. Neutrophil and endothelial adhesive function during human fetal ontogeny. *J Leukoc Biol*. (2013) 93:175–84. doi: 10.1189/jlb.0912468
42. Oh IY, Yoon CH, Hur J, Kim JH, Kim TY, Lee CS, et al. Involvement of E-selectin in recruitment of endothelial progenitor cells and angiogenesis in ischemic muscle. *Blood*. (2007) 110:3891–9. doi: 10.1182/blood-2006-10-048991
43. Brodowski L, Burlakov J, Hass S, von Kaisenberg C, von Versen-Hoynck F. Impaired functional capacity of fetal endothelial cells in preeclampsia. *PLoS ONE*. (2017) 12:e0178340. doi: 10.1371/journal.pone.0178340
44. Namsolleck P, Recarti C, Foulquier S, Steckelings UM, Unger T. AT(2) receptor and tissue injury: therapeutic implications. *Curr Hypertens Rep*. (2014) 16:416. doi: 10.1007/s11906-013-0416-6
45. Tayeh MA, Scicli AG. Angiotensin II and bradykinin regulate the expression of P-selectin on the surface of endothelial cells in culture. *Proc Assoc Am Physicians*. (1998) 110:412–21.
46. Luppi P, Tse H, Lain KY, Markovic N, Piganelli JD, DeLoia JA. Preeclampsia activates circulating immune cells with engagement of the NF-kappaB pathway. *Am J Reprod Immunol*. (2006) 56:135–44. doi: 10.1111/j.1600-0897.2006.00386.x
47. Sacks GP, Studena K, Sargent K, Redman CW. Normal pregnancy and preeclampsia both produce inflammatory changes in peripheral blood leukocytes akin to those of sepsis. *Am J Obstet Gynecol*. (1998) 179:80–6. doi: 10.1016/S0002-9378(98)70254-6
48. Haught WH, Mansour M, Rothlein R, Kishimoto TK, Mainolfi EA, Hendricks JB, et al. Alterations in circulating intercellular adhesion molecule-1 and L-selectin: further evidence for chronic inflammation in ischemic heart disease. *Am Heart J*. (1996) 132(1 Pt 1):1–8. doi: 10.1016/S0002-8703(96)90383-X
49. Kurlak LO, Hott-Ogalde MV, Scaife PJ, Escher G, Mistry HD. Maternal and placental selectins in women with pre-eclampsia; association with the renin-angiotensin system. *Placenta*. (2019) 83:e105. doi: 10.1016/j.placenta.2019.06.332

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Mistry, Ogalde, Broughton Pipkin, Escher and Kurlak. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.