Early View

Research letter

Age-dependent response of the human nasal epithelium to rhinovirus infection

Jakob Usemann, Marco P Alves, Nicole Ritz, Philipp Latzin, Loretta Müller

Please cite this article as: Usemann J, Alves MP, Ritz N, *et al.* Age-dependent response of the human nasal epithelium to rhinovirus infection. *Eur Respir J* 2020; in press (<https://doi.org/10.1183/13993003.00877-2020>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Copyright ©ERS 2020

Age-dependent response of the human nasal epithelium to rhinovirus infection

Jakob Usemann^{1,2,3}, Marco P Alves^{4,5}, Nicole Ritz², Philipp Latzin^{1,2,#}, Loretta Müller^{1,2,#,*}

¹Paediatric Respiratory Medicine, Department of Paediatrics, Inselspital, Bern University Hospital, University of Bern, Switzerland

²University Children's Hospital Basel (UKBB), Switzerland

³Division of Respiratory Medicine, University Children's Hospital Zurich, Zurich, Switzerland

⁴Institute of Virology and Immunology, Bern, Switzerland

⁵Department of Infectious Diseases and Pathobiology, Vetsuisse Faculty, University of Bern, Bern, Switzerland

#These authors contributed equally to this project.

***Corresponding Author:** loretta.mueller@insel.ch; Paediatric Pulmonology, Inselspital Bern, Murtenstrasse 50, 3008 Bern, Switzerland

Take home message (255 of 256 character incl. spaces): After rhinovirus (RV) infection, RV viral loads are higher and immune factor expressions lower in nasal epithelial cells from children versus adults in vitro. This may explain age-dependent severity of RV infections and association with asthma development.

Keywords

Airways, children, adults, air-liquid interface, respiratory infection

To the editor:

Rhinovirus (RV)-induced wheezing in early childhood is a major risk factor for later asthma (1, 2) and RV-infections have been identified as a main reason for asthma exacerbations (3-5). While RV-infections can manifest with a severe clinical course, RVs can also colonize the respiratory tract asymptotically (6-8). Thus, additional factors seem to determine which children develop a symptomatic RV-infection and are at increased risk for asthma development, and which remain asymptomatic. For example, there is a *genetic susceptibility*, since RV-induced wheezing *and* specific genetic variants increase the risk for asthma (9). *Environmental factors*, such as farming and air pollution, have been shown to influence the risk for asthma development (10). Further, the age of RV-infection differently affects the clinical course, suggesting different time windows of susceptibility (1). For example, RV-induced wheezing at ages 2-3 years is associated with a higher risk for asthma development compared to wheezing during the first year of life (1), indicating age-dependent immune responses. However, underlying mechanisms responsible for age-dependent severity of RV-infections have not yet been studied comprehensively. Therefore, we investigated age-dependency of anti-RV response in primary epithelial cells (NECs), which are the main target for RV-infections and act as a switchboard to initiate and regulate immune responses (11, 12). We hypothesize that the initial response of NECs to RVs is weaker at younger ages. To test this, we investigated the age-dependency of viral loads and the early epithelial immune response to *in vitro* infection with RV-16 and RV-1b of re-differentiated NECs of children and adults.

NECs were obtained via nasal brushings (interdental brush, Top Caredent) from healthy adult volunteers and healthy children undergoing elective surgery (13) (study approved by the Ethics Committee Northwest- und Zentralschweiz, Switzerland, reference number 250/13, written informed consent obtained from all donors). After cultivation and re-differentiation using the PneumCult Expansion Plus and ALI media (Stemcell Technologies), NECs were infected with RV-16 or RV-1b (multiplicity of infection (MOI) 1 and 4; 1h, 37°C; polyinosinic: polycytidylic acid (pI:C) (10µg/ml) and phosphate-buffered saline with Mg/Ca were used as controls). 20h post-infection, we harvested basolateral supernatants and cell lysates (using TRizol).

mRNA was isolated (RNA Clean & Concentrator-5 w/ Zymo-Spin IC Columns, Zymo Research) and converted into cDNA (GoScript Reverse Transcription System, Promega). Real-time RT-PCR was performed using the GoTaq qPCR Master Mix system (Promega). Protein concentrations were measured in the basolateral media using a Milliplex MAP kit (Merck).

We subdivided the paediatric population into three groups: infancy (0-12 months, group 1, mean age (range) 8.0 (2.5-12.7) months), early ($\geq$ 13-36 months, group 2, 26.4 (13.3-35.5) months), and late childhood (>36-93 months, group 3, 62.2 (36.3-92.7) months). Comparison of outcomes between children (N=49, (mean age (range)) 3.4 (0.21-7.7) years) versus adults (N=12, 35.1 (24.9-65.6) years) was done using the Wilcoxon rank-sum test. A p-value <0.05 defined statistical significance, analysis was performed with Stata[®], release 15.

Re-differentiated, air-liquid-interface (ALI) exposed NECs of children showed higher viral loads after RV-1b and RV-16 infection than NECs of adults. This finding was more pronounced in cells infected with RV-16 compared to RV-1b and those infected

at higher MOI (Fig.A). In the paediatric population, levels of viral loads after RV-16 infection increased with increasing age, albeit not significantly (Fig.B), RV-1b infection showed a similar pattern (data not shown). In children and adults separately, we could show a robust, dose-dependent anti-viral response induced by RV-infection (increase of *ICAM-1*, *LDLR*, *RIG-1*, *MDA5*, *TLR3*, β -*Def2*, *IFN- β* , *IFN- λ* (only for RV-16), *CXCL10*, *IL-1 β* (only for RV-16), *CXCL8* (only for RV-16), *IL-6* mRNA and CXCL10, CXCL8 and IL-6, but not IFN- γ protein, data not shown), and observed differences of the anti-RV response between adults children. Interestingly, those differences were already present at baseline (in uninfected controls). We found significantly lower mRNA levels of *ICAM-1*, *LDLR*, β -*DEF2*, *CXCL10*, *IL-1 β* , *IL-6* and *IFN- λ* (only borderline significant) in children compared to adults, while mRNA levels of *MDA5* were significantly higher in children compared to adults. We did not find any differences between children and adult NECs in mRNA levels of *RIG1*, *TRL3*, *IFN- β* , *IFN- λ* (Fig.C). These differences persisted after infection (data not shown). Compared to children, protein concentrations of IFN- γ and IL-6 were higher in adults at baseline, but not after infection. We did not see differences in CXCL10 or CXCL8 protein concentrations (Fig.D). Within the paediatric age groups, expression levels and protein release did not differ significantly.

This is the first study to investigate RV-1b and RV-16 infection in an ALI cell culture model. We found lower viral loads in cells infected with RV-1b compared to RV-16. Cell entry of RV-16 occurs via the ICAM-1 cell surface receptor, while RV-1b enters via the LDLR receptor (14). We found that compared to the highly expressed ICAM-1 levels, LDLR expression levels were much lower, possibly explaining the reduced RV-1b mRNA levels found in this study. Expression levels of *ICAM-1* and *LDLR*

mRNA levels were higher in adults compared to children. Association between age and the expression of *ICAM-1* or *LDLR* has scarcely been studied. Within a study including 29 adults (age 42-85 years), there was no association with age and *ICAM-1* levels (15). The here reported age-dependent expression of *ICAM-1* and *LDLR* could be relevant for epidemiological and experimental studies addressing a large age-range.

The childhood origins of asthma (COAST) study investigated the timing and specific viral aetiology of wheezing illnesses during early childhood and the impact on asthma development within a high-risk population. These data suggest that the age of RV-infection has prognostic value for subsequent asthma risk. Children who had RV-induced wheezing during their first year of life had a 2.7-fold asthma risk at six years, while RV-induced wheezing during the second year was associated with a 6.5-fold asthma risk, and RV-induced wheezing during the third year was associated with a dramatic 31.7-fold asthma risk (1). Interestingly, we found lower viral load in infants compared to older children potentially reflecting the age-dependent associations reported previously (1). Our findings, along with those previously reported (1), are important for future intervention studies addressing RV-infections as age-dependent differences in anti-viral responses in epithelial cells.

Our study is the first to investigate immune factors following RV-infection using an ALI cell model within a large population across a broad age range. Previous studies investigated either RV-16 or RV-1b, while we investigated both virus types in parallel, allowing comparison of immune response to both RV types. The study is limited by the cross-sectional assessment of the immune response, precluding the analysis of individual immune development or course of infection. Other cell types and viruses should be studied to investigate not only the anti-viral response to RV represented by

the respiratory epithelium, but more comprehensively the innate and late immune response.

We report on age-dependent anti-viral response in NECs after infection with major and minor group RVs. Children had higher viral loads than adults, likely the result of lower levels of immune factors. These findings may explain age-dependent severity of RV-infections and age-dependent associations between RV-induced wheezing and asthma development and indicate that early immune function priming is relevant for later chronic airway disease.

Figure. (A&B) Viral loads of RV-1b and RV-16 in NECs 20 h post-infection measured by quantitative real-time RT-PCR. (A) Comparison of viral loads in NECs of children (n=48) and adults (n=12). (B) Comparison of viral loads after RV-16 infection in different paediatric age groups and adults. Group 1 n=10, group 2 n=14, group 3 n=25, group 4 adults n=12. (C) mRNA levels of immune factors at baseline (without RV infection). mRNA levels are presented as dCt values, normalized to the house-keeping gene phosphoglycerate kinase 1. (D) Baseline

protein concentrations assessed using multiplex bead-based immunoassays.

Data are shown as box and whisker plot with line at median, interquartile range (box) and range (line). * $p < 0.05$, tested with Wilcoxon test.

Acknowledgments

We would like to thank all volunteers for donating nasal cells, Prof. Thomas Erb, Chief Physician Anesthesiology, University Children's Hospital Basel, and his team for performing the nasal brushings, Selina K Steiner and Andrea Zelmer (both University Children's Hospital Basel) for their excellent lab work, and Karine Bell Hugentobler (University Children's Hospital Basel) for proofreading the manuscript.

Funding

This project was supported by the Fondation Botnar Switzerland, the "Freiwillige Akademische Gesellschaft (FAG) Basel", "Swiss Life Jubiläumsstiftung", and the Fondation Johanna-Dürmüller Bol.

Conflict of interest

The authors have no conflicts of interest to disclose. Dr. Latzin reports personal fees from Gilead, Novartis, Polyphor, Roche, Santhera, Schwabe, Vertex, Vifor, and Zambon, outside the submitted work. Dr. Usemann reports personal fees from Vertex outside the submitted work.

Additional Information

Data availability

The datasets generated during the current study are available from the corresponding author on reasonable request.

Author contribution: JU performed RT-PCR measurements, analysed the data and wrote the main manuscript text. MPA provided stock of RV-16 and RV1B and helped with data interpretation. NR supported the cytokine analysis and helped with data

interpretation. PL developed the project idea and helped with data interpretation. LM designed the study, performed experiments and analysis, prepared the figures and the main manuscript text. All authors reviewed and approved the manuscript.

References

1. Jackson DJ, Gangnon RE, Evans MD, Roberg KA, Anderson EL, Pappas TE, Printz MC, Lee WM, Shult PA, Reisdorf E, Carlson-Dakes KT, Salazar LP, DaSilva DF, Tisler CJ, Gern JE, Lemanske RF, Jr. Wheezing rhinovirus illnesses in early life predict asthma development in high-risk children. *Am J Respir Crit Care Med* 2008; 178: 667-672.
2. Turi KN, Shankar J, Anderson LJ, Rajan D, Gaston K, Gebretsadik T, Das SR, Stone C, Larkin EK, Rosas-Salazar C, Brunwasser SM, Moore ML, Peebles RS, Jr., Hartert TV. Infant Viral Respiratory Infection Nasal Immune-Response Patterns and Their Association with Subsequent Childhood Recurrent Wheeze. *Am J Respir Crit Care Med* 2018; 198: 1064-1073.
3. Johnston SL, Pattemore PK, Sanderson G, Smith S, Lampe F, Josephs L, Symington P, O'Toole S, Myint SH, Tyrrell DA, et al. Community study of role of viral infections in exacerbations of asthma in 9-11 year old children. *BMJ* 1995; 310: 1225-1229.
4. Busse WW, Lemanske RF, Jr., Gern JE. Role of viral respiratory infections in asthma and asthma exacerbations. *Lancet* 2010; 376: 826-834.
5. Altman MC, Gill MA, Whalen E, Babineau DC, Shao B, Liu AH, Jepson B, Gruchalla RS, O'Connor GT, Pongracic JA, Kercksmar CM, Khurana Hershey GK, Zoratti EM, Johnson CC, Teach SJ, Kattan M, Bacharier LB, Beigelman A, Sigelman SM, Presnell S, Gern JE, Gergen PJ, Wheatley LM, Togias A,

- Busse WW, Jackson DJ. Transcriptome networks identify mechanisms of viral and nonviral asthma exacerbations in children. *Nat Immunol* 2019; 20: 637-651.
6. Müller L, Mack I, Tapparel C, Kaiser L, Alves MP, Kieninger E, Frey U, Regamey N, Latzin P. Human Rhinovirus Types and Association with Respiratory Symptoms During the First Year of Life. *Pediatr Infect Dis J* 2015; 34: 907-909.
7. Mack I, Kieninger E, Cangiano G, Tapparel C, Kuehni C, Spycher B, Kaiser L, Frey U, Regamey N, Latzin P. Rhinovirus Infections and Associated Respiratory Morbidity in Infants: A Prospective Cohort Study. *Pediatr Infect Dis J* 2016.
8. Kusel MM, de Klerk NH, Holt PG, Keadze T, Johnston SL, Sly PD. Role of respiratory viruses in acute upper and lower respiratory tract illness in the first year of life: a birth cohort study. *Pediatr Infect Dis J* 2006; 25: 680-686.
9. Caliskan M, Bochkov YA, Kreiner-Moller E, Bonnelykke K, Stein MM, Du G, Bisgaard H, Jackson DJ, Gern JE, Lemanske RF, Jr., Nicolae DL, Ober C. Rhinovirus wheezing illness and genetic risk of childhood-onset asthma. *N Engl J Med* 2013; 368: 1398-1407.
10. Stein MM, Hrusch CL, Gozdz J, Igartua C, Pivniouk V, Murray SE, Ledford JG, Marques Dos Santos M, Anderson RL, Metwali N, Neilson JW, Maier RM, Gilbert JA, Holbreich M, Thorne PS, Martinez FD, von Mutius E, Vercelli D, Ober C, Sperling AI. Innate Immunity and Asthma Risk in Amish and Hutterite Farm Children. *The New England journal of medicine* 2016; 375: 411-421.
11. Swamy M, Jamora C, Havran W, Hayday A. Epithelial decision makers: in search of the 'epimmunome'. *Nat Immunol* 2010; 11: 656-665.

12. Müller L, Jaspers I. Epithelial cells, the "switchboard" of respiratory immune defense responses: Effects of air pollutants. *Swiss Medical Weekly* 2012.
13. Stokes AB, Kieninger E, Schogler A, Kopf BS, Casaulta C, Geiser T, Regamey N, Alves MP. Comparison of three different brushing techniques to isolate and culture primary nasal epithelial cells from human subjects. *Exp Lung Res* 2014; 40: 327-332.
14. Uncapher CR, DeWitt CM, Colonno RJ. The major and minor group receptor families contain all but one human rhinovirus serotype. *Virology* 1991; 180: 814-817.
15. Shukla SD, Mahmood MQ, Weston S, Latham R, Muller HK, Sohal SS, Walters EH. The main rhinovirus respiratory tract adhesion site (ICAM-1) is upregulated in smokers and patients with increased with age chronic airflow limitation (CAL). *Respir Res* 2017; 18: 6.