Identification of growth mechanisms in metamorphic garnet by high-resolution trace element mapping with LA-ICP-TOFMS

Daniela Rubatto¹, Marcel Burger², Pierre Lanari¹, Bodo Hattendorf², Gunnar Schwarz², Christoph Neff², Peter Keresztes Schmidt², Jörg Hermann¹, Alice Vho¹, Detlef Günther²

1. Institute of Geological Sciences, University of Bern, Bern, CH-3012, Switzerland
2. Laboratory of Inorganic Chemistry, Department of Chemistry and Applied Biosciences, ETH, Zurich, CH- 8093, Switzerland

ORCID ID

Rubatto: 0000-0002-7425-7904

Burger: 0000-0002-9904-2547

Lanari: 0000-0001-8303-0771

Hattendorf: 0000-0001-9991-2410

Schwarz: 0000-0003-4449-7672

Neff: 0000-0002-9369-9580

Keresztes Schmidt: 0000-0002-4519-8861

Hermann: 0000-0001-8360-3592

Vho: 0000-0003-0158-9104

Günther: 0000-0001-7867-4310

Acknowledgements

We thank Yannick Bussweiler and Andy Smye for providing constructive comments that assisted in improving the manuscript. The careful editorial handling of Othmar Müntener is acknowledged.

D. Rubatto acknowledges the financial support of the Swiss National Science Foundation (project 200021_166280 and 206021_170722). D. Günther acknowledges financial support by ETH Zurich for the development of the LA-ICP-TOFMS system.

Keywords: garnet, trace element maps, metamorphism, Alps, metapelite, eclogite, LA-ICP-TOFMS

ABSTRACT

Garnet is one of the most robust and ubiquitous minerals that record element zoning during crustal metamorphism. In addition to major elements, zoning in trace elements can provide a wealth of information to document the changing conditions of garnet growth and modification. However, mapping trace elements at low concentrations, over large areas and with high resolution has remained a major challenge.

We present a comprehensive investigation of the TE distribution in garnet from three Alpine samples that underwent a complex evolution at different metamorphic conditions. The TE distribution in garnet grains is mapped in 2D in thin section with a novel approach using laser ablation inductively coupled plasma time of flight mass spectrometry (LA-ICP-TOFMS) to achieve a lateral resolution of 5 μm and limits of detection for the heavy rare earth elements (REE) down to 0.2 $\mu\text{g/g}$. Comparison with major element zoning measured by electron probe micro-analysis (EPMA) and trace elements measured by conventional LA-ICPMS spot analysis testifies to the accuracy of the measurements.

Garnet in an amphibolite-facies metapelite from Campolungo, Central Alps that recorded metamorphism to 600°C preserves Y+REE trace element zoning that closely matches that of Ca. In this sample, there is no notable diffusive modification for trace elements. Y+REE zoning is dominated by Rayleigh fractionation in the core and by the sporadic breakdown of accessory phases producing annuli in the rim of the garnet.

A granulite-facies garnet from Malenco, Eastern Central Alps, formed during subsolidus heating, followed by peritectic melting reactions up to temperatures of 800–850°C. Major and trace element zoning are decoupled indicating diffusional resetting of major elements, whereas trace elements still largely document the growth history. Enrichment of trace elements in the garnet mantle may be related to the consumption of biotite (V, Cr) and the dissolution of zircon (Zr) and monazite (Y+REE) in the melt. Diffusion of Y+HREE at the core–mantle boundary occurred over a length scale of $\sim 200\text{ }\mu\text{m}$.

Garnet in an eclogite from the Sesia Zone, Western Alps ($P \sim 2$ GPa, $T \sim 600^\circ\text{C}$) displays pronounced fluid-related veinlets, visible in FeO, MgO and MnO, which cross-cut the primary growth zoning. Surprisingly, complex Y+REE and Cr zoning is not affected by the veinlets, indicating that they did not form by a crack-seal mechanism but are rather related to a selective replacement process.

The trace element maps provide a detailed insight into the growth and modification of garnet and thus allow assessment of equilibrium versus disequilibrium processes, and assist in determination of P-T conditions, garnet dating, diffusion modelling as well as documenting fluid-induced modifications.

Introduction

Geological processes in the solid Earth are recorded in the structure and composition of minerals. Variations in composition within a mineral (zoning) reflect changes in the conditions of growth (pressure (P), temperature (T), reactive bulk composition (X), fluids, etc.), which in turn can be traced back to physical and chemical processes in the growth environment. Mineral zoning can be observed in major and trace elements (TE, mass fractions below $1000 \mu\text{g/g}$), and while the first is broadly understood and modelled to reconstruct P-T-X paths, the zoning in TE in a number of minerals is still underexplored and poorly comprehended.

Many TE have the advantage over major elements of i) being more robust to resetting by diffusional re-equilibration at high temperature and thus better preserving growth zoning (Hermann and Rubatto 2003), and ii) exhibiting large variations in concentrations during mineral formation and subsequent growth. The study of TE in minerals has allowed, for example, to reconstruct melt compositions (van Westrenen and Draper 2007), determine time scales of magmatic and metamorphic processes (e.g. Chakraborty 2006), calculate the temperature of mineral growth (e.g. Ferry and Watson 2007), recognize formation and breakdown of coexisting phases during mineral growth (Spear and Kohn 1996; Yang and Rivers 2002), and to establish mineral equilibrium for P-T-time determination (e.g. Rubatto 2002).

Garnet is one of the most robust minerals recording major and TE zoning. It is unsurpassed in its utility to reconstruct metamorphic P-T paths (Spear 1993; Caddick et al. 2010) on the basis of major element zoning. The investigation of TE in garnet is relatively mature (Hickmott et al. 1987; Spear and Kohn 1996; Otamendi et al. 2002; Hermann and Rubatto 2003; Skora et al. 2006; Konrad-Schmolke et al. 2008; Kelly et al. 2011; Moore et al. 2013; Ague and Axler 2016). The reasons for investigating TE distribution and zoning in garnet is multi-fold, from deducing metamorphic reactions (Spear and Kohn 1996; Pyle and Spear 1999; Moore et al. 2013), to the determination of geological rates based on element diffusivity (Jollands et al. 2018), to distribution of Sm and Lu to assist garnet geochronology (Anczkiewicz et al. 2007; Kelly et al. 2011; Anczkiewicz et al. 2014; Bloch et al. 2015; Guilmette et al. 2018), to equilibrium partitioning with zircon and monazite to reconstruct P-T-time paths (Hermann and Rubatto 2003; Rubatto and Hermann 2007; Taylor et al. 2014; Taylor et al. 2017).

Previous studies have proposed different processes to explain the observed TE distribution in garnet. Rayleigh-type fractionation during prograde garnet growth (at least for compatible elements such as Y and heavy rare earth elements – Y+HREE) should produce a monotonic decrease in such elements (i.e. “bell-shaped” mass fraction profile; Otamendi et al. 2002). This is however not always observed, particularly at subsolidus conditions, and alternative models to explain complex TE zoning have been proposed: (i) TE supply regulated by episodic breakdown of other TE-bearing phases, including accessory minerals (Hickmott et al. 1987; Pyle and Spear 1999; Yang and Rivers 2002; Konrad-Schmolke et al. 2008; Moore et al. 2013; Raimondo et al. 2017); (ii) differential supply of TE limited by diffusion in the matrix (Skora et al. 2006; Moore et al. 2013); (iii) inherited TE distribution from pre-existing phases in the site of garnet growth (Hirsch et al. 2003; Moore et al. 2013; George et al. 2018); (iv) solubility of TE in an intergranular fluid (Moore et al. 2013); and (v) possibly replacement of garnet along fractures and irregular domains by a second garnet generation, as shown for major elements (Pollok et al. 2008; Giuntoli et al. 2018). Understanding if key TE in garnet are distributed according to equilibrium processes, or to disequilibrium limited by diffusion and other phase stability, or if they are modified by secondary replacement is fundamental in the

interpretation of mineral dates, thermometry and thermodynamic modelling (Lanari and Engi 2017). Discerning between these processes in natural samples requires high-resolution imaging of the TE distribution within garnet grains.

Most commonly, TE mineral composition is determined using laser ablation inductively coupled plasma mass spectrometry (LA-ICPMS) in spot analyses and line profile mode, whereas mapping (here intended as 2D images where the ablation crater's depth is smaller than its diameter) is in its infancy (Ulrich et al. 2009; Raimondo et al. 2017; George et al. 2018; Hyppolito et al. 2019). This one-dimensional documentation has limited our understanding of how TE are distributed within garnet and which zoning can be considered typical for certain growth processes (Moore et al. 2013). In the same way that electron microprobe mapping revolutionized the understanding of mineral zoning for diffusion and thermodynamic studies, the mapping of TE zoning can make a substantial difference in the understanding of TE systematics as petrogenetic indicators and of mechanisms of mineral growth during metamorphism.

Certain TE are amenable to mapping by electron microprobe analysis at the 100 $\mu\text{g/g}$ level (P and Y, more rarely Cr and Sc) (Spear and Kohn 1996; Yang and Rivers 2002; Moore et al. 2013). Other studies have used secondary ion mass spectrometry for the analysis of partial or full sets of TE in garnet at the 0.1 $\mu\text{g/g}$ level (Hickmott et al. 1987; Green et al. 2000; Skora et al. 2006; Moore et al. 2013), but this method is time-consuming and relies heavily on appropriate matrix-matched standards. In the last 15 years, TE mapping by LA-ICPMS has been optimized for geological materials (Woodhead et al. 2007; Ulrich et al. 2009), but its application to garnet is still very limited (Ulrich et al. 2009; Raimondo et al. 2017; George et al. 2018; Guilmette et al. 2018). Even more recently, LA-ICP-TOFMS instrumentation has been developed to achieve rapid acquisition of composition maps for all elements at high lateral resolution (Gundlach-Graham et al. 2015; Bussweiler et al. 2017, Burger et al. 2017; Burger et al. 2019) and the first applications to geological materials have been published (Gundlach - Graham et al. 2018; Bussweiler et al. 2019; Ubide et al. 2019). The major advantage of LA-ICP-TOFMS is its quasi-simultaneous acquisition of complete mass spectra at high repetition rates. In combination with low-

dispersion aerosol transport systems (Wang et al. 2013), it allows for rapid and sensitive element mapping.

In this contribution we present mapping of the TE zoning in garnet using LA-ICP-TOFMS. We investigate the distribution of a large set of TE at a high lateral resolution of 5 μm in three garnet grains that represent examples of contrasting metamorphic conditions with respect to T, P and relation to major elements. (1) Garnet from an amphibolite-facies metapelite from the Central Alps grew during two orogenic cycles at T-time conditions at which diffusion of major elements is negligible ($<600^\circ\text{C}$), allowing direct comparison of major and TE zoning. (2) Garnet in granulite-facies metapelite from Malenco (Eastern Central Alps) records a single metamorphic cycle over at least 20 My reaching maximum T of 850°C , which has erased major element zoning, whereas TE zoning is preserved. (3) Garnet in a low temperature eclogite (1.8–2.3 GPa and $520\text{--}600^\circ\text{C}$) from the Sesia Zone in the Western Alps shows major element zoning with secondary garnet replacement cross-cutting the initial growth zoning. The record of different behaviour of TE in these three samples sheds light on the mechanism of garnet growth, equilibration, and replacement, which provides fundamental insights into crustal metamorphic processes.

Geological background and sample description

Sample CLM6 is from the Campolungo locality (N $46^\circ27'60''$, E $8^\circ42'54''$) in the Swiss Central Alps. The Mesozoic sediments at Campolungo document amphibolite-facies conditions during Barrovian metamorphism at T of $\sim 600^\circ\text{C}$ and P of 0.5–0.65 GPa at peak T (Todd and Engi 1997). The studied sample is from a paragneiss unit that is tectono-stratigraphically situated below the Mesozoic sediments. The sample was previously investigated by Boston et al. (2017) who reported major element zoning in the garnet during two distinct metamorphic cycles. Boston et al. (2017) obtained monazite ages of 332 ± 4 Ma for sample CLM6 and allanite ages between 32 and 28 Ma for a nearby Mesozoic metasediment. They concluded that the paragneiss unit underwent lower amphibolite-facies metamorphism during the Variscan orogeny and was again metamorphosed at amphibolite facies during the Alpine collision. On the

contrary, Gieré et al. (2011) interpreted the garnet growth in similar metapelites from the Campolungo area as solely grown during Alpine metamorphism, although without any geochronological constraints. The P-T-time reconstructed by Boston et al. (2017) suggests that the sample remained at T of 500-600 °C for ~15 My (from 33 to 18 Ma) during the Alpine orogeny. Fast cooling in the order of > 100°C/Ma occurred after 18 Ma (Boston et al. 2017). The duration of Barrovian metamorphism in this unit during Variscan orogeny is unknown but likely on the order of several My.

The sample mineral assemblage consists of quartz, white mica, biotite, plagioclase, garnet (15% in volume), staurolite and minor kyanite and graphite, with accessory monazite and zircon. The investigated garnet is euhedral and has a diameter of ~5 mm (Online Resource Fig. S1a). It is wrapped by a foliation defined by mica, quartz, plagioclase and graphite; it is mainly in direct contact with biotite, white mica and quartz. In the core, it contains an internal foliation defined by inclusions of fine-grained ilmenite, quartz and white mica; larger inclusions of ilmenite are in the rim, minor secondary chlorite is distributed along fractures.

Sample MG1 is from the granulite-facies metapelites of the Val Malenco area, Northern Italy (N 46°18'38.1", E 9°48'04.3"). These metapelites are part of a lower crustal sequence that experienced extensive partial melting and melt extraction (up to 50%, Hermann et al. 1997), leading to the formation of restitic rocks that are rich in garnet. The peak metamorphic conditions recorded are 1.0 GPa and 800-850°C (Hermann et al. 1997; Müntener et al. 2000). The timing of protracted high-T metamorphism and partial melting in the metapelites is constrained by U-Pb dating of monazite and zircon to 281–257 Ma (Hermann and Rubatto 2003). Hermann and Rubatto (2003) reported flat major element zoning in large garnet cores and a small overgrowth at the rim. In contrast to major elements, the garnet cores display marked Y and HREE zoning. Based on garnet composition and inclusion assemblages Hermann and Rubatto (2003) concluded that garnet growth may have started at subsolidus amphibolite facies (600°C), but most of the preserved garnet grew in a 20 My period at T of 700–

850°C in the presence of melt (their zone 1–2), with a minor Ca-rich rim (zone 4–5) forming during isobaric cooling and reaction with plagioclase.

Sample MG1 has a granoblastic texture with an assemblage of garnet (50% in volume), quartz, plagioclase (largely retrogressed to chlorite, clinozoisite and white mica), biotite (partly retrogressed to chlorite) and kyanite, with accessory rutile, zircon, monazite and pyrrhotite (Online Resource Fig. S1b). The analysed garnet is subhedral, 1.5 mm in diameter, and is surrounded by more or less altered plagioclase, with minor rutile. It contains inclusions of altered plagioclase and biotite; secondary mica fills the fractures.

Sample AV16021 (hereafter AV) is from the Ivozio Complex (Zucali et al. 2004; Delleani et al. 2018) at the eastern border of the Sesia Zone (Western Alps, Italy). The Sesia Zone consists of a basement that underwent Variscan metamorphism in the Carboniferous and that was intruded by Permian bimodal magmatism. This basement experienced eclogite facies metamorphism during Alpine subduction in the late Cretaceous (Rubatto et al. 2011). The banded texture in the eclogites and the overall abundance of mafic rocks in the Ivozio Complex suggests a gabbroic body as protolith (Zucali et al. 2004). The P-T path of the Ivozio metabasites during the Alpine metamorphism is characterized by a multi-stage evolution (Zucali and Spalla 2011). Garnet growth was protracted from ~ 1.1 GPa and 500°C to peak conditions of 1.8 – 2.3 GPa and 520 – 600 °C. Zircon dating in the surrounding micaschists (Rubatto et al. 1999) and allanite dating in a mafic rock (Vho 2020) constrains prograde to peak metamorphism of this unit to ~68-60 Ma.

This sample (N 45°32'09", E 07°50'46") contains a typical eclogite-facies assemblage and is characterized by (i) mafic layers rich in garnet (ca. 40%), omphacite and glaucophane, with minor rutile, white mica and pyrite, and (2) quartz-rich domains with garnet (20–30%), white mica, omphacite and glaucophane. The analysed garnet grain (Online Resource Fig. S1c) is 1.4 mm in diameter; it is surrounded mostly by quartz and minor omphacite, white mica and zoisite, and contains inclusions of quartz, zoisite, white mica and minor rutile. There is no previous study of the TE distribution in garnet in this unit.

In each sample, the investigated grains are large relative to other grains in the same sample, and are thus most likely to show the complete zoning sequence. Indeed, the selected garnet grains in CLM6 and MG1 show all the zones in major and TE that were identified by previous studies based on multiple garnet grains (Hermann and Rubatto 2003; Boston et al. 2017).

Methods

EPMA mapping

X-ray compositional maps were acquired by electron probe micro analysis (EPMA). They are presented as standardized mass fraction maps of oxide weight percentage obtained using the program XMAPTOOLS 3.2.2 (Lanari et al. 2014; Lanari et al. 2019). X-ray mapping was carried out using a JEOL JXA-8200 SuperProbe at the Institute of Geological Sciences of the University of Bern following the analytical procedure described in Lanari and Piccoli (2019). The maps for garnet CLM6 and MG1 were acquired after LA-ICP-TOFMS analysis and minor repolishing (a few μm) of the thin sections. Acquisition was at 15 keV accelerating voltage using a specimen current of 100 nA, dwell times of 210 ms (CLM6), 250 ms (MG1), a step size of 8 μm and a beam diameter of 1 μm . The spot analyses used as reference standards were acquired at 15 keV accelerating voltage using a specimen current of 20 nA, dwell times of 20 s and 2 x 10 s for peak and background with a beam diameter of 1 μm . Oxide mass fractions were calculated using the CITZAF routine and the following synthetic and natural standard reference materials: garnet (SiO_2 , Al_2O_3), olivine (MgO , FeO), anorthite (CaO), tephroite (MnO) and ilmenite (TiO_2).

The CLM6 garnet was imaged across an area of 8 x 8 mm using wavelength dispersive spectroscopy (WDS) for Mg, Ca, Mn, Al, Fe and energy dispersive spectroscopy (EDS) for P, S, Si, Al, Ti, Na, K and Ni. The composition maps of SiO_2 , TiO_2 , Al_2O_3 , FeO , MnO , MgO and CaO were obtained using 60 fully quantified spot analyses.

The MG1 garnet was mapped across an area of 5 x 6 mm using WDS for Mg, Ca, Mn, Al, Fe and EDS for P, S, Zr, Si, Al, Ti, Na and K. The compositional maps of

SiO₂, TiO₂, Al₂O₃, FeO, MnO, MgO, CaO were obtained using 45 fully quantified spot analyses.

The AV garnet grain mapped for major elements is not the same as that investigated for TE, but a very similar one from the same sample (as shown below). Acquisition conditions were similar to the other samples and with a dwell time of 150 ms. EPMA mapping covered an area of 2.40 x 2.19 mm using WDS for Si, Mg, Ca, Mn, Al, Fe, Ti Na, K and EDS for P, S, Ni, La, Ce, Zr and Au. The compositional maps of SiO₂, TiO₂, Al₂O₃, FeO, MnO, MgO and CaO were obtained using 27 fully quantified spot analyses.

LA-ICP-TOFMS mapping

Element mapping was carried out using a 193 nm ArF excimer laser ablation system (GeoLas C, Lambda Physik AG, Goettingen, Germany) coupled to an ICP-TOFMS (icpTOF, TOFWERK AG Thun, CH) equipped with a high capacity interface pump (UNO 065 D, Pfeiffer, DE) at ETH Zurich. Laser ablation sampling was performed in a custom-built, low-dispersion aerosol transport system that provided single pulse transient signal durations of less than 10 ms (FW0.01M) for laser spot diameters $\leq 10 \mu\text{m}$ (Gundlach-Graham et al. 2015). The data for the three garnets was acquired over a time span of almost two years and experiments differed considerably. To an extent, this also reflects the distinct steps in method development accomplished over this time.

Garnet CLM6 was mapped using a method described earlier (Burger et al. 2017). Briefly, 60 adjacent sections of 25 parallel lines each were ablated. The individual sections were vertically concatenated during data post-processing to generate the full image. The laser fluence was 15 J/cm², a repetition rate of 100 Hz, a circular ablation spot size 5 μm in diameter and a scan rate of 500 $\mu\text{m/s}$ were employed for edge-to-edge sampling across an area of 8 x 7.5 mm or 2.4 million pixels (Mpx). The total image acquisition lasted 12 hours. At this stage, laser ablation and ICP-TOFMS were not synchronized, and acquisition of individual sections was manually started. The ICP-TOFMS was operated with its multi-notch filter to attenuate the plasma background ions ⁴⁰Ar⁺ and ⁴⁰Ar₂⁺ to levels <400'000 cps. This was to avoid saturation of the micro-channel plate

detector, but it also affected the transmission of nearby isotopes of lower m/z in particular. During data processing, it was found that Al and Si were too strongly biased to be evaluable and excluded from the quantification (see below). A vertical tilt of the sample surface was compensated for by manually refocusing the laser at the beginning of each individual subsection. A correction of horizontal tilt was however not possible at that stage. However, the quantified images suggest that there is no substantial impact of lateral tilt on data quality with respect to the geological interpretation (see “Results” and “Discussion”).

For imaging the second garnet MG1, the icpTOF's reaction cell was pressurized with hydrogen gas to reduce argon-based background species and molecular ions (Burger et al. 2019) without having to employ the multi-notch filter. Thereby, lower limits of detection (LOD) could be achieved for Al, Si and Ca in particular. Further improvements over the previous analyses included triggered acquisition of individual subsections, which enabled considerable automation of the measurements. The sample was imaged across an area of 7.3 x 6.6 mm (1.9 Mpx) in 88 equally sized subsections at a laser repetition rate of 50 Hz. The dimensions of the displayed image are 6.25 x 6.25 mm because areas surrounding the garnet were of no particular interest and were therefore cropped during image processing. The fluence of the laser was 20 J/cm². Total acquisition lasted 15 hours. The reaction cell in the ICP-TOFMS was fed with an H₂ gas flow rate of 2.5 mL/min throughout. During the experiment, however, an accidental interruption of the carrier gas flow occurred. This caused 25 lines of the image to be lost (black lines at 2/3 of the garnet height). A tilt of the sample surface in the vertical direction was corrected for by constantly adjusting the laser focus between the beginning of the first and the last line of the image, respectively. No tilt correction was necessary in the horizontal direction.

Finally, element imaging for the AV garnet was performed using the ICP-TOFMS with extended drift tube (icpTOF 2R, ToFwerk, Thun, CH) providing higher mass resolving power. The ablation strategy involved a hole-drilling approach for which the isotope signals for each individual pixel were obtained by firing 25 laser pulses at the same position using a spot diameter of 5 µm and a frequency of 100 Hz, before moving the sample to the next, adjacent position. While this strategy decreases data acquisition speed per unit area, sensitivities and LOD can

be substantially improved. This is a direct consequence of the increase in the amount of material sampled. The garnet was mapped across an area of 1 x 2 mm (i.e. 80 kpx) and collected in 8 equally sized subsections arranged in a 2 x 4 grid, lasting 7.5 hours. The fluence of the laser was 15 J/cm². In addition to argon background species ⁴⁰Ar⁺, and ⁴⁰Ar₂⁺, also ⁵⁶Fe⁺ was attenuated via the multi-notch filter to prevent detector saturation.

The laser craters' depths were not measured directly. Considering the 50% lower ion signals compared to SRM NIST 610, however, we assume the depth ablation rate to be correspondingly lower. For SRM NIST 610, a depth ablation rate between 200 and 300 nm/pulse has been determined for a fluence of 15 – 25 J/cm² with the same laser ablation system (Horn et al. 2001). The material removed from garnets CLM6 and MG1 is thus considered to originate from a depth of less than 200 nm (1 pulse/pixel) and less than 5 µm for garnet AV (25 pulses).

The data acquisition protocol and data processing routines have been described in more detail in a previous publication (Burger et al. 2017). Briefly, the mass calibration was verified and mass spectra were baseline-subtracted using Tofware (versions 2.5.10 and 2.5.11, Tofwerk AG, Thun). In-house written MATLAB scripts (version R2015a, Mathworks, MA) were then used for calculation of gas-blank-subtracted time traces, integration of signal intensities for individual pixels and their assignment to sample position. For the processing of the AV garnet data, the MATLAB routines had been incorporated in an in-house written Python 3 software suite.

Quantification of each pixel was based on relative sensitivity factors established through external calibration with the standard reference material NIST SRM 610 and a normalization of the mass fractions of all determined element oxides (Liu et al. 2008). Because Al and Si were biased through the use of the multi-notch filter, quantification of garnet CLM6 was done by excluding the mass fractions of Al₂O₃ and SiO₂ from the normalization. Electron probe microanalyses from this sample had revealed a homogeneous distribution of both elements at mass fractions of 21.0% Al₂O₃ and 37.5 % SiO₂. Element-oxide mass fractions determined by LA-ICP-TOFMS were thus normalized to the remaining 41.5%.

The gas blank intensities were assumed to represent the instrumental background. LOD were then estimated using counting statistics and the mean sensitivity of the elements detected during the mapping. Sensitivities were typically found to decrease by 30% to 60% during the acquisition of a map. This drift is assumed to originate mainly from drift in the ICP-TOFMS system and changes in laser energy reaching the sample surface. The sensitivity drift affects the pixels' signal/noise ratios (S/N) while element mass fractions and their ratios in particular are not considered to be affected substantially.

Results

Major element zoning

LA-ICP-TOFMS maps were produced for all elements including major, minor and TE. In all samples Ca, Mg and Mn maps best describe garnet major element zoning (Fig. 1-3). The LA-ICP-TOFMS maps for major elements generally reproduce the EPMA maps.

In CLM6 garnet, the EPMA-CaO map shows a euhedral concentric zoning in the core and mantle, from Ca-rich core to Ca-poor mantle, truncated by an irregular boundary to a homogeneous, Ca-poor rim (Fig. 1d). Both the euhedral Ca-rich core and the irregular mantle-rim boundary are recognizable in the LA-ICP-TOFMS-CaO map. The EPMA-MnO map (Fig. 1a) shows a diffuse zoning at the core-mantle transition (less pronounced euhedral core) and does not identify the irregular mantle-rim boundary. The LA-ICP-TOFMS-MnO map (Fig. 1b) is remarkably similar to the EPMA-MnO map. A line profile across the grain in MnO in both maps shows their close correspondence (Fig. 1c).

For garnet CLM6, the LA-ICP-TOFMS major element maps show analytical artefacts in the form of parallel bands with less uniform mass fractions for CaO in the core and mantle in particular (Fig. 1e). This is a result of an imperfectly mounted thin section, causing the laser to become slightly de-focused during individual sections analysed in a sequence. Together with the comparably low S/N obtained for Ca during these analyses, the change in ablated mass led to an underestimation of the CaO content. After re-focusing at the start of a new

sequence, the S/N increased quantification improved. Although these artefacts are present in all maps including the TE, they are clearly distinguishable from the true element zoning and thus do not affect the interpretation of the TE maps substantially.

The major element zoning depicted by the EPMA maps is less complex for the high-T garnet MG1 (Fig. 2 and Online Resource Fig. S2), which is composed of two homogeneous zones with a sharp boundary: (i) a large core that is nearly homogeneous in CaO, MgO and MnO (a very weak CaO outwards decrease is visible, Fig. 2d); (ii) a thin, discontinuous rim (5-50 μm across, left side and upper right corner, Fig. 2a and d) that is rich in CaO and poor in MnO and MgO. The LA-ICP-TOFMS maps (Fig. 2b and e) for the same elements show comparable zoning to those obtained by EPMA.

In the eclogitic sample AV, the garnet grain mapped by EPMA (Fig. 3) is a nearby grain with a comparable composition and zoning to the one analysed by LA-ICP-TOFMS. Still, core, mantle and rim composition obtained for the two different grains agreed within 0.1 wt% for major element oxides. The grain is characterized by a primary zoning pattern truncated by secondary radial features (Fig. 3, also see comparable features in the LA-ICP-TOFMS Mn and Mg maps discussed below). CaO, MnO and FeO generally decrease from core to rim, whereas MgO increases (Fig. 3). The central part of the core, particularly rich in inclusions, is poor in FeO and MnO and richer in MgO with respect to the rest of the core and similar to the rim composition. However, this feature is not visible in CaO (Fig. 3a). A broad rim of $\sim 200\ \mu\text{m}$ is rather homogeneous in composition, with relatively higher MgO and lower CaO, FeO and MnO. This main core-rim zoning is parallel to the euhedral garnet shape indicating that it represents primary growth zoning. This main zoning pattern is truncated by irregular linear features (hereafter veinlets) from 10 to 100 μm in width that run radially from rim to core and are similar to “healed cracks”. The veinlets are marked by distinct compositional changes in MgO, MnO and FeO, which are not observed for CaO.

Trace element zoning

Different types of zoning patterns can be recognized among the TE. In the following description of TE zoning in garnet the different substitution mechanisms for TE in garnet should be kept in mind. In the general garnet structure $X_3Y_2Z_3O_{12}$ the main substitutions are (i) divalent cations such as Zn and Co on the eight-coordinated X-site for Mg and Fe; (ii) trivalent Y+REE on the X-site for Ca; (iii) Ti^{4+} and trivalent V and Cr on the six-coordinated (octahedral) Y-site for Al; (iv) P^{5+} and Ge^{4+} on the four-coordinated (tetrahedral) Z-site.

In the amphibolite-facies garnet of CLM6, the sharp zoning defined by Ca (euhedral core, oscillatory zoning in the mantle and sharp discordant rim, best visible in the EPMA-CaO map, Fig. 1d) is reflected in numerous elements such as Sc, Y and the detectable REE from Eu to Lu (Fig. 4). The match between the Y+HREE zoning and the CaO zoning of the EPMA map is remarkable: the sharp boundaries, the fine fractures and annuli defined by CaO zoning are reproduced in the Y+HREE images, both in shape and position. Among the Y+HREE there is no distinguishable difference in the location of sharp zoning features such as the euhedral inner core or the annulus in the outer core (Fig. 4 and 5).

Some of the zoning features, particularly the discordant mantle-rim boundary, are also visible in P (Fig. 4), which shows a weak inverse zoning with respect to Ca, Y and REE: the core is poorer in P than the mantle and rim. Notably, the diffuse zoning of Mg, Fe and Mn in the EPMA maps (Fig. 1 and Online Resource Fig. S2) is not mimicked by any of the detectable TE. Some of the TE that are below LOD in garnet (see Online Resource Table S1 for LOD), instead highlight inclusions and alteration along fractures. The metals Ni, Cu, As, Rh, Ba, Ta, Pt, Au, Bi and U (maps not shown) are systematically distributed along fractures and highlight a potential bias for these elements when fractures are intersected during spot or line analysis. Elements such as Cu, Zr, the light-REE (LREE), Hf, Pb, Th and U are relatively incompatible elements in garnet and well mark the micro-inclusions of sulphides, zircon and monazite/allanite. Around the largest inclusions some re-deposition of material from previous laser pulses is visible as a halo (see as an example Ce, plotted in logarithmic scale in Fig. 4).

The granulite-facies MG1 garnet has visible zoning in the core–mantle in most detectable elements, including some of the transition metals that were not detected in CLM6 garnet (Fig. 6). While there is a strong correlation between major and TE zoning in CLM6, MG1 garnet shows far more features in the TE than in the major element maps. A detailed zoning is defined by Y, which is enriched in the core, decreases outwards, and has an outer band of enrichment (hereafter, mantle) of 50-100 μm , before the truncating thin rim with low Y contents. This low-Y rim corresponds to the high-Ca and low-Mn rim in the major element map (Fig. 2). The mantle–rim boundary is particularly sharp, whereas the core to mantle zoning is more gradational in all elements. The full zoning is also shown by the REE Tb–Lu (Fig. 6); notably the HREE increasingly show a patchy zoning in the inner core.

A gradually decreasing concentric zoning across the core is well defined in Sc (Fig. 6) and weak in Ge (Online Resource Fig. S3), but the enrichment in the mantle is not evident. The opposite zoning with higher mass fractions in the mantle than in the core is shown in V and low concentrations detected for Cr and Zr (Fig. 6 and Online Resource Fig. S3). The mantle enrichment versus the core is also evident in the mid REE Gd and Tb, whereas HREE Ho–Lu have a higher mass fraction in the inner core than in the mantle.

The high-Ca thin rim, that is distinct in major elements, is depleted in most TE (mainly below LOD), but locally enriched in Ti, V and Cr (Fig. 6, left and top right of the crystal). Notably these elements are not equally present throughout the thin rim, but only in one corner that is in contact with altered biotite; in this area Ti, V and Cr show an additional zoning within the rim.

As for sample CLM6, some elements show particularly well the inclusions of zircon and monazite with halos: these are Zr, LREE and Ce (Fig. 6, but also e.g. La and Th not shown). Elements that are not detectable and are in low mass fraction in the garnet are instead concentrated in fractures (K, Ti, Cu, Zn, Rb, Rh, Au, Sn, Ba, La, Ce, Pt, Gd, Pb and Bi). This is likely due to later alteration by fluids and/or mineral replacement.

In the eclogitic AV garnet, Y and HREE define the principal zoning with a euhedral core and progressive depletion towards the outside (Fig. 7). An equally

euohedral rim is marked by an overall increase in Y and some REE with local oscillatory zoning. Most notably, the crosscutting radial veinlets characterized by higher Mn and lower Mg than the rim composition are also visible in TE such as Zn and Co, but do not affect the euohedral and oscillatory zoning of Y+HREE and Ti (see “Discussion”).

The inclusion-rich core has an inner part with more irregular zoning, mimicking the major elements in the nearby grains (see also Fig. 3). A thin halo depleted or enriched in elements can be seen around some of the inclusions in the core; in particular, V is enriched at the expenses of most other elements around epidote inclusions. The external part of the rim has the highest mass fraction in Y, Dy and Ho, whereas from Tm to Lu the inner core is richer than the rim. The rim has a visible oscillatory zoning in most Y+HREE, particularly in the outer part, and a ~ 50 μm thick annulus midway across marked by Y, Ho and Er (see red arrows in the Y and Er maps of Fig. 7). The euohedral, but discontinuous zoning in Y+HREE suggests that the core and rim represent two distinct stages of growth, but without signs of resorption between the core and rim. The transition metals, Sc, Cr, Co and Zn, have zoning roughly opposite to Mn, with flat and low mass fraction in the core of the garnet, and higher mass fractions in the rim. Co and Zn match particularly close the Mg zoning. Ti and Zr are the opposite, enriched in the core with respect to the rim. V and Cr have a more irregular distribution, and have patchy areas of enrichment.

Discussion

Spatial resolution, detection limits and accuracy

The LA-ICP-TOFMS setup delivers element maps that have a lateral resolution of 5 μm (pixel size) compared to the 15-17 μm previously reported for LA-ICPMS images (Ulrich et al. 2009; Raimondo et al. 2017; George et al. 2018). This increased resolution approaches that of EPMA major element imaging, with comparable or shorter acquisition times. The improved lateral resolution allows for a closer investigation of diffuse versus sharp compositional zoning and of thin features like mineral replacement veinlets (see below).

The achievable LOD are determined by the mass removed per pixel. Depending on the sampling mode, low abundant elements may not be detectable: (i) the approach used for the CLM6 garnet did not allow for detection of Zr (LOD 47 $\mu\text{g/g}$), except for several inclusions. Lower LOD could be achieved with the modes used for MG1 and AV garnets, which allowed for mapping the Zr distribution (LOD 20 and 3 $\mu\text{g/g}$, respectively), which is useful to identify events of zircon dissolution in aqueous fluids and melts during garnet growth. (ii) LREE La–Pr were always below LOD (10–0.4 $\mu\text{g/g}$ depending on setup), but these elements are not crucial for the petrogenesis of garnet. LREE La–Pr were not detectable even by LA-ICPMS analyses at 40 μm diameter laser spot size. (iii) Sm and Nd mass fractions are below LOD (50–2 $\mu\text{g/g}$ depending on setup), while analyses at 40 μm spot size yielded mass fractions of 0.1–10 and 0.2–4 $\mu\text{g/g}$ of Sm and Nd for CLM6 and MG1 garnet (Online Resource Table S2), respectively. The distribution of Sm and Nd would be relevant for the interpretation of Sm–Nd ages. Previous studies using conventional LA-ICPMS and a spot size of 15 μm have not reported LOD for elements, but present coarse maps of Zr, Sm and Nd (Raimondo et al. 2017; George et al. 2018).

Normalized REE patterns are a common way to analyse their relative occurrences and it is interesting to compare the REE patterns obtained with 5 μm pixel size LA-ICP-TOFMS data to those obtained using larger laser spots. The low sample mass analysed per 5 μm diameter pixel limits the attainable precision of the quantification and thus the REE pattern of individual pixels appears noisier than analyses using larger spot sizes (Raimondo et al. 2017). Nonetheless, the comprehensive element maps can be used to identify contiguous regions of similar composition and thus likely similar genesis. REE patterns from integrated pixel data within such areas show smooth trends that compare well with those generated by LA-ICPMS analyses using larger spot size. Figure 8 compares the data obtained for the main zones in MG1 garnet by LA-ICP-TOFMS using areas of ~50–600 pixels (~200–2400 μm^3 sample volume) with those from LA-ICPMS analyses using 60 μm spot size (~20 μm deep, ~56000 μm^3 , data in Online Resource Table S2). By averaging a larger sample mass of the LA-ICP-TOFMS analyses, the patterns become very similar. The

pattern for the M-HREE is relatively smooth and the individual features of the different regions are fully reproduced.

The reconstruction of REE composition from LA-ICP-TOFMS maps offers the advantage that, even for zones with low mass fractions, accurate REE patterns can be constructed, if a sufficiently large area can be identified. Using XMapTools, data from areas of variable size and shape can be extracted while avoiding areas with multiple inclusions (Lanari and Piccoli 2019). Additionally, the REE data represent the near surface composition and mitigates artefacts from unrecognized inclusions at sub-surface levels, which would be sampled in LA-ICPMS spot analyses at greater depths. This is demonstrated for sample AV, where the garnet's TE mass fractions in the core could not be obtained via conventional spot analysis because of the high number of inclusions, mainly epidote. Using the area sampling approach smooth REE pattern can be extracted for all domains (Fig. 9)

Diffusion versus growth zoning

Diffusion of major and TE can be assessed relative to each other by comparing maps and traverses across garnets CLM6 and MG1, for which better constraints for the temperature and duration of metamorphism exist.

In sample CLM6, Ca preserves pronounced zoning that matches that of Y and P (Fig. 1, 4 and 5), which are slow diffusing TE (Spear and Kohn 1996; Yang and Rivers 2002; Ague and Axler 2016; Jollands et al. 2018; Higashino et al. 2019); this is thus interpreted as growth zoning. This is in contrast with the diffusive blurring of Mg and Mn zoning at a scale of 100-200 μm (Fig. 1 and Online Resource Fig. S2), which is not observed in any of the detectable TE. This indicates that none of the detectable TE, including the transition metals Sc, V and Cr, display any diffusion equilibration at the conditions recorded by the CLM6 sample (max 600 °C and cooling over 15-30 My). The garnet core underwent two metamorphic cycles at amphibolite-facies conditions (Variscan and Alpine orogeny, see above), whereas the rim only recorded Alpine metamorphism and remained at 500–600°C for ~15 My (Boston et al. 2017). The duration of the Variscan metamorphism is unconstrained, but expected to be of the same order of magnitude.

Element diffusion is observed in the granulite-facies MG1 garnet in core and mantle: while the mantle–rim boundary is sharp in both major and trace elements, the TE show compositional zoning in the large garnet core and mantle where Ca and Mn are homogeneously distributed. The high lateral resolution of the LA-ICP-TOFMS maps allows assessing how sharp the TE zoning is. As for the CLM6 garnet, it is expected that the initial zoning of the major elements in the garnet core matches that of TE such as Y and HREE. In garnet MG1, these slow diffusing elements do not show sharp zoning at the microscale, for example between the core and the mantle (Fig. 10), as instead observed in CLM6 garnet for the same elements. While it cannot be excluded that the gradual zoning in Y+HREE in garnet MG1 is a growth feature, this pattern suggests diffusion.

Representative compositional profiles for Y and HREE between garnet core and mantle were extracted from the maps by averaging spots along core–mantle profiles (Fig. 10 and Online Resource Fig. S5-S8). The mass fraction change as a function of distance fits the solution of a diffusion law and permits calculating diffusion coefficients D (in m^2s^{-1} ; expressed in $\log D$). The $\log D$ values for each element were obtained by inversion based on single-component diffusion over a time interval of 10 My in a homogeneous medium along a one-dimensional semi-infinite profile (see details in Online Resource). The 10 My period is based on the P-T-time constrained by Hermann and Rubatto (2003) for this sample and represents a minimum estimate of the period of time that the sample remained above 750°C after the growth of the garnet mantle.

The extracted values of $\log D$ are between -22.8 and $-23.3 \text{ m}^2\text{s}^{-1}$ for Y and HREE (Fig. 10 and Online Resource Fig. S7 and S8). Lower diffusion coefficients would be obtained if (i) a longer residence time at these temperatures is used, (ii) the section is not perpendicular to the core/mantle boundary, and (iii) if part of the transitional zoning was due to growth. When compared to literature data calculated at 1 GPa, 750 °C and 800 °C (Fig. 10) our D values show a good match with the diffusion coefficients of Carlson (2012) obtained via numerical simulation on natural data, and with the experimental data of Bloch et al. (2015) for Lu and Dy. The diffusion coefficients for Y obtained in our study are instead

an order-of-magnitude lower than diffusion coefficients derived from experimental data by Cherniak (2005).

This modelling shows that the gradual zoning between mantle and core of granulite garnet MG1 is reconcilable with diffusion of Y and HREE across $\sim 200\ \mu\text{m}$ in 10 My at temperatures of 750–800°C. An advantage of the mapping technique is that the profiles can be perfectly placed so that they are perpendicular to the compositional interface, and avoid inclusions and heterogeneities that compromise the diffusion modelling.

Our data unfortunately do not allow a comparison of the diffusivity of Lu with that of Hf, or of Sm with Nd, which are mostly below LOD in MG1 garnet, and thus prevents us from making conclusions on relative diffusivities of the elements used for garnet geochronology (Kohn 2009; Bloch et al. 2015). Our data however confirm that Lu can diffuse over hundreds of microns in granulite facies garnet over geological times. This has implications for the interpretation of Lu-Hf garnet ages as they may not record garnet growth, but cooling below a certain T in granulite-facies rocks.

Garnet growth mechanisms recorded by trace element zoning

The similarities and differences in the TE zoning patterns observed in the three samples can be related to the different mechanism and conditions of garnet growth. Previous studies have identified different processes that can control TE incorporation and thus zoning in garnet (Pyle and Spear 1999; Otamendi et al. 2002; Skora et al. 2006; Moore et al. 2013; Ague and Axler 2016; George et al. 2018). The TE zoning images obtained can be used to identify which of these processes was dominant during garnet growth at the different conditions.

1) Equilibrium subsolidus growth

Equilibrium growth with a constant matrix composition and not limited by TE diffusion in the matrix produces simple zoning patterns typical of Rayleigh fractionation (Hollister 1966; Otamendi et al. 2002; Moore et al. 2013). The resulting zoning pattern in garnet is a gradual decrease from core to rim in the mass fractions of TE (bell shape) that are compatible in garnet. This is the dominant zoning type observed for Sc, Y and HREE in the core of the garnet

grains investigated, and particularly in garnet CLM6 and MG1. The establishment of equilibrium growth in TE in CLM6 and MG1 garnets lends support to the TE partitioning with other phases calculated in previous studies (Hermann and Rubatto 2003; Boston et al. 2017).

2) Peritectic garnet growth

In sample MG1, a part from the core, an additional zone with high TE content (mantle) is present (Fig. 6). The gradational transition from core to mantle is compatible with diffusional relaxation of a sharp boundary (see above). The mantle is much broader in size than thin annuli observed in garnet grown at lower temperatures (CLM6 and AV garnet, see discussion below). V, Cr, Zr, Y and the M-HREE are enriched in the mantle with respect to the outer core. Based on petrological, geochronological and inclusion evidence, Hermann and Rubatto (2003) related the garnet core to prograde metamorphism mainly at subsolidus conditions and the mantle to peritectic garnet growth during biotite melting. During the growth of the garnet mantle, V was likely liberated by muscovite and biotite melting (Yang and Rivers 2000), whereas Zr, Y and REE could be derived from dissolution of accessory phases, including xenotime, zircon and monazite. The Zr zoning should be particularly diagnostic for the transition of subsolidus to suprasolidus garnet growth as the solubility of Zr in the fluid phase increases over at least an order of magnitude from an aqueous fluid to a hydrous melt (Hermann and Rubatto 2009). Another feature of growth zoning in the presence of melt is that TE mass fractions can increase towards the garnet rim as seen in the REE. This is likely related to the increase of REE solubility with increasing temperature in the coexisting melt buffered by monazite (Stepanov et al. 2012) during prograde garnet growth. The embayed shape of the core–mantle boundary in most Y+REE may also indicate that the first melt resorbed a small amount of the core garnet.

Notably, Lu is enriched in this mantle domain, which is volumetrically significant. It implies that in this sample Lu-Hf ages would not preferentially date the core, as argued for most garnet in crustal rocks (e.g. Lapen et al. 2003; Skora et al. 2006), instead Lu-Hf ages in this sample would be intermediate between initial garnet growth in subsolidus and final growth during late anatexis. Lu enrichment

during melting has a similar effect to Lu enrichment in garnet during intense resorption as proposed by Kelly et al. (2011).

3) Diffusion-limited uptake of TE

This mechanism produces zoning patterns with a central peak and annular maxima (broad annulus where the TE mass fraction is higher) as modelled by Skora et al. (2006) and Moore et al. (2013). The model predicts a progressive shift in the position of the maximum REE mass fraction from core to rim for heavy to mid-REE (MREE), as the REE are incorporated progressively later with decreasing atomic number, due to the relatively slow diffusivity of the lighter REE. The resulting zoning can have many variations as it depends on relative diffusivities, T and duration of growth (Moore et al. 2013).

Models agree that diffusion-limited uptake during growth is more pronounced at lower T when the diffusivity is slow, and becomes negligible at high T. By comparing garnet zoning in metapelites from two localities, Moore et al. (2013) put the limit where diffusion-limited supply is dominant between 465 and 550°C. This feature has been observed in garnet from amphibolite-facies metapelites (Moore et al. 2013; George et al. 2018) and from eclogites (Skora et al. 2006) where prograde garnet growth is documented to a maximum T between 500–600°C. Amphibolite-facies garnet CLM6 does not exhibit broad annular maxima for Y and the REE, but these elements decrease constantly from the inner to the outer core (Fig 4 and 5). The identical shape in Y+REE zoning testifies to the fact that diffusion-limited supply and uptake of REE was not a significant mechanism during the growth of CLM6 garnet.

A diffusion-limited uptake of REE would fit, to some extent, the zoning observed in eclogitic AV garnet, whose centre/rim ratio mass fractions progressively decrease from Lu to Dy (Fig. 7). According to this mechanism, the core to rim growth in AV garnet would be continuous, which is however hard to reconcile with the very steep increase in Y, Dy, Ho and even Sc at the core–rim boundary. An alternative explanation for this type of zoning would be rim formation following an increase in MREE in the reactive bulk composition, e.g. by breakdown of a MREE-rich mineral (see mechanism 4 below). Therefore, we

conclude that the garnet core in sample AV shows no diffusion-limited uptake of REE.

Conforming with model predictions (Moore et al. 2013), annular maxima are not observed in granulite-facies MG1 garnet because at high temperature TE diffusion in the matrix is increasingly fast and can keep pace with garnet growth.

4) Trace element supply by the breakdown of major and accessory phases

The breakdown of major or accessory minerals during garnet growth has been recognized to form sharp annuli in garnet for specific TE such as MREE (controlled by titanite and epidote) or P (controlled by apatite) (Pyle and Spear 1999; Yang and Rivers 2002; Konrad-Schmolke et al. 2008; Gieré et al. 2011; Spandler et al. 2011; Raimondo et al. 2017). Another feature of reaction-controlled TE availability is the decoupling of equally compatible elements, such as for example a different zoning in Y vs. Mn, which are hosted in distinct minerals (Konrad-Schmolke et al. 2006).

This type of zoning is prominent in the external part of the core of amphibolite-facies CLM6 garnet, where a few annuli are present in Sc, Y and Yb. In CLM6 the annuli in Ca and Y do not match in the inner part, but partly correspond in the outer part of the core. The breakdown of accessory amounts of xenotime (48 wt% Y, 1.5-4 wt% Yb; e.g. Pyle and Spear 2000) or allanite (2000–5000 µg/g Y and 50–350 µg/g Yb; Boston et al. 2017) could provide significant Y to form minor annuli in garnet that roughly preserve the same Y/Yb ratio (1500 µg/g Y and 120 µg/g Yb). While xenotime and allanite were not observed in sample CLM6, xenotime has been reported as an included phase in garnet from metapelites in the area (Gieré et al. 2011) and allanite is present in other Campolungo metapelites (Boston et al. 2017). Likewise, Gieré et al. (2011) proposed that the Y and Yb annuli in similar garnet from this area were produced by the breakdown of allanite.

The oscillatory zoning in the rim of eclogitic garnet AV could be related to breakdown of accessory minerals, and because the oscillatory pattern is marked in different type of elements (Ti, Cr, Y and M-HREE) that would require the synchronous breakdown of multiple minerals. The REE patterns of the AV garnet (Fig. 9) display a pronounced increase in MREE from core (zone 1–4) to rim

(zone 5–7); this enrichment is in agreement with the breakdown of epidote/allanite and/or lawsonite that are a main repository for these elements in blueschist-facies rocks (Spandler et al. 2003). Indeed, Zucali et al. (2004) proposed that epidote and lawsonite were consumed between stage 1 and 2 of garnet growth. A milder enrichment in MREE also exists between zone 1 and 2 in AV garnet and this could correspond to the prograde breakdown of titanite to rutile.

5) Inherited TE from precursor and neighbouring phases

Overprinting zoning, inherited from the prior distribution of TE in precursor minerals, is predicted to produce small-scale irregular zoning patterns disconnected from major elements. This has been most commonly documented in garnet from amphibolite and eclogite-facies rocks for Ca, Cr and Ti (Spandler et al. 2011; Moore et al. 2013; Angiboust et al. 2014; Raimondo et al. 2017; George et al. 2018).

Zoning reconcilable with this mechanism is observed in granulite-facies MG1 garnet and eclogitic AV garnet. In MG1 garnet, the HREE Er–Lu (Fig. 6) show patchy zoning in the inner core and this could be related to inheritance from precursor phases, by which the prograde and relatively low temperature core grew before trace element equilibration was reached at the mm scale. Another zoning feature comparable to inheritance is the enrichment of Ti, Cr and V in parts of the thin rim of MG1 garnet (Fig. 4) in contact with retrogressed biotite. It is proposed that these elements were acquired from nearby biotite, but the process by which this happens is still unclear. In the AV garnet, the patchy distribution of V and Cr (Fig. 7) can also be explained as inherited from previous phases, most likely lawsonite and/or epidote.

6) Fluid related equilibration and replacement

Mineral replacement is a general term to address the process by which a mineral that comes into contact with a fluid out of equilibrium will tend to be replaced by a new mineral to reduce the free energy of the system (Putnis 2009). This replacement, also called interface coupled dissolution-precipitation (Putnis 2009), can also occur by garnet of a different composition that replaces pre-

existing garnet (e.g. Ague and Axler 2016). This process necessarily involves fluids and, when replacement is not complete, it commonly produces irregular zoning features, embayments, truncated zoning patterns and may be associated with porosity or hydrofractures. Such a process has been documented for secondary garnet replacing garnet in numerous cases from high-T granulites (Pollok et al. 2008; Ague and Axler 2016) to low-T eclogites (Erambert and Austrheim 1993; Kurz et al. 1998; Zack et al. 2002; Angiboust et al. 2011; Giuntoli et al. 2018) based on major element zoning.

The radial pattern of veinlets that cut across the main zoning in Mg and Mn in eclogitic garnet AV is interpreted as replacement (Fig. 7). The high lateral resolution of the LA-ICP-TOFMS TE maps allows resolving these thin radial features. Only the divalent cations Co and Zn, which have the same geochemical affinities as Mn and Mg, show the replacement features. None of the other detected TE are affected by garnet replacement. In the areas of intense replacement there is no significant correlation between Mg and other trace elements (see an example for Sc, V, Y and Er in Online Resource Fig. S4). This suggests that the veinlets do not represent a crack-seal mechanism in garnet as proposed by other studies that only looked at major elements (Angiboust et al. 2011; Giuntoli et al. 2018), but are rather related to a selective replacement that affected only some elements. This does not exclude that a micro-crack has acted as a fast fluid pathway in the centre of the veinlets, but replacement is the mechanism that controls garnet re-equilibration in this sample. The replacement can be partial and is likely controlled by the local bonding environment in the garnet, element mobility in the fluid and/or chemical potential gradients, by which the TE were not compatible in any of the nearby minerals forming at the time of garnet replacement.

Conclusions

We have demonstrated that LA-ICP-TOFMS can produce accurate and detailed images of TE zoning in garnet at a lateral resolution of 5 μm and with limits of detection suitable for mapping a wide range of elements at mass fractions down to sub $\mu\text{g/g}$ levels. Mapping of TE in garnet grains that formed in different

conditions from eclogite (500–600°C) to amphibolite (600°C) to granulite facies (850°C) provides insight into their growth mechanism and the distribution of TE during metamorphism in the presence of fluid and melt.

The method can generate TE maps that provide quantitative data to establish diffusivities of elements in garnet. The gradual zoning between mantle and core of the granulite-facies garnet is reconcilable with diffusion Y and HREE over ~200 µm in 10 My at temperatures of 750–800°C.

TE distribution in metapelite garnet grown at upper amphibolite to granulite facies is mainly controlled by an equilibrium Rayleigh fractionation process.

Peritectic garnet growth in the presence of melt is recognised in the granulite-facies sample, and marked by an increase in V, Cr, Zr, Y and the M-HREE. The redistribution for these elements in the melt is related to the melting of biotite (V and Cr), and the solubility of accessory minerals zircon and monazite.

Annuli or sharp enrichments in TE due to the breakdown of TE-rich minerals are confirmed to be typical of amphibolite-facies garnet, and also present in eclogitic garnet. The correlation between different TE can be used to speculate about the TE-rich phase that is breaking down and helps to correlate garnet growth with metamorphic reactions and accessory phase stability. Diffusion-limited uptake is not a dominant mechanism in any of the investigated garnet grains and is not observed even in the relatively low T eclogite garnet.

Overprinting zoning, inherited from precursor minerals, is identified in the core of granulite-facies garnet for HREE and in the eclogite-facies garnet for Cr and V, possibly in relation to epidote. In the granulite-facies metapelite, Ti, Cr and V can have enriched mass fraction when garnet grows in contact with (or in place of) high-T biotite.

Fluid-induced, secondary garnet replacement that affected the major elements in the eclogite-facies sample did not perturb the TE, with the exclusion of Zn and Co, divalent cations that substitute for Mg and Fe. This is in line with the low solubility of TE in aqueous fluids. The undisturbed TE distribution in the secondary replacement also indicates that this rather common garnet zoning in eclogites does not necessarily proceed by a crack-seal process.

References

- Ague JJ, Axler JA (2016) Interface coupled dissolution-reprecipitation in garnet from subducted granulites and ultrahigh-pressure rocks revealed by phosphorous, sodium, and titanium zonation. *Am Mineral* 101:1696-1699
- Anczkiewicz R, Chakraborty S, Dasgupta S, Mukhopadhyay D, Kołtonik K (2014) Timing, duration and inversion of prograde Barrovian metamorphism constrained by high resolution Lu–Hf garnet dating: A case study from the Sikkim Himalaya, NE India. *Earth Planet Sci Lett* 407:70-81
- Anczkiewicz R, Szczepanski J, Mazur S, Storey C, Crowley Q, Villa IM, Thirlwall MF, Jeffries TE (2007) Lu-Hf geochronology and trace element distribution in garnet: Implications for uplift and exhumation of ultra-high pressure granulites in the Sudetes, SW Poland. *Lithos* 95:363-380
- Angiboust S, Agard P, Raimbourg H, Yamato P, Huet B (2011) Subduction interface processes recorded by eclogite-facies shear zones (Monviso, W. Alps). *Lithos* 127:222-238
- Angiboust S, Pettke T, De Hoog JCM, Caron B, Oncken O (2014) Channelized fluid flow and eclogite-facies metasomatism along the subduction shear zone. *J Petrol* 55:883-916
- Bloch E, Ganguly J, Hervig R, Cheng W (2015) 176 Lu–176 Hf geochronology of garnet I: experimental determination of the diffusion kinetics of Lu 3+ and Hf 4+ in garnet, closure temperatures and geochronological implications. *Contrib Mineral Petrol* 169:12
- Boston K, Rubatto D, Hermann J, Engi M, Amelin Y (2017) Geochronology of accessory allanite and monazite in the Barrovian metamorphic sequence of the Central Alps, Switzerland *Lithos* 286-287:502-518
- Burger M, Hendriks L, Kaeslin J, Gundlach-Graham A, Hattendorf B, Günther D (2019) Characterization of inductively coupled plasma time-of-flight mass spectrometry in combination with collision/reaction cell technology – insights from highly time-resolved measurements *Journal of Analytical Atomic Spectrometry* 34:135-146
- Burger M, Schwarz G, Gundlach-Graham A, Käser D, Hattendorf B, Günther D (2017) Capabilities of laser ablation inductively coupled plasma time-of-flight mass spectrometry. *Journal of Analytical Atomic Spectrometry* 32:1946-1959
- Bussweiler Y, Borovinskaya O, Tanner M (2017) Laser Ablation and inductively coupled plasma-time-of-flight mass spectrometry - A powerful combination for high-speed multielemental imaging on the micrometer scale. *Spectroscopy* 32:20
- Bussweiler Y, Gervasoni F, Rittner M, Berndt J, Klemme S (2019) Trace element mapping of high-pressure, high-temperature experimental samples with laser ablation ICP time-of-flight mass spectrometry–Illuminating melt-rock reactions in the lithospheric mantle. *Lithos*:105282
- Caddick MJ, Konopásek J, Thompson AB (2010) Preservation of garnet growth zoning and the duration of prograde metamorphism. *J Petrol* 51:2327-2347
- Carlson WD (2012) Rates and mechanism of Y, REE, and Cr diffusion in garnet. *Am Mineral* 97:1598-1618
- Chakraborty S (2006) Diffusion modeling as a tool for constraining timescales of evolution of metamorphic rocks. *Mineral Petrol* 1-22
- Cherniak D (2005) Yb and Y diffusion in grossular garnet *Geochimica et Cosmochimica Acta Supplement* 69:405
- Cherniak DJ (2005) Yb and Y diffusion in grossular garnet. *Geochimica et Cosmochimica Acta, Supplement* 69:405
- Delleani F, Rebay G, Zucali M, Tiepolo M, Spalla MI (2018) Insight on Variscan geodynamics from the structural and geochemical characterization of a Devonian-

- Carboniferous gabbro from the Austroalpine domain (Western Alps) *Ofioliti* 43:23-39
- Erambert M, Austrheim H (1993) The effect of fluid and deformation on zoning and inclusion patterns in poly-metamorphic garnets. *Contrib Mineral Petrol* 115:204-214
- Ferry JM, Watson EB (2007) New thermodynamic models and revised calibrations for the Ti-in-zircon and Zr-in-rutile thermometers *Contrib Mineral Petrol* 154:429-437
- George FR, Gaidies F, Boucher B (2018) Population-wide garnet growth zoning revealed by LA-ICP-MS mapping: implications for trace element equilibration and syn-kinematic deformation during crystallisation. *Contrib Mineral Petrol* 173:74
- Gieré R, Rumble D, Günther D, Connolly J, Caddick MJ (2011) Correlation of Growth and Breakdown of Major and Accessory Minerals in Metapelites from Campolungo, Central Alps. *J Petrol* 52:2293-2334
- Giuntoli F, Lanari P, Engi M (2018) Deeply subducted continental fragments – Part 1: Fracturing, dissolution–precipitation, and diffusion processes recorded by garnet textures of the central Sesia Zone (western Italian Alps). *Solid Earth* 9:167-189
- Green TH, Blundy JD, Adam J, Yaxley GM (2000) SIMS determination of trace element partition coefficients between garnet, clinopyroxene and hydrous basaltic liquids at 2-7.5 GPa and 1080-1200°C. *Lithos* 53:165-187
- Guilmette C, Smit MA, van Hinsbergen DJJ, Gürer D, Corfu F, Charette B, Maffione M, Rabeau O, Savard D (2018) Forced subduction initiation recorded in the sole and crust of the Semail Ophiolite of Oman. *Nature Geoscience* 11:688-695
- Gundlach-Graham A, Burger M, Allner S, Schwarz G, Wang HAO, Gyr L, Grochimund D, Hattendorf B, Günther D (2015) High-Speed, High-Resolution, Multielemental Laser Ablation-Inductively Coupled Plasma-Time-of-Flight Mass Spectrometry Imaging: Part I. Instrumentation and Two-Dimensional Imaging of Geological Samples *Analytical Chemistry* 87:8250-8258
- Gundlach - Graham A, Garofalo PS, Schwarz G, Redi D, Günther D (2018) High - resolution, Quantitative Element Imaging of an Upper Crust, Low - angle Cataclasite (Zuccale Fault, Northern Apennines) by Laser Ablation ICP Time - of - Flight Mass Spectrometry. *Geostandards and Geoanalytical Research* 42:559-574
- Hermann J, Müntener O, Trommsdorff V, Hansmann W, Piccardo GB (1997) Fossil crust to mantle transition, Val Malenco (Italian Alps). *J Geophys Res* 102:20'123-120'132
- Hermann J, Rubatto D (2003) Relating zircon and monazite domains to garnet growth zones: age and duration of granulite facies metamorphism in the Val Malenco lower crust. *J Metamorphic Geol* 21:833-852
- Hermann J, Rubatto D (2009) Accessory phase control on the trace element signature of sediment melts in subduction zones. *Chem Geol* 265:512-526
- Hickmott DD, Shimizu N, Spear FS, Selverstone J (1987) Trace-element zoning in a metamorphic garnet. *Geology* 15:573-576
- Higashino F, Rubatto D, Kawakami T, Bouvier A-S, Baumgartner LP (2019) Oxygen isotope speedometry in granulite facies garnet recording fluid/melt-rock interaction (Sør Rondane Mountains, East Antarctica). *J Metamorphic Geol* 37:1037-1048
- Hirsch DM, Prior DJ, Carlson WD (2003) An overgrowth model to explain multiple, dispersed high-Mn regions in the cores of garnet porphyroblasts. *Am Mineral* 88:131-141
- Hollister LS (1966) Garnet zoning: an interpretation based on the Rayleigh fractionation model. *Science* 154:1647-1651
- Horn I, Guillon M, Günther D (2001) Wavelength dependant ablation rates for metals and silicate glasses using homogenized laser beam profiles—implications for LA-ICP-MS. *Applied Surface Science* 182:91-102
- Hyppolito T, Cambeses A, Angiboust S, Raimondo T, García-Casco A, Juliani C (2019) Rehydration of eclogites and garnet-replacement processes during exhumation in the amphibolite facies. *Geological Society, London, Special Publications* 478:217-239

- Jollands MC, Hanger BJ, Yaxley GM, Hermann J, Kilburn MR (2018) Timescales between mantle metasomatism and kimberlite ascent indicated by diffusion profiles in garnet crystals from peridotite xenoliths. *Earth Planet Sci Lett* 481:143-153
- Kelly ED, Carlson WD, Connelly JN (2011) Implications of garnet resorption for the Lu-Hf garnet geochronometer: an example from the contact aureole of the Makhavinekh Lake Pluton, Labrador. *J Metamorphic Geol* 29:901-916
- Kohn MJ (2009) Models of garnet differential geochronology. *Geochim Cosmochim Acta* 73:170-182
- Konrad-Schmolke M, Babist J, Handy MR, O'Brien PJ (2006) The Physico-Chemical Properties of a Subducted Slab from Garnet Zonation Patterns (Sesia Zone, Western Alps). *J Petrol* 47:2123-2148
- Konrad-Schmolke M, Zack T, O'Brien PJ, Jacob DE (2008) Combined thermodynamic and rare earth element modelling of garnet growth during subduction: Examples from ultrahigh-pressure eclogite of the Western Gneiss Region, Norway. *Earth Planet Sci Lett* 272:488-498
- Kurz W, Neubauer F, Dachs E (1998) Eclogite meso- and microfabrics: implications for the burial and exhumation history of eclogites in the Tauern Window (Eastern Alps) from P-T-d paths. *Tectonophysics* 285:183-209
- Lanari P, Engi M (2017) Local bulk compositional effects on metamorphic mineral assemblages. *Reviews in Mineralogy and Geochemistry* 83:55-102
- Lanari P, Piccoli F (2019) New horizons in quantitative compositional mapping – analytical conditions and data reduction using XMAPTOOLS. *Institute of Physics Conference Series: Materials Science and Engineering*, in press
- Lanari P, Vho A, Bovay T, Airaghi L, Centrella S (2019) Quantitative compositional mapping of mineral phases by electron probe micro-analyser. In: Ferrero S, Lanari P, Goncalves P, Grosch EG (eds) *Metamorphic Geology: Microscale to Mountain Belts* vol 478. Geological Society, Special Publications, London, pp 39-63
- Lanari P, Vidal O, De Andrade V, Dubacq B, Lewin E, Grosch EG, Schwartz S (2014) XMapTools: A MATLAB®-based program for electron microprobe X-ray image processing and geothermobarometry. *Computers & Geosciences* 62:227-240
- Lapen TJ, Johnson CM, Baumgartner LP, Mahlen NJ, Beard BL, Amato JM (2003) Burial rates during prograde metamorphism of an ultra-high-pressure terrane: an example from Lago di Cignana, western Alps, Italy. *Earth Planet Sci Lett* 215:57-72
- Liu Y, Hu Z, Gao S, Günther D, Xu J, Gao C, Chen H (2008) In situ analysis of major and trace elements of anhydrous minerals by LA-ICP-MS without applying an internal standard. *Chem Geol* 257:34-43
- Moore SJ, Carlson WD, Hesse MA (2013) Origins of yttrium and rare earth element distributions in metamorphic garnet. *J Metamorphic Geol* 31:663-689
- Müntener O, Hermann J, Trommsdorff V (2000) Cooling history and exhumation of lower crustal granulites and upper mantle (Malenco, Eastern Central Alps). *J Petrol* 41:175-200
- Otamendi JE, de la Rosa JD, Patiño Douce AE, Castro A (2002) Rayleigh fractionation of heavy rare earths and yttrium during metamorphic garnet growth. *Geology* 30:159-162
- Pollok K, Lloyd GE, Austrheim H, Putnis A (2008) Complex replacement patterns in garnets from Bergen Arcs eclogites: A combined EBSD and analytical TEM study. *Geochemistry* 68:177-191
- Putnis A (2009) Mineral Replacement Reactions. *Reviews in Mineralogy and Geochemistry* 70:87-124
- Pyle JM, Spear FS (1999) Yttrium zoning in garnet: Coupling of major and accessory phases during metamorphic reactions. *Geological Materials Research* 1:1-49
- Pyle JM, Spear FS (2000) An empirical garnet (YAG) – xenotime thermometer. *Contrib Mineral Petrol* 138:51-58

- Raimondo T, Payne J, Wade B, Lanari P, Clark C, Hand M (2017) Trace element mapping by LA-ICP-MS: assessing geochemical mobility in garnet. *Contrib Mineral Petrol* 172:17
- Rubatto D (2002) Zircon trace element geochemistry: distribution coefficients and the link between U-Pb ages and metamorphism. *Chem Geol* 184:123-138
- Rubatto D, Gebauer D, Compagnoni R (1999) Dating of eclogite-facies zircons: the age of Alpine metamorphism in the Sesia-Lanzo Zone (Western Alps). *Earth Planet Sci Lett* 167:141-158
- Rubatto D, Hermann J (2007) Experimental zircon/melt and zircon/garnet trace element partitioning and implications for the geochronology of crustal rocks. *Chem Geol* 241:62-87
- Rubatto D, Regis D, Hermann J, Boston K, Engi M, Beltrando M, McAlpine SRB (2011) Yo-Yo subduction recorded by accessory minerals in the Sesia Zone, Western Alps. *Nature Geoscience* 4:338-342
- Skora S, Baumgartner LP, Mahlen NJ, Johnson CM, Pilet S, Hellebrand E (2006) Diffusion-limited REE uptake by eclogite garnets and its consequences for Lu-Hf and Sm-Nd geochronology. *Contrib Mineral Petrol* 152:703-720
- Spandler C, Hermann J, Arculus R, Mavrogenes J (2003) Redistribution of trace elements during prograde metamorphism from lawsonite blueschist to eclogite facies; implications for deep subduction-zone processes. *Contrib Mineral Petrol* 146:205-222
- Spandler C, Pettke T, Rubatto D (2011) Internal and external fluid sources for eclogite-facies veins in the Monviso meta-ophiolite, Western Alps: Implications for fluid flow in subduction zones. *J Petrol* 52:1207-1236
- Spear FS (1993) *Metamorphic phase equilibria and pressure-temperature-time paths*. Mineralogical Society of America, Washington D.C.
- Spear FS, Kohn MJ (1996) Trace element zoning in garnet as a monitor of crustal melting. *Geology* 24:1099-1102
- Stepanov AS, Hermann J, Rubatto D, Rapp RP (2012) Experimental study of monazite/melt partitioning with implications for the REE, Th and U geochemistry of crustal rocks. *Chem Geol* 300-301:200-220
- Taylor RJM, Clark C, Harley SL, Kylander-Clark ARC, Hacker BR, Kinny PD (2017) Interpreting granulite facies events through rare earth element partitioning arrays. *J Metamorphic Geol* 35:759-775
- Taylor RJM, Harley SL, Hinton RW, Elphick S, Clark C, Kelly NM (2014) Experimental determination of REE partition coefficients between zircon, garnet and melt: A key to understanding high-T crustal processes. *J Metamorphic Geol*
- Todd CS, Engi M (1997) Metamorphic field gradients in the Central Alps. *J Metamorphic Geol* 15:513-530
- Ubide T, Caulfield J, Brandt C, Bussweiler Y, Mollo S, Di Stefano F, Nazzari M, Scarlato P (2019) Deep magma storage revealed by multi-method elemental mapping of clinopyroxene megacrysts at Stromboli volcano. *Frontiers in Earth Science* 7:239
- Ulrich T, Kamber BS, Jugo PJ, Tinkham DK (2009) Imaging element distribution patterns in minerals by Laser Ablation-Inductively Coupled Plasma-Mass Spectrometry (LA-ICP-MS). *Can Mineral*:1001-1012
- van Westrenen W, Draper DS (2007) Quantifying garnet-melt trace element partitioning using lattice-strain theory: New crystal-chemical and thermodynamic constraints. *Contrib Mineral Petrol* 154:717-730
- Vho A (2020) Fluid pathways in the subducting crust: insights from oxygen isotope records in metamorphic minerals, PhD thesis, Universität Bern, Bern
- Wang HAO, Grolimund D, Giesen C, Borca CN, Shaw-Stewart JRH, Bodenmiller B, Günther D (2013) Fast Chemical Imaging at High Spatial Resolution by Laser Ablation Inductively Coupled Plasma Mass Spectrometry. *Analytical Chemistry* 85:10107-10116

- Woodhead JD, Hellstrom J, Hergt JM, Greig A, Maas R (2007) Isotopic and Elemental Imaging of Geological Materials by Laser Ablation Inductively Coupled Plasma-Mass Spectrometry. *Geostandards and Geoanalytical Research* 31:331-343
- Yang P, Rivers T (2000) Trace element partitioning between coexisting biotite and muscovite from metamorphic rocks, Western Labrador: Structural, compositional and thermal controls. *Geochim Cosmochim Acta* 64:1451-1472
- Yang P, Rivers T (2002) The origin of Mn and Y annuli in garnet and the thermal dependence of P in garnet and Y in apatite in calc-pelite and pelite, Gagnon terrane, western Labrador. *Geological Materials Research* 4:1-35
- Zack T, Foley SF, Rivers T (2002) Equilibrium and Disequilibrium Trace Element Partitioning in Hydrous Eclogites (Trescolmen, Central Alps). *J Petrol* 43:1947-1974
- Zucali M, Spalla MI (2011) Prograde lawsonite during the flow of continental crust in the Alpine subduction: Strain vs. metamorphism partitioning, a field-analysis approach to infer tectonometamorphic evolutions (Sesia-Lanzo Zone, Western Italian Alps) *J Struct Geol* 33:381-398
- Zucali M, Spalla MI, Gosso G, Racchetti S, Zulbati F (2004) Prograde Lws-Ky transition during subduction of the Alpine continental crust of the Sesia-Lanzo Zone: the Ivozio Complex *Journal of Virtual Explorer* 16:1-21

Figures

Fig. 1 Comparison between LA-ICP-TOFMS and EPMA zoning in amphibolite-facies sample CLM6 for MnO (a-c) and CaO (d-f) both as images and profiles (c and f) along the line marked in (d). The slight difference in the MnO and CaO mass fractions along the profile is due to the different lateral resolution of the two methods. The horizontal parallel bands in the CaO LA-ICP-TOFMS map are an analytical artefact due to tilting of the thin section that caused the laser to become slightly de-focused during analysis.

Fig. 2 Comparison between LA-ICP-TOFMS and EPMA zoning in granulite-facies sample MG1 for MnO (a-c) and CaO (d-f) both as images and profiles (c and f) along the trace marked in (d). The slight difference in the MnO and CaO mass fractions along the profile is due to the different lateral resolution of the two methods. The black line at 2/3 of the garnet height in image (b) and (e) is due to an accidental interruption of the He gas flow. The black circles in the bottom-left quadrant of image (d) are LA-ICPMS spot analyses.

Fig 3 EPMA map of major elements in a garnet grain from eclogitic sample AV; the grain is not the same analysed by LA-ICP-TOFMS, but a nearby one with comparable zoning.

Fig. 4 LA-ICP-TOFMS element maps for amphibolite-facies garnet CLM6 for selected elements. The horizontal parallel bands are an analytical artefact due to tilting of the thin section that caused the laser to become slightly de-focused during analysis. Note that the Ce mass fraction is displayed in a logarithmic scale (*). Elements for which a median filter has been applied are marked with “mF”. For the median filtering, each pixel was recalculated as the median value in the 2-by-2 neighborhood around the corresponding pixel.

Fig. 5 Mass fraction profile for Y, Dy and Yb from rim to core of amphibolite-facies garnet CLM6 along the trace marked in the map. Note the good correspondence of the three elements: all three elements have the same rise at 1.6–2.0 mm, and Y and Yb mark another zone at 2.5 mm. The dotted red line represents the LOD.

Fig. 6 LA-ICP-TOFMS element maps for granulite-facies garnet MG1 for selected elements. The black line at 2/3 of the garnet height is due to an accidental interruption of the He gas flow. Ti and V mass fraction are displayed in a logarithmic scale (*). Maps for which a median filter has been applied are marked with “mF”. For the median filtering, each pixel was recalculated as the median value in the 2-by-2 neighborhood around the corresponding pixel.

Fig. 7 LA-ICP-TOFMS element maps for eclogitic garnet AV for selected elements. The red arrows in the Y and Er maps indicate the annulus. See discussion in text.

Fig. 8 Chondrite normalized REE patterns for different domains of granulite-facies garnet MG1 obtained using different instrumentation and setups. (a) REE patterns obtained sampling an area of ~ 50 – 600 pixels (~ 200 – $2400 \mu\text{m}^3$ sample volume) in the LA-ICP-TOFMS maps using the software XMapTools. (b) REE patterns obtained by routine LA-ICPMS spot analysis with a $60 \mu\text{m}$ diameter and $\sim 20 \mu\text{m}$ deep ($\sim 56000 \mu\text{m}^3$; data in Online Resource Table S2). See text for discussion.

Fig. 9 Chondrite normalized REE patterns for different domains of eclogitic garnet AV obtained by sampling the marked area in the LA-ICP-TOFMS maps shown in the inset using the software XMapTools.

Fig. 10 Results of diffusion modeling for Y, Lu and Ho (a-c) using a diffusion time-scale of 10 My. The red curve shows the original step function between mantle (left) and core (right). The fitted compositional profile is shown in blue and the obtained diffusion coefficient is reported in each plot as $\log D$. (d) comparison of the determined diffusion coefficients with literature data at $T = 750\text{ }^{\circ}\text{C}$ and $T = 800\text{ }^{\circ}\text{C}$ for a fixed pressure of 1 GPa. Abbreviations: C12: Carlson (2012); C05: from Cherniak (2005) published in Carlson (2012); B15: Bloch et al. (2015). See Online Resource for details on the modeling procedure.

Table S1. Average per-pixel LODs for the LA-ICP-TOFMS sessions.

CLM6 garnet		MG1 garnet		AV garnet	
Isotope measured	LOD in µg/g	Isotope measured	LOD in µg/g	Isotope measured	LOD in µg/g
²³ Na	5000	²³ Na	1000	²³ Na	200
²⁵ Mg	3000	²⁵ Mg	3000	²⁵ Mg	200
²⁷ Al	500	²⁷ Al	700	²⁷ Al	30
²⁹ Si	50000	²⁹ Si	40000	²⁹ Si	7000
³¹ P	600	³¹ P	2000	³¹ P	400
³⁴ S	n.d.	³⁴ S	10000	³⁴ S	3000
³⁹ K	1000	³⁹ K	90	³⁹ K	4000
⁴⁴ Ca	9000	⁴⁴ Ca	4000	⁴⁴ Ca	400
⁴⁵ Sc	200	⁴⁵ Sc	100	⁴⁵ Sc	10
⁴⁹ Ti	1000	⁴⁹ Ti	800	⁴⁹ Ti	100
⁵¹ V	100	⁵¹ V	80	⁵¹ V	6
⁵² Cr	300	⁵² Cr	100	⁵² Cr	10
⁵⁵ Mn	100	⁵⁵ Mn	100	⁵⁵ Mn	30
⁵⁷ Fe	4000	⁵⁷ Fe	3000	⁵⁷ Fe	900
⁵⁹ Co	100	⁵⁹ Co	100	⁵⁹ Co	5
⁶⁰ Ni	800	⁶⁰ Ni	400	⁶⁰ Ni	30
⁶⁵ Cu	300	⁶⁵ Cu	200	⁶⁵ Cu	20
⁶⁶ Zn	400	⁶⁶ Zn	300	⁶⁶ Zn	30
⁷¹ Ga	80	⁷¹ Ga	100	⁷¹ Ga	6
⁷⁴ Ge	100	⁷⁴ Ge	100	⁷⁴ Ge	7
⁷⁵ As	300	⁷⁵ As	300	⁷⁵ As	10
⁷⁷ Se	4000	⁷⁷ Se	2000	⁷⁷ Se	300
⁸⁵ Rb	70	⁸⁵ Rb	50	⁸⁵ Rb	20
⁸⁸ Sr	40	⁸⁸ Sr	20	⁸⁸ Sr	2
⁸⁹ Y	30	⁸⁹ Y	20	⁸⁹ Y	1
⁹⁰ Zr	70	⁹⁰ Zr	20	⁹⁰ Zr	3
⁹³ Nb	30	⁹³ Nb	20	⁹³ Nb	1
⁹⁵ Mo	200	⁹⁵ Mo	100	⁹⁵ Mo	7
¹⁰³ Rh	60	¹⁰³ Rh	20	¹⁰³ Rh	1
¹⁰⁵ Pd	20	¹⁰⁵ Pd	10	¹⁰⁵ Pd	2
¹⁰⁷ Ag	30	¹⁰⁷ Ag	40	¹⁰⁷ Ag	2
¹¹¹ Cd	200	¹¹¹ Cd	300	¹¹¹ Cd	9
¹¹⁵ In	20	¹¹⁵ In	10	¹¹⁵ In	0.6
¹¹⁸ Sn	70	¹¹⁸ Sn	50	¹¹⁸ Sn	4
¹²¹ Sb	40	¹²¹ Sb	40	¹²¹ Sb	2
¹²⁵ Te	2000	¹²⁵ Te	600	¹²⁵ Te	20
¹³³ Cs	n.d.	¹³³ Cs	10	¹³³ Cs	0.8
¹³⁷ Ba	80	¹³⁷ Ba	100	¹³⁷ Ba	5
¹³⁹ La	10	¹³⁹ La	9	¹³⁹ La	0.6
¹⁴⁰ Ce	10	¹⁴⁰ Ce	9	¹⁴⁰ Ce	0.6
¹⁴¹ Pr	9	¹⁴¹ Pr	7	¹⁴¹ Pr	0.4
¹⁴⁶ Nd	50	¹⁴⁶ Nd	40	¹⁴⁶ Nd	2
¹⁴⁷ Sm	50	¹⁴⁷ Sm	50	¹⁴⁷ Sm	3
¹⁵³ Eu	10	¹⁵³ Eu	10	¹⁵³ Eu	0.6
¹⁵⁷ Gd	40	¹⁵⁷ Gd	40	¹⁵⁷ Gd	2
¹⁵⁹ Tb	6	¹⁵⁹ Tb	6	¹⁵⁹ Tb	0.3
¹⁶³ Dy	20	¹⁶³ Dy	30	¹⁶³ Dy	1
¹⁶⁵ Ho	5	¹⁶⁵ Ho	6	¹⁶⁵ Ho	0.3
¹⁶⁶ Er	20	¹⁶⁶ Er	20	¹⁶⁶ Er	0.8
¹⁶⁹ Tm	5	¹⁶⁹ Tm	6	¹⁶⁹ Tm	0.3
¹⁷² Yb	20	¹⁷² Yb	30	¹⁷² Yb	1
¹⁷⁵ Lu	4	¹⁷⁵ Lu	5	¹⁷⁵ Lu	0.2
¹⁷⁸ Hf	10	¹⁷⁸ Hf	20	¹⁷⁸ Hf	1
¹⁸¹ Ta	3	¹⁸¹ Ta	4	¹⁸¹ Ta	0.4
¹⁸² W	10	¹⁸² W	20	¹⁸² W	1
¹⁸⁵ Re	10	¹⁸⁵ Re	10	¹⁸⁵ Re	0.7
¹⁹⁵ Pt	30	¹⁹⁵ Pt	20	¹⁹⁵ Pt	2
¹⁹⁷ Au	10	¹⁹⁷ Au	10	¹⁹⁷ Au	0.9
²⁰⁵ Tl	5	²⁰⁵ Tl	10	²⁰⁵ Tl	0.4
²⁰⁸ Pb	9	²⁰⁸ Pb	10	²⁰⁸ Pb	0.7
²⁰⁹ Bi	4	²⁰⁹ Bi	8	²⁰⁹ Bi	0.4
²³² Th	3	²³² Th	4	²³² Th	0.3
²³⁸ U	2	²³⁸ U	3	²³⁸ U	0.2

n.d. - not determined. Signal intensities for ³⁴S and ¹³³Cs were not detected above baseline levels within the section of the garnet image considered for evaluation.

Table S2.**LA-ICP-MS spot analyses of MG1 garnet. Mass fractions are given in µg/g.**

Label	position	Distance rim-core (µm)	La	Ce	Pr	Nd	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
MG1_109GTE15	inner core	1940	<0.01	<0.03	0.005	0.23	0.83	0.13	5.0	2.8	40	12	42	6.0	36	4.3
MG1_092GTE1	inner core	1660	<0.01	<0.03	0.014	0.41	0.78	0.13	5.2	2.9	38	11	34	4.6	27	3.5
MG1_095GTE3*	outer core	1270	<0.01	<0.03	0.008	0.42	1.2	0.11	5.4	2.0	15	2.4	4.3	0.40	1.9	0.20
MG1_098GTE6	outer core	880	<0.01	<0.03	0.015	0.59	1.7	0.10	9.1	2.7	15	2.0	3.5	0.30	1.4	0.18
MG1_103GTE10	mantle	470	<0.01	<0.03	0.037	1.08	3.9	0.15	17	4.5	35	7.5	23	3.5	24	3.6
MG1_106GTE13	rim	50	1.8	3.0	0.29	0.96	0.39	0.38	1.4	0.5	4.9	1.1	2.9	0.38	2.5	0.32

* from Hermann and Rubatto, 2003.

Trace element analyses were performed by Laser Ablation–ICP-MS at the Research School of Earth Sciences, Australian National University, using a pulsed 193 nm ArF Excimer laser with 100 mJ energy at a repetition rate of 5 Hz coupled to an Agilent 7500 quadrupole ICP-MS. During the time-resolved analysis of minerals, the contamination from inclusions, fractures and zones of different composition was detected by monitoring several elements and integrating only the relevant part of the signal. Spot size was 60 µm in diameter. External calibration was performed relative to NIST 612 glass using the concentrations given in Pearce et al. (1997). The internal standard was Ca mass fraction measured by EPMA. Accuracy was assessed to be better than 10% for all elements by analysing a Columbia River Basalt.

Pearce NJG, Perkins WT, Westgate JA, Gorton MP, Jackson SE, Neal CR, Chenery SP (1997) A compilation of new and published major and trace element data for NIST SRM 610 and NIST SRM 612 glass reference materials. *Geostandards Newsletter-the Journal of Geostandards and Geoanalysis* 21(1):115-144

Supplementary Material

Figures and diffusion

Identification of growth mechanisms in metamorphic garnet by high-resolution trace element mapping with LA-ICP-TOFMS

Daniela Rubatto, Marcel Burger, Pierre Lanari, Bodo Hattendorf, Gunnar Schwarz, Christoph Neff, Peter Keresztes Schmidt, Jörg Hermann, Alice Vho, Detlef Günther

2020 – Contributions to Mineralogy and Petrology

Fig. S1. Photomicrograph of the samples in plain polarized light: (a) CLM6 amphibolite-facies metapelite, (b) MG1 granulite-facies metapelite and (c) eclogite AV16021. The images show the main rock texture, mineral assemblage and the position of the analysed garnet.

CLM6

MG1

Fig. S2. EPMA MgO and FeO maps for garnet MG1 and CLM6.

Fig. S3. LA-ICP-TOFMS maps of garnet CLM6 for Cr, Ge and V, and map of garnet MG1 for Ge.

Fig. S4. Divariant diagrams showing the mass fraction of Sc, V, Y and Er versus that of Mg in an area of eclogitic garnet AV that is rich in secondary veinlets (area marked by dotted red line in the Mg map at the top). The lack of correlation between the trace elements and Mg is in favour of a selective replacement process (see text for discussion).

Diffusion models of Y+REE in MG1 garnet

Formalism

The diffusion of Y+REE was treated as single-component diffusion of elements present in trace concentrations in a homogeneous medium along a one-dimensional semi-infinite profile. As the characteristic diffusion distance (300–400 μm) in this example is much smaller than the grain diameter (~ 3 mm), the spherical geometry of the crystal was not considered. The model was kept simple in order to achieve order-of-magnitude estimates. The rate of accumulation (or depletion) was assumed to be dictated by the Fick's 2nd law of diffusion:

$$\frac{\partial C}{\partial t} = D \frac{\partial^2 C}{\partial x^2} \quad (1)$$

Here, the composition C is expressed in $\mu\text{g/g}$. The diffusion coefficient D (in m^2s^{-1}) is T and P dependent and defined as $\log_{10} D$ via

$$\log D = \log D_0 + \frac{-(E_a + P\Delta V)}{2.303RT} \quad (2)$$

with D_0 the pre-exponential factor (in m^2s^{-1}), E_a the activation energy for diffusion (in J mol^{-1}), P the pressure (in bar), ΔV the activation volume ($\text{J bar}^{-1}\text{mol}^{-1}$), R the gas constant (8.31 J mol^{-1}) and T the temperature (in Kelvin).

Fig. S5. Extraction of compositional profiles (e.g. for Y) for trace element diffusion modelling using the tool “integrated lines (strip)” in XMAPTOOLS. (a) Map showing the strip position (purple); (b) the individual profiles are shown in black and the average profile in purple.

Compositional profile, initial geometry and fitting procedure

Representative compositional profiles for diffusion modelling in garnet MG1 were obtained using the tool “integrated lines (strip)” implemented in XMAPTOOLS 3.3.1 (Lanari, et al. 2019; Lanari, et al. 2014). This technique is based on the integration of parallel compositional profiles defining a rectangle, which in this case was located perpendicular to the core-mantle interface (Figure S5a). The average profile is smoother than individual profiles and therefore more suitable for diffusion modelling (Figure S6). The average approach also lowers the LOD for each element compared to the single pixel LOD (Table S1). The compositional profiles of Y, Lu, Ho, Dy and Er were extracted from the same area.

The initial profile for diffusion was generated for each element using a step function; assuming that the compositions in the core and mantle were flat on both side of the interface (*initial profile* in Fig. S6). This jump in composition was centred on the position 342.5 μm set on the basis of Y. This value was kept constant for every element.

Fig. S6. Diffusion modelling over a period of 10 My for Y. The black spots are the points taken from the average profile (shown in Fig. S5a). The red curve shows the original profile for diffusion modelling defined as a step function between mantle (left) and core (right). The final compositional profile after 10 My is shown in blue; this profile was obtained using the diffusion coefficient expressed in $\log D$. * Fitted (see text).

The compositional profile after diffusion for a total duration of 10 My divided into 3000 time steps was obtained by solving Equation (1) using a finite difference scheme and a backward time, centred space implicit method. The value of D was obtained by non-linear optimization as the value minimizing the sum of the point-model distances for the n points of the compositional profile defined as

$$\text{minimize } \sum_{i=1}^n (C_i^{ref} - C_i^{model})^2 \quad (3)$$

with C_i^{ref} the observed composition and C_i^{model} the model compositions of point i expressed in ppm. As boundary conditions, the compositions at each end of the profile C_1^{model} and C_n^{model} were kept constant (red spots in Figure S2). The inversion allows a value of D to be determined for each element considered (Figure S3; other plots are given in the paper).

The same procedure was repeated for two additional profiles with different locations across the core-mantle boundary. Results are reported in in Table S3 and for Y in Figure S8. The obtained diffusion coefficients (in $\log D$) are within 0.1 log unit for Y, Dy and Er and within 0.3 log unit for Lu, Ho.

Fig. S7. Diffusion modeling over a period of 10 My for Dy and Er. Other elements are shown in the paper.

Fig. S8. Diffusion modeling over a period of 10 My for Y in two additional locations across the core-mantle boundary (from left to right in the profile) of garnet MG1.

	<i>Profile 1 (T2)</i>	<i>Profile 2 (T3s)</i>	<i>Profile 3 (T4)</i>
Y	-23.0 ($R^2=0.99$)	-23.0 (0.99)	-23.1 (0.99)
Lu	-22.8 (0.93)	-22.8 (0.87)	-23.1 (0.87)
Ho	-23.3 (0.96)	-23.1 (0.92)	-23.4 (0.95)
Dy	-23.1 (0.92)	-23.1 (0.86)	-23.2 (0.89)
Er	-23.1 (0.96)	-23.1 (0.91)	-23.2 (0.96)

Table S3. Diffusion coefficients expressed in $\log D$ and R^2 values obtained from three diffusion profiles. The reference given within parentheses for each profile is an internal reference corresponding to the fitting experiment.

References

- Lanari P, Vho A, Bovay T, Airaghi L, Centrella S (2019) Quantitative compositional mapping of mineral phases by electron probe micro-analyser. In: Ferrero S, Lanari P, Goncalves P, Grosch EG (eds) *Metamorphic Geology: Microscale to Mountain Belts*, vol 478. Geological Society of London, Special Publications, pp 39-63
- Lanari P, Vidal O, De Andrade V, Dubacq B, Lewin E, Grosch EG, Schwartz S (2014) XMapTools: A MATLAB©-based program for electron microprobe X-ray image processing and geothermobarometry. *Computers & Geosciences* 62:227-240 doi:10.1016/j.cageo.2013.08.010