

COVID-19 in Healthcare Workers: A Living Systematic Review and Meta-analysis of Prevalence, Risk Factors, Clinical Characteristics, and Outcomes

Sergio Alejandro Gómez-Ochoa, Oscar H. Franco, Lyda Z. Rojas, Peter Francis Raguindin, Zayne Milena Roa-Díaz, Beatrice Minder Wyssmann, Sandra Lucrecia Romero Guevara, Luis Eduardo Echeverría, Marija Glisic*, and Taulant Muka*

Correspondence to Dr. Taulant Muka, MD, MPH, PhD; Institute of Social and Preventive Medicine (ISPM), University of Bern, Mittelstrasse 43, 3012, Bern, Switzerland (e-mail: taulant.muka@ispm.unibe.ch)

Author affiliations:

Research Division, Cardiovascular Foundation of Colombia, Floridablanca, Colombia (Sergio Alejandro Gómez-Ochoa and Lyda Z. Rojas); Public Health and Epidemiological Studies Group, Cardiovascular Foundation of Colombia, Floridablanca, Colombia (Sergio Alejandro Gómez-Ochoa); Institute of Social and Preventive Medicine (ISPM), University of Bern, Bern, Switzerland (Oscar H. Franco, Peter Francis Raguindin, Zayne Milena Roa-Díaz, Marija Glisic and Taulant Muka); Institute of Child Health and Human Development (ICHHD), National Institutes of Health, University of the Philippines Manila, Manila, Philippines (Peter Francis Raguindin); Research Group and Development of Nursing Knowledge (GIDCEN-FCV), Research Institute, Cardiovascular Foundation of Colombia, Floridablanca, Santander, Colombia (Lyda Z. Rojas and Zayne Milena Roa-Díaz); Public Health & Primary Care Library, University Library of Bern, University of Bern, Bern, Switzerland (Beatrice Minder Wyssmann); Nursing School, Universidad Industrial de Santander, Bucaramanga, Colombia (Sandra Lucrecia Romero Guevara); Chief, Heart Failure Clinic, Cardiovascular Foundation of Colombia, Floridablanca, Colombia (Luis Eduardo Echeverría); Swiss Paraplegic Research, Nottwil, Switzerland (Marija Glisic).

*denotes equal contribution

Funding: None.

Conflict of interest: None declared.

Running head: COVID-19 in Health Care Workers

ABSTRACT

Health care workers (HCW) are at the frontline response to the new coronavirus disease 2019 (COVID-19), being at a higher risk of acquiring the disease, and subsequently, exposing patients and colleagues. Searches in eight bibliographic databases were performed to systematically review the evidence on the prevalence, risk factors, clinical characteristics, and prognosis of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection among HCW. Ninety-seven studies (All published in 2020), including 230,398 HCW, met the inclusion criteria. From the screened HCW using RT-PCR and the presence of antibodies, the estimated prevalence of SARS-CoV-2 infection was 11% (95%CI; 7%-15%) and 7% (95% CI; 4%-11%), respectively. The most frequently affected personnel were the nurses (48%. 95%CI; 41%-56%), while most of the COVID-19 positive medical personnel were working in hospitalization/non-emergency wards during the screening (43%, 95%CI;28%-59%). Anosmia, fever and myalgia were identified as the only symptoms associated with HCW SARS-CoV-2 positivity. Among RT-PCR positive HCW, 40% (95%CI;17%-65%) did not show symptoms at the time of diagnosis. Finally, 5% (95%CI;3%-8%) of the COVID-19 positive HCW developed severe clinical complications, and 0.5% (95% CI; 0.02%-1.3%) died. HCW suffer a significant burden from COVID-19, with HCW working in hospitalization/non-emergency wards and nurses being the most infected personnel.

Keywords: 2019-nCoV; SARS-CoV-2; COVID-19; Health Care Workers; Medical Workers

Abbreviations

COVID-19: Coronavirus disease 2019

SARS-CoV-2: Severe acute respiratory syndrome coronavirus 2

HCW: Healthcare workers

RT-PCR: Reverse transcription polymerase chain reaction

PPE: Personal protective equipment

HCQ: Hydroxychloroquine

INTRODUCTION

The pandemic of Coronavirus disease 2019 (COVID-19) caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) virus has already caused more than 14 million infections and 600 thousand deaths globally(1). Although SARS-CoV-2 infection has a lower mortality rate compared to infections caused by the severe acute respiratory syndrome (SARS) virus or Middle East respiratory syndrome (MERS) virus, its long incubation period and lower virulence have resulted in a large number of asymptomatic carriers(2). Several studies have shown that asymptomatic carriers contribute substantially to the spread of the virus, even by merely breathing in a room(3–5). Among asymptomatic carriers and individuals at risk due to asymptomatic coronavirus transmission, health care workers (HCW) represent an important but yet understudied population(6). HCW may experience an increased risk of SARS-CoV-2 infection due to their close contact with highly infectious patients, but also due to exposure to undiagnosed or subclinical infectious cases. This could be even more problematic, considering the poor access to personal protective equipment (PPE) worldwide(7). A recent report of the Centers for Disease Control and Prevention (CDC) shows that, as of April 9th, 2020, 9,282 known coronavirus disease 2019 (COVID-19) cases were labeled as HCW in the U.S, with numbers probably underestimated (8).

Currently, there is no clarity regarding the prevalence of SARS-CoV-2 infection among HCW according to specific clinical settings, limiting the possibility of designing effective preventive measures to limit the transmission of the virus within a hospital, and from hospitals to the community(9,10). Furthermore, it is unknown whether the clinical characteristics and outcomes of HCW may be different from those of the general population, considering that the repeated exposure to the virus may lead to higher SARS-CoV-2 viral load and therefore to worse clinical outcomes(11,12). Therefore, characterizing SARS-CoV-2 infection within health-care workers is critical for achieving optimal control of the pandemic. The present systematic review and meta-analysis aimed to identify, analyze, and quantify the prevalence, risk factors, clinical characteristics, and outcomes of COVID-19 among HCW.

METHODS

This systematic review and meta-analysis was conducted following a recently published guideline on how to perform a systematic review and reported following the PRISMA guidelines (**Web Table 1**) (13,14).

Data source and strategy

MEDLINE, EMBASE, LILACS, Cochrane, Web of Science, WHO COVID-19 database, Google scholar and 'Living Evidence on COVID-19', a database developed by the University of Bern (ISPM), were searched to identify relevant articles from inception until July 8th, 2020 without language restrictions. The following search terms related to the COVID-19 infection in HCW were used: coronavirus disease 2019, coronaviridae, SARS-

CoV-2, SARS coronavirus, 2019-nCov, prevalence, screening, clinical characteristics, clinical course, severity of illness, outcomes, among others. We limited our search to human studies, with no language restriction. The complete search strategy is described in the **Web Appendix 1**.

Study selection and eligibility criteria

All observational studies (e.g., cross-sectional, cohort, case-control studies, and case-series), except for case reports, were included. We included studies that reported the prevalence of COVID-19 in HCW by using either RT-PCR or a serum antibodies assay. We also included studies evaluating the risk factors for SARS-CoV-2 infection and those analyzing the clinical characteristics and outcomes of laboratory-confirmed COVID-19 among HCW. We excluded those articles that evaluated HCW with suspected but not laboratory-confirmed SARS-CoV-2 infection. Two independent reviewers screened the titles and abstracts according to the selection criteria.

Methods on Data extraction, Quality assessment and Living systematic review can be found in the Web Material (Web Appendix 2).

Data synthesis and analysis

Based on the extracted data of each study (performed by two independent investigators), we first estimated the global prevalence of SARS-CoV-2 infection by each test used for

screening (RT-PCR vs. antibodies tests). Before pooling, all proportions were transformed using the Freeman-Tukey Double Arcsine method. For dichotomous risk factors, results were expressed as ORs with 95% confidence intervals (CI). Heterogeneity of results was assessed using the I^2 measure of inconsistency; however, regardless of heterogeneity, random-effects models were chosen for all the analyses. Screening criteria, geographical location, HCW professions, the clinical setting, and the mean of daily new cases of SARS-CoV-2 per million inhabitants (obtained from the European Centers of Disease Control website. <https://www.ecdc.europa.eu/en>) in the country during the period in which the study was carried out were pre-specified as characteristics for assessment of heterogeneity and were evaluated using stratified analyses and univariate random-effects meta-regression. All analyses were conducted using STATA 15.1 (Statacorp, Texas, US, 2017). For main analysis, a p-value <0.05 was considered significant. To account for multiple testing, in the stratified analysis, we considered a p-value of 0.01 as significant.

RESULTS

The initial search yielded 4,107 studies, from which ninety-seven studies met the inclusion criteria (**Figure 1**). A total of 230,398 participants were evaluated in the included studies, mostly women (69.98%) and with a mean age of 40 ± 11 years (8,15–58). Of the 97 studies, 70 studies reported data regarding the prevalence of SARS-CoV-2 infection in HCW (total screened HCW: 96,813) (15–18,22,24–26,28,30,33–35,35–39,41,42,42,43,49–98), 38 studies analyzed the clinical characteristics of infected medical workers ($n=32,144$) (8,19–21,23,27,29–32,34,44–46,50,52,57,59,64,65,68,71,74,77–80,93,99,99–107) and, thirteen

studies evaluated risk factors for COVID-19 positivity among HCW(24,34,44,48,57,63,65,67,70,89,93,108,109).

SARS-CoV-2 infection prevalence in HCW using RT-PCR

Forty-six studies evaluated the prevalence of SARS-CoV-2 infection among HCW using RT-PCR. Of those, 31 studies were based in clinical facilities in Europe, nine in the USA, and six in Asia (15,16,18,22,24,26,28,30,34–36,38,39,41,43,44,50–54,57–59,62,64–68,70,71,74,75,78–84,86,87,92–95,110) (**Figure 2**). The prevalence of SARS-CoV-2 infection ranged from 0.4% in the study of *Olalla et al* among 498 Spanish HCW to 57,06% in the study of *Breazzano et al*, carried out in New York city. Among 75,859 HCW screened for COVID-19 using RT-PCR, the estimated pooled prevalence of SARS-CoV-2 infection was 11% (95% CI; 7%-15%; p-value for heterogeneity <0.001, I^2 : 98%) (**Figure 3, Web Figure 1**). Furthermore, the prevalence among symptomatic HCW was the highest (19%. 95% CI; 12%-28%; p-value for heterogeneity <0.001, I^2 : 99%), followed by the one observed in studies including both symptomatic and asymptomatic individuals (8%. 95% CI; 3%-16%; p-value for heterogeneity <0.001, I^2 : 99%). Finally, asymptomatic HCW showed the lowest prevalence of SARS-CoV-2 infection (5%. 95% CI; 1%-13%; p-value for heterogeneity <0.001, I^2 : 98%) (**Figure 3, Web Figure 1, Table 1**). Among HCW with positive results, 48% (95% CI; 41%-56%; p-value for heterogeneity <0.001, I^2 : 98%) were nurses, followed by physicians (25%, 95% CI; 16%-35%; p-value for heterogeneity <0.001, I^2 : 99%) and other HCW (23%, 95% CI; 12%-36%; p-value for heterogeneity <0.001, I^2 : 99%). Most of the SARS-CoV-2 positive personnel were working in hospitalization/non-emergency wards during the screenings (43 %, 95% CI; 28%-59%; p-value for

heterogeneity <0.001 . I^2 : 91%), followed by the operating rooms and surgery services (24%, 95% CI: 17% - 31%; p-value for heterogeneity: 0.05, I^2 : 60%) (15,16,20,22,24–27,38,39,41–44,47,49,50,52–54,59–63,66,67,73,74,80,82–84,94,103,105,106) (Web Figure 2 and Table 2).

Stratification analysis, supported by meta-regression analysis, showed that the symptoms criteria (p-value=0.002) for performing SARS-CoV-2 RT-PCR screenings was significantly associated with SARS-CoV-2 infection prevalence among HCW (Web Figure 3), while no role of other factors was observed (Web Figures 4 and Web Table 2).

Prevalence of antibodies against SARS-CoV-2 in HCW

Twenty-eight studies evaluated the prevalence of antibodies against SARS-COV-2 in HCW using serum antibody tests. The data regarding the sensitivity and specificity of the antibody detection kits used in each study are available in Web Table 3, with sensitivity ranging from 75% to 100%, and specificity being 80% and above. Among 27,445 HCW screened for the presence of antibodies, a pooled infection prevalence of 7% (95% CI: 4%-11%; p-value for heterogeneity < 0.001 . I^2 : 99%) was estimated. The prevalence of COVID-19 infection was similar after comparing studies regarding their screening criteria (p-value=0.543) and the mean of new daily cases of SARS-CoV-2 infection per million inhabitants in the country during the previous 2 weeks to the study initiation (p-value=0.787). (Web Figure 5).

Prevalence of COVID-19 in HCW after exposure to undiagnosed infected patients

Eight studies analyzed the scenario of direct exposure of HCW to an individual or a group of SARS-CoV-2 infected patients without knowing their infection status. The studies comprised a total of 1126 HCW screened after in-hospital exposure, highlighting the lack of PPE use in 57.37% (n=646) of the exposed workers. From the studies including HCW without proper PPE use, 4.7% (n=28) of the exposed individuals had a positive RT-PCR or antibody test result during the contact tracings. On the other hand, there was no single case of COVID-19 attributed to the exposure to the index case in the studies that included HCW with proper PPE use. In line with this trend, the study of *Chen et al* reported that adequate PPE use was associated with a reduced risk of seroconversion in HCW exposed to COVID-19 patients (OR, 0.127, 95% CI 0.017, 0.968)(42,49,60,63,69,71,75,91).

Clinical characteristics and outcomes of COVID-19 positive HCW

Thirty-seven studies, based on 31,866 COVID-19 positive HCW (69% women, mean age: 40.1 ± 12.33), evaluated the clinical characteristics or outcomes of HCW infected by the SARS-CoV-2(8,19,19–21,23,27,29–32,32,34,44–46,46,50,52,57,59,64,65,68,71,74,77–80,93,99–107). Among the 11,772 positive HCW with data regarding comorbidities, the pooled prevalence of hypertension, cardiovascular disease, type 2 diabetes and chronic obstructive pulmonary disease was 7% (95% CI; 4%-10%; p-value for heterogeneity: 0.35, I^2 : 10%), 3% (95% CI; 1%-8%; p-value for heterogeneity: 0.19, I^2 : 39%), 4% (95% CI; 2%-7%; p-value for heterogeneity: 0.01, I^2 : 63%) and 3% (95% CI; 1%-6%; p-value for heterogeneity: 0.01, I^2 : 77%), respectively. Furthermore, based on data from 15 studies,

including 12,089 HCW, the pooled prevalence of individuals diagnosed by RT-PCR that did not show symptoms at time of diagnosis was 40% (95% CI 17%-65%. p-value for heterogeneity < 0.001. I^2 : 99%)(16,24,38,50,51,59,62,71,74,80,83,84,102,110,111) (**Web Figure 6**).

Among symptomatic COVID-19 HCW, the most frequently reported symptoms were fever (57%, 95% CI; 50%-64%. 29 studies. p-value for heterogeneity < 0.001, I^2 : 96%) and dry cough (57%, 95% CI; 50%-65%. 26 studies. p-value for heterogeneity < 0.001, I^2 : 97%), followed by malaise (43%, 95% CI; 26%-61%. 1 studies. p-value for heterogeneity < 0.001, I^2 : 96%) and myalgia (48%, 95% CI; 35%-62%. 10 studies. p-value for heterogeneity < 0.001, I^2 : 92%) (**Table 3**). Finally, eight studies reported the severity of the disease, including ICU admission(8,20,23,27,29,32,45,50). Among these, a pooled prevalence of 5% (95% CI; 3%-8%. p-value for heterogeneity < 0.001, I^2 : 95%) for severe disease was estimated. Finally, eleven studies provided information regarding mortality in this population(8,19,23,32,50). An estimated 0.5% (95% CI; 0.02%-1.3%. p-value for heterogeneity < 0.001. I^2 : 96%) of the total HCW reported as infected by the SARS-CoV-2 died because of complications of the disease.

Factors associated with SARS-CoV-2 infection in HCW

Fifteen studies analyzed the factors potentially associated with SARS-CoV-2 infection in HCW. From these, four studies provided sufficient data from infected and non-infected HCW to perform a meta-analysis for anosmia and five for fever, finding a significantly

higher risk of COVID-19 with the presence of these symptoms (OR 28.37; 95%CI 9.45-85.16. p-value for heterogeneity = 0.002, I^2 : 79% for anosmia and OR 4.86; 95%CI 2.83-8.37 p-value for heterogeneity < 0.001, I^2 : 84% for fever) (**Web Table 4**). Furthermore, the study of *Rudberg et al* also found a significantly higher risk of a positive RT-PCR result in HCW with anosmia (OR 28.43; $p < 0.001$) or fever (OR 6.27; $p < 0.001$).

A similar result was observed for myalgia, a symptom that was significantly associated with SARS-CoV-2 infection (OR 3.06; 95% CI 1.24-7.56. p-value for heterogeneity = 0.001, I^2 : 86%. three studies). Finally, the study of *Clemency et al* evaluated the predictive value of different symptoms for COVID-19 diagnosis, highlighting the loss of taste and smell, with a positive predictive value (PPV) of 0.5 and a negative predictive value (NPV) of 0.85, fever, with a PPV of 0.31 and a NPV of 0.83, and myalgia, with a PPV of 0.27 and a NPV of 0.80. On the other hand, no significant associations were found for fatigue (5 studies. OR 2.41. 95% CI; 0.92, 6.27. p-value for heterogeneity < 0.001, I^2 : 92%)(24,44,57) and sore throat (4 studies. OR 0.55. 95% CI; 0.30, 1.01. p-value for heterogeneity: 0.02, I^2 : 75%) (**Web Table 4**)(44,57).

Ran et al. found that, among seventy-two HCWs in Wuhan, China, unqualified hand-washing (OR 2.64; 1.04-6.71), suboptimal hand hygiene before, patient contact (OR 3.10; 1.43-6.73.), and inadequate PPE (OR 2.82; 1.11-7.18) were risk factors for SARS-CoV-2 infection(34). Similarly, the study of *Wang X et al.* reported that in a sample of 493 HCW in Wuhan, China, the risk of COVID-19 in HCW using medical masks was significantly

higher when compared to those using N95 respirators (OR: 464.82, (95% CI:97.73-infinite), despite this last group had a significantly higher exposure to infected patients (48). *Chatterjee et al* reported similar results, highlighting a higher risk of SARS-CoV-2 infection in HCW that never used PPE compared to those with usual protection usage (OR 3.72; 95% CI 2.12-6.52).

Moreover, the risk of aerosol-generating procedures for HCW was analyzed by the prospective multi-center study by *El-Boghdadly et al*, which evaluated 1718 HCW participating in tracheal intubation of patients with suspected or confirmed COVID-19. The overall incidence of the composite outcome (new laboratory-confirmed COVID-19 or new COVID-19 symptoms requiring self-isolation or hospitalization) was 10.7% over a median follow-up of 32 (18–48) days. Furthermore, 10% of the HCW involved in tracheal intubation procedures in these cohort had subsequently the composite outcome(109).

The potential source of infection was studied by *García Basteiro et al*, observing that HCW with a larger household size tended to have more frequently detectable antibodies (IgM or IgA) against SARS-CoV-2; albeit non-significant ($p=0.093$)(24). The study of *Kluytmans et al*. reported that among 84 SARS-CoV-2 positive HCW in two hospitals in The Netherlands, only three percent reported having been exposed to an inpatient with confirmed COVID-19 before symptoms onset(30). Finally, *Sikkema et al.*, by investigating the genome sequences from infected HCW and patients, reported that the obtained patterns

were consistent with multiple introductions into the hospitals through community-acquired infections and local amplification of the viral disease in the community context(93).

Interestingly, two studies evaluated the benefit of pharmacological prophylaxis with hydroxychloroquine (HCQ) to prevent SARS-CoV-2 infection among high-risk HCW, with both studies showing that the history of having taken maintenance doses of HCQ was associated with a significantly lower risk of COVID-19(70,108).

Study Quality

The majority of studies were of moderate (n=61 [62.9%]) quality, with 29.9% (n=29) being high quality, while of the rest of low quality (n=7, 7.2%). Web **Tables 5, 6, and 7** present a summary of the studies' quality evaluation.

Discussion

The current evidence shows that around a tenth of the total HCW in the screened hospitals have a diagnosis of acute SARS-CoV-2 infection, with half being nurses. To date, only 7% of the HCW have resulted to be positive for the presence of antibodies indicating SARS-CoV-2 infection. Furthermore, most of the SARS-CoV-2 positive personnel were working in hospitalization/non-emergency wards during the laboratory screenings. Fever, anosmia and myalgia were the main associated factors for SARS-CoV-2 infection in the meta-analysis. From the fifteen studies that screened HCW irrespective of their symptoms and reported the clinical features of SARS-CoV-2 positive individuals, 40% did not report any symptom compatible with COVID-19 during the screenings. Furthermore, we observed a

pooled prevalence of severe COVID-19 of 5% among HCW, while 0.5% of the infected HCW died because of complications of the disease.

Our findings show a higher prevalence of SARS-CoV-2 infection among HCW when compared to the data from the general population reported in the literature(112). This difference may be attributable to workplace exposures of the HCW; nevertheless, only a few studies analyzed the potential source of infection in this population, limiting the possibility of evaluating the impact of nosocomial vs. community-acquired infection. The study of *Folgueira et al.* reported that there were no significant differences in the infection rates between the groups of HCW working in high, intermediate, and low exposure risk settings(22). Furthermore, the study of *Hunter et al.* found no differences in the proportion of infected HCW when comparing the ones with patient-facing roles with those without this exposure(26). In addition, the results of the studies of *García-Basteiro et al*, *Kluytmans et al*, and *Sikkema et al* provided evidence suggesting a relevant role of community transmission of the disease in HCW infections. These results may suggest that household contacts may play a significant role in SARS-CoV-2 infection in HCW, mainly due to the rapid circulation of the virus in the community. Another reason could be the infection from asymptomatic carriers, considering that about a half of the SARS-CoV-2 infected HCW were asymptomatic during the screenings. However, the importance of nosocomial transmission needs to be analyzed in light of the use of PPE and other measures designed to reduce the exposure of HCW in the work settings, but that are not usually applied by the health personnel in low-risk settings or the community context(113).

As new strategies have been designed and implemented for the “re-opening” of the economic activities in different countries, the understanding of the role of asymptomatic transmission of SARS-CoV-2 is essential, especially in a clinical setting. According to WHO, at least 50% of patients dying from COVID-19 were residents in hospitals or nursing homes, highlighting the need to control the spread of infection in a health care setting. Our findings suggest that almost half of SARS-CoV-2 infected HCW are asymptomatic during the screenings. However, the contribution of asymptomatic carriers in the transmission of SARS-CoV-2 infection is still not clear. A recent study showed similar viral loads in symptomatic patients compared to asymptomatic individuals, highlighting the transmission potential of SARS-CoV-2 carriers despite their clinical status(114). On the other hand, the study of *Gao et al.* suggested potentially low infectivity of asymptomatic SARS-CoV-2 carriers, as none of the 455 contacts who were exposed to an asymptomatic COVID-19 virus carrier had a positive test(73). These results were in-line with the studies performed by *Ng et al.* and *Canova et al.*, in which none of the exposed HCW had a subsequent positive SARS-CoV-2 test(42,49). Nevertheless, the lack of infection among HCW in these studies could also be due to adequate use of PPE, hand hygiene, and other standard procedures. Indeed, all asymptomatic patients investigated in the study by *Gao et al.* were wearing a mask, reducing further the spread of the infection. Nonetheless, even if this low infectivity is confirmed by further studies, the potential of silent transmission still represents an enormous issue that needs to be addressed efficiently, especially in low and middle-income countries lacking medical resources (e.g. PPE and diagnostic capacity). Considering the results of the present study, in which almost half of the positive HCW by RT-PCR were asymptomatic, there is an urgent need to promote a process of continuous, systematic screening of all HCW in high-risk settings, the use of adequate PPE and other

standard procedures. Moreover, a low threshold for suspicion of infection in low-risk settings is also needed to promote early isolation to avoid cross-infection(57).

To improve the screening performance and early detection of SARS-CoV-2 infection among HCW, the analysis of risk factors for COVID-19 positivity is of importance. In the present review, fifteen studies analyzed the symptoms and signs associated with SARS-CoV-2 infection among HCW. Fever (OR: 4.86, 95% CI: 2.38-8.37, I^2 : 84%), anosmia (OR: 28.37; 95%CI 9.45-85.16 I^2 : 79%) and myalgia (OR 3.06; 95% CI 1.24-7.56. I^2 : 86%) proved to be associated with higher odds of COVID-19 in symptomatic HCW; however, relying on these specific but not sensitive symptoms to define screening criteria may lead to an important proportion of missed COVID-19 positive cases. This was the conclusion of the study of *Yombi et al.*, which assessed the impact of using fever as a predictor for the positivity of SARS-CoV-2 RT-PCR. Their results showed that fever might have a positive impact on the yield of RT-PCR for SARS-CoV-2; nonetheless, when this symptom was required as a criterion for testing, an important number of positive cases were missed(57). Similar results were reported by *Chow et al.* in a cohort of HCW in King County, Washington. In this study, screening only for fever, cough, shortness of breath, or sore throat might have missed 17% of symptomatic HCW at the time of illness onset. The authors also mention that expanding criteria for symptoms screening to include chills and myalgia may still have missed 10%(21). According to the results of the present systematic review and meta-analysis, malaise (48%) and headache (36%) represent additional common symptoms with a high prevalence among positive HCW (**Table 3**). Therefore, the inclusion of these symptoms in the screening criteria for testing may improve the identification of

SARS-CoV-2 positive individuals and prevent further transmission(21,57). Specifically, screening for these symptoms could be useful in low- and middle-income countries with limited testing capacity. However, screening HCW by symptoms may still miss a significant proportion of COVID-19 cases; therefore, universal screening for all exposed HCW regardless of symptoms should be the standard strategy to reduce transmission of SARS-CoV-2 in a hospital setting.

The high number of nurses being SARS-CoV-2 positive in our study could be explained by the larger time staff nurses usually spend with direct patient care involving tasks performed at the bedside, drug administration, and being the first line of response in case of any patient complications. The fact that more infected HCW were observed in hospitalization/non-emergency wards in the current study may suggest a difference in PPE use across the settings, being the compliance to this measures higher in the emergency departments and ICUs (where the subjective risk perception is higher) compared to non-COVID-19 wards(115–117). However, most of the studies did not report the total number of screened HCW per area/professional category, limiting the representativeness of the studies and precluding the analysis of the prevalence of SARS-CoV-2 infection by unit or per profession. For instance, nurses being the highest number of employees in the health care setting could explain the high nurse SARS-CoV-2 positivity among other positive HCW. If future studies confirm nurses being the most affected personnel and hospitalization/non-emergency wards are associated with higher risk of SARS-CoV-2 infection, then the findings would have important implications for policy makers and

hospital administrators in better planning of resources to reduce SARS-CoV-2 transmission in hospitals.

Finally, our findings highlight the risk of clinical complications and mortality among HCW. We found that COVID-19 among HCW is associated with approximately a 0.5% proportion of fatal cases. Although the mortality among HCW is lower compared to the rate of mortality in the general population reported in the literature, better overall health and care among HCW, different age demography, and other factors could explain these differences. Yet, due to exposure to numerous infected individuals exposures, HCW, if infected, could be characterized by higher viral load which is associated with worst clinical outcomes(12).

Strengths and limitations

To our knowledge, this is the first systematic review quantifying the burden of COVID-19 among HCW. However, limitations in the current study merit careful consideration. First, in our analysis, we included evidence mainly deriving from preprint publications, which are not peer-reviewed. Nevertheless, we assessed their quality, highlighting the potential limitations of each study. Second, the heterogeneity of the included studies represented a challenge when pooling the results; thus, we aimed to overcome this limitation by performing different sub-group analyses when estimating the prevalence. Third, an important limitation of the study was the lack of reporting of the test quality (sensitivity and specificity) in most of the included articles, and therefore the prevalence reported from each study could be under or overestimated depending on the applied test. Fourth, clinical settings and HCW professional categories were determined mainly based on employment, which may not necessarily represent specific exposure levels to COVID-19 positive

patients. Fifth, the potential source of SARS-CoV-2 infection in positive HCW was poorly studied in the included articles, therefore limiting the possibility of analyzing the impact of nosocomial vs. community transmission. Last, most of the studies did not report the level of adherence to preventive measures and PPE use, which is an essential factor affecting the transmission of the virus.

To conclude, HCW represent a population with a significant burden from COVID-19. HCW exhibit a high prevalence of SARS-CoV-2 infection, with a significant proportion of the infected HCW being asymptomatic carriers, a condition that favors silent transmission both in the clinical and community contexts if preventive measures and other standard procedures are not implemented.

Declaration of interests

None to declare

Funding

Not received

References

1. Progressier. The Coronavirus App. <https://coronavirus.app/map>. Published February 1, 2020. Accessed July 8, 2020.
2. Mahase E. COVID-19 has killed more people than SARS and MERS combined, despite lower case fatality rate. *BMJ*. 2020;368.
3. Yu X, Yang R. COVID-19 transmission through asymptomatic carriers is a challenge to containment. *Influenza Other Respir Viruses*. 2020;14(4):474-475
4. Riediker M, Tsai D-H. Estimation of SARS-CoV-2 emissions from non-symptomatic cases. *medRxiv*. 2020. Accessed July 8, 2020.
5. Oran DP, Topol EJ. Prevalence of Asymptomatic SARS-CoV-2 Infection. *Ann Intern Med*. 2020:M20-3012. Available in: <https://www.acpjournals.org/doi/10.7326/M20-3012>
6. Kursumovic E, Lennane S, Cook TM. Deaths in healthcare workers due to COVID-19: the need for robust data and analysis. *Anaesthesia*. 2020;75(8):989-992.
7. Wang X, Zhang X, He J. Challenges to the system of reserve medical supplies for public health emergencies: reflections on the outbreak of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) epidemic in China. *Biosci Trends*. 2020;14(1):3-8.
8. CDC COVID-19 Response Team. Characteristics of Health Care Personnel with COVID-19 - United States, February 12-April 9, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(15):477-81.
9. Wang J, Zhou M, Liu F. Reasons for healthcare workers becoming infected with novel coronavirus disease 2019 (COVID-19) in China. *J Hosp Infect*. 2020;105(1):100-101.
10. The Lancet. COVID-19: protecting health-care workers. *Lancet Lond Engl*. 2020;395(10228):922.
11. Yu X, Sun S, Shi Y, Wang H, Zhao R, Sheng J. SARS-CoV-2 viral load in sputum correlates with risk of COVID-19 progression. *Crit Care*. 2020;24(1):170.
12. Liu Y, Yan L-M, Wan L, Xiang T-X, Le A, Liu J-M, et al. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis*. 2020;20(6):656-7.
13. Moher D, Liberati A, Tetzlaff J, Altman DG, Group TP. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLOS Med*. 2009;6(7):e1000097.
14. Muka T, Glisic M, Milic J, Verhoog S, Bohlius J, Bramer W, et al. A 24-step guide on how to design, conduct, and successfully publish a systematic review and meta-analysis in medical research. *Eur J Epidemiol*. 2020;35(1):49-60.

15. Bai Y, Wang X, Huang Q, Wang H, Gurarie D, Ndeffo-Mbah M, et al. SARS-CoV-2 infection in health care workers: a retrospective analysis and a model study. medRxiv. 2020. doi: 10.1101/2020.03.29.20047159
16. Barrett ES, Horton DB, Roy J, Gennaro ML, Brooks A, Tischfield J, et al. Prevalence of SARS-CoV-2 infection in previously undiagnosed health care workers at the onset of the U.S. COVID-19 epidemic. medRxiv. 2020. doi: 10.1101/2020.04.20.20072470
17. Behrens GM, Cossmann A, Stankov MV, Witte T, Ernst D, Happle C, et al. Perceived versus proven SARS-CoV-2 specific immune responses in health care workers. *Infection*. 2020; 48(4): 631–634.
18. Breazzano MP, Shen J, Abdelhakim AH, Glass LRD, Horowitz JD, Xie SX, et al. Resident physician exposure to novel coronavirus (2019-nCoV, SARS-CoV-2) within New York City during exponential phase of COVID-19 pandemic: Report of the New York City Residency Program Directors COVID-19 Research Group. medRxiv. 2020. doi:10.1101/2020.04.23.20074310
19. Team TNCPERE. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. *Chin J Epidemiol*. 2020;41(02):145-51.
20. Chu J, Yang N, Wei Y, Yue H, Zhang F, Zhao J, et al. Clinical Characteristics of 54 medical staff with COVID-19: A retrospective study in a single center in Wuhan, China. *J Med Virol*. 2020; 92(7):807-813.
21. Chow EJ, Schwartz NG, Tobolowsky FA, Zacks RLT, Huntington-Frazier M, Reddy SC, et al. Symptom Screening at Illness Onset of Health Care Personnel With SARS-CoV-2 Infection in King County, Washington. *JAMA*. 2020;323(20):2087-2089.
22. Folgueira MD, Munoz-Ruipérez C, Alonso-Lopez MA, Delgado R. SARS-CoV-2 infection in Health Care Workers in a large public hospital in Madrid, Spain, during March 2020. medRxiv. 27 de abril de 2020. doi: 10.1101/2020.04.07.20055723
23. Gao W, Sanna M, Tsai MK, Wen CP. Geo-temporal distribution of 1,688 Chinese healthcare workers infected with COVID-19 in severe conditions—A secondary data analysis. *PLOS ONE*. 2020;15(5):e0233255.
24. Garcia-Basteiro AL, Moncunill G, Tortajada M, Vidal M, Guinovart C, Jimenez A, et al. Seroprevalence of antibodies against SARS-CoV-2 among health care workers in a large Spanish reference hospital. *Nat Commun* 11, 3500 (2020).
25. Hains DS, Schwaderer AL, Carroll AE, Starr MC, Wilson AC, Amanat F, et al. Asymptomatic Seroconversion of Immunoglobulins to SARS-CoV-2 in a Pediatric Dialysis Unit. *JAMA*. 2020;323(23): 2424–2425.
26. Hunter E, Price DA, Murphy E, van der Loeff IS, Baker KF, Lendrem D, et al. First experience of COVID-19 screening of health-care workers in England. *Lancet Lond Engl*. 2020;395(10234):e77-8.

27. Jin Y-H, Huang Q, Wang Y-Y, Zeng X-T, Luo L-S, Pan Z-Y, et al. Perceived infection transmission routes, infection control practices, psychosocial changes, and management of COVID-19 infected healthcare workers in a tertiary acute care hospital in Wuhan: a cross-sectional survey. *Mil Med Res.* 2020;7(1):24.
28. Keeley AJ, Evans C, Colton H, Ankcorn M, Cope A, State A, et al. Roll-out of SARS-CoV-2 testing for healthcare workers at a large NHS Foundation Trust in the United Kingdom, March 2020. *Eurosurveillance.* 2020;25(14):2000433.
29. Liu J, Ouyang L, Guo P, Wu H sheng, Fu P, Chen Y liang, et al. Epidemiological, Clinical Characteristics and Outcome of Medical Staff Infected with COVID-19 in Wuhan, China: A Retrospective Case Series Analysis. *medRxiv.* 2020. doi: 10.1101/2020.03.09.20033118.
30. Kluytmans M, Buiting A, Pas S, Bentvelsen R, Bijlgaardt W van den, Oudheusden A van, et al. SARS-CoV-2 infection in 86 healthcare workers in two Dutch hospitals in March 2020. *medRxiv.* 2020. doi: 10.1101/2020.03.23.20041913
31. McMichael TM, Currie DW, Clark S, Pogojans S, Kay M, Schwartz NG, et al. Epidemiology of Covid-19 in a Long-Term Care Facility in King County, Washington. *N Engl J Med.* 2020;382(21):2005-2011.
32. Liu M, He P, Liu HG, Wang XJ, Li FJ, Chen S, et al. [Clinical characteristics of 30 medical workers infected with new coronavirus pneumonia]. *Zhonghua Jie He He Hu Xi Za Zhi Zhonghua Jiehe He Huxi Zazhi Chin J Tuberc Respir Dis.* 2020;43(3):209-14.
33. Paradiso AV, Summa simona D, Silvestris N, Tommasi S, Tufaro A, Palma GD, et al. COVID-19 SCREENING AND MONITORING OF ASYMPTOMATIC HEALTH WORKERS WITH A RAPID SEROLOGICAL TEST. *medRxiv.* 2020. doi: 10.1101/2020.05.05.20086017
34. Ran L, Chen X, Wang Y, Wu W, Zhang L, Tan X. Risk Factors of Healthcare Workers with Corona Virus Disease 2019: A Retrospective Cohort Study in a Designated Hospital of Wuhan in China [available online ahead of print August 18, 2020]. *Clin Infect Dis.* 2020; ciaa287 .doi: 10.1093/cid/ciaa287
35. Reusken CB, Buiting A, Bleeker-Rovers C, Diederens B, Hooiveld M, Friesema I, et al. Rapid assessment of regional SARS-CoV-2 community transmission through a convenience sample of healthcare workers, the Netherlands, March 2020. *Eurosurveillance.* 2020;25(12).
36. Rivett L, Sridhar S, Sparkes D, Routledge M, Jones NK, Forrest S, et al. Screening of healthcare workers for SARS-CoV-2 highlights the role of asymptomatic carriage in COVID-19 transmission. *eLife.* 2020;9:e58728..
37. Guery R, Delaye C, Brule N, Nael V, Castain L, Raffi F, et al. Limited effectiveness of systematic screening by nasopharyngeal RT-PCR of medicalized nursing home staff

after a first case of COVID-19 in a resident [available online ahead of print July 8 2020]. *Med Mal Infect.* 2020;S0399-077X(20)30126-8. doi: 10.1016/j.medmal.2020.04.020.

38. Roxby AC. Detection of SARS-CoV-2 Among Residents and Staff Members of an Independent and Assisted Living Community for Older Adults — Seattle, Washington, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(14):416–418
39. Solodky ML, Galvez C, Russias B, Detourbet P, N’Guyen-Bonin V, Herr A-L, et al. Lower detection rates of SARS-COV2 antibodies in cancer patients vs healthcare workers after symptomatic COVID-19. *Ann Oncol Off J Eur Soc Med Oncol.* 2020;31(8):1087-1088.
40. Takita M, Matsumura T, Yamamoto K, Yamashita E, Hosoda K, Hamaki T, et al. Preliminary Results of Seroprevalence of SARS-CoV-2 at Community Clinics in Tokyo. *medRxiv.* 2020. doi: 10.1101/2020.04.29.20085449
41. Treibel TA, Manisty C, Burton M, McKnight Á, Lambourne J, Augusto JB, et al. COVID-19: PCR screening of asymptomatic health-care workers at London hospital. *The Lancet.* 2020;395(10237):1608-10.
42. Canova V, Lederer Schläpfer H, Piso RJ, Droll A, Fenner L, Hoffmann T, et al. Transmission risk of SARS-CoV-2 to healthcare workers -observational results of a primary care hospital contact tracing. *Swiss Med Wkly.* 2020; 25(12): 2000334.
43. Tosato F, Pelloso M, Gallo N, Giraudo C, Llanaj G, Cosma C, et al. Severe Acute Respiratory Syndrome Coronavirus 2 Serology in Asymptomatic Healthcare Professionals: Preliminary Experience of a Tertiary Italian Academic Center. *medRxiv.* 2020. doi: 10.1101/2020.04.27.20073858v1
44. Tostmann A, Bradley J, Bousema T, Yiek W-K, Holwerda M, Bleeker-Rovers C, et al. Strong associations and moderate predictive value of early symptoms for SARS-CoV-2 test positivity among healthcare workers, the Netherlands, March 2020. *Euro Surveill Bull Eur Sur Mal Transm Eur Commun Dis Bull.* 2020;25(16):2000508.
45. Wang X, Liu W, Zhao J, Lu Y, Wang X, Yu C, et al. Clinical characteristics of 80 hospitalized frontline medical workers infected with COVID-19 in Wuhan, China. *J Hosp Infect.* 2020;105(3):399-403
46. Wei X-S, Wang X-R, Zhang J-C, Yang W-B, Ma W-L, Yang B-H, et al. A cluster of health care workers with COVID-19 pneumonia caused by SARS-CoV-2 [available online ahead of print April 1, 2012]. *J Microbiol Immunol Infect Wei Mian Yu Gan Ran Za Zhi.* 2020. doi: 10.1016/j.jmii.2020.04.013
47. Zheng L, Wang X, Zhou C, Liu Q, Li S, Sun Q, et al. Analysis of the infection status of the health care workers in Wuhan during the COVID-19 outbreak: A cross-sectional study [available online ahead of print July 8 2020]. *Clin Infect Dis.* 2020. doi: 10.1093/cid/ciaa588/5837357

48. Wang X, Pan Z, Cheng Z. Association between 2019-nCoV transmission and N95 respirator use. *J Hosp Infect.* 2020;105(1):104-5.
49. Ng K, Poon BH, Kiat Puar TH, Shan Quah JL, Loh WJ, Wong YJ, et al. COVID-19 and the Risk to Health Care Workers: A Case Report [available online ahead of print July 8 2020]. *Ann Intern Med.* 2020. doi: 10.7326/L20-0175
50. Lai X, Wang M, Qin C, Tan L, Ran L, Chen D, et al. Coronavirus Disease 2019 (COVID-2019) Infection Among Health Care Workers and Implications for Prevention Measures in a Tertiary Hospital in Wuhan, China. *JAMA Netw Open.* 2020;3(5):e209666-e209666.
51. Khalil A, Hill R, Ladhani S, Pattisson K, O'Brien P. COVID-19 screening of health-care workers in a London maternity hospital. *Lancet Infect Dis.* 2020;S1473-3099(20)30403-5
52. Contejean A, Leporrier J, Canoui E, Alby-Laurent F, Lafont E, Beaudeau L, et al. Comparing dynamics and determinants of SARS-CoV-2 transmissions among health care workers of adult and pediatric settings in central Paris. *medRxiv.* 2020. doi: 10.1101/2020.05.19.20106427
53. Graham N, Junghans C, Downes R, Sendall C, Lai H, Mckirdy A, et al. SARS-CoV-2 infection, clinical features and outcome of COVID-19 in United Kingdom nursing homes. *Journal of Infection.* 2020;81(3):411-419.
54. Vahidy F, Sostman HD, Bernard D, Boom ML, Drews AL, Christensen PA, et al. Prevalence of SARS-CoV-2 infection among asymptomatic healthcare workers in greater Houston: a cross-sectional analysis of surveillance data from a large healthcare system. *JAMA Netw Open.* 2020;3(7):e2016451.
55. Fujita K, Kada S, Kanai O, Hata H, Odagaki T, Satoh-Asahara N, et al. Quantitative SARS-CoV-2 antibody screening of healthcare workers in the southern part of Kyoto city during the COVID-19 peri-pandemic period. *medRxiv.* 2020. doi: 10.1101/2020.05.12.20098962
56. Sikora K, Barwick I, Hamilton C. Serological prevalence of antibodies to SARS CoV-2 amongst cancer centre staff. *medRxiv.* 2020. doi: 10.1101/2020.05.16.20099408.
57. Yombi JC, De Greef J, Marsin A-S, Simon A, Rodriguez-Villalobos H, Penaloza A, et al. Symptom-based screening for COVID-19 in health care workers: The importance of fever. *J Hosp Infect.* 2020;105(3): 428–429.
58. Olalla J, Correa AM, Martin-Escalante MD, Hortas ML, Martin-Sendarrubias MJ, Fuentes V, et al. Search for asymptomatic carriers of SARS-CoV-2 in healthcare workers during the pandemic: a Spanish experience [available online ahead of print July 8th 2020]. *QJM: An International Journal of Medicine.* doi: 10.1093/qjmed/hcaa238

59. Antonio-Villa NE, Bello-Chavolla OY, Vargas-Vazquez A, Fermin-Martinez CA, Marquez-Salinas A, Bahena-Lopez JP. Health-care workers with COVID-19 living in Mexico City: clinical characterization and related outcomes. medRxiv. 2020. doi: 10.1101/2020.07.02.20145169.
60. Baker M.A., Rhee C., Fiumara K., Bennett-Rizzo C., Tucker R., Williams S.A., et al. COVID-19 infections among healthcare workers exposed to a patient with a delayed diagnosis of COVID-19. Infect Control Hosp Epidemiol. 2020: 1–2. doi: 10.1017/ice.2020.256
61. Fernández-Rivas G, Quirant-Sánchez B, González V, Doladé M, Martínez-Caceres E, Piña M, et al. Seroprevalence of SARS-CoV-2 IgG Specific Antibodies among Healthcare Workers in the Northern Metropolitan Area of Barcelona, Spain, after the first pandemic wave. medRxiv. doi: 10.1101/2020.06.24.20135673.
62. Borrás-Bermejo B, Martínez-Gómez X, San Miguel MG, Esperalba J, Antón A, Martín E, et al. Asymptomatic SARS-CoV-2 Infection in Nursing Homes, Barcelona, Spain, April 2020 [available online ahead of print July 8 2020]. Emerg Infect Dis. 2020;26(9). doi: 10.3201/eid2609.202603.
63. Chen Y., Tong X., Wang J., Huang W., Yin S., Huang R., et al. High SARS-CoV-2 antibody prevalence among healthcare workers exposed to COVID-19 patients. J Infect. 2020; 81(3): 420–426.
64. Cho J.-H., Kang S.H., Park H.C., Kim D.K., Lee S.-H., Do J.Y., et al. Hemodialysis with Cohort Isolation to Prevent Secondary Transmission during a COVID-19 Outbreak in Korea. J Am Soc Nephrol. 2020;31(7):1398-1408.
65. Clemency BM, Varughese R, Scheafer DK, Ludwig B, Welch JV, McCormack RF, et al. Symptom Criteria for COVID-19 Testing of Health Care Workers. Acad Emerg Med. 2020;27(6):469-74.
66. Fusco FM, Pisaturo M, Iodice V, Bellopede R, Tambaro O, Parrella G, et al. COVID-19 infections among Healthcare Workers in an Infectious Diseases specialized setting in Naples, Southern Italy: results of a cross-sectional surveillance study. J Hosp Infect. 2020; 105(4): 596–600.
67. Galan I, Velasco M, Casas ML, Goyanes MJ, Rodríguez-Caravaca G, Losa JE, et al. SARS-CoV-2 SEROPREVALENCE AMONG ALL WORKERS IN A TEACHING HOSPITAL IN SPAIN: UNMASKING THE RISK. medRxiv. 2020. doi: 10.1101/2020.05.29.20116731.
68. Albalade M, Arribas P, Torres E, Cintra M, Alcázar R, Puerta M, et al. High prevalence of asymptomatic COVID-19 in hemodialysis: Daily learning during first month of COVID-19 pandemic. Nefrol Engl Ed. 2020; 40(3)279-286.
69. Bays DJ, Nguyen M-VH, Cohen SH, Waldman S, Martin CS, Thompson GR, et al. Investigation of Nosocomial SARS-CoV-2 Transmission from Two Patients to Health Care

Workers Identifies Close Contact but not Airborne Transmission Events [available online ahead of print July 8 2020]. *Infect Control Hosp Epidemiol.* 2020:1-22.

70. Bhattacharya R, Chowdhury S, Mukherjee R, Nandi A, Kulshrestha M, Ghosh R, et al. Pre exposure Hydroxychloroquine use is associated with reduced COVID19 risk in healthcare workers - a Retrospective cohort. *medRxiv.* 2020. doi: 10.1101/2020.06.09.20116806.

71. Brandstetter S, Roth S, Harner S, Buntrock-Döpke H, Toncheva AA, Borchers N, et al. Symptoms and immunoglobulin development in hospital staff exposed to a SARS-CoV-2 outbreak [available online ahead of print July 8 2020]. *Pediatr Allergy Immunol.* 2020. doi: 10.1111/pai.13278

72. Mansour M, Leven E, Muellers K, Stone K, Mendu DR, Wajnberg A. Prevalence of SARS-CoV-2 Antibodies Among Healthcare Workers at a Tertiary Academic Hospital in New York City. *J Gen Intern Med.* 2020; 35(8): 2485–2486.

73. Gao M, Yang L, Chen X, Deng Y, Yang S, Xu H, et al. A study on infectivity of asymptomatic SARS-CoV-2 carriers. *Respir Med.* 2020;169:106026. doi: 10.1016/j.rmed.2020.106026

74. Houlihan C, Vora N, Byrne T, Lewer D, Heaney J, Moore DA, et al. SARS-CoV-2 virus and antibodies in front-line Health Care Workers in an acute hospital in London: preliminary results from a longitudinal study. *medRxiv.* 2020. doi: 10.1101/2020.06.08.20120584.

75. Jeon YW, Park ES, Jung SJ, Kim Y, Choi JY, Kim HC. Protection of Healthcare Workers Against COVID-19 at a Large Teaching Hospital in Seoul, Korea. *Yonsei Med J.* 2020;61(7):631-4.

76. Kammon AM, El-Arabi AA, Erhouma EA, Mehemed TM, Mohamed OA. Seroprevalence of antibodies against SARS-CoV-2 among public community and health-care workers in Alzintan City of Libya. *medRxiv.* 2020. doi: 10.1101/2020.05.25.20109470.

77. Korth J, Wilde B, Dolff S, Anastasiou OE, Krawczyk A, Jahn M, et al. SARS-CoV-2-specific antibody detection in healthcare workers in Germany with direct contact to COVID-19 patients. *J Clin Virol.* 2020;128:104437-104437.

78. Lahner E, Dilaghi E, Prestigiacomo C, Alessio G, Marcellini L, Simmaco M, et al. Prevalence of Sars-Cov-2 Infection in Health Workers (HWs) and Diagnostic Test Performance: The Experience of a Teaching Hospital in Central Italy. *Int J Env Res Public Health Online.* 2020;17(12): 4417.

79. Lan F-Y, Filler R, Mathew S, Buley J, Iliaki E, Bruno-Murtha LA, et al. COVID-19 symptoms predictive of healthcare workers' SARS-CoV-2 PCR results. *PLoS One.* 2020;15(6):e0235460-e0235460.

80. Lombardi A, Consonni D, Carugno M, Bozzi G, Mangioni D, Muscatello A, et al. Characteristics of 1573 healthcare workers who underwent nasopharyngeal swab testing for SARS-CoV-2 in Milan, Lombardy, Italy [available online ahead of print July 8 2020]. *Clin Microbiol Infect.* 2020. doi: 10.1016/j.cmi.2020.06.013
81. Ma Y, Diao B, Lv X, Zhu J, Chen C, Liu L, et al. Epidemiological, clinical, and immunological features of a cluster of COVID-19 contracted hemodialysis patients. *Kidney Int Rep.* 2020; 5(8): 1333–1341.
82. Mani NS, Budak JZ, Lan KF, Bryson-Cahn C, Zelikoff A, Barker GEC, et al. Prevalence of COVID-19 Infection and Outcomes Among Symptomatic Healthcare Workers in Seattle, Washington [available online ahead of print July 8 2020]. *Clin Infect Dis.* 2020:ciaa761. doi: 10.1093/cid/ciaa761
83. Jones NK, Rivett L, Sparkes D, Forrest S, Sridhar S, Young J, et al. Effective control of SARS-CoV-2 transmission between healthcare workers during a period of diminished community prevalence of COVID-19. *eLife* 2020;9:e59391 DOI: 10.7554/eLife.59391
84. Martin C, Montesinos I, Dauby N, Gilles C, Dahma H, Van Den Wijngaert S, et al. Dynamic of SARS-CoV-2 RT-PCR positivity and seroprevalence among high-risk health care workers and hospital staff. *J Hosp Infect.* 2020;102-106.
85. Nakamura A, Sato R, Ando S, Oana N, Nozaki E, Endo H, et al. Seroprevalence of Antibodies to SARS-CoV-2 in Healthcare Workers in Non-epidemic Region: A Hospital Report in Iwate Prefecture, Japan. *medRxiv.* 2020. doi: 10.1101/2020.06.15.20132316.
86. Paderno A., Fior M., Berretti G., Schreiber A., Grammatica A., Mattavelli D., et al. SARS-CoV-2 Infection in Health Care Workers: Cross-sectional Analysis of an Otolaryngology Unit. *Otolaryngol Head Neck Surg.* 2020;194599820932162. doi: 10.1177/0194599820932162.
87. Parcell B, Brechin K, Allstaff S, Park M, Third W, Bean S, et al. Drive-through testing for SARS-CoV-2 in symptomatic health and social care workers and household members: an observational cohort study in Tayside, Scotland. *medRxiv.* 2020. doi: 10.1101/2020.05.08.20078386.
88. Psychogiou M, Karabinis A, Pavlopoulou ID, Basoulis D, Petsios K, Roussos S, et al. Antibodies against SARS-CoV-2 among health care workers in a country with low burden of COVID-19. *medRxiv.* 2020. doi: 10.1101/2020.06.23.20137620
89. Rudberg A-S, Havervall S, Manberg A, Falk AJ, Aguilera K, Ng H, et al. SARS-CoV-2 exposure, symptoms and seroprevalence in health care workers. *medRxiv.* 2020. doi: 10.1101/2020.06.22.20137646.
90. Schmidt SB, Grüter L, Boltzmann M, Rollnik JD. Prevalence of serum IgG antibodies against SARS-CoV-2 among clinic staff. *PLoS One.* 2020;15(6):e0235417-e0235417.

91. Schneider KN, Correa-Martínez CL, Gosheger G, Rickert C, Schorn D, Mellmann A, et al. Assessing the spreading potential of an undetected case of COVID-19 in orthopaedic surgery [available online ahead of print July 8 2020]. *Arch Orthop Trauma Surg.* 2020: 1–7. doi: 10.1007/s00402-020-03516-1
92. Shields AM, Faustini SE, Perez-Toledo M, Jossi S, Aldera EL, Allen JD, et al. SARS-CoV-2 seroconversion in health care workers. *medRxiv.* 2020. doi: 10.1101/2020.05.18.20105197
93. Sikkema RS, Pas SD, Nieuwenhuijse DF, O'Toole Á, Verweij J, Linden A van der, et al. COVID-19 in health-care workers in three hospitals in the south of the Netherlands: a cross-sectional study [available online ahead of print July 8 2020]. *Lancet Infect Dis.* 2020. doi: 10.1016/S1473-3099(20)30527-2
94. Bird P, Badhwar V, Fallon K, Kwok KO, Tang JW. High SARS-CoV-2 infection rates in respiratory staff nurses and correlation of COVID-19 symptom patterns with PCR positivity and relative viral loads. *J Infect.* 2020; 81(3): 452–482.
95. Wee LE, Sim XYJ, Conceicao EP, Aung MK, Goh JQ, Yeo DWT, et al. Containment of COVID-19 cases among healthcare workers: The role of surveillance, early detection, and outbreak management [available online ahead of print July 8 2020]. *Infect Control Hosp Epidemiol.* 2020;1-7. doi: 10.1017/ice.2020.219.
96. Xiong S, Guo C, Dittmer U, Zheng X, Wang B. The prevalence of antibodies to SARS-CoV-2 in asymptomatic healthcare workers with intensive exposure to COVID-19. *medRxiv.* 2020. doi: 10.1101/2020.05.28.20110767
97. Xu X, Sun J, Nie S, Li H, Kong Y, Liang M, et al. Seroprevalence of immunoglobulin M and G antibodies against SARS-CoV-2 in China. *Nat Med.* 2020;26:1193–1195.
98. Zhao R, Li M, Song H, Chen J, Ren W, Feng Y, et al. Early detection of SARS-CoV-2 antibodies in COVID-19 patients as a serologic marker of infection [available online ahead of print July 8 2020]. *Clin Infect Dis.* 2020. doi: 10.1093/cid/ciaa523
99. Jary A, Flandre P, Chabouis A, Nguyen S, Marot S, Burrel S, et al. Clinical presentation of Covid-19 in health care workers from a French University Hospital. *J Infect.* 2020; 81(3): e61–e63.
100. Ji M., Yuan L., Shen W., Lv J., Li Y., Li M., et al. Characteristics of Disease Progress in Patients with Coronavirus Disease 2019 in Wuhan, China [available online ahead of print July 8 2020]. *Epidemiol Infect.* 2020;148: e94.
101. Malani A, Mohanan M, Kumar C, Kramer J, Tandel V. Prevalence of SARS-CoV-2 among workers returning to Bihar gives snapshot of COVID across India. *medRxiv.* 2020. doi: 10.1101/2020.06.26.20138545.

102. Mandić-Rajčević S, Masci F, Crespi E, Franchetti S, Longo A, Bollina I, et al. Contact tracing and isolation of asymptomatic spreaders to successfully control the COVID-19 epidemic among healthcare workers in Milan (Italy). medRxiv. 2020. doi: 10.1101/2020.06.26.20138545. Accessed July 8, 2020.
103. Caseiro MM, Mazzurana M, Barreiro B, Barbosa EC, Barbosa AP. Positivity of SARS-CoV-2, by RT-PCR among workers of a Public Hospital in the city of Santos, SP, Brazil 2020. medRxiv. 2020. doi: 10.1101/2020.06.30.20143529. Accessed July 8, 2020.
104. Ortiz-Brizuela E, Villanueva-Reza M, González-Lara MF, Tamez-Torres KM, Román-Montes CM, Díaz-Mejía BA, et al. CLINICAL AND EPIDEMIOLOGICAL CHARACTERISTICS OF PATIENTS DIAGNOSED WITH COVID-19 IN A TERTIARY CARE CENTER IN MEXICO CITY: A PROSPECTIVE COHORT STUDY. Rev Invest Clin. 2020;72(3):165-77.
105. Schwartz KL, Achonu C, Buchan SA, Brown KA, Lee B, Whelan M, et al. COVID-19 infections among Healthcare Workers and Transmission within Households. medRxiv. 2020. doi: 10.1101/2020.06.12.20129619. Accessed July 8, 2020.
106. Wu F, Guo M, Zhang S, Ma Y, Ma P, Duan L, et al. Transmission of 2019-nCoV to Health-Care Workers in the Early Epidemic. SSRN. doi: 10.2139/ssrn.3542155. Accessed July 8, 2020.
107. Li YK, Peng S, Li LQ, Wang Q, Ping W, Zhang N, et al. Clinical and Transmission Characteristics of Covid-19 - A Retrospective Study of 25 Cases from a Single Thoracic Surgery Department [available online ahead of print July 8 2020]. Curr Med Sci. 2020:1-6. doi: 10.1007/s11596-020-2176-2.
108. Chatterjee P, Anand T, Singh KJ, Rasaily R, Singh R, Das S, et al. Healthcare workers & SARS-CoV-2 infection in India: A case-control investigation in the time of COVID-19. Indian J Med Res. 2020;151(5):459-67.
109. El-Boghdadly K, Wong DJN, Owen R, Neuman MD, Pocock S, Carlisle JB, et al. Risks to healthcare workers following tracheal intubation of patients with COVID-19: a prospective international multicentre cohort study[available online ahead of print July 8 2020]. Anaesthesia. 2020. doi: 10.1111/anae.15170.
110. Guery R, Cécile D, Noëlle B, Virginie N, Louise C, François R, et al. Limited effectiveness of systematic screening by nasopharyngeal RT-PCR of medicalized nursing home staff after a first case of COVID-19 in a resident[available online ahead of print July 8 2020]. Med Mal Infect. 2020;S0399-077X(20)30126-8. doi: 10.1016/j.medmal.2020.04.020.
111. Alsafyan Y.M., Althunayyan S.M., Khan A.A., Hakawi A.M., Assiri A.M. Clinical characteristics of COVID-19 in Saudi Arabia: A national retrospective study. J Infect Public Health. 2020;13:920-925.

112. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*.2020;323(13):1239-42.
113. Eurosurveillance editorial team. Updated rapid risk assessment from ECDC on coronavirus disease 2019 (COVID-19) pandemic: increased transmission in the EU/EEA and the UK. *Eurosurveillance*. 2020;25(12).
114. Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected Patients. *N Engl J Med*. 2020;382(12):1177-9.
115. Felice C., Di Tanna G.L., Zanusi G., Grossi U. Impact of COVID-19 Outbreak on Healthcare Workers in Italy: Results from a National E-Survey. *J Community Health*. 2020;45(4):675-683.
116. Tabah A, Ramanan M, Laupland KB, Buetti N, Cortegiani A, Mellingerhoff J, et al. Personal protective equipment and intensive care unit healthcare worker safety in the COVID-19 era (PPE-SAFE): An international survey. *J Crit Care*. 2020;59:70-5.
117. Iqbal M.R., Chaudhuri A. COVID-19: Results of a national survey of United Kingdom healthcare professionals' perceptions of current management strategy – A cross-sectional questionnaire study. *Int J Surg*. 2020;79: 156–161.

Table 1. Characterization of studies describing SARS-CoV-2 prevalence in HCW using RT-PCR

First Author, Year (Reference No.)	Number of participants	Setting	Selection criteria	City, Country
<i>Kluytmans et al (2020)(30)</i>	86	Hospital, all services	HCW who fever or (mild) respiratory symptoms	Breda, The Netherlands
<i>García-Basteiro et al (2020) (24)</i>	583	Hospital, all services	All HCW who deliver care and services to patients, either directly as physicians or nurses, or indirectly as assistants, technicians, stretcher-bearers or other support staff (administrative officers, cleaning, kitchen, laundry, maintenance, etc.)	Barcelona, Spain
<i>Breazzano et al (2020) (18)</i>	2088	Hospitals at the city level. Data reported by each postgraduate director	Medical residents in universities of NYC. Screened according to symptoms.	New York City, NY, USA
<i>Tosato et al (2020) (43)</i>	133	Laboratory Department	All workers in the laboratory department.	Padova, Italy
<i>Keeley et al (2020) (28)</i>	1533	Hospital, all services	All HCW	Sheffield, UK
<i>Tostmann et al (2020) (44)</i>	803	Hospital, all services	HCW with symptoms suggestive of COVID-19	Nijmegen, The Netherlands
<i>Reusken et al (2020) (35)</i>	1097	Hospital, all services	HCW with respiratory any symptom, even mild respiratory complaints.	Noord-Brabant Province, The Netherlands
<i>Hunter et al (2020) (26)</i>	1654	Hospital, all services including ambulance service staff	Staff with compatible symptoms (ie, new continuous cough or fever)	Newcastle upon Tyne, UK
<i>Ran et al (2020) (34)</i>	72	Hospital, all services	All HCW	Wuhan, China
<i>Solodky et al (2020) (39)</i>	244	Hospital, Cancer Institute	Voluntary HCW	Lyon, France

<i>Guery et al (2020) (37)</i>	136	Long-term care facility	All HCW	Nantes, France
<i>Treibel et al (2020) (41)</i>	400	Hospitals at a city level	All HCW	London, UK
<i>Roxby et al (2020) (38)</i>	62	Long-term care facility	All HCW	Seattle, WA, USA
<i>Lai et al (2020) (50)</i>	335	Hospital, all services	All HCW	Wuhan, China
<i>Graham et al (2020) (53)</i>	70	Long-term care facility	Asymptomatic HCW	London, UK
<i>Vahidy et al (2020) (54)</i>	2887	Hospital, all services	Asymptomatic HCW	Houston, TX, USA
<i>Olalla et al (2020) (58)</i>	498	Hospital, all services	Asymptomatic HCW	Marbella, Spain
<i>Yombi et al (2020) (57)</i>	536	Hospital, all services	Symptomatic HCW	Brussels, Belgium
<i>Khalil et al (2020) (51)</i>	266	Hospital, all services	All HCW	London, UK
<i>Contejean et al (2020) (52)</i>	1344	Hospital, all services	Symptomatic HCW	Paris, France
<i>Barrett et al (2020) (16)</i>	546	Hospital, all services	All HCW that reported: (1) ≥ 20 hours of hospital work weekly; (2) occupations with regular patient exposure (e.g., residents, fellows, attending physicians, dentists, nurse practitioners, physician assistants, registered nurses, technicians, respiratory therapists, physical therapists); and (3) regular direct patient contact (≥ 3 patients/shift) expected in the next three months.	Newark, NJ, USA
<i>Bai et al (2020) (15)</i>	118	Hospital, Department Neurosurgery	All HCW	Wuhan, China

<i>Folgueira et al (2020) (22)</i>	1438	Hospital, City-level	All HCW	Madrid, Spain
<i>Antonio-Villa et al (2020) (59)</i>	34263	Hospital, all services	All HCW	Mexico City, Mexico
<i>Borras-Bermejo et al (2020) (62)</i>	2655	Long-term care facility	All HCW	Catalonia, Spain
<i>Cho JH et al (2020) (64)</i>	278	Hospital, all services	Symptomatic HCW	South Korea
<i>Clemency BM et al (2020) (65)</i>	961	Hospital, all services	Symptomatic HCW	New York City, NY, USA
<i>Fusco FM et al (2020) (66)</i>	115	Hospital, all services	Asymptomatic HCW	Naples, Italy
<i>Galán MI et al (2020) (67)</i>	2590	Hospital, all services	HCW with positive IgG and symptoms in the past 14 days	Madrid, Spain
<i>Albalade M. et al (2020) (68)</i>	14	Hospital, all services	Symptomatic HCW	Madrid, Spain
<i>Bhattacharya et al (2020) (70)</i>	106	Hospital, all services	All HCW	Kolkata, India
<i>Houlihan et al (2020) (74)</i>	200	Hospital, all services	All HCW	London, UK
<i>Lahner et al (2020) (78)</i>	2115	Hospital, all services	All HCW	Rome, Italy
<i>Lan et al (2020) (79)</i>	592	Hospital, all services	All HCW	Massachusetts, USA
<i>Lombardi et al (2020) (80)</i>	1573	Hospital, all services	All HCW	Milan, Italy
<i>Ma et al (2020) (81)</i>	33	Hemodialysis unit	All HCW	Wuhan, China

<i>Mani et al (2020) (82)</i>	3477	Hospital, all services	Symptomatic HCW	Seattle, WA, USA
<i>Jones et al (2020) (83)</i>	4800	Hospital, all services	Symptomatic HCW	London, UK
<i>Martin et al (2020) (84)</i>	326	Hospital, all services	Symptomatic HCW	Brussels, Belgium
<i>Paderno et al (2020) (86)</i>	58	Otorrinolaringology Clinic	Symptomatic HCW	Brescia, Italy
<i>Parcell et al (2020) (87)</i>	1173	Hospital, all services	Symptomatic HCW	Tayside, UK
<i>Shields et al (2020) (92)</i>	554	Hospital, all services	All HCW	Birmingham, UK
<i>Sikkema et al (2020) (93)</i>	1796	Hospital, all services	Symptomatic HCW	Breda and tilburg, Netherlands
<i>Bird et al (2020) (94)</i>	152	Hospital, all services	Symptomatic HCW	Leicester, UK
<i>Wee et al (2020) (95)</i>	1642	Hospital, all services	Symptomatic HCW	Singapore
<i>Brandstetter et al (2020) (71)</i>	201	Hospital, all services	All HCW	Regensburg, Germany

HCW: Health care workers; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

Table 2. Areas in which COVID-19 positive health care workers were laboring during RT-PCR screenings

Area/Setting	Number of studies	Proportion, (%)	95% CI	I ²
Clinics/Wards	5	43	28% - 59%	91%
Operating Room	4	24	17% - 31%	60%
Others	4	29	13% - 48%	91%
Emergency Room	5	16	6% - 29%	91%
ICU	5	9	4% - 15%	68%

COVID-19: Coronavirus disease 2019; ICU: Intensive care unit.

Table 3. Prevalence of symptoms in COVID-19 positive health care workers

Symptom	Number of studies	Pooled Prevalence, (%)	95% CI	I ²
Fever	29	57	50% - 64%	96%
Cough	26	57	50% - 65%	96%
Malaise	10	48	35% - 62%	89%
Myalgia	22	44	36% - 52%	96%
Headache	22	36	27% - 46%	97%
Sore Throat	15	32	23% - 42%	97%
Shortness of Breath	21	22	17% - 28%	95%
Diarrhea	21	18	14% - 22%	87%
Nausea	7	9	6% - 14%	73%
Chest Pain	6	8	1% - 18%	86%

COVID-19: Coronavirus disease 2019

Figure 1. Flowchart summarizing the study search and selection process.

Legend. The PRISMA 2009 Flow Diagram, obtained from “Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097”


Figure 2. Geographical distribution of studies that reported data on SARS-CoV-2 prevalence in healthcare workers.

Legend: SARS-CoV-2, Severe acute respiratory syndrome coronavirus 2; %, percent.

Figure 3. SARS-CoV-2 infection prevalence in healthcare workers using RT-PCR


Legend: The results are presented as fractions. The overall summary estimates presented are calculated using random-effects models. HCW, health care workers; LTCF, long-term care facility; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; %, percent.

ORIGINAL UNEDITED MANUSCRIPT





B)



c)

