**Original Research Article** 

# Air conditioning and heat-related mortality: a multi-country longitudinal study

Francesco Sera<sup>1\*,2</sup>, Masahiro Hashizume<sup>3</sup>, Yasushi Honda<sup>4</sup>, Eric Lavigne<sup>5,6</sup>, Joel Schwartz<sup>7</sup>, Antonella Zanobetti<sup>7</sup>, Aurelio Tobias<sup>8</sup>, Carmen Iñiguez<sup>9,10</sup>, Ana M. Vicedo-Cabrera<sup>1,11,12</sup>, Marta Blangiardo<sup>13</sup>, Ben Armstrong<sup>1,2</sup>, Antonio Gasparrini<sup>1,2,14</sup>

<sup>1</sup>Department of Public Health Environments and Society, London School of Hygiene & Tropical Medicine, London UK

<sup>2</sup>Centre for Statistical Methodology, London School of Hygiene & Tropical Medicine, London UK

<sup>3</sup>Department of Global Health Policy, School of International Health, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

<sup>4</sup>Faculty of Health and Sport Sciences, University of Tsukuba, Tsukuba, Japan

<sup>5</sup>Air Health Science Division, Health Canada, Ottawa, Canada

<sup>6</sup>School of Epidemiology & Public Health, University of Ottawa, Ottawa, Canada

<sup>7</sup>Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA

<sup>8</sup>Institute of Environmental Assessment and Water Research, Spanish Council for Scientific Research, Barcelona, Spain

<sup>9</sup>Department of Statistics and Computational Research, Universitat de València, Spain <sup>10</sup>CIBERESP, Madrid, Spain

<sup>11</sup>Institute of Social and Preventive Medicine. University of Bern, Bern, Switzerland
<sup>12</sup>Oeschger Center for Climate Change Research. University of Bern, Bern, Switzerland
<sup>13</sup>Department of Epidemiology and Biostatistics, Imperial College London, London, United Kingdom

<sup>14</sup>Centre on Climate Change and Planetary Health, London School of Hygiene & Tropical Medicine, London UK

\*Correspondence to: Francesco Sera, London School of Hygiene & Tropical Medicine, 15-17 Tavistock Place, London WC1H 9SH, UK. Telephone: 0044 (0)20 79272992. E-mail: <u>francesco.sera@lshtm.ac.uk</u>.

Running title: A longitudinal study on air conditioning and mortality

**Funding**: This work was supported by the Medical Research Council-UK [Grant ID: Grant ID: MR/R013349/1], by the Natural Environment Research Council UK (Grant ID: NE/R009384/1) and by the European Union Horizon 2020 programme (Grant ID: 820655)

**Competing financial interests**: The author declares he has no actual or potential conflict of interest.

**Data and code**: The computer code used to conduct analyses for this paper is available from the first author upon request. The mortality data have been obtained through a restricted data use agreement with each national institute (Statistics Canada for Canada, Ministry of Health, Labour and Welfare for Japan, Spain National Institute of Statistics for Spain, and National Center for Health Statistics (NCHS) for USA), and are therefore not available for public dissemination.

#### ABSTRACT

#### Background

Air conditioning has been proposed as one of the key factors explaining reductions of heatrelated mortality risks observed in the last decades. However, direct evidence is still limited.

#### Methods

We used a multi-country, multi-city, longitudinal design to quantify the independent role of air conditioning in reported attenuation in risk. We collected daily time series of mortality, mean temperature, and yearly air conditioning prevalence for 311 locations in Canada, Japan, Spain, and the USA between 1972 and 2009. For each city and sub-period, we fitted a quasi-Poisson regression combined with distributed lag non-linear models to estimate summer-only temperature–mortality associations. At the second stage, we used a novel multilevel, multivariate spatio-temporal meta-regression model to evaluate effect modification of air conditioning on heat–mortality associations. We computed relative risks and fractions of heat-attributable excess deaths under observed and fixed air conditioning prevalences.

#### Results

Results show an independent association between increased air conditioning prevalence and lower heat-related mortality risk. Excess deaths due to heat decreased during the study periods from 1.40% to 0.80% in Canada, 3.57% to 1.10% in Japan, 3.54% to 2.78% in Spain, and 1.70% to 0.53% in the US. However, increased air conditioning explains only part of the observed attenuation, corresponding to 16.7% in Canada, 20.0% in Japan, 14.3% in Spain, and 16.7% in the US.

### Conclusions

Our findings are consistent with the hypothesis that air conditioning represents an effective heat adaptation strategy, but suggests that other factors have played an equal or more important role in increasing the resilience of populations.

**Keywords**: air conditioning; temperature; adaptation; longitudinal; meta-analysis; multilevel.

#### INTRODUCTION

Epidemiologic studies in various countries have provided evidence of a decrease in mortality risks associated to exposure to heat over the last decades.<sup>1,2</sup> Several mechanisms have been suggested as potential drivers of such attenuation, including physiologic (referred to as acclimatization), behavioral (e.g., clothing), infrastructural (green spaces), and technological (heat warning system).<sup>2-4</sup> However, evidence is still limited, and an appropriate characterization of factors responsible for the attenuation of heat-related risks is still lacking. This information is nonetheless critical for planning effective public health and climate policies.<sup>1-3</sup>

Air conditioning is one of the most straightforward strategies to reduce heat stress, and previous investigations have assessed its role in modifying mortality risks associated to exposure to high temperature using both individual- or aggregated-level designs, although with conflicting results.<sup>5-13</sup> These studies adopted either a cross-sectional and/or longitudinal design, comparing risks at different air conditioning prevalence between individuals/locations or at different times. However, they faced a number of methodologic challenges. Analyses based on the cross-sectional comparison of subjects or cities with different air conditioning use and prevalence are prone to bias, as other characteristics, such as socio-economic or climatic conditions, can be responsible for differences in health risks. Longitudinal designs can address this issue, but they need data consistently collected over a long period of time to allow for substantial variation in air conditioning use within each location. More importantly, these studies can be affected by temporal confounding due to concurrent changes in other modifying factors, such as infrastructural changes and access to health care. Finally, the complexity of exposure–response relationships,

characterised by non-linearity and temporally delayed effects, presents additional problems in modelling temperature—mortality associations. A recent investigation by Nordio and colleagues<sup>10</sup> partly addressed these issues by comparing estimates from several USA cities over five decades, while using flexible exposure—response functions and adjusting for underlying trends. However, that study was performed in a single country, and its estimates of the role of air conditioning can be affected by the lack of separation between spatial and temporal contrasts.

In this contribution, we extend the assessment to a multi-country setting and adopting sophisticated longitudinal designs to control for spatial and temporal confounding. Specifically, the analysis makes use of a unique dataset with time series data from 331 locations in four countries (US, Japan, Canada, and Spain) in the period 1972-2019, and applies novel two-stage methods based on multilevel multivariate spatio-temporal metaregression models.

#### METHODS

#### Data

We collated data on mortality, temperature, and air conditioning prevalence from multiple locations in the four countries (see eTable 1). For each location the data consist of daily counts of all-cause (Canada, Japan, and Spain) or non-accidental (US) mortality and temperature series in summer months (June to September), and air conditioning prevalence from survey data in multiple years within the study period. Table 1 lists the study locations, the observation period as well as the air conditioning variable and surveys used to derive air conditioning prevalences in the four countries included in this study. Across countries air conditioning prevalence data comes from different surveys with different frequency of reporting (see eAppendix). More detailed information on the data collected in each country are reported in the eAppendix.

#### Statistical methods

The analytical strategy was based on three steps, briefly summarized here and described in detail below. In the first step, each country-specific study interval was split into multiple periods. Then, we fitted separate regression models to obtain estimates of heat–mortality associations for each location and period. In addition, we reconstructed location-specific air conditioning trends and assigned prevalence estimates to each location or period unit. In the second step, we pooled the set of coefficients defining the associations to evaluate changes in heat-related mortality risks by calendar year and air conditioning prevalence, accounting for both within- and between-city variations. Finally, in the third and last step, we used the coefficients of the meta-regression models to derive trends in relative risk (RR)

and attributable fractions (AF%) predicted using observed and alternative scenarios of air conditioning prevalence trends.

#### Step 1: Estimating location and period-specific air conditioning prevalence and risks

In the first step, for each location, we divided the observation time was divided into specific time intervals. The number and the different periods for each country are reported in eTable 2. Time intervals have a length of 4 or 5 years. The length of time intervals was chosen a priori in order to provide enough statistical power to derive period-specific estimates, and enough time points to detect changes over time. For each country and locations, using the original air conditioning data, which was assessed intermittently, we estimated the air conditioning prevalence for each period, as described in the eAppendix. Briefly, for the US, Canada, and Spain, we fitted a linear mixed-effects model with a B-spline parameterization of the time variable (years), and city as grouping level. We used best linear unbiased prediction estimates were used to predict yearly air conditioning prevalence in mid-summer (1st of July) in each city of the three countries. For Japan, we used the original yearly data, and assigned it to mid-summer. To assessed if changes in reporting air conditioning prevalence over time affected the predicted trends we performed a sensitivity analysis including an indicator that defines pre- and post-periods corresponding to implementation of the new reporting methods (see eAppendix).

We estimated the location and period-specific temperature–mortality associations through quasi-Poisson regression<sup>14</sup> with distributed lag non-linear models (DLNMs).<sup>15</sup> Based on previous work<sup>16</sup>, we specified the cross-basis function of daily mean temperature using a quadratic B-spline function for the temperature dimension, with two internal knots at the 50th and 90th percentiles of the location and period-specific summer temperature

distributions, and unconstrained parameterization over lag 0-2. To control for long-term trends and residual seasonality, we included interaction terms between a natural cubic Bspline function with 4 degrees of freedom (df) of the day of the year and indicators of year, along with an indicator of day of the week. We tested these modelling choices in a sensitivity analysis.

#### Step 2: modelling spatial and temporal variation in risk

The location and period-specific estimates obtained from the quasi-Poisson model in Step 1 were then combined using multilevel multivariate spatio-temporal models that consider possible non-independence of estimates within each location.<sup>17</sup> For each location i =1, ..., m and year  $t = 1, ..., T_i$  (defined as mid-points of periods), we obtained a k = 4 length column vector of spline coefficients  $\theta_{it}$  representing the temperature–mortality association cumulated over lag 0-2 in location i and period t, and associated  $k \times k$  estimated (co)variance matrix  $S_{it}$ . The multilevel multivariate spatio-temporal meta-regression model for the multivariate vector response  $\theta_{it}$  can be written as:

$$\theta_{it} = X_{it}\beta + Z_ib_i + \varepsilon_{it} \tag{1}$$

with 
$$b_i \sim N(0, \Psi_1)$$
, and  $\varepsilon_{it} \sim N(0, S_{it})$ .

The matrix *X<sub>it</sub>* in the meta-regression model in (1) included fixed-effect predictors, represented by indicators of country, calendar year, period specific average and interquartile range (IQR) of daily mean temperature, in addition to air conditioning prevalence. Temperature variables were selected following previous evidence of their role in modifying heat-related mortality risks, while a linear term for calendar year was included to control for underlying variations in risk unrelated to air conditioning use. We compared the role of different fixed-effect predictors through likelihood ratio (LR) test in models fitted with a maximum likelihood (ML) estimator. We included random terms at city or prefecture level, represented by indicators  $Z_i$  with random coefficients  $b_i$ . The random coefficients have unstructured (co)variance matrices  $\Psi_1$ . The term  $S_{it}$  represents the estimation error within location/period combinations. A restricted maximum likelihood estimator was used for the final model.

This modeling approach allows investigation of the independent effect of changes over time in air conditioning prevalence on the temperature—mortality association, while adjusting for country and location-specific trends. Using random terms at location level allows the use of information both within and between locations.

#### Step 3: quantifying heat-related risks and AC contribution

The estimated fixed-effects coefficients  $\hat{\beta}$  from the multilevel multivariate spatio-temporal meta-regression model (1) fitted in Step 2 can be used to predict a set of spline coefficients  $\hat{\theta}_{ct}$  that represent pooled heat–mortality association curves for any combination of country, year, and air conditioning prevalence. Specifically, associations were predicted longitudinally or at the end of country-specific study periods, either using observed values of meta-predictors or under specific scenarios of air conditioning prevalence. Results were first reported in terms of country-averaged relative risk (RR), using country-specific temperature distributions and minimum mortality temperature as references. In addition, we also derived summaries corresponding to estimated mortality fractions (in percentage) attributed to summer heat for each country/sub-period, following a procedure described elsewhere.<sup>18</sup> In brief, we computed the mortality attributable to heat first by summing the temperature-related deaths occurring in days with temperatures higher than the location

total number of deaths. We calculated empirical standard error (SE) using Monte Carlo simulations, assuming a multivariate normal distribution of the fixed-effects coefficients estimated in Step 2.<sup>18</sup>

#### RESULTS

#### **Data description**

During the study period, more than 23 million deaths were registered in the 331 locations assessed in the four countries. On average, air conditioning prevalence increased in all countries (Figure 1), with the highest prevalence at the end of the study period observed in Japan (89.2%), followed by the USA (82.8%), Canada (48.8%), and Spain (26.9%).

#### Multilevel multivariate spatio-temporal meta-regression model

The results of meta-regression models with different fixed-effects specifications are shown in eTable 3. In the final specification of the multilevel multivariate spatio-temporal metaregression model, air conditioning prevalence shows an independent association with heatrelated risks (p-value = 0.011), while accounting for country-specific trends and adjusting also by locations and period-specific average and interquartile range of mean temperature. We did not find strong evidence of a differential effect of air conditioning prevalence between countries (p-value = 0.084). Inspection of distribution of the residuals and their scatter plot versus time and air conditioning prevalence suggested a good fit of the model (see eFigure 3).

#### Quantification of the heat-related risk and its trend

Figure 2 represents the changes in the heat–mortality association curves predicted by spatio-temporal meta-regression, at the beginning and end of the study periods in the four countries. Japan showed a strong attenuation in risk, with a decline of the RRs across almost all the summer temperature range. The US and Spain also displayed a decrease in risk,

although more evident at highest temperature percentiles. Canada showed little evidence of a reduction in heat-related RR over the observed period.

Table 2 presents air conditioning prevalence, estimated RR at 99<sup>th</sup> percentile of the temperature distribution versus minimum mortality temperature, and estimated excess mortality by country and calendar year. The trend is consistent with the attenuation in risk, especially in Japan where the RR declined from 1.32 to 1.08 during the period 1975-2007. In the same period, the heat-related excess deaths reduced from 3.57% to 1.10%. A reduction in RR is also evident in the USA and Spain, with a reduction of excess deaths due to heat from 0.54% to 2.78% in Spain, and 1.70% to 0.53% in the USA. In Canada, there was no evidence of reduction of the RR corresponding to the 99<sup>th</sup> temperature percentile, but we observed a decrease in mortality fraction attributable to heat, from 1.40% to 0.80%, due to an attenuation in risk at lower temperature percentiles (90<sup>th</sup> and 50<sup>th</sup>), as shown in eFigure 2.

Temporal changes in temperature-related risks are generated by both variation in air conditioning prevalence and underlying trends due to other factors. In order to quantify the role of air conditioning, we fixed the calendar year at the end of the study period and calculated the RR at 99<sup>th</sup> temperature percentile and heat-related mortality fraction for different levels of air conditioning prevalence (Table 3). Results indicate that increasing the AC prevalence from 30% to 80% would be associated with important reduction in heat-related death: 30.2% in the US, 24.9% in Canada, 20.3% in Japan, and 8.8% in Spain.

Finally, in order to separate and quantify the contribution of air conditioning prevalence from other time-varying factors in attenuating heat-related risks, we compared the excess mortality under scenarios of observed increase or no change in air conditioning prevalence

(Figure 3). The dark and light blue bars represent the excess mortality fraction calculated at the beginning and at the end of the study periods, using the actual air conditioning prevalences, with figures reported in Table 2. The middle blue bar represent instead the excess mortality fraction at the end of study period assuming no change in air conditioning prevalence: the comparison indicates that an increased air conditioning prevalence is responsible for only part of the observed attenuation, corresponding approximately to 16.7% in Canada, 20.0% in Japan, 14.3% in Spain, and 16.7% in the US. These results suggest that other adaptation factors can be equally and, in some cases, more important for explaining the decreasing trend (see eTable 4).

#### DISCUSSION

Our results on air conditioning prevalence in Japan, the US, Canada, and Spain are consistent with the hypothesis that air conditioning reduces heat-related mortality. This reduction occurs on top of variations in heat-related health risks possibly associated with planned and unplanned adaptation processes other than air conditioning use. These independent adaptation pathways were quantified and compared using alternative scenarios of air conditioning prevalence and underlying temporal trends. These scenarios indicate that while the increase in air conditioning use is associated with a reduction in heatrelated mortality, this only explains a part of the decline in risk experienced in some countries, and other adaptation pathways have had a more important role in reducing the health burden.

Our results are consistent with published epidemiological investigations that have reported a substantial attenuation of heat-related health risk.<sup>1,2,19</sup> In particular, similar declining trends were observed in the US<sup>6,7,10,16,20-24</sup>, Japan<sup>8,9,25</sup>, Spain<sup>26</sup>, and Canada<sup>16</sup>. Similar

declining trends were also observed in Sweden<sup>27</sup>, Austria<sup>28</sup>, UK<sup>29,30</sup>, Netherlands<sup>31</sup>, nine European cities<sup>32</sup>, and Korea<sup>33,34</sup>, but not in China.<sup>35</sup>

Previous studies have evaluated the protective effect of air conditioning on heat-related risks. Some assessments used cohort<sup>12</sup> and case–control study designs<sup>13</sup>, and suggested a role of AC in reducing the heat-related mortality risks in the USA. These studies were followed by two-stage studies in which the first-stage estimates obtained through caseonly<sup>11</sup> or time-series analyses<sup>5</sup> in multiple cities were combined using meta-regression models with air conditioning prevalence as a contextual variable. These studies confirmed the protective effect of air conditioning in the USA, but were prone to ecologic confounding as the selected cities can differ by other unmeasured characteristics (e.g., demographic, socio-economic, and infrastructural) related to health risk. More recent studies in the US and Japan used a longitudinal design to disentangle the effect of air conditioning as behavioural adaptive measure. In the US, two studies found an independent protective effect of air conditioning<sup>5,6</sup>, but Bobb and colleagues observed no evidence of protective effect.<sup>7</sup> The longitudinal study of Nordio and colleagues<sup>10</sup> reported independent protective effects of air conditioning while controlling for region, time trend, and mean summer temperature, using spline models in individual cities and a meta-regression approach. The two longitudinal studies conducted in Japan did not find evidence consistent with an independent protective effect of air conditioning over the declining heat-related risk trend.<sup>8,9</sup> Differences on previous studies results can be partly explained by low statistical power, as these investigations were conducted in a single country and/or the temperaturemortality curve was summarized using simplified indices. Moreover, these studies did not jointly consider the longitudinal and spatial structure of the data, and the nonindependence of the observations within locations.

Our study has several strengths. First, we used distributed lag non-linear modeling techniques to estimate the heat-mortality association. This modeling framework helps avoid biases due to simplification of the exposure–response association and considers possible lagged effects of heat on mortality.<sup>15</sup> Second, we were able to collect mortality, temperature and air conditioning data for 331 locations in four countries for a period of four decades. This provided large variability in air conditioning prevalence both within and across locations, offering sufficient statistical power to isolate the impact on modifying heatmortality relationships. Third, we used a study design based on both spatial and longitudinal comparison, reducing the chance of ecologic bias and temporal confounding due to concurrent changes in other modifying factors, such as socio-economic conditions and access to health care. The spatial component provides increased variability in response and exposure, while the longitudinal design compares variations in risk within a location. Finally, we used novel multilevel multivariate spatio-temporal meta-regression models that allow disentangling of the reduction in heat-related risk associated to the increase in air conditioning prevalence from underlying trends due to other adaptation pathways, while at the same time correctly accounting for correlations between repeated measures taken within the same location.<sup>17</sup>

We must acknowledge some limitations. First, the results of our study refer to developed countries with predominantly temperate or continental climates. Caution should be used when extrapolating results to low-income countries, which are characterized by different climatic, socio-demographic, and development conditions, and where technology based adaptation measures, such as increasing air conditioning use, may be problematic as many low-income countries already experience chronic shortages of power.<sup>2</sup> Second, we reconstructed air conditioning prevalence along the past decades by applying smoothing

techniques to irregular survey data from multiple sources. However, additional analyses described in the eAppendix show that results are robust to this filling-up procedure. The results of the sensitivity analysis suggest that the smoothing process could have introduced some error, although it is unlikely that this is correlated with the estimated period-specific risk, and therefore can probably be assumed as random. Third, our air conditioning variable is defined as presence of air conditioning units or central air conditioning at home, but does not capture its actual use. Moreover, this measure is not informative about air conditioning use in other environments, such as on public transport, stores, workplaces, and public areas. This may induce some additional problems in the interpretation of the results.

The analysis of factors related to changes in susceptibility to temperature-related mortality is critical to inform health and climate policies. Air conditioning is a solution to regulate ambient indoor temperatures and lower the heat stress imposed on the human thermoregulatory function<sup>36</sup>, and it represents one of the most cited behavioural adaptation strategy to climate change.<sup>37</sup> The results of our analysis confirm that air conditioning is an effective adaptive measure and have contributed to reduce the burden of heat-related mortality. According to our estimates in the USA and Japan, nearly 0.09% and 0.32% of deaths during summer months were delayed by increasing the air conditioning market seems to have reached a plateau, but the heat-related mortality is still substantial. However, the quantitative comparison of the contribution of increase in air conditioning prevalence, and the independent attenuation of the risk reported in Figure 3, suggest that other adaptation pathways can be equally or even more effective in reducing the health burden. In Spain and Canada, the delayed deaths during summer months were both 0.05%, suggesting a further margin on reduction of heat-related mortality, especially in Spain where the reported air

conditioning prevalence reaches only 30% in 2009. In addition, increasing air conditioning use has also important negative consequences, including capital and energy cost, carbon and pollution-generating energy demand, and contribution to the heat-island effect.<sup>2</sup> However, the current rapid transition of electricity generation to carbon zero sources is likely to ameliorate the pollution impact in the next few decades. A quantitative assessment of health and economic impacts of this and other adaptive changes is critical for generating plausible scenarios of potential mitigation and adaptation benefit and costs.

In conclusion, in this study we found a reduction over time of the heat-related health risk in Japan, the USA, and Spain. Air conditioning prevalence was factor that independently explained part of the decrease in heat-related deaths, although we estimated that other adaptive strategies accounted for a larger proportion of the attenuation. These results can be used to inform policy measures based at individual, community, and international level, and to improve and extend projections of future heat impacts on human health.

# FIGURES



Figure 1. Air conditioning (AC) prevalence (%) by year in Canada, Japan, Spain and the US.

**Figure 2**. Country-average exposure—response curves (in relative risk, RR) predicted at the beginning and end of the study periods in Canada, Japan, Spain, and the US. The x-axis represents relative temperatures in percentiles, but rescaled using the average distribution of absolute temperature across cities in each country.



**Figure 3**. Excess mortality associated to heat reported as attributable fraction (AF%) estimated at the beginning (Baseline, dark blue) and end of the study period assuming no change (End-Study period with fixed air conditioning, medium blue) or with the observed change (End-Study period, light blue) in air conditioning (AC) prevalence.



## REFERENCES

- 1. Arbuthnott K, Hajat S, Heaviside C, Vardoulakis S. Changes in population susceptibility to heat and cold over time: assessing adaptation to climate change. *Environmental Health* 2016;**15**(1):S33.
- 2. Kinney PL. Temporal Trends in Heat-Related Mortality: Implications for Future Projections. *Atmosphere* 2018;**9**(10):409.
- 3. Hondula DM, Balling RC, Vanos JK, Georgescu M. Rising temperatures, human health, and the role of adaptation. *Current Climate Change Reports* 2015;**1**(3):144-154.
- 4. II IPoCCWG. *Climate change 2014: Impacts, adaptation, and vulnerability* IPCC Working Group II, 2014.
- 5. Anderson BG, Bell ML. Weather-related mortality: how heat, cold, and heat waves affect mortality in the United States. *Epidemiology* 2009;**20**(2):205.
- 6. Barreca A, Clay K, Deschenes O, Greenstone M, Shapiro JS. Adapting to climate change: The remarkable decline in the US temperature–mortality relationship over the twentieth century. *Journal of Political Economy* 2016;**124**(1):105-159.
- 7. Bobb JF, Peng RD, Bell ML, Dominici F. Heat-related mortality and adaptation to heat in the United States. *Environmental health perspectives* 2014;**122**(8):811-816.
- Chung Y, Yang D, Gasparrini A, Vicedo-Cabrera AM, Fook Sheng Ng C, Kim Y, Honda Y, Hashizume M. Changing Susceptibility to Non-Optimum Temperatures in Japan, 1972-2012: The Role of Climate, Demographic, and Socioeconomic Factors. *Environ Health Perspect* 2018;**126**(5):057002.
- 9. Ng CFS, Boeckmann M, Ueda K, Zeeb H, Nitta H, Watanabe C, Honda Y. Heat-related mortality: effect modification and adaptation in Japan from 1972 to 2010. *Global environmental change* 2016;**39**:234-243.
- 10. Nordio F, Zanobetti A, Colicino E, Kloog I, Schwartz J. Changing patterns of the temperature–mortality association by time and location in the US, and implications for climate change. *Environment international* 2015;**81**:80-86.
- 11. Ostro B, Rauch S, Green R, Malig B, Basu R. The effects of temperature and use of air conditioning on hospitalizations. *Am J Epidemiol* 2010;**172**(9):1053-61.
- 12. Rogot E, Sorlie PD, Backlund E. Air-conditioning and mortality in hot weather. *Am J Epidemiol* 1992;**136**(1):106-16.
- 13. Semenza JC, Rubin CH, Falter KH, Selanikio JD, Flanders WD, Howe HL, Wilhelm JL. Heat-related deaths during the July 1995 heat wave in Chicago. *N Engl J Med* 1996;**335**(2):84-90.
- 14. Bhaskaran K, Gasparrini A, Hajat S, Smeeth L, Armstrong B. Time series regression studies in environmental epidemiology. *International journal of epidemiology* 2013;**42**(4):1187-1195.
- 15. Gasparrini A. Modeling exposure–lag–response associations with distributed lag nonlinear models. *Statistics in medicine* 2014;**33**(5):881-899.
- 16. Gasparrini A, Guo Y, Hashizume M, Kinney PL, Petkova EP, Lavigne E, Zanobetti A, Schwartz JD, Tobias A, Leone M. Temporal variation in heat–mortality associations: a multicountry study. *Environmental health perspectives* 2015;**123**(11):1200-1207.
- 17. Sera F, Armstrong B, Blangiardo M, Gasparrini A. An extended mixed-effects framework for meta-analysis. *Stat Med* 2019;**38**(29):5429-5444.
- 18. Gasparrini A, Leone M. Attributable risk from distributed lag models. *BMC medical research methodology* 2014;**14**(1):55.

- 19. Sheridan SC, Allen MJ. Temporal trends in human vulnerability to excessive heat. *Environmental research letters* 2018;**13**(4):043001.
- 20. Barnett AG. Temperature and cardiovascular deaths in the US elderly: changes over time. *Epidemiology* 2007;**18**(3):369-372.
- 21. Davis RE, Knappenberger PC, Michaels PJ, Novicoff WM. Changing heat-related mortality in the United States. *Environmental health perspectives* 2003;**111**(14):1712-1718.
- 22. Donaldson G, Keatinge W, Näyhä S. Changes in summer temperature and heatrelated mortality since 1971 in North Carolina, South Finland, and Southeast England. *Environmental research* 2003;**91**(1):1-7.
- 23. Guo Y, Barnett AG, Tong S. High temperatures-related elderly mortality varied greatly from year to year: important information for heat-warning systems. *Scientific reports* 2012;**2**:830.
- 24. Petkova EP, Gasparrini A, Kinney PL. Heat and mortality in New York City since the beginning of the 20th century. *Epidemiology (Cambridge, Mass.)* 2014;**25**(4):554.
- 25. Onozuka D, Hagihara A. Variation in vulnerability to extreme-temperature-related mortality in Japan: A 40-year time-series analysis. *Environmental research* 2015;**140**:177-184.
- 26. Martínez-Solanas È, Basagaña X. Temporal changes in temperature-related mortality in Spain and effect of the implementation of a Heat Health Prevention Plan. *Environmental research* 2019;**169**:102-113.
- 27. Åström DO, Forsberg B, Edvinsson S, Rocklöv J. Acute fatal effects of short-lasting extreme temperatures in Stockholm, Sweden: evidence across a century of change. *Epidemiology* 2013;**24**(6):820-829.
- 28. Matzarakis A, Muthers S, Koch E. Human biometeorological evaluation of heatrelated mortality in Vienna. *Theoretical and applied climatology* 2011;**105**(1-2):1-10.
- 29. Carson C, Hajat S, Armstrong B, Wilkinson P. Declining vulnerability to temperaturerelated mortality in London over the 20th century. *American journal of epidemiology* 2006;**164**(1):77-84.
- 30. Christidis N, Donaldson GC, Stott PA. Causes for the recent changes in cold-and heatrelated mortality in England and Wales. *Climatic Change* 2010;**102**(3-4):539-553.
- 31. Ekamper P, Van Poppel F, Van Duin C, Garssen J. 150 Years of temperature-related excess mortality in the Netherlands. *Demographic Research* 2009;**21**:385-426.
- de'Donato F, Leone M, Scortichini M, De Sario M, Katsouyanni K, Lanki T, Basagaña X, Ballester F, Åström C, Paldy A. Changes in the effect of heat on mortality in the last 20 years in nine European cities. Results from the PHASE project. *International journal of environmental research and public health* 2015;**12**(12):15567-15583.
- 33. Ha J, Kim H. Changes in the association between summer temperature and mortality in Seoul, South Korea. *International journal of biometeorology* 2013;**57**(4):535-544.
- 34. Lee W, Choi HM, Kim D, Honda Y, Guo Y-LL, Kim H. Temporal changes in mortality attributed to heat extremes for 57 cities in Northeast Asia. *Science of the total environment* 2018;**616**:703-709.
- 35. Yang C, Meng X, Chen R, Cai J, Zhao Z, Wan Y, Kan H. Long-term variations in the association between ambient temperature and daily cardiovascular mortality in Shanghai, China. *Science of the Total Environment* 2015;**538**:524-530.
- 36. Deschenes O. Temperature, human health, and adaptation: A review of the empirical literature. *Energy Economics* 2014;**46**:606-619.

37. O'Neill MS. Air conditioning and heat-related health effects. *Appl Environ Sci Public Health* 2003;**1**(1):9-12.

## TABLES

**Table 1**. Geographical boundaries, observation period, and definition of air conditioning prevalence in each country.

Country	Locations	Period	Air conditioning variable	Survey
Canada	20 census	1991-	Proportion of dwellings	-Survey of Household
	metropolitan	2009	with an air conditioning	& Energy Use (SHEU) <sup>1</sup>
	areas + city of		system (central or with a	-Households and
	Hamilton		window or room	Environment Survey
			mounted air	(HES) <sup>2</sup>
			conditioning system)	
Japan	47 prefectures	1972-	Proportion of	-Regional statistics
		2009	households with two or	database <sup>3</sup>
			more occupants with air	
			conditioning	
Spain	52 capital cities	1990-	Proportion of family	-Population and
		2009	homes with	Housing Census <sup>4</sup>
			"refrigeration"; and	-"Life Conditions"
			from 2007 Proportion of	Survey <sup>5</sup>
			"homes with air	
			conditioning"	
USA	211	1973-	Proportion of	-Census of
	metropolitan	2006	households in each	Population <sup>6</sup>
	areas		metropolitan area with	-American Housing
			central air conditioning	Survey (AHS) <sup>7</sup>
				- Residential Energy
				Consumption Survey <sup>8</sup>

<sup>1</sup>Estimates at regional level in years 1993, 1997, 2003

<sup>2</sup> Estimates at city level in years 2006, 2007, 2009

<sup>3</sup> Asahi Newspaper Publishing.2015

 $^{\rm 4}\,{\rm Estimates}$  at city level in years 1991 and 2001

<sup>5</sup> Estimates at regional level in 2007

<sup>6</sup> Estimates before 1985 at city level

<sup>7</sup> AHS use a rotation sampling of cities; data available yearly from 1985

<sup>8</sup> Used to estimate air conditioning prevalence in northern New England cities

**Table 2**. Reconstructed air conditioning (AC) prevalence, relative risk (RR) at 99th percentile of the temperature distribution versus minimum mortality temperature, and attributed mortality fraction AF% by country and year.

	N/	1.00/		
Country	Year	AC%	99 <sup>111</sup> RR 95%CI	AF% 95%CI
Canada	1994	30.1	1.13 (1.09, 1.17)	1.40 (1.23 ;1.55)
	1998	35.5	1.12 (1.08; 1.16)	1.33 (1.20; 1.44)
	2003	41.9	1.11 (1.08; 1.14)	1.22 (1.05; 1.38)
	2008	48.8	1.11 (1.07; 1.16)	0.80 (0.59; 0.98)
Japan	1975	15.9	1.32 (1.29; 1.34)	3.57 (3.53; 3.61)
	1979	31.1	1.28 (1.26; 1.30)	3.13 (3.10; 3.17)
	1983	41.3	1.24 (1.23; 1.26)	2.83 (2.79; 2.86)
	1987	52.3	1.21 (1.19; 1.22)	2.52 (2.49; 2.56)
	1991	64.1	1.18 (1.16; 1.19)	2.24 (2.20; 2.28)
	1995	73.7	1.15 (1.13; 1.16)	1.90 (1.86; 1.94)
	1999	81.3	1.12 (1.11; 1.14)	1.70 (1.66; 1.75)
	2003	86.0	1.10 (1.08; 1.11)	1.43 (1.39; 1.46)
	2007	89.2	1.08 (1.06; 1.10)	1.10 (1.05; 1.14)
Spain	1993	9.0	1.37 (1.32; 1.42)	3.54 (3.38; 3.69)
	1998	12.9	1.42 (1.37; 1.46)	3.54 (3.42; 3.65)
	2003	19.2	1.35 (1.32; 1.39)	3.51 (3.41; 3.60)
	2007	26.9	1.26 (1.22; 1.31)	2.78 (2.63; 2.92)
USA	1975	49.4	1.14 (1.13; 1.15)	1.70 (1.67; 1.73)
	1979	56.5	1.13 (1.12; 1.14)	1.56 (1.54; 1.58)
	1984	64.1	1.11 (1.10; 1.12)	1.32 (1.30; 1.33)
	1989	71.0	1.09 (1.08; 1.10)	1.09 (1.07; 1.10)
	1994	76.8	1.08 (1.07; 1.09)	0.88 (0.87; 0.90)
	1999	80.7	1.06 (1.05; 1.07)	0.67 (0.65; 0.68)
	2004	82.8	1.05 (1.04; 1.06)	0.53 (0.51; 0.55)

**Table 3**. Predicted relative risk (RR) at 99th temperature percentile, and attributed mortality fraction (AF%) with 95% confidence intervals (CI) calculated at the end of the study period for four scenarios of air conditioning prevalence levels (30%, 55%, 80% and 100%) in Canada, Japan, Spain, and the USA

Country; Year	AC%	RR	AF%
Canada (2008)	30%	1.12 (1.07;	0.93 (0.75;
		1.17)	1.10)
	55%	1.11 (1.06;	0.82 (0.63;
		1.15)	1.00)
	80%	1.09 (1.05;	0.70 (0.51;
		1.14)	0.89)
	100%	1.08 (1.03;	0.61 (0.40;
		1.13)	0.80)
Japan (2007)	30%	1.12 (1.09;	1.48 (1.41;
		1.14)	1.54)
	55%	1.10 (1.08;	1.33 (1.28;
		1.12)	1.37)
	80%	1.08 (1.07;	1.18 (1.13;
		1.10)	1.22)
	100%	1.07 (1.06;	1.06 (1.01;
		1.09)	1.10)
Spain (2007)	30%	1.26 (1.22;	2.86 (2.70;
		1.31)	2.99)
	55%	1.24 (1.20;	2.73 (2.58;
		1.29)	2.87)
	80%	1.23(1.18; 1.28)	2.61 (2.45;
			2.77)
	100%	1.21 (1.16;	2.50 (2.32;
		1.27)	2.66)
USA (2004)	30%	1.07 (1.05;	0.82 (0.79;
		1.09)	0.84)
	55%	1.06 (1.05;	0.69 (0.67;
		1.07)	0.71)
	80%	1.05 (1.04;	0.57 (0.55;
		1.06)	0.59)
	100%	1.04 (1.03;	0.47 (0.45;
		1.05)	0.49)