Journal Pre-proof

Reported symptoms differentiate diagnoses in children with exercise-induced respiratory problems: findings from the Swiss Paediatric Airway Cohort (SPAC)

Eva SL. Pedersen, PhD., Carmen CM. de Jong, MD, Cristina Ardura-Garcia, MD PhD, Maria Christina Mallet, MBChB, Juerg Barben, MD, Carmen Casaulta, MD, Karin Hoyler, MD, Anja Jochmann, MD, Alexander Moeller, MD, Dominik Mueller-Suter, MD, Nicolas Regamey, MD, Florian Singer, MD PhD, Myrofora Goutaki, MD PhD, Claudia E. Kuehni, MD PhD

PII: S2213-2198(20)30959-4

DOI: <https://doi.org/10.1016/j.jaip.2020.09.012>

Reference: JAIP 3121

To appear in: *The Journal of Allergy and Clinical Immunology: In Practice*

Received Date: 12 March 2020

Revised Date: 8 September 2020

Accepted Date: 9 September 2020

Please cite this article as: Pedersen ES, de Jong CC, Ardura-Garcia C, Mallet MC, Barben J, Casaulta C, Hoyler K, Jochmann A, Moeller A, Mueller-Suter D, Regamey N, Singer F, Goutaki M, Kuehni CE, Reported symptoms differentiate diagnoses in children with exercise-induced respiratory problems: findings from the Swiss Paediatric Airway Cohort (SPAC), *The Journal of Allergy and Clinical Immunology: In Practice* (2020), doi: <https://doi.org/10.1016/j.jaip.2020.09.012>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Inc. on behalf of the American Academy of Allergy, Asthma & Immunology

Reported symptoms differentiate diagnoses in children with exercise-induced respiratory problems: findings from the Swiss Paediatric Airway Cohort (SPAC)

Authors

Eva SL Pedersen (eva.pedersen@ispm.unibe.ch), PhD.¹, Carmen CM de Jong (carmen.dejong@ispm.unibe.ch) MD¹, Cristina Ardura-Garcia (Cristina.ardura@ispm.unibe.ch) MD PhD¹, Maria Christina Mallet (maria.mallet@ispm.unibe.ch) MBChB¹, Juerg Barben (juerg.barben@kispi.sg.ch) MD², Carmen Casaulta (CarmenAnnemarie.Casaulta@ksgr.ch) MD³, Karin Hoyler (karin.hoyler@swissonline.ch) MD⁴, Anja Jochmann (anja.jochmann@ukbb.ch) MD⁵, Alexander Moeller (alexander.moeller@kispi.uzh.ch) MD⁶, Dominik Mueller-Suter (Dominik.Mueller-Suter@ksa.ch) MD⁷, Nicolas Regamey (nicolas.regamey@luks.ch) MD⁸, Florian Singer (florian.singer@insel.ch) MD PhD^{9, 10}, Myrofora Goutaki (Myrofora.goutaki@ispm.unibe.ch) MD PhD^{1,9}, Claudia E. Kuehni (Claudia.kuehni@ispm.unibe.ch) MD PhD^{1,9}

Corresponding Author

Claudia E Kuehni, Institute of Social and Preventive Medicine, University of Bern, Mittelstrasse 43, 3012 Bern, Tel: +41 31 631 35 07
E-mail: Claudia.kuehni@ispm.unibe.ch

Affiliations

¹ Institute of Social and Preventive Medicine, University of Bern, Switzerland

² Division of Paediatric Pulmonology, Children's Hospital St. Gallen, Switzerland

³ Division of Paediatric Pulmonology, Children's Hospital Chur, Chur, Switzerland

⁴ Paediatric Pulmonology Practice Horgen, Horgen, Switzerland

⁵ Division of Paediatric Pulmonology, University Children's Hospital, University of Basel, Switzerland

⁶ Division of Paediatric Pulmonology, University Children's Hospital Zurich, Switzerland

⁷ Division of Paediatric Pulmonology, Kantonsspital Aarau, Switzerland

⁸ Division of Paediatric Pulmonology, Children's Hospital Lucerne, Switzerland

⁹ Paediatric Respiratory Medicine, Children's University Hospital of Bern, University of Bern, Switzerland

¹⁰ PedNet, Children's University Hospital of Bern, University of Bern, Switzerland

35

36

37 Disclosure statement

38 Dr. Singer reports personal fees from Novartis, personal fees from Vertex, outside the
39 submitted work. All other authors declare that they have no competing interests.

40

41 Funding Sources

42 This work was funded by the Swiss National Science Foundation (SNSF 32003B_162820) and
43 the Swiss Lung Association (2019-02 641670). Further funding to develop the SPAC cohort
44 came from the Allergiestiftung U. Müller-Gierok and the Lung League St. Gallen.

45

46 Word count

47 Abstract: 246 words

48 Manuscript: 2592 words

Abstract (246 words)

Background: Exercise-induced breathing problems with similar clinical presentations can have different etiologies. This makes distinguishing common diagnoses such as asthma, extrathoracic and thoracic dysfunctional breathing (DB), insufficient fitness, and chronic cough difficult.

Objective: We studied which parent-reported, exercise-induced symptoms can help distinguish diagnoses in children seen in respiratory outpatient clinics.

Methods: This study was nested in the Swiss Paediatric Airway Cohort (SPAC), an observational study of children aged 0-17 years referred to pediatric respiratory outpatient clinics in Switzerland. We studied children aged 6-17 years and compared information on exercise-induced symptoms from parent-completed questionnaires between children with different diagnoses. We used multinomial regression to analyze whether parent-reported symptoms differed between diagnoses (asthma as base).

Results: Among 1109 children, exercise-induced symptoms were reported for 732 (66%) (mean age 11 years, 318 of 732 [43%] female). Among the symptoms, dyspnea best distinguished thoracic DB (relative risk ratio [RRR] 5.4, 95%CI 1.3-22) from asthma. Among exercise triggers, swimming best distinguished thoracic DB (RRR 2.4, 95%CI 1.3-6.2) and asthma plus DB (RRR 1.8, 95%CI 0.9-3.4) from asthma only. Late onset of symptoms was less common for extrathoracic DB (RRR 0.1, 95%CI 0.03-0.5) and thoracic DB (RRR 0.4, 95%CI 0.1-1.2) compared with asthma. Localization of dyspnea (throat vs. chest) differed between extrathoracic DB (RRR 2.3, 95%CI 0.9-5.8) and asthma. Reported respiration phase (inspiration or expiration) did not help distinguish diagnoses.

71 **Conclusion:** Parent-reported symptoms help distinguish different diagnoses in children with
72 exercise-induced symptoms. This highlights the importance of physicians obtaining detailed
73 patient histories.

Highlights box

1. What is already known about this topic?

Experts suggest that information about the symptoms and their onset and duration can assist accurate diagnosis of children with exercise-induced respiratory problems, but no original studies have tested this.

2. What does this article add to our knowledge?

Exercise-induced symptoms reported by parents and further information about their onset, triggers, and effects of treatment help differentiate diagnoses in children with exercise-induced respiratory problems.

3. How does this study impact current management guidelines?

Our results emphasize the importance of taking detailed symptom histories of children with exercise-induced problems, and suggest which questions are most helpful.

Key words

Exercise-induced, ILO, asthma, childhood, adolescents, dysfunctional breathing

89 **List of abbreviations**

90	BMI	Body mass index
91	DB	Dysfunctional breathing
92	FeNO	Fractional exhaled nitric oxide
93	ILO	Inducible laryngeal obstruction
94	RRR	Relative risk ratio
95	SPAC	Swiss Paediatric Airway Cohort

Manuscript: 2592 words

Introduction

Exercise-induced respiratory symptoms are common in childhood. But their underlying causes can be difficult to identify because the clinical presentation of exercise-induced symptoms of different etiologies can overlap (1-3). Exercise-induced symptoms are most often caused by asthma, extrathoracic or thoracic dysfunctional breathing (DB), insufficient fitness, and nonspecific chronic cough (4, 5). Even though clinical presentations can be similar, certain symptoms are typically associated with specific diagnoses (6-8). Knowing which symptoms are particularly characteristic of different underlying causes may help physicians to make a diagnosis, in addition to formal exercise testing (4, 6, 9). For example, expiratory wheeze, cough, and shortness of breath are typical for asthma with symptoms lasting from minutes to hours that usually peak after exercise (10, 11). Inspiratory problems with stridor, throat tightness, and shortness of breath are more typical for extrathoracic DB with symptoms that last only a few minutes and peak during exercise (10, 12). Typical symptoms for thoracic DB are shortness of breath, sighing, dizziness, and symptoms can last from minutes to hours and peak during exercise (9). Tingling in fingers or lips is typical for thoracic DB with hyperventilation.

A few studies have examined the association of diagnoses with typical symptoms. However, exercise-induced symptoms have been reported only for specific diagnostic groups such as children with asthma (13), or no more than two diagnostic groups have been compared (14-16). If we better knew which symptoms most usefully distinguish diagnoses, then misdiagnoses in children with exercise-induced symptoms, such as extrathoracic DB misdiagnosed as asthma, might be reduced (17-19). We therefore studied children visiting

pediatric respiratory outpatient clinics in Switzerland to investigate which symptoms reported by parents are most useful to distinguish different diagnoses of exercise-induced symptoms.

Method

Study design

We used cross-sectional data from the Swiss Paediatric Airway Cohort (SPAC), a multicenter study of children referred to pediatric respiratory outpatient clinics in Switzerland (20). The SPAC study includes children aged 0-17 years who were referred for respiratory problems such as wheeze, cough, dyspnea, or symptoms related to sleep or exercise, and who spoke sufficient German to participate. At the time of the visit, the physicians explained the SPAC study to the families. Parents completed a questionnaire before or shortly after the visit that inquired about symptoms, medication, environment, and health behaviors. After the visit, the SPAC study team collected the outpatient clinic letters that had been sent back to the referring pediatrician with information on diagnosis, diagnostic investigations, and treatment. We entered questionnaire responses and information from outpatient clinic letters into a Research Electronic Data Capture (REDCap) database (21). Recruitment for SPAC started in July 2017 and is ongoing. At the time we extracted data for this analysis, eight pediatric respiratory outpatient clinics in Switzerland were participating. Among 2971 children invited, 1838 (62%) agreed to participate (December 1, 2019).

The SPAC study was approved by the Bern Cantonal Ethics Committee (Kantonale Ethikkommission Bern 2016-02176). Written informed consent was obtained from parents and patients older than 13 years. This article follows the STROBE reporting recommendations (22).

144

145 Inclusion criteria

146 We included children aged 6-17 years with a completed questionnaire and an available
147 outpatient clinic letter with information on diagnosis. We restricted the population to
148 schoolchildren because nearly all children referred for exercise-induced symptoms to
149 respiratory outpatient clinics are older than 5 years. The question used to identify children
150 with exercise-induced symptoms was “Does your child sometimes experience breathing
151 problems during exercise?”

152

153 Parent reported exercise-induced symptoms (EIS)

154 The questionnaire was completed by parents at the first visit to the clinic and inquired about
155 symptoms that included exercise-induced wheeze, cough, dyspnea, tingling sensations in
156 fingertips/lips, and other symptoms that could be reported in a free-text field. It also asked
157 about characteristics of symptoms that included trigger factors (running, bicycle riding,
158 intensive sport games, swimming), localization of dyspnea (chest, throat, or both),
159 respiration phase (inspiration, expiration), onset of symptoms (during or after exercise),
160 duration of symptoms, and whether a short-acting bronchodilator helped to relieve
161 symptoms. **Table E1** reproduces English translations of the questions about exercise-induced
162 symptoms in the German language questionnaire. Parental questionnaires were not
163 disclosed to outpatient physicians.

164

165 Diagnosis

166 Diagnosis was taken from the outpatient clinic letter that the hospital pulmonologists sent
167 back to the referring physician. We distinguished six diagnoses: asthma, extrathoracic DB,

thoracic DB, asthma plus DB, chronic cough, and other (including insufficient fitness level, exercise-induced symptoms of unknown etiology, allergic rhinoconjunctivitis, recurrent respiratory infections, and rare pulmonary diseases). Exact definitions of diagnoses are in **Table E2**. The diagnosis given at the clinic was made by the attending pediatric pulmonologist and supported by at least one pulmonary function test such as spirometry, bodyplethysmography, measurement of fractional exhaled nitric oxide (FeNO), direct or indirect bronchial provocation tests. Some children were seen more than once in the outpatient clinic, and we took the diagnosis from the outpatient clinic with the latest date. If a child had more than one diagnosis listed in the letter, we used the diagnosis listed first, except in children who had asthma and any type of DB. In these children we created a separate category (asthma plus DB) because we believed that symptoms might differ between children with asthma, DB, and both occurring together. Outpatient clinic physicians were blinded to the parent completed questionnaire when giving the final diagnosis.

Other variables

Age, sex, height, and weight were taken from the outpatient clinic letter. We calculated body mass index (BMI) as weight/height^2 (kg/m^2) and calculated age-adjusted BMI z-scores based on Swiss reference values (23), defining overweight as BMI z-score > 1 . We obtained information on symptoms not induced by exercise from the questionnaire including parental education, environmental factors, and physical activity.

Statistical methods

We compared proportions of exercise-induced symptoms by diagnosis categories: asthma, extrathoracic DB, thoracic DB, asthma plus DB, chronic cough, and others using chi-square

and Fisher's exact tests. We studied which symptoms were most useful to distinguish diagnoses using multinomial logistic regression. We defined diagnosis as outcome and asthma as the reference category, and studied each explanatory variable in turn, adjusted for age and sex. For the multinomial regression, due to the sample size we grouped chronic cough with other diagnoses. Overall, we had little missing information in the questionnaire replies (<7%) apart from the question about the respiration phase when symptoms are worst (inspiration or expiration) for which 14% were missing. Children with missing data were excluded. We used STATA version 14 for statistical analysis.

Results

Of the 1109 children aged 6-17 whose parents completed the questionnaire and for whom we had information about diagnosis, 732 (66%) reported exercise-induced symptoms in the questionnaire (**Figure E1**). On average, children were 11 years old (SD 3.2), 318 (43%) were female (**table 1**). Children with reported exercise-induced symptoms were older and more often female than children without reported exercise-induced symptoms. Among these children with exercise-induced symptoms, 549 (75%) were diagnosed with asthma, 38 (5%) with extrathoracic DB, 30 (4%) with thoracic DB, 43 (6%) with asthma plus DB, 21 (3%) with chronic cough, and 51 (7%) received other diagnoses. Overall, more boys than girls were diagnosed with asthma (63% boys) whereas more girls than boys were diagnosed with extrathoracic DB (62% girls), thoracic DB (59% girls), and asthma + DB (61% girls), data not shown.

Symptoms differed between diagnostic groups (**figure 1, table 2, figure 2**). Results from our multinomial regression analysis (adjusted for age and sex) showed that wheeze was

reported less often for children with other diagnoses (relative risk ratio [RRR] 0.2, 95% CI 0.1-0.4) than for children with asthma. Cough was less common in children with thoracic DB (RRR 0.3, 95% CI 0.2-0.7) and asthma plus DB (RRR 0.3, 95% CI 0.2-0.6) than in children with asthma alone. Dyspnea was reported more often for children with thoracic DB (RRR 5.4, 95% CI 1.3-23.1) and asthma plus DB (RRR 4.9, 95% CI 1.5-16.2) than in children with asthma alone. A tingling feeling in fingertips or lips was more common in children with thoracic DB (RRR 3.0, 95% CI 1.2-7.3) than in children with asthma.

The type of physical activity reported to trigger exercise-induced symptoms differed between diagnostic groups (**table 2, figure 3**). Compared to children with asthma, swimming was more commonly reported as trigger in children with thoracic DB (RRR 2.9, 95%CI 1.3-6.2), asthma plus DB (RRR 1.8, 95%CI 0.9-3.4), and other diagnoses (RRR 2.1, 95%CI 1.2-3.4). Bicycle riding was reported more often for children with extrathoracic DB (RRR 2.0, 95%CI 1.0-4.2), and intensive sports games were more often reported for children with asthma plus DB (RRR 2.4, 95%CI 1.0-5.8).

The characteristics of exercise-induced symptoms differed between diagnostic groups (**table 2, figure 4**). Late onset (after exercise) of symptoms was rarely reported for extrathoracic DB (RRR 0.1, 95% CI 0.03-0.5) compared to asthma. A long duration of symptoms (more than 10 minutes) was reported more often for thoracic DB (RRR 4.8, 95% CI 1.4-16.8) than asthma. For localization of dyspnea, throat was reported more often than chest for children with extrathoracic DB (RRR 2.3, 95% CI 0.9-5.8) than for children with asthma. Respiration phase (inspiration or expiration) did not differ between diagnostic groups. Use of a bronchodilator

made symptoms disappear in 172 (43%) children with asthma in contrast to 2 (14%) children with extrathoracic DB and 1 (8%) child with asthma plus DB (**table 2**).

Discussion

Parent-reported symptoms can distinguish different diagnoses in children with exercise-induced symptoms referred to pediatric outpatient clinics. We observed that especially reported exercise-induced cough, dyspnea, and tingling sensation in fingers or lips differed between diagnostic groups. Of the physical activities triggering symptoms, intensive sport games and swimming best distinguished diagnosis groups. Additionally, onset of symptoms, duration of symptoms, and effect of a short-acting bronchodilator differed between the diagnostic groups. Respiration phase (inspiration or expiration) was less helpful.

Strengths and limitations

Information about exercise-induced symptoms and activities that trigger them has not previously been reported in such detail. The comparison of questionnaire-reported exercise-induced symptoms and diagnostic groups, which included asthma, extrathoracic DB, thoracic DB, and the combination of asthma and DB, is this study's major strength. The level of detail afforded examination of how activities trigger different exercise-induced problems. In addition, our study was nested in SPAC, a real-life prospective observational clinical cohort which is representative of children referred to pediatric respiratory outpatient clinics for respiratory problems. We therefore believe our findings can be broadly generalized to children seen by respiratory physicians for exercise-induced symptoms.

A limitation of the study is that we did not analyze if results from diagnostic tests can help distinguish diagnoses additionally to reported symptoms. The SPAC study is embedded in routine care, and therefore some tests (e.g., exercise challenge tests) were performed by indication and therefore not done in all children, and including these results in our analyses would have introduced selection bias. Another limitation is that the questionnaire was addressed to the parents rather than the children themselves. However, we encouraged parents to fill in the questionnaire together with their child, which increases validity of reported symptoms (24, 25). Our questionnaire included the set of questions usually asked by physicians during the consultation. However, physicians might have worded some questions differently, addressed them to the child rather than the mother, simulated respiratory sounds such as wheeze or stridor, or provided additional explanations. So, although collected at the same time, the replies in the parental questionnaire, used for the analysis, might not always mirror the information retrieved by the physician who took the history. Our limited sample size for some diagnostic categories (thoracic DB, n=30) led to wide confidence intervals, and we could not investigate combinations of reported symptoms. Still, our study is the largest of its kind. A further limitation is that the final diagnosis was made by different pulmonologists and not based on a standardized, predefined diagnostic algorithm specific for this study. However, all pulmonologists were board-certified and diagnoses were based on clinical history and standardized objective diagnostic test results representative of typical situations in a tertiary care clinic.

Comparison with other studies

A few studies have presented questionnaire-reported symptoms for children or adolescents with exercise-induced symptoms. A Swedish population based study in children aged 12-13

years reported exercise-induced symptoms for 128 children with an asthma diagnosis (13). Exercise-induced wheeze was reported for 76 (59%), cough for 81 (63%), and chest tightness for 56 (44%); throat tightness also was reported for 63 (49%), and inspiratory stridor for 47 (37%). We saw higher prevalence of symptoms overall because our study included respiratory outpatients and not children from the general population.

In a case series study of 12 adolescent athletes seen for suspected exercise-induced laryngeal obstruction (EILO) (15), dyspnea during inspiration was reported by all (100%) and dyspnea during expiration by 8 (67%), and throat tightness was reported more frequently (50%) than chest tightness (25%). A Danish study that compared 42 adolescents with EILO with 16 adolescents diagnosed with airway hyper-responsiveness (AHR) similarly found that all reported wheeze and stridor, but those with EILO mostly reported cough and throat tightness while those with AHR reported mostly dyspnea (14). Our results and those from previous studies emphasize that no symptom is uniquely reported for single diagnostic groups among children with exercise-induced symptoms, but some symptoms are reported more frequently for certain diagnoses than others.

Interpretation

Cough, dyspnea, and tingling sensation in fingers or lips better distinguished thoracic DB from asthma than extrathoracic DB from asthma. This partly explains why extrathoracic DB can be misdiagnosed as asthma (17, 19). Onset of symptoms during exercise was strongly associated with extrathoracic DB, while onset after exercise was associated with asthma. This finding is in line with the literature and could help physicians distinguish extrathoracic DB from asthma (7, 26, 27). We found that of the different triggers, swimming best distinguished diagnoses. Swimming has been reported as a trigger of bronchoconstriction in

children with asthma (28). It was therefore surprising that children with thoracic DB more often reported swimming as a trigger of symptoms than children with asthma. We found no other studies that reported on triggers of exercise-induced symptoms in children with dysfunctional breathing. An explanation could be that children with thoracic DB have difficulties with their breathing patterns and might therefore find swimming especially difficult as correct breathing is a requirement during swimming. We found no evidence that the distinction between inspiration and expiration helped to distinguish diagnoses. Most parents reported that their child's symptoms occur during inspiration (n=298, 47%) or during inspiration and expiration (n=276, 44%) rather than during expiration alone (n=54, 9%). Results from other studies confirm that adolescents with exercise-induced symptoms rarely only report symptoms during expiration alone but usually report symptoms during inspiration or both inspiration and expiration (14, 15). Therefore, although asthma is associated with expiratory airway obstruction, most adolescents report symptoms during inspiration and expiration. While physicians are trained to distinguish inspiratory sounds from expiratory sounds during auscultation, this might be more difficult for parents and children (7, 29). This feature might therefore be useful for clinical examination but not for interpretation of patient-reported symptoms. We also did not see any difference in the duration of symptoms between diagnostic groups. It might be difficult for parents in a stressful moment to judge whether the child's symptoms lasted a few minutes, between 5-10 minutes, or longer unless they observed their child at the time of an attack.

Diagnosing children with exercise-induced symptoms is not easy and requires a thorough diagnostic work up including objective diagnostic tests. Our study confirms that parent-reported symptoms can help to distinguish different diagnoses in children with exercise-

334 induced symptoms. This highlights the importance of physicians taking detailed symptom
335 histories.

336

337

Journal Pre-proof

Ethics approval and consent to participate

The SPAC study was approved by the Bern Cantonal Ethics Committee (Kantonale Ethikkommission Bern 2016-02176). Written informed consent was obtained from patients' parents and directly from patients older than 13 years.

Author's contributions

EP and CK made substantial contributions to the study conception and design. EP, CdJ, and MCM collected and prepared data from the SPAC study. EP drafted the manuscript. EP, CdJ, CA, MCM, JB, CC, KH, AJ, AM, DM, NR, FS, MG, and CK critically revised and approved the manuscript.

Acknowledgements

We would like to thank the families who took part in the SPAC study. We would also like to thank the outpatient clinic assistants, nurses, and doctors for recruiting patients.

Availability of data and material

The SPAC dataset is available on reasonable request by contacting Claudia Kuehni by email: Claudia.kuehni@ispm.unibe.ch.

References

1. Connett GJ, Thomas M. Dysfunctional Breathing in Children and Adults With Asthma. *Frontiers in pediatrics*. 2018;6:406.
2. Johansson H, Norlander K, Berglund L, Janson C, Malinovschi A, Nordvall L, et al. Prevalence of exercise-induced bronchoconstriction and exercise-induced laryngeal obstruction in a general adolescent population. *Thorax*. 2015;70(1):57-63.
3. Fretzayas A, Moustaki M, Loukou I, Douros K. Differentiating vocal cord dysfunction from asthma. *Journal of Asthma and Allergy*. 2017;10:277-83.
4. Barker N, Everard ML. Getting to grips with 'dysfunctional breathing'. *Paediatric respiratory reviews*. 2015;16(1):53-61.
5. Depiazzi J, Everard ML. Dysfunctional breathing and reaching one's physiological limit as causes of exercise-induced dyspnoea. *Breathe (Sheffield, England)*. 2016;12(2):120-9.
6. Kenn K, Hess MM. Vocal Cord Dysfunction: An Important Differential Diagnosis of Bronchial Asthma. *Deutsches Ärzteblatt International*. 2008;105(41):699-704.
7. Roksund OD, Heimdal JH, Clemm H, Vollsaeter M, Halvorsen T. Exercise inducible laryngeal obstruction: diagnostics and management. *Paediatric respiratory reviews*. 2017;21:86-94.
8. Roksund OD, Heimdal JH, Olofsson J, Maat RC, Halvorsen T. Larynx during exercise: the unexplored bottleneck of the airways. *European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery*. 2015;272(9):2101-9.
9. Niggemann B. How to diagnose psychogenic and functional breathing disorders in children and adolescents. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology*. 2010;21(6):895-9.
10. Liyanagedara S, McLeod R, Elhassan HA. Exercise induced laryngeal obstruction: a review of diagnosis and management. *European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery*. 2017;274(4):1781-9.
11. Chiang T, Marcinow AM, deSilva BW, Ence BN, Lindsey SE, Forrest LA. Exercise-induced paradoxical vocal fold motion disorder: diagnosis and management. *The Laryngoscope*. 2013;123(3):727-31.
12. Olin JT. Exercise-Induced Laryngeal Obstruction: When Pediatric Exertional Dyspnea Does not Respond to Bronchodilators. *Frontiers in pediatrics*. 2019;7:52.
13. Johansson H, Norlander K, Hedenstrom H, Janson C, Nordang L, Nordvall L, et al. Exercise-induced dyspnea is a problem among the general adolescent population. *Respiratory medicine*. 2014;108(6):852-8.
14. Christensen PM, Thomsen SF, Rasmussen N, Backer V. Exercise-induced laryngeal obstructions: prevalence and symptoms in the general public. *European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery*. 2011;268(9):1313-9.
15. Shembel AC, Hartnick CJ, Bunting G, Ballif C, Shaiman S, de Guzman V, et al. Perceptual Clinical Features in Exercise-Induced Laryngeal Obstruction (EILO): Toward

- Improved Diagnostic Approaches. *Journal of voice : official journal of the Voice Foundation*. 2018.
16. Nielsen EW, Hull JH, Backer V. High prevalence of exercise-induced laryngeal obstruction in athletes. *Medicine and science in sports and exercise*. 2013;45(11):2030-5.
 17. Abu-Hasan M, Tannous B, Weinberger M. Exercise-induced dyspnea in children and adolescents: if not asthma then what? *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology*. 2005;94(3):366-71.
 18. Khan DA. Exercise-induced bronchoconstriction: burden and prevalence. *Allergy and asthma proceedings*. 2012;33(1):1-6.
 19. Seear M, Wensley D, West N. How accurate is the diagnosis of exercise induced asthma among Vancouver schoolchildren? *Archives of disease in childhood*. 2005;90(9):898-902.
 20. Pedersen ESL, de Jong CCM, Ardura-Garcia C, Barben J, Casaulta C, Frey U, et al. The Swiss Paediatric Airway Cohort (SPAC). *ERJ open research*. 2018;4(4).
 21. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of biomedical informatics*. 2009;42(2):377-81.
 22. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Journal of clinical epidemiology*. 2008;61(4):344-9.
 23. Van den Broeck J, Willie D, Younger N. The World Health Organization child growth standards: expected implications for clinical and epidemiological research. *European journal of pediatrics*. 2009;168(2):247-51.
 24. Braun-Fahrlander C, Gassner M, Grize L, Minder CE, Varonier HS, Vuille JC, et al. Comparison of responses to an asthma symptom questionnaire (ISAAC core questions) completed by adolescents and their parents. SCARPOL-Team. Swiss Study on Childhood Allergy and Respiratory Symptoms with respect to Air Pollution. *Pediatric pulmonology*. 1998;25(3):159-66.
 25. Olson LM, Radecki L, Frintner MP, Weiss KB, Korfmacher J, Siegel RM. At what age can children report dependably on their asthma health status? *Pediatrics*. 2007;119(1):e93-102.
 26. Griffin SA, Walsted ES, Hull JH. Breathless athlete: exercise-induced laryngeal obstruction. *British journal of sports medicine*. 2018;52(18):1211-2.
 27. Hall A, Thomas M, Sandhu G, Hull JH. Exercise-induced laryngeal obstruction: a common and overlooked cause of exertional breathlessness. *The British journal of general practice : the journal of the Royal College of General Practitioners*. 2016;66(650):e683-5.
 28. Romberg K, Tufvesson E, Bjermer L. Asthma symptoms, mannitol reactivity and exercise-induced bronchoconstriction in adolescent swimmers versus tennis players. *Journal of Asthma and Allergy*. 2017;10:249-60.
 29. Liyanagedera S, McLeod R, Elhassan HA. Exercise induced laryngeal obstruction: a review of diagnosis and management. *European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery*. 2017;274(4):1781-9.

449

450

451

452

Journal Pre-proof

Figure legends

Figure 1: Type of exercise-induced symptoms in children referred to paediatric respiratory outpatient clinics, displayed by diagnosis group (n=732)

Figure 2 Comparison of type of exercise-induced symptoms between diagnosis groups using multinomial regression models adjusted for age and sex

Figure 3 Comparison of activities triggering exercise-induced symptoms between diagnosis groups using multinomial regression models adjusted for age and sex

Figure 4 Comparison of exercise-induced symptom characteristics (localization of dyspnea, respiratory phase, onset, and duration of symptoms) between diagnosis groups using multinomial regression models adjusted for age and sex

Table 1 Comparison of characteristics, respiratory symptoms and diagnoses between included patients with and without exercise-induced symptoms (EIS) (N=1109)

	Yes to EIS in questionnaire N=732 n(%)	No to EIS in questionnaire N=377 n(%)
Characteristics		
Demographic and socioeconomic characteristics		
Age (years), mean (SD)	11.0 (3.2)	9.5 (3.1)
Sex (female)	318 (43)	132 (35)
BMI z-score, mean (SD) (n=1091)	0.3 (1.2)	0.2 (1.2)
Overweight (BMI z-score >1) (n=1094)	177 (26)	76 (23)
Sports apart from at school (n=1056)	566 (78)	257 (70)
Swiss nationality	609 (83)	309 (82)
Parental education		
Mother, tertiary ^a (n=1071)	259 (37)	114 (31)
Father, tertiary ^a (n=1056)	315 (45)	142 (39)
Parental smoking		
Mother, current smoking (n=1090)	114 (16)	49 (14)
Father, current smoking (n=1046)	174 (25)	80 (23)
Respiratory symptoms in the past 12 months		
Cough apart from colds, yes often (n=1096)	88 (12)	55 (15)
Cough at night apart from colds (n=1079)	329 (46)	151 (41)
Wheeze (n=1086)	452 (63)	168 (46)
>3 attacks of wheeze (n=1086)	216 (30)	48 (13)
Rhinitis apart from colds (n=1100)	479 (66)	213 (57)
Eczema ever (n=1090)	215 (30)	102 (28)
Diagnosis given at outpatient clinic		
Asthma	549 (75)	276 (73)
Extrathoracic dysfunctional breathing	38 (5)	1 (0)
Thoracic dysfunctional breathing	30 (4)	7 (2)
Asthma + any DB	43 (6)	1 (0)
Chronic cough	21 (3)	35 (9)
Other	51 (7)	57 (15)

^a Degree from university of applied sciences or university. Abbreviations: EIS, exercise induced symptoms

Table 2 Reported exercise-induced symptoms (EIS) by diagnosis group in children who reported exercise-induced respiratory symptoms in the questionnaire (n=732)

Baseline questionnaire	Asthma N=549 N (%)	DB extra- thoracic N=38 N (%)	DB thoracic N=30 N (%)	Asthma + any DB N=43 N (%)	Cough N=21 N (%)	Other N=51 N (%)	P-value
Type of EIS (n=732) ^a							
Wheeze	329 (60)	20 (53)	15 (52)	27 (63)	9 (43)	9 (18)	<0.001
Cough	390 (71)	21 (55)	13 (43)	17 (40)	18 (86)	32 (63)	<0.001
Dyspnea	376 (68)	32 (84)	28 (93)	40 (93)	10 (48)	29 (57)	<0.001
Tingling feelings in finger or lips	53 (11)	6 (16)	8 (30)	10 (24)	0 (0)	10 (21)	0.002
Other symptoms	50 (9)	5 (13)	7 (23)	7 (16)	4 (19)	8 (16)	0.016
Trigger activities (n=697)							
Run short (50-100 m)	327 (63)	27 (71)	19 (66)	29 (69)	14 (70)	34 (71)	0.730
Run far (>1 km)	402 (77)	32 (84)	22 (76)	35 (83)	13 (65)	35 (73)	0.522
Cycle	254 (50)	27 (75)	19 (68)	23 (59)	9 (50)	24 (53)	0.034
Intensive sport games [#]	396 (75)	27 (71)	23 (77)	37 (86)	13 (65)	34 (71)	0.453
Swim	162 (31)	13 (34)	17 (59)	20 (48)	8 (40)	24 (50)	0.002
Localization of dyspnea (n=496 of 515 with dyspnea) ^b							
Chest	189 (52)	15 (47)	13 (46)	19 (50)	3 (30)	19 (70)	0.187
Throat	47 (13)	9 (28)	6 (21)	6 (16)	3 (30)	4 (15)	
Chest and Throat	125 (35)	8 (25)	9 (32)	13 (34)	4 (40)	4 (15)	
Respiration phase ^c (n=628)							
Inspiration	214 (46)	19 (51)	12 (44)	27 (64)	9 (60)	17 (44)	0.271
Expiration	45 (10)	0	2 (7)	2 (5)	0	5 (13)	
Inspiration and Expiration	209 (44)	18 (49)	13 (48)	13 (31)	6 (40)	17 (44)	
EIS start ^d (n=677)							
During exercise	344 (69)	36 (95)	24 (86)	33 (81)	9 (43)	29 (60)	<0.001
After ending exercise	156 (31)	2 (5)	4 (14)	8 (19)	12 (57)	19 (40)	
Duration of EIS ^e (n=648)							
1-2 minutes	189 (37)	13 (34)	5 (20)	14 (34)	8 (38)	21 (45)	0.503
5-10 minutes	268 (53)	22 (58)	14 (56)	23 (56)	10 (48)	22 (47)	
Longer than 10 min	48 (10)	3 (8)	6 (24)	4 (10)	3 (14)	4 (9)	
Used asthma-spray before or during exercise? ^g (n=712)	41 (77)	15 (39)	14 (47)	37 (88)	13 (62)	21 (43)	<0.001
Effect of asthma-spray ^h (n=494 of 511 who used asthma spray)							
EIS disappear	172 (43)	2 (14)	2 (14)	9 (25)	1 (8)	3 (17)	*
EIS are reduced	204 (51)	8 (57)	9 (64)	18 (50)	8 (62)	11 (61)	
No effect	23 (6)	4 (29)	3 (21)	9 (25)	4 (31)	4 (22)	

This table displays n(%) with column percentages.

EIS: exercise-induced symptoms

^aWhich symptoms does your child have during exercise?

^bIf reported dyspnea: Where is the sensation of symptoms felt the strongest?

^cWhen are the symptoms worst?

^dWhen do the symptoms begin?

^eAfter ending the exercise, how long do the symptoms usually stay?

^fDoes your child sometimes get a tingling sensation in fingertips or around the mouth during the EIS?

^gHas your child ever used an asthma inhaler during exercise?

484 ^hHow well does this asthma inhaler help?
485 *Too few observations in single cells to calculate Fisher's exact
486

487

488

Journal Pre-proof

Figure 2 Comparison of type of exercise-induced symptoms between diagnosis groups using multinomial regression models adjusted for age and sex

The graphs represent relative risk ratios from multinomial regression analysis with diagnosis categories as outcome (asthma as base variable) and types of symptom (wheeze, cough, dyspnea, tingling sensation in fingertips/lips, other symptoms) adjusted for age and sex. RRR, relative risk ratio; 95%CI, 95% confidence interval; DB, dysfunctional breathing

Figure 3 Comparison of activities triggering exercise-induced between diagnosis groups using multinomial regression models adjusted for age and sex

The graphs represent relative risk ratios from multinomial regression analysis with diagnosis categories as outcome (asthma as base variable) and trigger activities (run, cycle, intensive sport games, swim) adjusted for age and sex. RRR, relative risk ratio; 95%CI, 95% confidence interval; m, meter; km, kilometer; DB, dysfunctional breathing

Figure 4: Comparison of exercise-induced symptom characteristics (localization of dyspnea, respiratory phase, onset, and duration of symptoms) between diagnosis groups using multinomial regression models adjusted for age and sex

The graphs represent relative risk ratios from multinomial regression analysis with diagnosis categories as outcome (asthma as base variable) and characterizations of exercise-induced symptoms (localization of dyspnea, respiration phase, and duration) as explanatory variables. RRR, relative risk ratio; 95%CI, 95% confidence interval; DB, dysfunctional breathing

*Too few observations to perform multinomial regression analysis