

RESEARCH ARTICLE

A polymorphism in the cachexia-associated gene *INHBA* predicts efficacy of regorafenib in patients with refractory metastatic colorectal cancer

Yuji Miyamoto¹, Marta Schirripa¹, Mitsukuni Suenaga¹, Shu Cao², Wu Zhang¹, Satoshi Okazaki¹, Martin D. Berger¹, Satoshi Matsusaka¹, Dongyun Yang², Yan Ning¹, Hideo Baba³, Fotios Loupakis⁴, Sara Lonardi⁴, Filippo Pietrantonio⁵, Beatrice Borelli⁶, Chiara Cremolini⁶, Toshiharu Yamaguchi⁷, Heinz-Josef Lenz^{1*}

1 Division of Medical Oncology, Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, United States of America, **2** Department of Preventive Medicine, Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, United States of America, **3** Department of Gastroenterological Surgery, Graduate School of Medical Sciences, Kumamoto University, Kumamoto, Japan, **4** Unit of Medical Oncology 1, Department of Clinical and Experimental Oncology, Istituto Oncologico Veneto, IRCCS, Padua, Italy, **5** Medical Oncology Department, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy, **6** Polo Oncologico, Azienda Ospedaliero-Universitaria Pisana, Pisa, Italy, **7** Department of Gastroenterological Surgery, Cancer Institute Hospital, Japanese Foundation for Cancer Research, Tokyo, Japan

* lenz@usc.edu



OPEN ACCESS

Citation: Miyamoto Y, Schirripa M, Suenaga M, Cao S, Zhang W, Okazaki S, et al. (2020) A polymorphism in the cachexia-associated gene *INHBA* predicts efficacy of regorafenib in patients with refractory metastatic colorectal cancer. PLoS ONE 15(9): e0239439. <https://doi.org/10.1371/journal.pone.0239439>

Editor: Jason Chia-Hsun Hsieh, Chang Gung Memorial Hospital at Linkou, TAIWAN

Received: March 2, 2020

Accepted: September 5, 2020

Published: September 24, 2020

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0239439>

Copyright: © 2020 Miyamoto et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Abstract

Activin/myostatin signaling has a critical role not only in cachexia but also in tumor angiogenesis. Cachexia is a frequent complication among patients with advanced cancer and heavily pretreated patients. We aimed to evaluate the prognostic significance of cachexia-associated genetic variants in refractory metastatic colorectal cancer (mCRC) patients treated with regorafenib. Associations between twelve single nucleotide polymorphisms in 8 genes (*INHBA*, *MSTN*, *ALK4*, *TGFBR1*, *ALK7*, *ACVR2B*, *SMAD2*, *FOXO3*) and clinical outcome were evaluated in mCRC patients of three cohorts: a discovery cohort of 150 patients receiving regorafenib, a validation cohort of 80 patients receiving regorafenib and a control cohort of 128 receiving TAS-102. In the discovery cohort, patients with any G variant in *FOXO3* rs12212067 had a significantly lower response rate ($P = 0.031$) and overall survival (OS) than those with a T/T in univariate analysis (4.5 vs. 7.6 months, hazard ratio [HR] = 1.63, 95% confidence interval [CI] = 1.09–2.46, $P = 0.012$). Among female patients, those with any G variant in *INHBA* rs2237432 had a significantly longer OS than those with an A/A in both univariate (7.6 vs. 4.3 months, HR = 0.57, 95%CI = 0.34–0.95, $P = 0.021$) and multivariable (HR = 0.53, 95%CI = 0.29–0.94, adjusted $P = 0.031$) analysis. This association was confirmed in female patients of the validation cohort, though without statistical significance ($P = 0.059$). Conversely, female patients with any G allele in the control group receiving TAS-102 did not show a longer OS. This was the first study evaluating the associations between polymorphisms in cachexia-associated genes and outcomes in refractory mCRC

Funding: This work was supported by the National Institute of Health (P30CA014089); the Gloria Borges Wunderglo Project; the Dhont Family Foundation; the Daniel Butler Research Fund; and the Call to Cure Research Fund. YM received a grant from Japan Society for the Promotion of Science (S2606). MS is the recipient of Takashi Tsuruo Memorial Fund. MDB received a grant from the Werner and Hedy Berger-Janser Foundation for cancer research and the Swiss Cancer League (BIL KLS-3334-02-2014). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have read the journal's policy and the authors have the following competing interests: HJ Lenz has received honoraria from Merck Serono, Roche, Celgene, Bayer, and Boehringer Ingelheim. HB received honoraria from Chugai Pharma, Bayer, Taiho Pharmaceutical and Merck Serono. There are no patents, products in development or marketed products to declare. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

patients treated with regorafenib. Further studies should be conducted to confirm these associations.

Introduction

Regorafenib is a small molecule multikinase inhibitor that blocks protein kinases involved in tumor angiogenesis, oncogenesis and the tumor microenvironment [1]. The benefit of regorafenib on overall survival (OS) in patients with metastatic colorectal cancer (mCRC) was demonstrated in two phase III randomized controlled trials, the CORRECT [2] and CONCUR [3] trials. Therefore, regorafenib is now established as an additional line of therapy for patients with mCRC refractory to previous chemotherapy as well as for best supportive care [4, 5]. Several investigators have attempted to identify molecular markers that predict the activity of regorafenib for the individualized treatment of patients with mCRC. For example, expression levels of biomarkers such as *VEGF* and *CCL5* [6] or plasma circulating cell-free DNA [7] may represent potential predictive biomarkers of regorafenib treatment, although these results have not been sufficiently validated.

Cancer cachexia is defined as an ongoing loss of skeletal muscle mass and is a more common complication in heavily pretreated cancer patients [8], leading to progressive impairment of physical function and quality of life as well as resistance to chemotherapy or radiotherapy [9, 10]. Skeletal muscle mass is dynamically regulated by various extracellular signals, which activate distinct intracellular signaling processes [11]. In particular, *INHBA* and *MSTN* are potent negative regulators of muscle mass [12]. The binding of *INHBA* and *MSTN* to membrane receptors (*ACVR1B*, *C*, and *ACVR2B*) leads to the activation of *SMAD*-mediated signal transduction, promoting muscle protein degradation [13]. *INHBA* or *MSTN* expression is associated with several types of human cancers, and CRC patients with high *INHBA* expression showed poorer OS than those with low *INHBA* expression [14]. In addition, accumulating evidence suggests that activin/myostatin signaling, like other members of the *TGF-beta* superfamily, can regulate angiogenesis. *MSTN* blockade reduced the tumor expression of genes involved in angiogenesis (e.g. *VEGF-A*, *HIF-1α*) [15]. Similarly, *INHBA* demonstrated both pro- [16] and anti-angiogenic [17] properties in different systems. Recently, we reported that germline variants within the cancer cachexia pathway are associated with outcome in mCRC patients treated with bevacizumab-based chemotherapy [18].

Based on the clinical importance of cachexia signaling being potentially involved in angiogenesis, we evaluated the prognostic and predictive significance of cachexia-associated genetic variants in refractory mCRC patients treated with regorafenib chemotherapy. A previous report indicated that gender differences may influence skeletal muscle changes after chemotherapy [19]. We therefore determined whether such associations were influenced by gender.

Materials and methods

Study design and patients

This study was a retrospective exploratory study in three independent cohorts of patients with refractory mCRC: a discovery cohort of 150 patients receiving regorafenib at Azienda Ospedaliero-Universitaria Pisana, Istituto Oncologico Veneto (Padova, Italy); a validation cohort of 80 patients receiving regorafenib at the Cancer Institute Hospital of the Japanese Foundation for Cancer Research (Japan); and a control cohort of 128 patients receiving TAS-102 at Azienda Ospedaliero-Universitaria Pisana, Istituto Oncologico Veneto (Padova, Italy) and

Istituto Nazionale Tumori (Milano, Italy). Patients with histologically verified colorectal adenocarcinoma, measurable metastatic disease according to Response Evaluation Criteria in Solid Tumors (RECIST) v1.1, and a history of previous standard chemotherapy with 5-FU, L-OHP, CPT-11, bevacizumab, and cetuximab or panitumumab were eligible. Patients received regorafenib 160 mg per body once daily from days 1–21, every 4 weeks, or TAS-102 35 mg per m² twice daily on days 1–5 and 8–12, every 4 weeks. Treatment was administered until disease progression, intolerable toxicities, or patient withdrawal occurred. All patients provided written informed consent, including consent for all medical record which were fully anonymized before we assessed, blood or tumor tissue to be used to explore relevant molecular parameters. This study was conducted according to the REporting recommendations for tumor MARKer prognostic studies (REMARK) [20]. The tissue analysis protocol was approved by the University of Southern California (USC) Institutional Review Board of Medical Sciences and conducted at the USC/Norris Comprehensive Cancer Center in accordance with the Declaration of Helsinki and Good Clinical Practice guidelines.

Selection of candidate single-nucleotide polymorphisms

The 12 candidate single nucleotide polymorphisms (SNPs) in the cachexia pathway examined in this study were *INHBA*, *MSTN*, *ALK4*, *TGFBRI*, *ALK7*, *ACVR2B*, *SMAD2*, and *FOXO3*, which were selected using one of the following criteria: i) SNP with potential biological significance based on the published literature or F-SNP database* (<http://compbio.cs.queensu.ca/F-SNP/>); or ii) minor allele frequency $\geq 10\%$ in both white and East Asians in the Ensembl Genome Browser. The characteristics of the selected polymorphisms are shown in [S1 Table](#).

DNA extraction and genotyping

Genomic DNA was extracted from patients' peripheral blood using a QIAmp Kit (Qiagen, Valencia, CA, USA) according to the manufacturer's protocol. The candidate SNPs were examined by PCR-based direct DNA sequence analysis using an ABI 3100A Capillary Genetic Analyzer and Sequencing Scanner v1.0 (Applied Biosystems, Foster City, CA, USA). The primers for amplification of extracted DNA are listed in [S1 Table](#). For quality control purposes, 10% of samples were randomly selected and analyzed by direct DNA sequencing for each SNP. The genotype concordance rate was found to be $\geq 99\%$. The investigators analyzing SNPs were blinded to the clinical data.

Statistical analysis

The primary endpoint in this study was progression-free survival (PFS), and the secondary endpoints were OS and disease control rate (DCR). PFS was defined as the period between the date of starting treatment and the date of confirmed disease progression or death. OS was calculated from the date of starting treatment until the date of death from any cause. If the event was not observed by the last follow up date, the patient was recorded as censored. In patients lost to follow-up, data were censored at the date of last follow up. According to RECIST v1.1, DCR was defined as the proportion of patients who achieved stable disease (SD), partial response (PR), or progressive disease (PD). Chi-square tests were used to examine the difference in baseline patient characteristics between the three cohorts. Allelic distribution of polymorphisms was tested for deviation from the Hardy–Weinberg equilibrium using the exact test. Linkage disequilibrium among SNPs was evaluated using D' and r^2 values and haplotype frequencies of genes were inferred using HaploView version 4.2 (<http://www.broad.mit.edu/mpg/haploview>). High linkage disequilibrium was defined as $r^2 > 0.7$. Fisher's exact test was applied to examine the associations between SNPs and DCR. Associations between candidate

SNPs and PFS or OS were analyzed by the Kaplan–Meier method and log-rank test in the univariable analysis and reevaluated using a Cox proportional hazards model and Wald test with predictive or prognostic baseline factors included. The baseline demographic and clinical characteristics statistically significantly associated with PFS and OS in multivariable analyses were included in the final models. We used codominant, dominant, or recessive genetic models where appropriate for the candidate SNPs, because the true modes were not yet established in the analyses. The minimum detectable hazard ratios of 1.61–1.82 corresponded to the minor allele frequency of 0.1–0.4 in the association between an SNP and PFS in the discovery cohort ($n = 150$, PFS events = 149), considering a dominant model and using a two-sided 0.05-level log-rank test with 80% power. In the validation cohort ($n = 80$, PFS events = 79), the power was 54% using the same model. All analyses were carried out with SAS 9.4 (SAS Institute, Cary, NC, USA). All tests were two-sided at a significance level of 0.050. *P*-act method, a modified multiple testing method, was applied for adjusting the *P* values for all SNPs when the linkage disequilibrium between candidate SNPs and different modes of inheritance was considered.

Results

Baseline characteristics

The baseline characteristics of enrolled cohorts are summarized in [Table 1](#). Gender, performance status, adjuvant treatment history, and the number of prior chemotherapy regimens were distributed differently between the cohorts. The median PFS, OS, and follow-up time were 2.1, 6.0, and 36.4 months in the discovery cohort; 2.0, 8.0, and 15.3 months in the validation cohort; and 2.0, 5.4, and 5.3 months in the control cohort. Genotyping was successful in at least 90% of cases for each polymorphism analyzed. The allelic frequencies for all SNPs were within the probability limits of the Hardy–Weinberg equilibrium ($P > 0.050$). High linkage disequilibrium was observed between *ACVR2B* rs13072731 and *ACVR2B* rs2268753 in the discovery cohort, with $D' = 0.98$ and $r^2 = 0.70$. No other high linkage disequilibrium was observed between the SNPs found in each cohort.

Associations between cachexia SNPs and outcome in the discovery and validation cohorts

Associations between candidate SNPs and clinical outcome were examined in the regorafenib discovery cohort. Patients with any G allele in *FOXO3* rs12212067 had significantly shorter PFS and OS and worse DCR than those with a T/T variant in univariate analysis (PFS: 1.8 vs. 2.1 months, hazard ratio [HR] 1.44, 95% confidence interval [CI] 0.98–2.12, $P = 0.056$; OS: 4.5 vs. 7.6 months, HR 1.63, 95% CI 1.09–2.46, $P = 0.012$; DCR: $P = 0.031$) ([Table 2](#)). However, in the validation cohort, patients with any G allele in *FOXO3* rs12212067 had longer PFS and OS than those with a T/T variant in univariate analysis (PFS: 2.0 vs. 2.5 months, HR 0.56, 95% CI 0.32–0.97, $P = 0.027$; OS: 7.6 vs. 15.3 months, HR 0.49, 95% CI 0.23–1.04, $P = 0.054$) ([S2 Table](#)). However, these effects were not significant in the multivariable model and after multiple testing.

Associations between cachexia SNPs and outcome stratified by sex in the discovery and validation cohorts

Among female patients in the discovery cohort, patients with any G allele in *INHBA* rs2237432 showed a significantly longer OS than those with the A/A allele in both univariate (7.6 vs. 4.3 months, HR 0.57, 95% CI 0.34–0.95, $P = 0.021$) and multivariable analysis (HR 0.53, 95% CI

Table 1. Baseline clinical characteristics of patients in the discovery, validation, and control cohorts.

Characteristics	Discovery cohort (n = 150)		Validation cohort (n = 80)		Control cohort (n = 128)		P value ^a
	N	(%)	N	(%)	N	(%)	
Age (years)							0.97
≤65	94	(63)	49	(61)	79	(62)	
>65	56	(37)	31	(39)	49	(38)	
Gender							0.49
Male	81	(54)	38	(48)	61	(48)	
Female	69	(46)	42	(53)	67	(52)	
Primary tumor site							0.80
Right	49	(33)	23	(29)	39	(30)	
Left	99	(66)	57	(71)	85	(66)	
Unknown ^b					4	(3)	
Primary tumor resected							0.52
Yes	127	(85)	70	(88)	109	(85)	
No	23	(15)	10	(12)	13	(10)	
Unknown ^b					6	(5)	
Adjuvant treatment							0.087
Yes	37	(25)	27	(34)	47	(37)	
No	112	(75)	53	(66)	81	(63)	
Unknown ^b	1	(0)					
Liver metastasis							0.11
Yes	120	(80)	54	(68)	96	(75)	
No	30	(20)	26	(33)	32	(25)	
Lung metastasis							0.096
Yes	109	(73)	47	(59)	88	(69)	
No	41	(27)	33	(41)	40	(31)	
LN metastasis							0.47
Yes	75	(50)	41	(51)	56	(44)	
No	75	(50)	39	(49)	72	(56)	
Peritoneal involvement							0.67
Yes	43	(29)	20	(25)	31	(24)	
No	107	(71)	60	(75)	97	(76)	
Number of metastases							0.001
1	16	(11)	24	(30)	21	(16)	
> 1	134	(89)	56	(70)	107	(84)	
Number of treatment regimens							< 0.001
≤ 3	108	(72)	72	(90)	80	(63)	
> 3	42	(28)	8	(10)	48	(38)	
Performance status							< 0.001
ECOG 0	117	(78)	45	(56)	72	(56)	
ECOG 1	33	(22)	35	(44)	56	(44)	
RAS status							0.41
Wild	52	(35)	-		47	(37)	
Mutant	93	(62)	-		68	(53)	
Unknown ^b	5	(3)	-		13	(10)	

Abbreviations: LN, lymph node; ECOG, Eastern Cooperative Oncology Group.

^a Based on the χ^2 test.

^b Not included in the test.

<https://doi.org/10.1371/journal.pone.0239439.t001>

Table 2. Association between cachexia-related gene polymorphism and clinical outcomes in the discovery cohort.

Genotype	N	Tumor response			Progression-free survival					Overall survival								
		PR+SD	PD	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*				
INHBA rs2237432				0.91					0.92					0.59				0.87
A/A	78	27 (37%)	46 (63%)		2.1 (1.8, 2.8)	1 (Reference)		1 (Reference)		6.5 (4.5, 8.7)	1 (Reference)		1 (Reference)					
A/G	56	18 (33%)	37 (67%)		2.1 (1.8, 2.3)	0.94 (0.66, 1.34)		0.99 (0.68, 1.43)		5.6 (4.3, 7.6)	0.91 (0.63, 1.31)		1.00 (0.68, 1.48)					
G/G	16	5 (33%)	10 (67%)		2.1 (1.7, 4.0)	1.02 (0.59, 1.74)		1.44 (0.82, 2.54)		9.5 (2.2, 13.9)	0.75 (0.42, 1.34)		1.17 (0.64, 2.12)					
INHBA rs17776182				1.00					0.78					0.21				0.47
G/G	122	41 (35%)	76 (65%)		2.1 (1.9, 2.3)	1 (Reference)		1 (Reference)		5.9 (4.5, 7.8)	1 (Reference)		1 (Reference)					
G/A ^a	25	8 (35%)	15 (65%)		1.9 (1.8, 3.6)	0.94 (0.62, 1.43)		1.09 (0.71, 1.69)		8.7 (5.5, 12.4)	0.77 (0.50, 1.18)		0.85 (0.54, 1.33)					
A/A ^a	3	1 (33%)	2 (67%)															
MSTN rs7570532				0.46					0.67					0.96				0.98
A/A	89	33 (39%)	51 (61%)		2.3 (1.9, 3.0)	1 (Reference)		1 (Reference)		7.8 (5.7, 9.4)	1 (Reference)		1 (Reference)					
A/G ^a	54	15 (29%)	37 (71%)		1.9 (1.8, 2.1)	1.07 (0.77, 1.49)		1.06 (0.76, 1.48)		5.3 (3.6, 7.6)	1.01 (0.71, 1.42)		1.00 (0.70, 1.41)					
G/G ^a	7	2 (29%)	5 (71%)															
ALK4 rs2854464				0.90					0.95					0.48				0.17
A/A	78	25 (34%)	49 (66%)		2.1 (1.8, 2.8)	1 (Reference)		1 (Reference)		6.0 (5.0, 8.9)	1 (Reference)		1 (Reference)					
A/G	56	18 (34%)	35 (66%)		2.1 (1.8, 2.3)	1.05 (0.74, 1.49)		1.13 (0.79, 1.61)		6.3 (4.4, 8.0)	1.09 (0.76, 1.57)		1.14 (0.78, 1.66)					
G/G	15	6 (40%)	9 (60%)		2.2 (1.4, 3.8)	1.06 (0.61, 1.85)		1.14 (0.65, 2.02)		5.7 (2.0, 10.0)	1.41 (0.80, 2.47)		1.74 (0.97, 3.10)					
TGFBR1 rs10760673				0.57					0.60					0.72				0.28
G/G	92	32 (37%)	54 (63%)		2.0 (1.8, 2.8)	1 (Reference)		1 (Reference)		7.7 (5.4, 9.1)	1 (Reference)		1 (Reference)					
G/A ^a	48	17 (35%)	31 (65%)		2.1 (1.9, 2.4)	0.92 (0.65, 1.29)		0.70 (0.49, 1.00)		5.6 (4.4, 7.8)	1.06 (0.75, 1.51)		0.82 (0.57, 1.18)					
A/A ^a	8	1 (14%)	6 (86%)															
ALK7 rs13010956				0.94					0.62					0.95				0.68
T/T	45	15 (34%)	29 (66%)		1.9 (1.8, 2.3)	1 (Reference)		1 (Reference)		5.4 (3.5, 7.9)	1 (Reference)		1 (Reference)					
T/C	80	26 (34%)	51 (66%)		2.1 (1.9, 2.3)	0.84 (0.58, 1.22)		0.83 (0.57, 1.20)		6.5 (5.5, 8.9)	0.99 (0.67, 1.46)		0.93 (0.62, 1.38)					
C/C	24	8 (38%)	13 (62%)		1.9 (1.8, 4.1)	0.84 (0.51, 1.39)		1.02 (0.60, 1.71)		7.4 (3.5, 9.7)	1.07 (0.64, 1.79)		1.16 (0.68, 1.97)					
ACVR2B rs13072731				0.30					0.89					0.49				0.33

(Continued)

Table 2. (Continued)

Genotype	N	Tumor response			Progression-free survival					Overall survival				
		PR+SD	PD	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*
C/C	52	20 (40%)	30 (60%)		2.1 (1.8, 3.1)	1 (Reference)		1 (Reference)		6.0 (4.4, 10.5)	1 (Reference)		1 (Reference)	
C/A	73	25 (37%)	43 (63%)		2.2 (1.9, 2.8)	1.03 (0.72, 1.47)		1.12 (0.78, 1.62)		6.4 (4.7, 8.0)	1.24 (0.85, 1.81)		1.35 (0.91, 1.99)	
A/A	23	5 (22%)	18 (78%)		1.8 (1.8, 2.1)	0.92 (0.55, 1.55)		0.89 (0.52, 1.53)		7.6 (2.7, 9.7)	1.05 (0.63, 1.76)		1.15 (0.68, 1.93)	
ACVR2B rs2268753				0.56			0.80		0.70			0.85		0.67
T/T	38	15 (41%)	22 (59%)		2.1 (1.8, 3.1)	1 (Reference)		1 (Reference)		6.2 (4.4, 10.1)	1 (Reference)		1 (Reference)	
T/C	81	27 (35%)	50 (65%)		2.2 (1.9, 2.7)	0.88 (0.60, 1.30)		0.96 (0.65, 1.43)		6.0 (4.7, 8.0)	1.12 (0.74, 1.69)		1.21 (0.80, 1.84)	
C/C	31	8 (28%)	21 (72%)		2.0 (1.8, 2.3)	0.89 (0.54, 1.46)		0.81 (0.49, 1.35)		7.6 (3.3, 9.1)	1.13 (0.69, 1.85)		1.15 (0.70, 1.89)	
SMAD2 rs1792671				0.52			0.18		0.44			0.15		0.58
G/G	55	18 (34%)	35 (66%)		2.1 (1.8, 2.5)	1 (Reference)		1 (Reference)		7.6 (5.1, 9.4)	1 (Reference)		1 (Reference)	
G/A	64	24 (40%)	36 (60%)		2.3 (1.8, 3.7)	0.78 (0.54, 1.13)		0.80 (0.54, 1.16)		7.0 (4.5, 9.6)	0.88 (0.60, 1.28)		0.93 (0.63, 1.37)	
A/A	30	8 (28%)	21 (72%)		1.9 (1.8, 2.2)	1.12 (0.71, 1.76)		0.99 (0.62, 1.59)		4.9 (3.1, 8.0)	1.36 (0.85, 2.16)		1.21 (0.75, 1.97)	
SMAD2 rs1792689				0.49			0.49		0.80			0.58		0.50
C/C	109	35 (34%)	69 (66%)		2.0 (1.8, 2.3)	1 (Reference)		1 (Reference)		6.5 (5.3, 8.7)	1 (Reference)		1 (Reference)	
C/T ^a	38	13 (36%)	23 (64%)		2.2 (1.8, 3.4)	0.88 (0.62, 1.27)		0.95 (0.66, 1.38)		5.9 (3.7, 8.9)	1.11 (0.76, 1.61)		1.14 (0.78, 1.68)	
T/T ^a	3	2 (67%)	1 (33%)											
FOXO3 rs12212067				0.031			0.056		0.19			0.012		0.094
T/T	114	42 (38%)	68 (62%)		2.1 (1.9, 2.5)	1 (Reference)		1 (Reference)		7.6 (5.9, 9.0)	1 (Reference)		1 (Reference)	
T/G ^a	34	6 (19%)	25 (81%)		1.8 (1.8, 2.3)	1.44 (0.98, 2.12)		1.31 (0.87, 1.96)		4.5 (2.7, 5.5)	1.63 (1.09, 2.46)		1.43 (0.94, 2.17)	
G/G ^a	1	1 (100%)	0 (0%)											
FOXO3 rs4946935				0.051			0.29		0.26			0.30		0.36
G/G	87	31 (38%)	51 (62%)		2.1 (1.9, 2.4)	1 (Reference)		1 (Reference)		6.5 (5.5, 9.1)	1 (Reference)		1 (Reference)	
G/A ^a	55	15 (28%)	39 (72%)		2.1 (1.8, 2.7)	1.19 (0.85, 1.66)		1.21 (0.87, 1.69)		5.4 (3.6, 8.7)	1.20 (0.85, 1.69)		1.18 (0.83, 1.67)	

(Continued)

Table 2. (Continued)

Genotype	N	Tumor response			Progression-free survival					Overall survival				
		PR+SD	PD	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*
A/A ^a	6	4 (80%)	1 (20%)											

Abbreviations: PR, partial response; SD, stable disease; PD, progressive disease; HR, hazard ratio; CI, confidence interval.

* P value based on Fisher's exact test for tumor response; log-rank test for progression-free survival (PFS) and overall survival (OS) in the univariate analysis (†); and Wald test for PFS and OS in the multivariable Cox regression model adjusted for time to start of regorafenib treatment (<18 vs. ≥18 months), ECOG performance status (0 vs. 1 or 2), primary tumor resection (yes vs. no), and Kohne score (low-intermediate vs. high) (‡). P values < 0.050 are shown in bold text.

^a In the dominant model. + Estimates not yet reached.

<https://doi.org/10.1371/journal.pone.0239439.t002>

0.29–0.94, adjusted $P = 0.031$) (Table 3 and Fig 1A); in addition, *SMAD2* rs1792671 showed significant association with PFS in both univariate and multivariable analyses ($P = 0.025$, adjusted $P = 0.047$) (Table 3). Similarly, female patients in the validation cohort with any G allele in *INHBA* rs2237432 showed longer OS which was marginally significant in multivariable analysis (adjusted $P = 0.059$) (Table 4 and Fig 1B). After *P*-act multiple testing, the effects were not significant.

In the discovery cohort, male patients with any G allele in *FOXO3* rs12212067 had a significantly shorter PFS and OS than those with T/T allele in both univariate and multivariable model (PFS: $P = 0.025$, adjusted $P = 0.009$; OS: $P = 0.015$, adjusted $P = 0.006$) (Table 3 and Fig 2A). After *P*-act multiple testing, the effects remained significant for both PFS and OS (P -act = 0.035 and 0.024, respectively). In the validation cohort, male patients carrying a *FOXO3* rs12212067 T/G allele had a significant longer OS ($P = 0.040$, adjusted $P = 0.069$) (Table 4 and Fig 2B).

Associations between cachexia SNPs and outcome in the control cohort

Within the control cohort, no significant associations were observed between the cachexia SNPs and outcome in female patients (Table 5).

Discussion

Our findings present the first evidence that germline variations in the cancer cachexia pathway are associated with outcome in chemorefractory mCRC patients treated with regorafenib. Furthermore, these associations may depend on gender. We analyzed data from 230 patients receiving regorafenib treatment in two cohorts. Among female patients in the Italian regorafenib discovery cohort, those with any G allele in *INHBA* rs2237432 had significantly better OS than those with an A/A variant. A similar association was confirmed in the Japanese regorafenib validation cohort.

Activin A (*INHBA*), a member of the *TGF-beta* superfamily, is a homodimer formed from two inhibin betaA chains [21] which is produced by several cell types and is involved in several physiologic functions, including embryogenesis, cell growth, differentiation, immune response, and angiogenesis [22]. Activins act via heteromeric complexes of two related transmembrane type I (*ACVR1B*, C) and type II (*ACVR2B*) serine/threonine kinase receptors to activate the downstream *SMAD* signaling pathway [13]. Circulating activin A level is associated with cachexia syndrome, and increased concentrations in cancer cachectic patients may contribute to the development of this condition [12]. In addition, a model of activin A overexpression in

Table 3. Association between cachexia-related gene polymorphism and clinical outcome by gender subgroup in the discovery cohort.

Genotype	N	Tumor response			Progression-free survival					Overall survival				
		PR+SD	PD	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*
Female														
INHBA rs2237432				0.79				0.77		0.59			0.021	0.031
A/A	37	9 (27%)	24 (73%)		1.8 (1.7, 2.3)	1 (Reference)	1 (Reference)		4.3 (2.9, 6.5)	1 (Reference)		1 (Reference)		
A/G ^a	24	7 (29%)	17 (71%)		2.0 (1.8, 2.3)	0.93 (0.57, 1.51)	1.16 (0.67, 2.02)		7.6 (5.4, 10.2)	0.57 (0.34, 0.95)		0.53 (0.29, 0.94)		
G/G ^a	8	1 (13%)	7 (88%)											
MSTN rs7570532				0.82				0.29		0.59			0.51	0.74
A/A	41	11 (30%)	26 (70%)		2.1 (1.8, 2.3)	1 (Reference)	1 (Reference)		6.0 (4.3, 9.0)	1 (Reference)		1 (Reference)		
A/G ^a	23	5 (22%)	18 (78%)		1.8 (1.7, 1.9)	1.29 (0.79, 2.11)	1.17 (0.67, 2.03)		4.7 (2.6, 7.6)	1.18 (0.71, 1.96)		1.10 (0.62, 1.97)		
G/G ^a	5	1 (20%)	4 (80%)											
SMAD2 rs1792671				0.39				0.025		0.047			0.98	0.95
G/G	29	5 (18%)	23 (82%)		1.9 (1.7, 2.1)	1 (Reference)	1 (Reference)		5.9 (3.7, 9.0)	1 (Reference)		1 (Reference)		
G/A ^a	30	9 (33%)	18 (67%)		1.9 (1.8, 3.0)	0.61 (0.35, 1.03)	0.56 (0.32, 0.99)		5.6 (3.4, 7.9)	0.99 (0.60, 1.64)		0.98 (0.59, 1.64)		
A/A ^a	10	3 (30%)	7 (70%)											
FOXO3 rs12212067				0.71				0.62		0.71			0.32	0.82
T/T	56	15 (28%)	39 (72%)		1.9 (1.8, 2.1)	1 (Reference)	1 (Reference)		6.0 (4.1, 8.0)	1 (Reference)		1 (Reference)		
T/G	13	2 (18%)	9 (82%)		1.8 (1.0, 3.0)	1.17 (0.62, 2.20)	0.87 (0.40, 1.86)		4.5 (2.6, 5.8)	1.38 (0.71, 2.70)		1.09 (0.51, 2.33)		
Male														
INHBA rs2237432				0.48				0.93		0.84			0.43	0.21
A/A	41	18 (45%)	22 (55%)		2.3 (2.0, 3.8)	1 (Reference)	1 (Reference)		9.4 (6.4, 12.0)	1 (Reference)		1 (Reference)		
A/G ^a	32	11 (35%)	20 (65%)		2.2 (1.8, 3.1)	0.98 (0.63, 1.53)	0.95 (0.60, 1.51)		5.4 (3.1, 8.9)	1.19 (0.75, 1.90)		1.37 (0.84, 2.24)		
G/G ^a	8	4 (57%)	3 (43%)											
MSTN rs7570532				0.56				0.87		0.51			0.62	0.55
A/A	48	22 (47%)	25 (53%)		2.5 (1.9, 3.8)	1 (Reference)	1 (Reference)		8.7 (5.6, 10.7)	1 (Reference)		1 (Reference)		
A/G ^a	31	10 (34%)	19 (66%)		2.1 (1.8, 2.8)	0.97 (0.61, 1.52)	1.18 (0.72, 1.94)		5.3 (3.6, 9.6)	0.89 (0.56, 1.44)		1.17 (0.70, 1.95)		
G/G ^a	2	1 (50%)	1 (50%)											
SMAD2 rs1792671				0.22				0.17		0.61			0.19	0.39

(Continued)

Table 3. (Continued)

Genotype	N	Tumor response			Progression-free survival					Overall survival				
		PR+SD	PD	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*
G/G	26	13 (52%)	12 (48%)		2.8 (1.9, 3.8)	1 (Reference)		1 (Reference)		9.1 (4.4, 11.9)	1 (Reference)		1 (Reference)	
G/A	34	15 (45%)	18 (55%)		2.8 (1.8, 3.9)	1.04 (0.62, 1.75)		1.07 (0.63, 1.83)		8.3 (5.0, 13.9)	0.88 (0.51, 1.50)		0.85 (0.49, 1.48)	
A/A	20	5 (26%)	14 (74%)		1.9 (1.8, 2.3)	1.63 (0.89, 2.99)		1.40 (0.71, 2.76)		4.9 (2.3, 10.1)	1.46 (0.79, 2.68)		1.35 (0.70, 2.58)	
FOXO3 rs12212067				0.026					0.025					0.009
T/T	58	27 (48%)	29 (52%)		2.6 (2.1, 3.9)	1 (Reference)		1 (Reference)		9.1 (6.3, 10.7)	1 (Reference)		1 (Reference)	
T/G ^a	21	4 (20%)	16 (80%)		2.0 (1.6, 2.3)	1.71 (1.03, 2.84)		1.99 (1.19, 3.34)		4.3 (2.3, 8.7)	1.83 (1.08, 3.09)		2.17 (1.25, 3.75)	
G/G ^a	1	1 (100%)	0 (0%)											

Abbreviations: PR, partial response; SD, stable disease; PD, progressive disease; HR, hazard ratio; CI, confidence interval.

* P value based on Fisher’s exact test for tumor response; log-rank test for progression-free survival (PFS) and overall survival (OS) in the univariate analysis (†); and Wald test for PFS and OS in the multivariable Cox regression model adjusted for time to start of regorafenib treatment (<18 vs. ≥18 months), ECOG performance status (0 vs. 1 or 2), primary tumor resection (yes vs. no), and Kohne score (low-intermediate vs. high) (‡). P values < 0.050 are shown in bold text.

^a In the dominant model.

+ Estimates not yet reached.

<https://doi.org/10.1371/journal.pone.0239439.t003>

muscle showed upregulation of muscle *Fn14* during muscle wasting [23]. Furthermore, almost all patients with stage IV CRC (93%) have enhanced tumor expression of activin compared with only 40% of patients with stage I CRC [24]. These data indicate that activin expression is higher in more advanced CRC. *INHBA* rs2237432 is reported to have a significant association with fertility [25], although the clinical significance in cancer remains unknown.




Fig 1. Kaplan–Meier cumulative overall survival probability curves stratified by *INHBA* rs2237432 in female patients in (A) the discovery cohort and (B) the validation cohort.

<https://doi.org/10.1371/journal.pone.0239439.g001>

Table 4. Association between cachexia-related gene polymorphism and clinical outcome by gender subgroup in the validation cohort.

Genotype	N	Tumor response			Progression-free survival					Overall survival				
		SD	PD	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*
Female														
INHBA rs2237432				0.51				0.55		0.84			0.15	0.059
A/A	23	7 (39%)	11 (61%)		1.8 (1.8, 3.3)	1 (Reference)		1 (Reference)		8.1 (4.0, 12.6)	1 (Reference)		1 (Reference)	
A/G ^a	15	5 (50%)	5 (50%)		1.8 (1.1, 2.5)	1.20 (0.64, 2.25)		1.07 (0.55, 2.08)		9.6 (5.2, 29.9+)	0.58 (0.27, 1.25)		0.46 (0.21, 1.03)	
G/G ^a	3	0 (0%)	2 (100%)											
MSTN rs7570532				0.56				0.42		0.83			0.81	0.46
A/A	22	6 (40%)	9 (60%)		1.8 (1.1, 3.3)	1 (Reference)		1 (Reference)		9.6 (4.0, 13.9)	1 (Reference)		1 (Reference)	
A/G ^a	18	5 (36%)	9 (64%)		2.0 (1.6, 3.3)	0.79 (0.42, 1.49)		0.93 (0.48, 1.82)		8.1 (3.6, 12.9)	1.09 (0.53, 2.28)		1.32 (0.62, 2.81)	
G/G ^a	1	1 (100%)	0 (0%)											
SMAD2 rs1792671				1.00				0.81		0.68			0.91	0.68
G/G	31	8 (38%)	13 (62%)		1.8 (1.5, 2.5)	1 (Reference)		1 (Reference)		9.6 (4.0, 12.9)	1 (Reference)		1 (Reference)	
G/A	10	4 (44%)	5 (56%)		1.9 (0.9, 3.7)	1.09 (0.51, 2.29)		0.84 (0.36, 1.93)		7.8 (4.0, 15.3)	1.05 (0.46, 2.37)		0.83 (0.33, 2.04)	
FOXO3 rs12212067				0.61				0.072		0.30			0.56	0.83
T/T	35	9 (36%)	16 (64%)		1.8 (1.7, 2.0)	1 (Reference)		1 (Reference)		8.1 (5.8, 12.0)	1 (Reference)		1 (Reference)	
T/G ^a	6	2 (40%)	3 (60%)		3.3 (0.5, 11.9)	0.53 (0.21, 1.30)		0.58 (0.21, 1.62)		13.9 (1.7, 16.9+)	0.76 (0.28, 2.02)		0.90 (0.33, 2.44)	
G/G ^a	1	1 (100%)	0 (0%)											
Male														
INHBA rs2237432				0.52				0.43		0.24			0.73	0.62
A/A	16	8 (53%)	7 (47%)		2.3 (1.7, 4.2)	1 (Reference)		1 (Reference)		7.6 (4.1, 27.7+)	1 (Reference)		1 (Reference)	
A/G ^a	14	9 (69%)	4 (31%)		2.8 (1.7, 4.6)	0.77 (0.39, 1.51)		0.66 (0.33, 1.33)		10.3 (4.0, 27.2+)	0.86 (0.36, 2.08)		1.27 (0.50, 3.23)	
G/G ^a	6	2 (40%)	3 (60%)											
MSTN rs7570532				0.45				0.091		0.61			0.023	0.13
A/A	19	8 (47%)	9 (53%)		2.0 (1.3, 3.0)	1 (Reference)		1 (Reference)		6.3 (4.0, 12.9)	1 (Reference)		1 (Reference)	
A/G ^a	12	7 (64%)	4 (36%)		3.3 (2.0, 6.2)	0.58 (0.29, 1.14)		0.83 (0.41, 1.70)		26.7+ (6.5, 26.7+)	0.37 (0.15, 0.92)		0.47 (0.18, 1.24)	
G/G ^a	5	4 (80%)	1 (20%)											
SMAD2 rs1792671				0.35				0.38		0.19			0.82	0.18

(Continued)

Table 4. (Continued)

Genotype	N	Tumor response			Progression-free survival					Overall survival				
		SD	PD	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*
G/G	26	14 (61%)	9 (39%)		2.3 (1.8, 3.2)	1 (Reference)		1 (Reference)		11.8 (5.1, 27.7+)	1 (Reference)		1 (Reference)	
G/A ^a	8	5 (63%)	3 (38%)		3.8 (1.7, 4.7)	0.72 (0.33, 1.54)		0.58 (0.26, 1.30)		8.7 (3.1, 27.2+)	1.11 (0.43, 2.88)		2.09 (0.71, 6.16)	
A/A ^a	2	0 (0%)	2 (100%)											
FOXO3 rs12212067				0.24				0.18	0.47			0.040		0.069
T/T	28	12 (48%)	13 (52%)		2.3 (1.7, 3.0)	1 (Reference)		1 (Reference)		6.3 (4.0, 10.3)	1 (Reference)		1 (Reference)	
T/G	10	7 (78%)	2 (22%)		3.7 (1.9, 7.2)	0.61 (0.29, 1.29)		0.75 (0.34, 1.65)		27.2+ (1.9, 27.2+)	0.30 (0.09, 1.02)		0.30 (0.08, 1.10)	

Abbreviations: SD, stable disease; PD, progressive disease; HR, hazard ratio; CI, confidence interval.

* P value based on Fisher’s exact test for tumor response, log-rank test for progression-free survival (PFS) and overall survival (OS) in the univariate analysis (†), and Wald test for PFS and OS in the multivariable Cox regression model adjusted for liver metastasis and lymph node involvement (§). P values < 0.1 are shown in bold text.

^a In the dominant model.

+ Estimates not yet reached.

<https://doi.org/10.1371/journal.pone.0239439.t004>

Activin has been associated with angiogenesis, but unlike the positive correlation between activin overexpression and cancer cachexia, several studies have reported conflicting data on the relationship of activin overexpression with angiogenesis in various tissue types. Activin A increases *VEGF* expression via the physical interaction of *SMAD2* with the *MAPK*-regulated transcription factor *SP1* in hepatocellular carcinoma [26]. In contrast, activin A acts as a tumor suppressor in neuroblastoma [27] and gastric cancer [28] cells via the inhibition of




Fig 2. Kaplan–Meier cumulative overall survival probability curves stratified by FOXO3 rs12212067 in male patients in (A) the discovery cohort and (B) the validation cohort.

<https://doi.org/10.1371/journal.pone.0239439.g002>

Table 5. Association between *INHBA* rs2237432 and clinical outcome in female patients in the control cohort (Italian TAS-102 cohort).

Genotype	N	Tumor response			Progression-free survival					Overall survival				
		SD	PD	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*	Median, months (95%CI)	HR (95%CI) †	P value*	HR (95%CI) ‡	P value*
<i>INHBA</i> rs2237432				1.00			0.28		0.85			0.097		0.24
A/A	26	7 (27%)	19 (73%)		2.3 (1.9, 2.6)	1 (Reference)		1 (Reference)		7.3+ (3.7, 7.3+)	1 (Reference)		1 (Reference)	
A/G ^a	22	6 (27%)	16 (73%)		2.0 (1.7, 2.4)	1.38 (0.73, 2.60)		1.07 (0.55, 2.06)		4.1 (2.7, 5.5+)	2.03 (0.85, 4.86)		1.77 (0.69, 4.56)	
G/G ^a	1	0 (0%)	1 (100%)											

Abbreviation: SD, stable disease; PD, progressive disease; HR, hazard ratio; CI, confidence interval.

* P value based on Fisher's exact test for response; log-rank test in the univariate analysis (†); and Wald test in the multivariate analysis within Cox regression model adjusted for age group (<61 vs ≥61), liver metastasis, ECOG performance status, previous anti-EGFR therapy (‡). P values < 0.050 are shown in bold text.

^a In the dominant model.

+ Estimates not yet reached.

<https://doi.org/10.1371/journal.pone.0239439.t005>

VEGF mediated-angiogenesis. These findings suggest that activin has dual proinflammatory and anti-inflammatory roles, depending on the cell type and stage of cancer development. Another potential activin-related mechanism is *CCL/CCR*-dependent angiogenesis. *CCL2* binds its receptor *CCR2* to promote angiogenesis by recruiting macrophages [29, 30]. Activin has a critical role in controlling the expression of *CCL2/CCR2* in macrophages by increasing *CCR2* expression while inhibiting *CCL2* expression [31]. Regorafenib is a small molecule that inhibits various intracellular kinases involved in tumor angiogenesis, metastasis, oncogenesis, and tumor immunity. Our results especially found a correlation with tumor angiogenesis. Regorafenib inhibits tumor angiogenesis through inhibiting VEGFR1-3 and TIE2. A preclinical study indicated that *INHBA* exerts diverse effects on the VEGF pathway, including upregulation of the ligand, VEGF, as well as VEGF receptors [16]. Considering these data together, *INHBA* polymorphism may be associated with the effect of regorafenib through exerting its actions via VEGFR.

INHBA rs2237432 is an intronic SNP that is classified as a synonymous SNP. Generally, non-synonymous SNPs are considered to affect gene behavior even more considerably than synonymous SNPs. However, some intronic SNPs may affect gene splicing or expression, and such SNPs may have an effect on the function of a gene [32, 33]. Indeed, prediction tools revealed that *INHBA* rs2237432 might have a role as a strong enhancer of *INHBA* expression [33]. These suggest that rs2237432 is associated with the expression of *INHBA*.

In this study, the association between *INHBA* rs2237432 and clinical outcome was demonstrated only in female patients. Activin is an important modulator of follicle-stimulating synthesis and secretion of hormones such as estrogen and progesterone [34, 35]. Several studies have demonstrated the importance of activin and estrogen crosstalk during cancer initiation [36–38]. In addition, estrogen is reported to suppress activin subunit gene promoter activities [39], suggesting that activin activities differ by gender. These results may explain why the association between activin polymorphism and clinical outcome was dependent on gender. Unfortunately, however, because of the lack of samples we were unable to evaluate estrogen levels.

We included 128 patients who were treated with TAS-102 as the control cohort. TAS-102 is an oral drug that combines trifluridine and thymidine phosphorylase inhibitor [40]. The main antitumor effect of TAS-102 is due to DNA dysfunction by trifluridine incorporation into DNA

[41]. In the TAS-102 cohort, contrary to the regorafenib cohort, *INHBA* rs2237432 at any G allele showed a trend toward worse PFS and OS compared with that at the A/A allele. However, these differences did not reach statistical significance, indicating that *INHBA* rs2237432 has a specific association with regorafenib efficacy. A recent retrospective study showed comparable efficacy between regorafenib and TAS-102 [42]. However, a systemic review demonstrated that regorafenib was associated with more toxicity compared with TAS-102 [43]. On the basis of these results, female patients with the *INHBA* rs2237432 A/A allele should avoid regorafenib treatment and be treated with TAS-102 or best supportive care. Such a biomarker-based strategy will identify patients who are eligible for regorafenib treatment, resulting in improved clinical outcomes and quality of life for all patients treated with regorafenib.

Our study also indicated that the impact of *FOXO3* rs12212067 on OS differed significantly between the discovery and validation cohorts. This finding may result from etiological differences between Japanese and Italian populations. Several studies have shown that *FOXO3* rs12212067 is associated with the clinical course of inflammatory diseases such as Crohn's disease or rheumatoid arthritis [44, 45]. *FOXO3* has also been linked to the regulation of immune responses using systems biology [46] and knockout mouse models [47]. Furthermore, the *FOXO3* rs12212067 T/T allele is significantly associated with increased inflammatory cytokine production by monocytes (*IL-6*, *IL-8*, *IL-1beta* and *TNF-alfa*) compared with the G/G variant [48]. The pathogenesis of cachexia may be influenced by various factors, including genetic predisposition, inflammatory cytokines, and hormonal aspects. Especially, SNPs within inflammatory cytokine genes can affect cytokine levels and the degree of inflammation, and these SNP functions are reported to differ according to ethnicity [49].

This study had some limitations, such as the sample size and the retrospective design. In addition, we were unable to correlate the *INHBA* and *FOXO3* polymorphisms with intratumoral or serum expression levels, which may have clarified the mechanisms of regorafenib resistance. We were also unable to determine the relationship between the polymorphisms and skeletal muscle mass. In addition, this study presents no information on *RAS* mutation status in the Japanese cohort. We did not confirm our previous results showing that the *ACVR2B* rs2268753 genotype was associated with survival in *RAS* mutant mCRC patients receiving first-line anti-VEGF therapy, which warrants further investigation.

In conclusion, we evaluated for the first time the association of genetic variations in cancer cachexia-associated genes with clinical outcome in mCRC patients treated with regorafenib. We found that *INHBA* rs2237432 was significantly associated with clinical outcomes in female mCRC patients treated with regorafenib. Our findings may contribute to the identification of predictive or prognostic biomarkers of regorafenib therapy and potential drug targets in mCRC patients with cancer cachexia. Further studies are required, however, to fully elucidate the underlying biological mechanisms of the cachexia disease pathway.

Supporting information

S1 Table. Polymorphisms and primers.

(DOCX)

S2 Table. Association between cachexia-related gene polymorphism and clinical outcome in the validation cohort (Japanese regorafenib cohort).

(DOCX)

Acknowledgments

We thank Clare Cox, PhD, and H. Nikki March, PhD, from Edanz Group (www.edanzediting.com/ac) for editing a draft of this manuscript.

Author Contributions

Conceptualization: Yuji Miyamoto, Heinz-Josef Lenz.

Data curation: Marta Schirripa, Mitsukuni Suenaga, Shu Cao, Fotios Loupakis.

Formal analysis: Shu Cao, Dongyun Yang.

Investigation: Yuji Miyamoto, Marta Schirripa, Mitsukuni Suenaga, Satoshi Okazaki, Martin D. Berger, Satoshi Matsusaka.

Resources: Marta Schirripa, Mitsukuni Suenaga, Fotios Loupakis, Sara Lonardi, Filippo Pietrantonio, Beatrice Borelli, Chiara Cremolini, Toshiharu Yamaguchi.

Supervision: Wu Zhang, Satoshi Okazaki, Martin D. Berger, Yan Ning, Hideo Baba, Heinz-Josef Lenz.

Writing – original draft: Yuji Miyamoto.

Writing – review & editing: Wu Zhang, Hideo Baba, Heinz-Josef Lenz.

References

1. Wilhelm SM, Dumas J, Adnane L, Lynch M, Carter CA, Schutz G, et al. Regorafenib (BAY 73–4506): a new oral multikinase inhibitor of angiogenic, stromal and oncogenic receptor tyrosine kinases with potent preclinical antitumor activity. *Int J cancer*. 2011; 129: 245–255. <https://doi.org/10.1002/ijc.25864> PMID: 21170960
2. Grothey A, Van Cutsem E, Sobrero A, Siena S, Falcone A, Ychou M, et al. Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT): An international, multicentre, randomised, placebo-controlled, phase 3 trial. *Lancet*. 2013; 381: 303–312. [https://doi.org/10.1016/S0140-6736\(12\)61900-X](https://doi.org/10.1016/S0140-6736(12)61900-X) PMID: 23177514
3. Li J, Qin S, Xu R, Yau TCC, Ma B, Pan H, et al. Regorafenib plus best supportive care versus placebo plus best supportive care in Asian patients with previously treated metastatic colorectal cancer (CONCUR): A randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol*. 2015; 16: 619–629. [https://doi.org/10.1016/S1470-2045\(15\)70156-7](https://doi.org/10.1016/S1470-2045(15)70156-7) PMID: 25981818
4. Network NCC. National Comprehensive Cancer Network (NCCN). Guidel Version 1 2016 Colon Cancer. 2016. Available: <http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0001308/>
5. Van Cutsem E, Cervantes A, Adam R, Sobrero A, Van Krieken JH, Aderka D, et al. ESMO consensus guidelines for the management of patients with metastatic colorectal cancer. *Ann Oncol*. 2016;0: mdw235. <https://doi.org/10.1093/annonc/mdw235> PMID: 27380959
6. Suenaga M, Mashima T, Kawata N, Wakatsuki T, Horiike Y, Matsusaka S, et al. Serum VEGF-A and CCL5 levels as candidate biomarkers for efficacy and toxicity of regorafenib in patients with metastatic colorectal cancer. *Oncotarget*. 2016; 7. <https://doi.org/10.18632/oncotarget.9187> PMID: 27166185
7. Tabernero J, Lenz H-J, Siena S, Sobrero A, Falcone A, Ychou M, et al. Analysis of circulating DNA and protein biomarkers to predict the clinical activity of regorafenib and assess prognosis in patients with metastatic colorectal cancer: a retrospective, exploratory analysis of the CORRECT trial. *Lancet Oncol*. 2015; 16: 937–948. [https://doi.org/10.1016/S1470-2045\(15\)00138-2](https://doi.org/10.1016/S1470-2045(15)00138-2) PMID: 26184520
8. Fearon K, Strasser F, Anker SD, Bosaeus I, Bruera E, Fainsinger RL, et al. Definition and classification of cancer cachexia: An international consensus. *Lancet Oncol*. 2011; 12: 489–495. [https://doi.org/10.1016/S1470-2045\(10\)70218-7](https://doi.org/10.1016/S1470-2045(10)70218-7) PMID: 21296615
9. Morishita S, Kaida K, Tanaka T, Itani Y, Ikegame K, Okada M, et al. Prevalence of sarcopenia and relevance of body composition, physiological function, fatigue, and health-related quality of life in patients before allogeneic hematopoietic stem cell transplantation. *Support Care Cancer*. 2012; 20: 3161–3168. <https://doi.org/10.1007/s00520-012-1460-5> PMID: 22526152
10. Andreyev HJ, Norman AR, Oates J, Cunningham D. Why do patients with weight loss have a worse outcome when undergoing chemotherapy for gastrointestinal malignancies? *Eur J Cancer*. 1998; 34: 503–509. [https://doi.org/10.1016/s0959-8049\(97\)10090-9](https://doi.org/10.1016/s0959-8049(97)10090-9) PMID: 9713300
11. Han HQ, Zhou X, Mitch WE, Goldberg AL. Myostatin/activin pathway antagonism: Molecular basis and therapeutic potential. *Int J Biochem Cell Biol*. 2013; 45: 2333–2347. <https://doi.org/10.1016/j.biocel.2013.05.019> PMID: 23721881

12. Loumaye A, De Barys M, Nachit M, Lause P, Frateur L, Van Maanen A, et al. Role of activin A and myostatin in human cancer cachexia. *J Clin Endocrinol Metab.* 2015; 100: 2030–2038. <https://doi.org/10.1210/jc.2014-4318> PMID: 25751105
13. Miyamoto Y, Hanna DL, Zhang W, Baba H, Lenz H-J. Molecular Pathways: Cachexia Signaling—A Targeted Approach to Cancer Treatment. *Clin cancer Res.* 2016; 22: 3999–4004. <https://doi.org/10.1158/1078-0432.CCR-16-0495> PMID: 27340276
14. Okano M, Yamamoto H, Ohkuma H, Kano Y, Kim H, Nishikawa S, et al. Significance of INHBA expression in human colorectal cancer. *Oncol Rep.* 2013; 30: 2903–2908. <https://doi.org/10.3892/or.2013.2761> PMID: 24085226
15. Gallot YS, Durieux AC, Castells J, Desgeorges MM, Vernus B, Plantureux L, et al. Myostatin gene inactivation prevents skeletal muscle wasting in cancer. *Cancer Res.* 2014; 74: 7344–7356. <https://doi.org/10.1158/0008-5472.CAN-14-0057> PMID: 25336187
16. Maeshima K, Maeshima A, Hayashi Y, Kishi S, Kojima I. Crucial role of activin a in tubulogenesis of endothelial cells induced by vascular endothelial growth factor. *Endocrinology.* 2004; 145: 3739–3745. <https://doi.org/10.1210/en.2004-0213> PMID: 15117880
17. Breit S, Ashman K, Wilting J, Rossler J, Hatzl E, Fotsis T, et al. The N-myc oncogene in human neuroblastoma cells: down-regulation of an angiogenesis inhibitor identified as activin A. *Cancer Res.* 2000; 60: 4596–4601. PMID: 10969812
18. Miyamoto Y, Stintzing S, Loupakis F, Zhang W, Cao S, Ning Y, et al. Genetic variations associated with cancer cachexia pathways to predict survival in metastatic colorectal cancer (mCRC): Results from FIRE-3 and TRIBE. *J Clin Oncol.* 2016; 34: (suppl; abstr 3590).
19. Miyamoto Y, Baba Y, Sakamoto Y, Ohuchi M, Tokunaga R, Kurashige J, et al. Negative Impact of Skeletal Muscle Loss after Systemic Chemotherapy in Patients with Unresectable Colorectal Cancer. *PLoS One.* 2015; 10: e0129742. <https://doi.org/10.1371/journal.pone.0129742> PMID: 26069972
20. McShane LM, Altman DG, Sauerbrei W, Taube SE, Gion M, Clark GM. REporting recommendations for tumor MARKer prognostic studies (REMARK). *Nat Clin Pract Oncol.* 2005; 2: 416–422. PMID: 16130938
21. Chen YG, Lui HM, Lin SL, Lee JM, Ying SY. Regulation of cell proliferation, apoptosis, and carcinogenesis by activin. *Exp Biol Med.* 2002; 227: 75–87. Available: <http://www.ncbi.nlm.nih.gov/pubmed/11815670>
22. Chen Y-G, Wang Q, Lin S-L, Chang CD, Chuang J, Ying S-Y. Activin signaling and its role in regulation of cell proliferation, apoptosis, and carcinogenesis. *Exp Biol Med (Maywood).* 2006; 231: 534–544. doi:231/5/534 [pii]
23. Johnston AJ, Murphy KT, Jenkinson L, Laine D, Emmrich K, Faou P, et al. Targeting of Fn14 Prevents Cancer-Induced Cachexia and Prolongs Survival. *Cell.* 2015; 162: 1365–1378. <https://doi.org/10.1016/j.cell.2015.08.031> PMID: 26359988
24. Wildi S, Kleeff J, Maruyama H, Maurer CA, Buchler MW, Korc M. Overexpression of activin A in stage IV colorectal cancer. *Gut.* 2001; 49: 409–417. <https://doi.org/10.1136/gut.49.3.409> PMID: 11511564
25. Kuningas M, Altmäe S, Uitterlinden AG, Hofman A, Van Duijn CM, Tiemeier H. The relationship between fertility and lifespan in humans. *Age (Omaha).* 2011; 33: 615–622. <https://doi.org/10.1007/s11357-010-9202-4> PMID: 21222045
26. Wagner K, Peters M, Scholz A, Benckert C, Ruderisch HS, Wiedenmann B, et al. Activin A stimulates vascular endothelial growth factor gene transcription in human hepatocellular carcinoma cells. *Gastroenterology.* 2004; 126: 1828–1843. <https://doi.org/10.1053/j.gastro.2004.03.011> PMID: 15188178
27. Panopoulou E, Murphy C, Rasmussen H, Bagli E, Rofstad EK, Fotsis T. Activin A suppresses neuroblastoma xenograft tumor growth via antimitotic and antiangiogenic mechanisms. *Cancer Res.* 2005; 65: 1877–1886. <https://doi.org/10.1158/0008-5472.CAN-04-2828> PMID: 15753386
28. Togashi Y, Kogita A, Sakamoto H, Hayashi H, Terashima M, de Velasco MA, et al. Activin signal promotes cancer progression and is involved in cachexia in a subset of pancreatic cancer. *Cancer Lett.* 2015; 356: 819–827. <https://doi.org/10.1016/j.canlet.2014.10.037> PMID: 25449777
29. Salcedo R, Ponce ML, Young HA, Wasserman K, Ward JM, Kleinman HK, et al. Human endothelial cells express CCR2 and respond to MCP-1: direct role of MCP-1 in angiogenesis and tumor progression. *Blood.* 2000; 96: 34–40. PMID: 10891427
30. Goede V, Brogelli L, Ziche M, Augustin HG. Induction of inflammatory angiogenesis by monocyte chemoattractant protein-1. *Int J cancer.* 1999; 82: 765–770. [https://doi.org/10.1002/\(sici\)1097-0215\(19990827\)82:5<765::aid-ijc23>3.0.co;2-f](https://doi.org/10.1002/(sici)1097-0215(19990827)82:5<765::aid-ijc23>3.0.co;2-f) PMID: 10417778
31. Sierra-Filardi E, Nieto C, Dominguez-Soto A, Barroso R, Sanchez-Mateos P, Puig-Kroger A, et al. CCL2 shapes macrophage polarization by GM-CSF and M-CSF: identification of CCL2/CCR2-dependent gene expression profile. *J Immunol.* 2014; 192: 3858–3867. <https://doi.org/10.4049/jimmunol.1302821> PMID: 24639350

32. Goto Y, Yue L, Yokoi A, Nishimura R, Uehara T, Koizumi S, et al. A novel single-nucleotide polymorphism in the 3'-untranslated region of the human dihydrofolate reductase gene with enhanced expression. *Clin cancer Res an Off J Am Assoc Cancer Res*. 2001; 7: 1952–1956.
33. Millar DS, Horan M, Chuzhanova NA, Cooper DN. Characterisation of a functional intronic polymorphism in the human growth hormone (GH1) gene. *Hum Genomics*. 2010; 4: 289–301. <https://doi.org/10.1186/1479-7364-4-5-289> PMID: 20650818
34. Hsueh AJ, Bicsak TA, Jia XC, Dahl KD, Fauser BC, Galway AB, et al. Granulosa cells as hormone targets: the role of biologically active follicle-stimulating hormone in reproduction. *Recent Prog Horm Res*. 1989; 45: 207–209.
35. Risbridger GP, Schmitt JF, Robertson DM. Activins and inhibins in endocrine and other tumors. *Endocr Rev*. 2001; 22: 836–858. <https://doi.org/10.1210/edrv.22.6.0450> PMID: 11739336
36. Drummond AE, Fuller PJ. Activin and inhibin, estrogens and NFkappaB, play roles in ovarian tumorigenesis is there crosstalk? *Mol Cell Endocrinol*. 2012; 359: 85–91. <https://doi.org/10.1016/j.mce.2011.07.033> PMID: 21839804
37. Bloise E, Couto HL, Massai L, Ciarmela P, Mencarelli M, Borges LE, et al. Differential expression of follistatin and FLRG in human breast proliferative disorders. *BMC Cancer*. 2009; 9: 320. <https://doi.org/10.1186/1471-2407-9-320> PMID: 19740438
38. Burdette JE, Woodruff TK. Activin and estrogen crosstalk regulates transcription in human breast cancer cells. *Endocr Relat Cancer*. 2007; 14: 679–689. <https://doi.org/10.1677/ERC-07-0054> PMID: 17914098
39. Kipp JL, Kilen SM, Bristol-Gould S, Woodruff TK, Mayo KE. Neonatal exposure to estrogens suppresses activin expression and signaling in the mouse ovary. *Endocrinology*. 2007; 148: 1968–1976. <https://doi.org/10.1210/en.2006-1083> PMID: 17255206
40. Mayer RJ, Van Cutsem E, Falcone A, Yoshino T, Garcia-Carbonero R, Mizunuma N, et al. Randomized trial of TAS-102 for refractory metastatic colorectal cancer. *N Engl J Med*. 2015; 372: 1909–1919. <https://doi.org/10.1056/NEJMoa1414325> PMID: 25970050
41. Miyamoto Y, Lenz H-J, Baba H. A novel antimetabolite: TAS-102 for metastatic colorectal cancer. *Expert Rev Clin Pharmacol*. 2016; 9: 355–365. <https://doi.org/10.1586/17512433.2016.1133285> PMID: 26677869
42. Moriwaki T, Fukuoka S, Taniguchi H, Takashima A, Kumekawa Y, Kajiwara T, et al. Propensity Score Analysis of Regorafenib Versus Trifluridine/Tipiracil in Patients with Metastatic Colorectal Cancer Refractory to Standard Chemotherapy (REGOTAS): A Japanese Society for Cancer of the Colon and Rectum Multicenter Observational Study. *Oncologist*. 2018; 23: 7–15. <https://doi.org/10.1634/theoncologist.2017-0275> PMID: 28894015
43. Abrahao ABK, Ko YJ, Berry S, Chan KKW. A Comparison of Regorafenib and TAS-102 for Metastatic Colorectal Cancer: A Systematic Review and Network Meta-analysis. *Clin Colorectal Cancer*. 2018; 17: 113–120. <https://doi.org/10.1016/j.clcc.2017.10.016> PMID: 29174481
44. Marlow G, Han DY, Triggs CM, Ferguson LR. Food Intolerance: Associations with the rs12212067 Polymorphism of FOXO3 in Crohn's Disease Patients in New Zealand. *J Nutrigenet Nutrigenomics*. 2015; 8: 70–80. <https://doi.org/10.1159/000435783> PMID: 26226934
45. Viatte S, Lee JC, Fu B, Espeli M, Lunt M, De Wolf JNE, et al. Association Between Genetic Variation in FOXO3 and Reductions in Inflammation and Disease Activity in Inflammatory Polyarthritides. *Arthritis Rheumatol (Hoboken, NJ)*. 2016; 68: 2629–2636. <https://doi.org/10.1002/art.39760> PMID: 27214848
46. Litvak V, Ratushny A V, Lampano AE, Schmitz F, Huang AC, Raman A, et al. A FOXO3-IRF7 gene regulatory circuit limits inflammatory sequelae of antiviral responses. *Nature*. 2012; 490: 421–425. <https://doi.org/10.1038/nature11428> PMID: 22982991
47. Dejean AS, Beisner DR, Ch'en IL, Kerdiles YM, Babour A, Arden KC, et al. Transcription factor Foxo3 controls the magnitude of T cell immune responses by modulating the function of dendritic cells. *Nat Immunol*. 2009; 10: 504–513. <https://doi.org/10.1038/ni.1729> PMID: 19363483
48. Lee JC, Espéli M, Anderson CA, Linterman MA, Pocock JM, Williams NJ, et al. Human SNP links differential outcomes in inflammatory and infectious disease to a FOXO3-regulated pathway. *Cell*. 2013; 155: 57–69. <https://doi.org/10.1016/j.cell.2013.08.034> PMID: 24035192
49. Zabaleta J, Schneider BG, Ryckman K, Hooper PF, Camargo MC, Piazuelo MB, et al. Ethnic differences in cytokine gene polymorphisms: Potential implications for cancer development. *Cancer Immunol Immunother*. 2008; 57: 107–114. <https://doi.org/10.1007/s00262-007-0358-4> PMID: 17618436