Response to Mahir Gachabayov, et al, ‘Methodological Biases May Render a Clinical Study Underpowered’

Manuel Jakob^{1,2}, Guido Beldi, MD¹

¹Department of Visceral Surgery and Medicine, University Hospital, Bern, Switzerland

²Department of Microbiology, Infectious Diseases and Immunology,

Charité Universitätsmedizin Berlin, Germany

Corresponding author and address for reprints: Guido Beldi, MD. Department of Visceral Surgery and Medicine, Bern University Hospital, Bern, Switzerland. Office: 0041 316328275. guido.beldi@insel.ch

E-mail addresses of authors: Manuel Jakob: manuel.jakob@charite.de, Guido Beldi: guido.beldi@insel.ch

Disclosure: The authors of this manuscript have no conflict of interest to disclose.

Financial support: The authors received financial support by StratticeTM, Allergan

To the editorial board of Journal of Trauma

We acknowledge the letter to the editor and the potential conflicts of interests of the authors.

To avoid selection bias, the inclusion or exclusion of patients was carried out by an independent emergency department physician who included patients at the emergency department (please refer to page 42 of the trial protocol (1)). The ASA scores reported in the manuscript correspond to the scores as recorded by anesthesiologists at a later time point after the inclusion of study participants. Indeed, in some patients, the ASA score may have deteriorated after the study inclusion. Furthermore, the severity of comorbidities may have been estimated differently by physicians at study inclusion and anesthesiologists, leading to higher recorded ASA scores by the latter. Of note no patient rated as ASA 5 died within 30 days of the operation.

We acknowledge that many surgical facilities may not be familiar with intra-abdominal placement of mesh and this lack of experience may be detrimental to outcome. However, at the Bern University Hospital, placement of intra-abdominal meshes is performed routinely in both emergency and elective procedures. The implantation of intra-abdominal mesh has not been shown to be associated with specific complications in previous studies published by our department (2-6)~~[1-5]~~. Furthermore, the described complications in patients following mesh placement, such as mesh dissolution, non-integration into the abdominal wall or late onset mesh infection are unlikely to be related to the surgical technique.

Our study was designed and powered for incisional hernia as its primary outcome. However, detection bias as described in the letter to the editor concerns a safety outcome and not the primary outcome parameter. We agree with the authors that the meshes should not affect anastomotic healing and patient deterioration. However, patients #3 and #10 both had severe mesh-associated complications such as torn mesh, necrosis or hematoma of the abdominal wall that are highly likely to become clinically apparent. In addition, case #12, had a type 2 surgical site infection according to CDC criteria (7)~~[6]~~.

Based on the letter to the editor by Gachabayov and Latifi, we recalculated the p-value in question using SPSS and received the same result as reported in our publication. The corresponding SPSS output has been made available to the editorial board.

This study has been designed to assess incisional herniation in an at risk patient cohort. However, the significantly higher rate of complications seen with the implantation of Strattice meshes did not allow the completion of the study and to reach the primary endpoint. Therefore this study is obviously underpowered to identify differences for the primary endpoint. However, the main statement of the publication is the description of safety related events, which were significantly different. Thus, we disagree with the authors of this letter that no robust conclusion can be drawn from this study. Based on the results of this study, the prophylactic use of biologic Strattice meshes cannot be recommended in patients undergoing emergency abdominal surgery because of safety concerns.

Your Sincerely,

Manuel Jakob, MD

Guido Beldi, MD

References:

1. Jakob MO, Haltmeier T, Candinas D, Beldi G. Biologic mesh implantation is associated with serious abdominal wall complications in patients undergoing emergency abdominal surgery - a randomized-controlled clinical trial. *J Trauma Acute Care Surg*. 2020.
2. Jakob MO, Schwarz C, Haltmeier T, Zindel J, Pinworasarn T, Candinas D, Starlinger P, Beldi G. Mesh-augmented versus direct abdominal closure in patients undergoing open abdomen treatment. *Hernia*. 2018;22(5):785-92.
3. Jakob MO, Spari D, Zindel J, Pinworasarn T, Candinas D, Beldi G. Prophylactic, Synthetic Intraperitoneal Mesh Versus No Mesh Implantation in Patients with Fascial Dehiscence. *J Gastrointest Surg*. 2018.
4. Kohler A, Lavanchy JL, Lenoir U, Kurmann A, Candinas D, Beldi G. Effectiveness of Prophylactic Intraperitoneal Mesh Implantation for Prevention of Incisional Hernia in Patients Undergoing Open Abdominal Surgery: A Randomized Clinical Trial. *JAMA Surg*. 2019;154(2):109-15.
5. Scholtes M, Kurmann A, Seiler CA, Candinas D, Beldi G. Intraperitoneal mesh implantation for fascial dehiscence and open abdomen. *World J Surg*. 2012;36(7):1557-61.

6. Kurmann A, Barnetta C, Candinas D, Beldi G. Implantation of prophylactic nonabsorbable intraperitoneal mesh in patients with peritonitis is safe and feasible. *World J Surg.* 2013;37(7):1656-60.
7. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for Prevention of Surgical Site Infection, 1999. Centers for Disease Control and Prevention (CDC) Hospital Infection Control Practices Advisory Committee. *Am J Infect Control.* 1999;27(2):97-132; quiz 3-4; discussion 96.

Disclosure: The authors of this manuscript have no conflict of interest to disclose.

Financial support: The authors received financial support by Strattice™, Allergan (1).

Authors contributions: Jakob M.O., Beldi G.: Writing of the manuscript.

Supplemental Material: SPSS output HPACS (<http://links.lww.com/TA/B826>)

CROSSTABS

/TABLES=Strattice BY Dindo_greater_2_abdominal_wall_complication

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ

/CELLS=COUNT ROW

/COUNT ROUND CELL

/METHOD=EXACT TIMER(5) .

Crosstabs

Notes

Output Created		16-SEP-2020 08:15:34
Comments		
Input	Data	\\FILER300\USERS3002\0073998\Desktop\HPACS\HPACS SPSS\HPACS interim 7.sav
	Active Dataset	DataSet1
	Filter	Abdominal wall complication
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	13
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Syntax		CROSSTABS /TABLES=Strattice BY Dindo_greater_2_abdominal_wall_complication /FORMAT=AVALUE TABLES /STATISTICS=CHISQ /CELLS=COUNT ROW /COUNT ROUND CELL /METHOD=EXACT TIMER(5).
Resources	Processor Time	00:00:00.05
	Elapsed Time	00:00:00.06

Notes

Dimensions Requested	2
Cells Available	524245
Time for Exact Statistics	0:00:00.01

Case Processing Summary

	Valid		Cases Missing		Total	
	N	Percent	N	Percent	N	Percent
Strattice * Dindo_greater_2_abdominal_wall_complication	13	100.0%	0	0.0%	13	100.0%

Strattice * Dindo_greater_2_abdominal_wall_complication Crosstabulation

			Dindo_greater_2_abdominal_wall_complication		
			0	1	Total
Strattice	0	Count	6	1	7
		% within Strattice	85.7%	14.3%	100.0%
	1	Count	1	5	6
		% within Strattice	16.7%	83.3%	100.0%
Total	Count	7	6	13	
	% within Strattice	53.8%	46.2%	100.0%	

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	6.198 ^a	1	.013	.029	.025
Continuity Correction ^b	3.731	1	.053		
Likelihood Ratio	6.796	1	.009	.029	.025
Fisher's Exact Test				.029	.025
Linear-by-Linear Association	5.721 ^c	1	.017	.029	.025
N of Valid Cases	13				

Chi-Square Tests

	Point Probability
Pearson Chi-Square	
Continuity Correction ^b	
Likelihood Ratio	
Fisher's Exact Test	
Linear-by-Linear Association	.024
N of Valid Cases	

- a. 4 cells (100.0%) have expected count less than 5. The minimum expected count is 2.77.
- b. Computed only for a 2x2 table
- c. The standardized statistic is 2.392.