


Journal Pre-proof

Clinical recommendations for in-hospital airway management during aerosol-transmitting procedures in the setting of a viral pandemic

Alexander Fuchs, MD, Daniele Lanzi, MD, Christian M. Beilstein, MD, Thomas Riva, MD, Richard D. Urman, MD MBA, Markus M. Luedi, MD, MBA, Matthias Braun, MD



PII: S1521-6896(20)30126-9

DOI: <https://doi.org/10.1016/j.bpa.2020.12.002>

Reference: YBEAN 1147

To appear in: *Best Practice & Research Clinical Anaesthesiology*

Received Date: 26 November 2020

Accepted Date: 3 December 2020

Please cite this article as: Fuchs A, Lanzi D, Beilstein CM, Riva T, Urman RD, Luedi MM, Braun M, Clinical recommendations for in-hospital airway management during aerosol-transmitting procedures in the setting of a viral pandemic, *Best Practice & Research Clinical Anaesthesiology*, <https://doi.org/10.1016/j.bpa.2020.12.002>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier Ltd. All rights reserved.

1 **Title:** **Clinical recommendations for in-hospital airway management**
2 **during aerosol-transmitting procedures in the setting of a viral**
3 **pandemic**

4 **Authors:**

5 **1. Alexander Fuchs, MD**

6 Affiliation: Department of Anaesthesiology and Pain Medicine, Inselspital, Bern
7 University Hospital, University of Bern, Bern, Switzerland

8 Address: Inselspital, Bern University Hospital
9 Freiburgstrasse
10 CH-3010 Bern, Switzerland

11 Email: Alexander.Fuchs@insel.ch

12 Conflicts: This author reported no conflicts of interest

13 Attestation: This author has seen, reviewed and approved the final manuscript

14 **2. Daniele Lanzi, MD**

15 Affiliation: Department of Anaesthesiology and Pain Medicine, Inselspital, Bern
16 University Hospital, University of Bern, Bern, Switzerland

17 Address: Inselspital, Bern University Hospital
18 Freiburgstrasse
19 CH-3010 Bern, Switzerland

20 Email: Daniele.Lanzi@insel.ch

21 Conflicts: This author reported no conflicts of interest

22 Attestation: This author has seen, reviewed and approved the final manuscript

23 **3. Christian M. Beilstein, MD**

24 Affiliation: Department of Anaesthesiology and Pain Medicine, Inselspital, Bern
25 University Hospital, University of Bern, Bern, Switzerland

26 Address: Inselspital, Bern University Hospital
27 Freiburgstrasse

28 CH-3010 Bern, Switzerland
29 Email: Christian.Beilstein@insel.ch
30 Conflicts: This author reported no conflicts of interest
31 Attestation: This author has seen, reviewed and approved the final manuscript

32 **4. Thomas Riva, MD**

33 Affiliation: Department of Anaesthesiology and Pain Medicine, Inselspital, Bern
34 University Hospital, University of Bern, Bern, Switzerland
35 Address: Inselspital, Bern University Hospital
36 Freiburgstrasse
37 CH-3010 Bern, Switzerland
38 Email: Thomas.Riva@insel.ch
39 Conflicts: This author reported no conflicts of interest
40 Attestation: This author has seen, reviewed and approved the final manuscript

41 **5. Richard D. Urman, MD, MBA**

42 Affiliation: Department of Anesthesiology, Perioperative and Pain Medicine,
43 Brigham and Women's Hospital, Harvard Medical School, Boston, MA,
44 USA
45 Address: 75 Francis Street
46 Boston, MA 02115, USA
47 Email: rurman@bwh.harvard.edu
48 Conflicts: This author reported no conflicts of interest
49 Attestation: This author has seen, reviewed and approved the final manuscript

50 **6. Markus M. Luedi, MD, MBA**

51 Affiliation: Department of Anaesthesiology and Pain Medicine, Inselspital, Bern
52 University Hospital, University of Bern, Bern, Switzerland
53 Address: Inselspital, Bern University Hospital
54 Freiburgstrasse

55 CH-3010 Bern, Switzerland
56 Email: Markus.Luedi2@insel.ch
57 Conflicts: This author reported no conflicts of interest
58 Attestation: This author has seen, reviewed and approved the final manuscript

59 **7. Matthias Braun, MD**

60 Affiliation: Department of Anaesthesiology and Pain Medicine, Inselspital, Bern
61 University Hospital, University of Bern, Bern, Switzerland
62 Address: Inselspital, Bern University Hospital
63 Freiburgstrasse
64 CH-3010 Bern, Switzerland
65 Email: Matthias.Braun@insel.ch
66 Conflicts: This author reported no conflicts of interest
67 Attestation: This author has seen, reviewed and approved the final manuscript

68
69 **Word Counts** Abstract 146 words, Main text 4450 words, summary 275 words, 2
70 tables, 93 references

71 **Contributions** Alexander Fuchs helped to write the article
72 Daniele Lanzi helped to write the article
73 Markus M. Luedi helped to write the article
74 Christian M. Beilstein helped to write the article
75 Thomas Riva helped to write the article
76 Richard D. Urman helped to write the article
77 Matthias Braun helped to write the article

78 **Correspondence** Matthias Braun, MD, Department of Anaesthesiology and Pain
79 Medicine, Inselspital, Bern University Hospital, University of Bern,

80 Freiburgstrasse, 3010 Bern, Switzerland, Phone: +41 31 632 24 83,
81 Email: Matthias.Braun@insel.ch

82

83 **Abstract (146 words)**

84 The coronavirus disease 2019 (COVID-19) pandemic, caused by severe acute respiratory
85 syndrome coronavirus 2 (SARS-CoV-2), can lead to severe pneumonia and multi-organ
86 failure. While most of the infected patients develop no or only mild symptoms, some need
87 respiratory support or even invasive ventilation.

88 The exact route of transmission is currently under investigation. While droplet exposure and
89 direct contact seem to be the most significant ways of transmitting the disease, aerosol
90 transmission appears to be possible under circumstances favored by high viral load. Despite
91 the use of personal protective equipment (PPE), this situation potentially puts healthcare
92 workers at risk of infection, especially if they are involved in airway management.

93 Various recommendations and international guidelines aim to protect healthcare workers,
94 although evidence-based research confirming the benefits of these approaches is still
95 scarce. In this article, we summarize the current literature and recommendations for airway
96 management of COVID-19 patients.

97

98

99 **Keywords:** COVID-19, aerosol-transmitted viral diseases, airway management, intubation,
100 critical care, narrative review

101

102

103 **Introduction**

104 As of mid-November 2020, COVID-19—the illness caused by the novel severe acute
105 respiratory syndrome-coronavirus-2 (SARS-CoV-2)—has been responsible for more than 54
106 million confirmed infections and 1,300,000 deaths worldwide[1]. Due to the highly infectious
107 nature of the virus — with a high viral load in human airways — establishing precise and safe
108 airway management guidelines is essential to protect healthcare workers and bystanders.

109 SARS-CoV-2 is mainly transmitted by droplet spread and by contact with infected patients
110 and contaminated surfaces, but infection is also possible through airborne spread, especially
111 after aerosol-generating procedures (AGP)[2]. Droplet and airborne spread differ in the size
112 of the suspended particles. Aerosols are airborne particles that evaporate but remain in the
113 air longer before settling on surfaces (airborne transmission). In particular, particles that are
114 5 µm or smaller can survive in the air for a long time, and directly enter the lower respiratory
115 tract of a person [3]. Although there is no clear cutoff between large and small droplets[4-6],
116 SARS-CoV-2 can be transmitted either way, so taking precautions to prevent contact and
117 airborne spread is crucial[7, 8].

118 COVID-19 primarily causes severe viral hypoxemic pneumonia, with or without multiorgan
119 dysfunction. Critical care is needed for around 5% of the patients, and mortality varies
120 considerably between countries, with the case fatality ratio worldwide having reached 2.4 by
121 November 2020[1]. Most of the patients—especially those who are young (under the age of
122 65) and 80-90% of those without comorbidities—develop only mild symptoms or no
123 symptoms at all [9-11]. This makes it more difficult to track and isolate potentially infectious
124 subjects and to slow down or stop the spread of the virus.

125 Various societies involved in airway management have published guidelines and
126 recommendations for the treatment of COVID-19 patients [12-21]. We aim to summarize how
127 airway management recommendations have evolved over the course of the present
128 pandemic, and to suggest how clinicians can benefit from new guidelines to be prepared for

129 future aerosol-transmitted diseases[22, 23], with a focus on aerosol-generating procedures
 130 (AGP) and their risk to healthcare workers (Table 1).

131

132 **Risk for health care workers**

133 The odds ratios for infection of healthcare workers involved in the care of patients with the
 134 severe acute respiratory syndrome (SARS) in Table 1 are based on a review published in
 135 2012 containing low-grade to very low-grade evidence [23]. A number of procedures are
 136 associated with potential infection.

137 The unknown number of asymptomatic COVID-19-positive patients was probably
 138 underestimated, especially at the beginning of the pandemic [24]. Elective procedures are
 139 being performed again, and the discipline required for testing, maintaining distance from
 140 contacts, and limiting social interaction may drop.

141 An international prospective multicenter study found that performing an intubation places the
 142 involved healthcare workers at high risk of infection (up to 10%)[25].

Table 1. Aerosol-generating procedures (AGP) identified by the US Centers for Disease Control and Prevention [26] and odds ratios (OR) for the risk of SARS transmission for healthcare workers exposed vs. not-exposed to SARS (Tran et al.[23])

Aerosol-generating procedure (AGP)	Estimate OR
Tracheal intubation and extubation	6.6
Manual ventilation	1.3 - 2.8
Tracheotomy or tracheostomy procedures (insertion or removal)	4.2
Bronchoscopy	1.9
Non-invasive ventilation (NIV)	3.1
High-flow nasal cannula (HFNC)	0.4
High-frequency oscillatory ventilation (HFOV)	0.7
Induction of sputum using nebulized saline	0.9
Respiratory tract suctioning (before and after intubation)	1.3 - 3.5

Chest compressions during cardiopulmonary resuscitation (CPR)	1.4
Abbreviations: SARS: severe acute respiratory syndrome	

143

144

145 Risks of airway-related procedures

146 The risk of transmission or infection is increased for healthcare workers during aerosol-
147 generating procedures (AGPs) [7, 27, 23] due to the high virus concentration in the patient's
148 upper airway and sputum [28]. Protection of workers involved in airway management must
149 be a high priority, and personal protective equipment (PPE) should be available for the whole
150 team involved in treating suspected or confirmed COVID-19 patients. To minimize the risk of
151 multiple transmissions, the number of healthcare personnel involved in an aerosol-generating
152 procedure should be reduced to the minimum, while maintaining the patient's safety at all
153 times. All of these procedures are best performed in rooms with negative pressure or with air
154 exchange rates of up to 6-12 times per hour [29, 12, 13]. Some authors even suggest
155 avoiding rooms with positive pressure (e.g., operating rooms (OR) for airway procedures
156 [13]. However, the recommendation seems to be difficult to apply due to lack of availability of
157 emergency rooms (ERs), ORs and intensive care units (ICUs). A preoperative huddle
158 consisting of all staff involved in the care of the patient is important to improve teamwork and
159 reduce unnecessary exposure [30].

160

161 Tracheal intubation

162 A severe COVID-19 infection can lead to acute respiratory failure and therefore requires
163 emergency intubation. Such patients are characterized by very high viral spread[31]. The
164 point at which intubation is indicated may differ from institution to institution and also may be
165 influenced by the existing resources in an acute pandemic situation, such as the availability
166 of beds with mechanical ventilation and ICU care, or the possibility to relocate an intubated

167 patient. Evidence is lacking with regard to the best time for intubation, but the current
168 recommendations tend to avoid emergency intubation of unstable patients, instead providing
169 early intubation in order to protect the involved employees [32, 13, 12]. Adapting an
170 institution's existing airway management algorithms to the pandemic situation seems to be
171 the most appropriate approach, due to prior experience and high acceptance [13, 33-37].

172 Some published consensus guidelines [12, 21, 13] suggested the creation of dedicated
173 intubation teams. These teams should be composed of an experienced airway manager and
174 1-2 assistants per patient, regardless of whether COVID-19 is confirmed or suspected.
175 Reducing the number of potentially exposed persons and the exposure time during an AGP
176 can potentially shorten workers' exposure to a high viral load and could be one of the keys to
177 minimizing new infections. Team members should be familiar with the airway tools,
178 monitoring equipment, ventilators and drugs used to safely perform tracheal intubation.
179 Simulation training of airway management and PPE can significantly increase adherence to
180 standards and therefore the safety of patients and healthcare providers [21]. The method of
181 communication has to be established before the procedure starts, and should result in an
182 "airway plan". Clear the procedure for every team member, and try to communicate directly
183 and in a closed-loop manner, both within your team and with any teams from other
184 disciplines [12].

185 Preoxygenation should be performed for 3-5 minutes [32, 17, 38] with a good sealed mask
186 and rapid sequence induction [12]. There is evidence of decreased desaturation rates with
187 low-flow apneic oxygenation (oxygen 1-5 l/min) provided by a conventional nasal cannula
188 during intubation [39, 40]. Although strong evidence is lacking, it seems unlikely that such
189 procedures produce a relevant amount of aerosol [12]. There is a strong consensus that
190 video laryngoscopy (VL) should be preferred [13, 29, 12] as it offers some distance between
191 the healthcare professional and a potentially extremely high virus load in comparison with
192 direct laryngoscopy (DL)[31]. In patients with a difficult airway, VL improves glottic view,
193 reduces airway trauma [41] and can increase first-pass intubation success[42]. In general,

194 the use of a standard blade is recommended, with a hyperangulated blade as a backup. The
195 choice of equipment should always be adapted to the skill and clinical judgement of the care
196 provider performing airway management [13]. In case of the unavailability of VL or
197 unfamiliarity with this tool, well-known standard procedures should be performed instead of
198 experiments with unfamiliar ones [33, 12].

199 Practicing rapid sequence induction (RSI) in patients with very limited pulmonary function
200 and almost nonexistent respiratory reserve is already challenging. In this context, induction
201 medications with rapid effect and a deep neuromuscular block to suppress coughing should
202 be mandatory. The recommendations for neuromuscular blocking agents (NMBAs) favor
203 rocuronium [12] in a dosage between 1.2 mg/kg and 1.5 mg/kg ideal body weight (IBW)[13]
204 and sugammadex available in case of an unexpected difficult airway[43]. If succinylcholine is
205 used as the NMBA, the recommended dosage is 1 mg/kg [43] up to 1.5 mg/kg total body
206 weight (TBW)[12, 13]. An international consensus on how to perform rapid sequence
207 induction is still lacking [44].

208 Induction of anesthesia for intubation exposes the patient to an increased risk of
209 hemodynamic instability or even cardiac arrest. In patients susceptible to significant
210 hemodynamic fluctuations, consider ketamine at a dosage of 1-2 mg/kg for a
211 hemodynamically stable induction and have vasopressors ready to use during the intubation
212 phase [12]. An easy way to administer norepinephrine in such situations is 10 mcg/ml as a
213 push dose.

214

215 A common concern of healthcare workers which should be taken into consideration is that
216 massive exposure to COVID-19 can occur during intubation. A recent study showed that
217 tracheal intubation—including facemask ventilation—produced rather low quantities of
218 aerosolized particles compared to extubation and much less than a coughing patient[45].
219 Therefore, there is strong agreement that coughing should be avoided whenever possible.
220 This underlines the importance of a deep neuromuscular block during airway procedures.

221 After intubation the cuff should be inflated immediately and a viral filter should be connected
222 at the end of the tube before starting positive pressure ventilation [13, 12, 29]. Expiratory
223 capnography has to be monitored continuously [12]. Auscultation of the chest can be
224 challenging when wearing a full PPE, and provides a potential risk of contamination;
225 therefore it may not be feasible. If correct placement of the tube needs to be verified, a chest
226 radiograph should be considered after potential central lines or catheters have been installed
227 [13]. Alternatively, point-of-care ultrasound (POCUS)[46] can be used to assist in determining
228 endotracheal tube depth and to rule out a pneumothorax, if needed [29].

229

230 ***Bag–mask ventilation***

231 Nowadays, careful bag-mask ventilation is acceptable while performing a rapid sequence
232 induction [47]. Nevertheless, as a recognized aerosol-generating procedure it should be
233 avoided whenever feasible in suspected or confirmed COVID-19 patients, unless it is a
234 rescue maneuver to treat an unexpected difficult airway [48, 13, 29].

235 These patients present with a higher risk of hypoxia, and mask ventilation should be
236 considered in some cases [10]. If indicated and applied, sealing of the mask is crucial, and a
237 filter needs to be placed directly after the mask to minimize the dispersion of aerosols. To
238 achieve a good seal, the two-handed V-E grip[49] can be used with a two-person-technique.
239 Oropharyngeal airways such as Guedel's or Wendel's may be used to ensure an open
240 airway [50], and minimal positive pressure and low oxygen flow should be applied.
241 Monitoring the bag-mask ventilation with continuous wave capnography is important in order
242 to detect possible leaks [51]. Some authors suggest placing two wet tissues between face
243 and mask to achieve a better seal [43], but there is no evidence supporting this technique. If
244 it is not possible to achieve a good seal with the mask, and careful ventilation seems
245 unattainable, a supraglottic airway device (SAD) should be inserted. This is considered to be
246 safer and produces less aerosol [13].

247

248 Airway manipulation

249 If disconnection of the tracheal tube of an intubated patient is necessary, proper preparation
250 helps to minimize the amount of aerosol generated and the amount of time it takes to
251 disperse. To help avoid a coughing patient, consider deep sedation and profound muscle
252 relaxation before the procedure starts. If no direct access to the airway is necessary, the
253 tracheal tube should be clamped between the filter and the patient after the patient has
254 inhaled, in order to maintain the positive end expiratory pressure (PEEP) generated and
255 therefore to avoid atelectasis during this maneuver [12]. Open suctioning, bronchoscopy, and
256 disconnection of the ventilator circuit should be avoided unless necessary, since these can
257 generate aerosols. If possible, a closed suctioning system should be installed [52, 43].

258

259 Extubation

260 Extubation is considered a high-risk AGP due to the high likelihood of coughing and possible
261 agitation while the endotracheal tube is being removed. Extubation produces up to 15 times
262 more aerosols than intubation[45]. A number of techniques have been developed to reduce
263 aerosol production and droplet spread during extubation. Physical protection—such as
264 consequent use of PPE and early application of a surgical mask to the patient's face—seems
265 to best protect staff from being contaminated [53, 54, 12]. The number of healthcare workers
266 involved should be reduced to the minimum. Extubation under deep sedation is not
267 recommended due to possible absence of spontaneous ventilation, prolonged time with an
268 unsecured airway, increased risk of aspiration and therefore increased risk of requiring bag-
269 mask ventilation and re-intubation. Medications suppressing coughing, such as opioids,
270 lidocaine or dexmedetomidine, may be considered as preventive measures [55] [12, 13].

271

272 Non-invasive ventilation

273 Indications for non-invasive ventilation (NIV) in patients with acute respiratory distress
274 syndrome (ARDS) or COVID-19 are beyond the scope of this review. In general, NIV and
275 humidified application of aerosolized (nebulized) medications should be avoided [13, 29] in
276 aerosol-transmitted viral diseases, to protect healthcare workers until evidence is available
277 from randomized controlled trials. A review article published in 2014 in the context of
278 influenza A H1N1 reported transmission to staff caring for patients treated with NIV in one
279 study out of 22 [56]. A recently published review article reports that the risk of transmission of
280 COVID-19 to healthcare workers may be increased [57]. However, if NIV is used, it is
281 recommended that the treating practitioners wear full PPE and that isolated areas be used to
282 protect the staff and other patients [13, 29].

283

284 ***Supraglottic airway device***

285 It still remains unclear if and how much aerosol is generated by inserting—and removing—a
286 SAD[33]. If a SAD is leaking or patients are coughing at its removal, it is very likely that
287 aerosols will be generated. In the management of an airway due to pulmonary exacerbation
288 in symptomatic COVID-19 patients who will be intubated anyway for long-term ventilation,
289 SAD might play a role as a rescue tool in an unanticipated difficult airway. If one is using
290 SAD for general anesthesia, the provider should keep the risk of the AGP in mind. The leak
291 may be smaller with a spontaneously breathing patient, but if the anesthesia is too “light”
292 there may be an increased risk of coughing [33]. A SAD of the second generation is believed
293 to produce less aerosol than bag-mask ventilation [13].

294

295 ***High-flow nasal cannula***

296 The use of high-flow nasal cannulas (HFNCs) is controversial. A retrospective analysis by
297 Patel et al. [58] postulated that they reduce the incidence of intubation and lead to better
298 outcomes in the case of severe COVID-19 infections. Some earlier randomized controlled

299 trials have shown benefits of high-flow nasal cannula therapy in the context of acute
300 respiratory failure— caused by pneumonia and without hypercapnia—compared to
301 conventional oxygen therapy to prevent NIV and invasive mechanical ventilation [59, 60].
302 Nevertheless, in some cases necessary intubation may be delayed [12]. These findings have
303 not been decisively demonstrated [61-63].

304 Although the use of HFNCs is highly suspected of generating aerosols, the amount still
305 remains unclear, and is thought to be smaller with newer models [12]. As HFNCs are on the
306 list of aerosol-generating procedures[64], there might be an increased risk of significant
307 aerosol exposure for healthcare workers, even if there is low evidence. Instead of completely
308 banning the cannulas, however, hospitals should evaluate their risks and benefits [13, 33,
309 65]. There are differences between using HFNCs to avoid desaturation during airway
310 instrumentation or with the intention of delaying or preventing an intubation that could lead
311 patients to a long period of mechanical ventilation.

312 If the use of HFNCs is considered, it should be subject to the same safety precautions as
313 NIV. This means that the healthcare workers providing the treatment need to wear PPE, and
314 treatment should only be provided in areas with isolation of airborne particles [13]. In
315 addition, during a pandemic, there is a need to conserve resources as much as possible. In
316 fact, even oxygen supply may be scarce [12], and the use of HFNCs may contribute to
317 depleted reserves.

318

319 ***Emergency front-of-neck airway (eFONA) and tracheostomy***

320 In “can't intubate, can't oxygenate” (CICO) situations, a surgical emergency front-of-neck
321 airway (eFONA) created with scalpel and bougie may be preferred over a needle
322 technique[12]. Attempts to oxygenate via bag–mask ventilation during the procedure should
323 be avoided to minimize the risk of generating aerosols [13, 12].

324 Tracheostomy is common for patients who need long-term ventilation, but it is considered a
325 high aerosol-generating procedure [66]. Even though there is evidence of better patient
326 outcomes in early compared to late tracheostomy [67], those findings are not specific to
327 COVID-19 patients, and the best time to perform a tracheostomy in these patients remains
328 controversial. One taskforce recommended that extended endotracheal intubation be
329 considered in order to protect healthcare workers[68], but a recent cohort study showed that
330 tracheostomy can be a safe procedure if performed by an experienced team wearing PPE
331 [69]. Deep neuromuscular blockade is recommended to prevent the patients from coughing
332 while tracheostomy is performed[70].

333

334 ***Awake tracheal intubation and expected difficult airway***

335 Awake tracheal intubation (ATI) is a procedure performed on patients with expected or
336 known difficult airways [71]. It is usually performed with a flexible endoscope, but can also be
337 performed with video laryngoscopes [72] or rigid optics such as the C-MAC VS (Karl Storz,
338 Tübingen, Germany). ATI is an aerosol-generating procedure, which is performed while in
339 close proximity to the spontaneously breathing patient.

340 Due to reductions in elective surgeries in a pandemic situation, the clinical load of such
341 cases —more likely to be found in ENT surgery—will decrease [33]. Nevertheless, there will
342 be intubations for patients with an expected or known difficult airway, for those needing
343 emergency surgery, and for those needing intubation due to COVID-19 infection.

344 ATIs should only be performed if there is a strong indication and no other alternative is
345 deemed safe. Coughing should be suppressed as much as possible. This can be achieved
346 with topical anesthesia and short-acting intravenous opioids (e.g., remifentanyl). Disposable
347 devices should be used if available, and the operator should be experienced in this technique
348 [17, 32]. Intratracheal application of local anesthetics should be avoided due to the cough
349 stimulus [33]. Some authors recommend primarily nasal intubation via an endoscopic mask,

350 and only switching to oral intubation in case of failure [17]. The authors of this review believe
351 that in a complex and difficult airway management situation, using the techniques and
352 materials one is most familiar with will be the most successful approach.

353 The choice of a small endotracheal tube reduces the cough generated during insertion [32].
354 Unfortunately, a small tube causes more resistance and therefore generates increased
355 airway pressures. Especially for long-term mechanical ventilation of a COVID-19 patient with
356 ARDS, it seems to be more reasonable to have a rather large endotracheal tube. Some
357 authors suggest using an endotracheal tube size 7 or 8 for women and a size 8 or 9 for men
358 [12]. Both indication and possible duration—a short emergency operation vs. long-term
359 intubation—should be considered in advance. In the event of ATI failure, the ENT surgeon
360 should decide early on whether to perform a tracheotomy [17]. Although the application of
361 HFNCs is controversial (see above), ATI could possibly be a good indication. HFNCs may
362 allow deeper sedation and therefore lead to less irritation and coughing.

363

364 **Personal protective equipment (PPE)**

365 In times of crisis, shortages of healthcare personnel are a central and crucial problem.
366 Although staffing adjustments that focus on epidemiological factors may reduce this shortage
367 [73], keeping healthcare workers healthy and safe must remain a central concern.

368 Availability and use of personal protective equipment is key to protecting the healthcare
369 workforce. In general, staff members who are involved in airway management of a patient
370 with suspected or proven COVID-19 infection should follow available recommendations. This
371 includes correct hand disinfection and single-use airborne PPE, consisting of a mask
372 (whenever possible N95, KN95 or filtering face piece class 2 (FFP2) or higher), protective
373 goggles, a hat, a gown, and gloves (optionally 2 pairs); a practical overview was recently
374 provided by Cook et al.[74]. This protective gear should be worn for all airway-related

375 procedures, as well as while caring for COVID-19 patients [12, 13, 29] and especially during
376 AGP.

377 Of equal importance is the procedure to be followed when doffing the PPE, as errors are
378 associated with a potential risk of infection. Simulation of donning and doffing can improve
379 safety in handling [75, 12], as can a “buddy system” with checklists followed by a specialized
380 supervisor for donning and doffing PPE [12, 13]. At the very least, supervision of the removal
381 process should be introduced, and proper hand hygiene after removal is mandatory [29].
382 Additionally, the environment should be decontaminated for at least 20 minutes [12] after
383 AGP or depending on the air exchange capacity of the room. The virus can be detectable for
384 up to 72 hours, depending on the surface material [20]

385

386 **Pediatric airway management**

387 Children who test positive are often asymptomatic (18-22%) or present with mild symptoms
388 like fever and general respiratory symptoms [76, 77]. Hospitalizations and intensive care unit
389 admissions are rare [78].

390 We will briefly discuss the challenges to be considered when anesthetizing children in the
391 specific context of the COVID-19 pandemic. As in cases involving adults, protection of
392 healthcare workers should be prioritized, with no exceptions, and PPE must be worn for all
393 risky procedures.

394 There are some precautions to be taken preoperatively that could differ from the daily routine
395 of a pediatric anesthesiologist. Because inhalational induction increases the risk of exposure
396 to respiratory droplets and aerosols, intravenous induction should be the first choice in the
397 case of COVID-positive children [79]. To reduce the child’s anxiety as well as crying during
398 intravenous (IV) line placement, the administration of premedication combined with patches
399 for topical anesthesia of the puncture site is highly recommended. Because of the risk of
400 sneezing or coughing, nasal premedication should be avoided and oral or rectal

401 premedication should be preferred [16]. To minimize the potential risk of transmission of
402 SARS-CoV-2 to staff and to preserve PPE, during induction the presence of parents who
403 have close contact with the child (and are considered potentially infected) is not
404 recommended. Nevertheless, if the parents are asymptomatic and wearing correct PPE, this
405 may be adapted to the specific situation, because a calm child with parents is safer than an
406 agitated, crying and coughing child[16]. The parents should leave before any aerosol-
407 generating procedure starts.

408 As in the case of performing anesthesia for an adult patient, endotracheal intubation (sealing
409 of the airway) with a VL using a modified RSI should be performed, and low-flow nasal
410 oxygen with a conventional cannula for apneic oxygenation (e.g., 0.2 l/kg/min) should be
411 considered during the procedure to prolong the time until desaturation and to increase safety
412 [80]. For the same reason, due to the risk of extreme desaturation during airway
413 management, the classical RSI technique should not be performed. In this case, the mask
414 should be kept sealed to minimize aerosolization. To minimize the duration of intubation and
415 the number of attempts needed, the use of apneic oxygenation should be considered and the
416 most experienced person should perform the laryngoscopy. In addition, a cuffed
417 endotracheal tube should be used whenever the situation allows and the weight of the child
418 is over 3 kg. In a situation where the placement of an IV catheter is difficult and the child is
419 agitated and combative, exposure to respiratory droplets may increase considerably. In this
420 case, we recommend an ultrasound-guided venous puncture to facilitate venous access or
421 an inhalational induction with the precaution of keeping the mask sealed and using the
422 lowest possible flow rate [16]. HFNC can be useful in elective endoscopic airway surgery
423 procedures [81], but should be carefully evaluated as it is potentially an AGP. Anesthesia can
424 be maintained following institutional routine. Emergence from anesthesia in deep sedation is
425 indicated to minimize dispersion of droplets due to coughing [82]; children with confirmed or
426 potential COVID-19 should be extubated before leaving the operating room to avoid a stay in
427 the pediatric post-anesthesia care unit [16]. Children who require a postoperative stay in the
428 pediatric intensive care unit (PICU) should be extubated in the PICU.

429 In situations where a difficult airway is encountered, the principles already mentioned for the
430 airway should be applied. In addition, task fixation and prolonged attempts at intubation
431 should be avoided in order to avoid increased aerosolization of the virus. In children with
432 COVID-19 the first choice for intubation is videolaryngoscopy. In case of failure, an early
433 change to more advanced intubation techniques such as intubation via fibroscopy through
434 SGA is desirable. In addition, even for children with COVID-19, anatomical and functional
435 airway obstructions must be recognized and treated in order to avoid CICO situations[83,
436 16]. In very rare cases these situations can degenerate and a surgical airway will be
437 required[84].

438 eFONA in children is very rare and is associated with poor outcomes[85]. As in adults, a
439 surgical eFONA with scalpel is recommended: in the absence of an ENT surgeon, rapid
440 sequence surgical tracheostomy should be preferred to access the trachea for emergencies
441 in children under 8 years of age [86]. For children older than 8 years, we prefer a scalpel
442 bougie technique over a needle technique [87, 88, 86]. There are no other special
443 recommendations for changes in technique in children with aerosol-transmitted viral
444 disease.

445

446 **Developing new protection tools**

447 The risk of exposing healthcare workers to the highly contagious SARS-CoV-2 virus and the
448 threat of shortages of PPE pushed clinicians to develop new types of physical protection
449 devices to use during aerosol-generating procedures. Boxes, drapes, sheets and shields
450 made of a variety of materials (mostly plastic) were tested and used [33-37], but there was
451 no validation and no randomized trials of these devices[89]. Systematic analyses do not
452 recommend the use of such tools, not only due to a lack of evidence, but also because most
453 of them have proven to be ineffective, impeding a rapid intubation through delays and
454 making airway management unsafe [90]. In fact, the concentration of aerosol in the
455 containment box could even be higher than without it, exposing the airway manager to a

456 greater risk of infection [91]. Reviews published until now therefore suggest avoiding the use
457 of protective aids and focusing more on correct handling of PPE and proper ventilation,
458 which workers are already familiar with in their daily practice [91-93].

459

460 **Airway guidelines and recommendations for the COVID-19 pandemic**

461 Many professional societies involved in airway management promptly published consensus
462 papers and guidelines based on expert opinions and experience from the field. The
463 algorithms often focus on PPE and avoiding or at least reducing the duration of aerosol-
464 generating procedures. Most of these guidelines are modifications of already existing
465 evidence-based guidelines on airway or difficult airway management, and there remains a
466 dearth of high-quality scientific evidence regarding the recommendations. Regardless,
467 protecting healthcare workers is a key goal. A summary of existing airway management
468 recommendations is presented in Table 2.

Table 2. Airway management recommendations and consensus

<i>First author</i> <i>Date</i> <i>Country or Society</i>	PPE	Intubation	Extubation	NIV & HFNC	eFONA & Tracheostomy	Bag Mask Ventilation & SAD	Medication	Key points
<i>Brewster et al.</i> <i>01.06.20</i> <i>Australian and New Zealand College of Anesthetists/Safe Airway Society</i>	PPE: minimum: impervious gown, theater hat, N95 mask, face shield, eye protection, double gloves "Buddy system": guided by specially trained and designated staff member acting as "spotter"	RSI, Indirect VL (video screen) maximizing distance between airway and operator; Macintosh or hyperangulated blade; place the tube to correct depth; inflation of the cuff before positive pressure ventilation; viral filter to end of the tube; cuff pressure monitoring	Face mask ready; 2 staff members with PPE (same as intubation); don't encourage the patient to cough; minimize coughing by the use of intravenous opioids, lidocaine or dexmedetomidine. Consider plastic sheets in case of coughing, place oxygen mask immediately after; oral suctioning	No evidence Should be assumed that NIV & HFNC are aerosol-generating procedures; airborne isolation rooms; protective PPE (including N95/FFP2 masks)	eFONA (CICO): Scalpel-bougie technique (to minimize the risk of high pressure oxygen insufflation via a small-bore cannula). No attempts to deliver oxygen from above during procedure (avoid aerosolization) Tracheostomy: N/A	Avoid BMV. If needed: use a vice (V-E) grip; minimize ventilation pressure through ramping and/or early use of an oropharyngeal airway with low gas flows; filter between mask and bag SAD: likely to protect better than BMV	Initial NMB: rocuronium (>1.5 mg/kg IBW) or suxamethonium (1.5 mg/kg TBW). Generous dosing for rapid onset and minimizes the risk of coughing. To avoid coughing during extubation: intravenous opioids, lidocaine or dexmedetomidine	Follow existing guidelines; modify them for COVID-19; early intubation; significant institutional preparation; principles for airway management should be same for all COVID-19 patients; safe, simple, familiar, reliable and robust practices should be adopted
<i>Wax et al.</i>	Fluid-resistant gown, gloves, eye protection, full face	VL; RSI; only essential team members; airborne	n/a	No evidence HFNC limited to	n/a	Bag-mask ventilation can generate aerosols (avoid when	Use of TIVA for anesthesia, avoid gas	n/a

12.02.2020	shield, fit-tested N95 mask, hair covers or hoods; longer sleeved gloves; consider powered air purifying respirator (PAPR); scrub suits or full coveralls under PPE; hand hygiene after PPE use; remove PPE under supervision of an infection control coach using checklist	isolation room; end tidalCO ₂ ; all exhaled gas from the ventilator should be filtered Rule out Pneumothorax in sudden respiratory deterioration (Ultrasound on bedside)		patients in appropriate airborne isolation. Avoid NIV (CPAP/BiPAP) use outside of appropriate airborne/droplet isolation. Avoid nebulization of medications		possible); filter between mask and bag		
------------	---	--	--	--	--	--	--	--

<p><i>Cook et al.</i></p> <p>17.03.2020</p> <p><i>UK / Difficult Airway Society, Association of Anesthetists, the Intensive Care Society, Faculty of Intensive Care Medicine, Royal College of Anesthetists</i></p>	<p>PPE; mask (FFP3), simple to remove; avoid complex systems; cover the whole upper body; dispose, appropriately immediately after "doffing". "Buddy system" (observer); checklists; double-gloving for endotracheal intubation; use anti-fog for goggles/eyewear; training and practicing PPE use; negative pressure rooms with good rates of air exchange (> 12 times/h)</p>	<p>Specific intubation team (not part of the risk groups); most experienced airway manager; simulation; single-use equipment; rather early than late intubation; limit team to 2 persons performing intubation inside + 1 runner (outside), prepare and communicate before intubation; airway strategy (primary plan and the rescue plans) avoid AGPs; good preoxygenation with sealed face mask (3-5 min), RSI, VL; intubation checklists; dedicated intubation trolley, aim to achieve first attempt success; no test of new techniques</p>	<p>Delayed extubation; minimize coughing; appropriate physiotherapy, tracheal and oral suction as normal before extubation; prepare for mask or low flow nasal oxygen delivery before extubation; after extubation, place a facemask; SAD may be considered as a bridge to extubation to minimize coughing; a second procedure and the possibility of airway difficulty, unlikely to be a first-line procedure; use of an airway exchange catheter is relatively contra-indicated; use drugs</p>	<p>No evidence</p> <p>HFNC recommendation debated: delays intubation, needs much O2 (empty tanks)</p>	<p>Scalpel cricothyroidotomy in CICO situations wearing full PPE; Closed suction</p>	<p>BMV: 2-handed V-E grip</p> <p>SGA: second generation as rescue airway also to improve seal</p>	<p>Intubation: Consider Ketamine 1-2 mg/kg; deep neuromuscular relaxation with rocuronium 1.2mg/kg IBW or succinylcholine 1.5mg/kg TBW</p> <p>Extubation: dexmedetomidine, lidocaine and opioids</p>	<p>Safe, accurate and swift airway management</p>
---	---	---	--	---	--	---	--	---

			to suppress coughing					
<p><i>Sorbello et al.</i></p> <p>27.03.2020</p> <p><i>Società Italiana di Anestesia Analgesia Rianimazione e Terapia Intensiva</i></p> <p><i>European Airway Management Society</i></p>	<p>PPE: PAPR, with helmet, protective total body suite, double gloves;</p> <p>If no PAPR available: goggles/face shield, FFP3/2 or N95 mask, waterproof gown, overshoes;</p> <p>Dedicated donning/doffing area</p>	<p>Preoxygenation with or without CPAP and PEEP;</p> <p>RSI technique;</p> <p>Nasal O2 1-3 L/min during apnea;</p> <p>VL (with separate screen) + introducer;</p> <p>second generation SDA if failed intubation;</p> <p>Early cricothyroidotomy if</p>	n/a	NIV, HFNC should not delay an early elective intubation	Cricothyrotomy in CICO situations	<p>Avoid BMV</p> <p>SADs only as rescue, 2nd generation to intubate through</p>	<p>Rocuronium 1.2 mg/kg IBW or Suxamethonium 1 mg/kg TBW</p>	<p>Full airborne protection for every phase of airway management;</p> <p>Training, planning, anticipation; Maximize first-pass attempt</p>

		CICO; ATI only if mandatory						
<i>Patwa et al.</i> 23.05.2020 <i>All India Difficult Airway Association</i>	Hand hygiene Full PPE: waterproof gown, long shoe covers, a cap, goggles, a fit-tested N95 mask, double layer of gloves, and a head hood or full face shield; Correct donning and supervised doffing; covering the patient with a plastic sheet or intubation box.	Preoxygenation with a 2-hands 2 persons technique; continuous capnography (leakage monitoring); Low-flow O ₂ (< 5 L/min) nasal during apnea; RSI; most experienced clinician; Ventilation after cuffing Closed suction system; Consider ATI only in high selected cases with anticipated difficult airway	Same protection as for intubation; Suction only if necessary; Prevent coughing, agitation and emesis; Avoid any manipulation; Defer extubation if there are concerns	NIV and HFNC not recommended	Avoid cannula or needle cricothyrotomy with jet ventilation Surgical cricothyrotomy in case of complete failure of ventilation	Avoid BMV Consider surgery with SAD if safe, or awakening the patient with SAD in place	Suxamethonium or rocuronium for anesthesia induction	Modified AIDAA algorithm for airway management during COVID-19 pandemic
<i>Al Harbi et al.</i> 17.04.2020 <i>Saudi Anesthesia Society</i>	Hand hygiene Disposable N-95 masks, goggles, footwear, water-proof gowns and gloves (consider double glove technique); PAPR for high-risk	Intubation by the most experienced clinician; Standard ASA monitoring; VL (single-use blade); Ventilation after inflation of the cuff; Lowest gas flow	n/a	n/a	n/a	n/a	n/a	Adherence and correct usage of PPE; Ad interim Guideline (COVID pandemic still outbreaking)

	AGP.	acceptable						
<i>Matava et al.</i> 13.04.2020 <i>Society for Pediatric Anesthesia's Pediatric Difficult Intubation Collaborative / Canadian Pediatric Anesthesia Society</i>	Not specified PPEs; teams reduced to the minimum to preserve PPE, importance of correctly donning/doffing PPEs (with coaches). High risk clinicians should not be involved.	RSI, VL; Parents may be present until airway management.	Deep sedation (see medication) to avoid coughing Closed suction system Extubation in the OR	HFNC to be avoided if possible	n/a	Avoid BMV and Mask Induction Consider 2nd generation SADs (good seal, low airway pressures).	Premedication not nasal, oral or rectal should be preferred Consider dexmedetomidine, TIVA for extubation	Protection of healthcare workers is priority; adapt guidelines to institutional protocols
<i>Chen et al.</i> 29.07.2020 <i>Chinese Society of Anesthesiology / Chinese Association of Anesthesiologists</i>	Hospital scrubs inside and protective coveralls outside; medical protective mask, disposable surgical cap, goggles/face shield; wear disposable medical latex gloves and boot covers.	Airway team (experienced), patient's mouth covered with two wet gauze during preoxygenation, RSI, VL or Bronchoscope/Fibers cope (airway manager is familiar and brings distance to the airway), filter between tube, no auscultation	n/a	If patient under HFNC or NIV before intubation use caution for aerosol and droplets	n/a	Two wet gauzes, rather avoid BMV	Consider midazolam 2-5mg, etomidate 10-20mg, propofol (if stable), succinylcholine 1mg/kg, if rocuronium have sugammadex nearby for a CICO	Protection of healthcare workers

		Respirator only for COVID-19 patients after use even with filter or need to be disinfected; closed suction system						
<p>Abbreviations: AGP: aerosol-generating procedure; ASA: American Society of Anesthesiologists; ATI: awake tracheal intubation; CICO: “can't intubate, can't oxygenate”; CPAP: continuous positive airway pressure; eFONA: emergency front-of-neck airway; HFNC: High-flow nasal cannula; IBW: ideal body weight; NIV: non-invasive ventilation; NMB: neuromuscular blockade; OR: operating room; PAPR: powered air-purifying respirator; PEEP: positive end expiratory pressure; PPE: personal protective equipment; RSI: rapid sequence induction; SGA: supraglottic airway device; TBW: total body weight; TIVA: total intravenous anesthesia; VL: video laryngoscopy; WHO: World Health Organization.</p>								

Summary

In order to further improve safety during airway management, it is important to clearly define an aerosol-generating procedure (AGP), how much aerosol is produced during one, how performing an AGP affects healthcare workers, and how much workers are put at risk of infection. A range of measures are under consideration for use in treating COVID-19 patients, among them proper hand hygiene and correct donning and doffing of PPE; simulation training for airway management involving PPE; the use of highly experienced “Airway teams” in which the person with the most experience performs the procedure; preparation for situations with an unexpected difficult airway; and suppression of coughing in patients undergoing airway-related procedures.

It is important to protect both high risk patients and healthcare workers and not to experiment with new techniques and tools that could lead to increased exposures. As was true before the onset of COVID-19, any AGP without strong indications should not be performed.

The care of COVID-19 patients is challenging due to many factors, not the least the reduced capacity of beds, ventilators, and personnel. Institutional requirements and resources need to be evaluated before the airway management starts. Triage adapted to the individual institution and situation is useful and should be discussed in advance, especially in times of low capacity. Overall, preparation and planning are even more essential.

In this review we discuss the management of the airway and the precautions that need to be taken in detail, but just as important is the realization that our knowledge will evolve over time as we learn more about the COVID-19 virus. The pandemic is ongoing, and we will be confronted with it for a while.

Practice Points

- There is limited evidence for the impact of aerosol-generating procedures and their influence on infection in healthcare workers;

- Airway preparation and management should be performed by experienced staff who do not belong to a high risk group;
- To avoid or suppress coughing in COVID patients, use a deep neuromuscular block and rapid-sequence induction for intubation and airway management;
- Hand disinfection and adequate PPE with a “buddy system” are essential for the protection of healthcare staff;
- Adapt and use algorithms and equipment that healthcare workers are already familiar with;
- Use simulation training—especially for airway management and usage of PPE—to improve adherence and safety.

Research Agenda

- Define an aerosol-generating procedure (AGP),
- Better estimate how much aerosol is produced during an AGP,
- How performing an AGP affects healthcare workers,
- How much workers are put at risk of infection during AGP procedures,
- Most effective ways to protect healthcare workers

Conflicts of Interest

This research did not receive specific support from funding agencies in the public, commercial, or not-for-profit sectors.

Richard D. Urman reports research funding/fees from Merck, Medtronic/Covidien, AcelRx, Takeda, Pfizer. All other authors declare no conflicts of interest.

Acknowledgements

We thank Jeannie Wurz for her careful proofreading of this manuscript.

References

1. WHO Coronavirus Disease (COVID-19) Dashboard. Geneva: World Health Organization, 2020. Available online: <https://covid19.who.int/>. Last viewed 11.24.2020. 2020.
2. Lotfi M, Hamblin MR, Rezaei N. COVID-19: Transmission, prevention, and potential therapeutic opportunities. *Clin Chim Acta*. 2020;508:254-66. doi:10.1016/j.cca.2020.05.044.
3. **Fennelly KP. Particle sizes of infectious aerosols: implications for infection control. *Lancet Respir Med*. 2020;8(9):914-24. doi:10.1016/s2213-2600(20)30323-4.**
4. Tellier R. Review of aerosol transmission of influenza A virus. *Emerg Infect Dis*. 2006;12(11):1657-62. doi:10.3201/eid1211.060426.
5. Judson SD, Munster VJ. Nosocomial transmission of emerging viruses via aerosol-generating medical procedures. *Viruses*. 2019;11(10). doi:10.3390/v11100940.
6. Fennelly KP. Particle sizes of infectious aerosols: implications for infection control. *Lancet Respir Med*. 2020;8(9):914-24. doi:10.1016/s2213-2600(20)30323-4.
7. Scales DC, Green K, Chan AK, Poutanen SM, Foster D, Nowak K et al. Illness in intensive care staff after brief exposure to severe acute respiratory syndrome. *Emerg Infect Dis*. 2003;9(10):1205-10. doi:10.3201/eid0910.030525.
8. **Verbeek JH, Rajamaki B, Ijaz S, Tikka C, Ruotsalainen JH, Edmond MB et al. Personal protective equipment for preventing highly infectious diseases due to exposure to contaminated body fluids in healthcare staff. *The Cochrane Database Syst Rev*. 2019;7(7):Cd011621. doi:10.1002/14651858.CD011621.pub3.**
9. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020;382(18):1708-20. doi:10.1056/NEJMoa2002032.
10. Tang LY, Wang J. Anesthesia and COVID-19: What we should know and what we should do. *Semin Cardiothorac Vasc Anesth*. 2020;24(2):127-37. doi:10.1177/1089253220921590.
11. Bulut C, Kato Y. Epidemiology of COVID-19. *Turk J Med Sci*. 2020;50(SI-1):563-70. doi:10.3906/sag-2004-172.
12. **Cook TM, El-Boghdadly K, McGuire B, McNarry AF, Patel A, Higgs A. Consensus guidelines for managing the airway in patients with COVID-19: Guidelines from the Difficult Airway Society, the Association of Anaesthetists, the Intensive Care Society, the Faculty of Intensive Care Medicine and the Royal College of Anaesthetists. *Anaesthesia*. 2020;75(6):785-99. doi:10.1111/anae.15054.**
13. Brewster DJ, Chrimes N, Do TB, Fraser K, Groombridge CJ, Higgs A et al. Consensus statement: Safe Airway Society principles of airway management and tracheal intubation specific to the COVID-19 adult patient group. *Medical J Australia*. 2020;212(10):472-81. doi:10.5694/mja2.50598.
14. Al Harbi MK, Abdulmomen A, Qurashi FA, Kattan M, Ahmad AE, Eldawlatly A. Airway management and COVID-19 patient -Saudi Anesthesia Society guidelines. *Saudi J Anaesth*. 2020;14(3):387-9. doi:10.4103/sja.SJA_265_20.
15. Pritchett MA, Oberg CL, Belanger A, De Cardenas J, Cheng G, Nacheli GC et al. Society for Advanced Bronchoscopy Consensus Statement and Guidelines for bronchoscopy and airway management amid the COVID-19 pandemic. *J Thorac Dis*. 2020;12(5):1781-98. doi:10.21037/jtd.2020.04.32.
16. **Matava CT, Kovatsis PG, Lee JK, Castro P, Denning S, Yu J et al. Pediatric airway management in COVID-19 patients: Consensus guidelines from the Society for Pediatric Anesthesia's Pediatric Difficult Intubation Collaborative and the Canadian Pediatric Anesthesia Society. *Anesth Analg*. 2020;131(1):61-73. doi:10.1213/ane.0000000000004872.**
17. Zuo MZ, Huang YG, Ma WH, Xue ZG, Zhang JQ, Gong YH et al. Expert recommendations for tracheal intubation in critically ill patients with novel Coronavirus disease 2019. *Chin Med Sci J*. 2020;35(2):105-9. doi:10.24920/003724.
18. Velly L, Gayat E, Quintard H, Weiss E, De Jong A, Cuillon P et al. Guidelines: anaesthesia in the context of COVID-19 pandemic. *Anaesth Crit Care Pain Med*. 2020;39(3):395-415. doi:10.1016/j.accpm.2020.05.012.

19. Takhar A, Walker A, Tricklebank S, Wyncoll D, Hart N, Jacob T et al. Recommendation of a practical guideline for safe tracheostomy during the COVID-19 pandemic. *Eur Arch Otorhinolaryngol* 2020;277(8):2173-84. doi:10.1007/s00405-020-05993-x.
20. Shamsoddin E. A COVID-19 pandemic guideline in evidence-based medicine. *Evidence-based Dentistry*. 2020;21(2):71-3. doi:10.1038/s41432-020-0105-7.
21. Sullivan EH, Gibson LE, Berra L, Chang MG, Bittner EA. In-hospital airway management of COVID-19 patients. *Critical Care*. 2020;24(1):292. doi:10.1186/s13054-020-03018-x.
22. Caputo KM, Byrick R, Chapman MG, Orser BJ, Orser BA. Intubation of SARS patients: infection and perspectives of healthcare workers. *Can J Anaesth*. 2006;53(2):122-9. doi:10.1007/bf03021815.
23. Tran K, Cimon K, Severn M, Pessoa-Silva C, Conly J. Aerosol-generating procedures and risk of transmission of acute respiratory infections: a systematic review. *CADTH Technol Overv*. 2013;3(1):e3201.
24. Yanes-Lane M, Winters N, Fregonese F, Bastos M, Perlman-Arrow S, Campbell JR et al. Proportion of asymptomatic infection among COVID-19 positive persons and their transmission potential: A systematic review and meta-analysis. *PloS one*. 2020;15(11):e0241536. doi:10.1371/journal.pone.0241536.
- 25. El-Boghdady K, Wong DJN, Owen R, Neuman MD, Pocock S, Carlisle JB et al. Risks to healthcare workers following tracheal intubation of patients with COVID-19: a prospective international multicentre cohort study. *Anaesthesia*. 2020;75(11):1437-47. doi:10.1111/anae.15170.**
26. Lee ZX, Ng KT, Ang E, Wang CY, Binti S, II. Effect of perioperative regional anesthesia on cancer recurrence: A meta-analysis of randomized controlled trials. *Int J Surg*. 2020;82:192-9. doi:10.1016/j.ijssu.2020.08.034.
- 27. Cheung JC, Ho LT, Cheng JV, Cham EYK, Lam KN. Staff safety during emergency airway management for COVID-19 in Hong Kong. *Lancet Respir Med*. 2020;8(4):e19. doi:10.1016/s2213-2600(20)30084-9.**
28. Wang W, Xu Y, Gao R, Lu R, Han K, Wu G et al. Detection of SARS-CoV-2 in different types of clinical specimens. *JAMA*. 2020;323(18):1843-4. doi:10.1001/jama.2020.3786.
29. Wax RS, Christian MD. Practical recommendations for critical care and anesthesiology teams caring for novel coronavirus (2019-nCoV) patients. *Can J Anaesth*. 2020;67(5):568-76. doi:10.1007/s12630-020-01591-x.
30. Pimentel MPT, Pimentel CB, Wheeler K, Dehmer E, Vacanti JC, Urman RD. Using a pre-procedure COVID-19 huddle to improve operating room safety. *J Clin Anesth*. 2020;65:109875. doi:10.1016/j.jclinane.2020.109875.
31. Yao W, Wang T, Jiang B, Gao F, Wang L, Zheng H et al. Emergency tracheal intubation in 202 patients with COVID-19 in Wuhan, China: lessons learnt and international expert recommendations. *Br J Anaesth*. 2020;125(1):e28-e37. doi:10.1016/j.bja.2020.03.026.
32. Sorbello M, El-Boghdady K, Di Giacinto I, Cataldo R, Esposito C, Falcetta S et al. The Italian coronavirus disease 2019 outbreak: recommendations from clinical practice. *Anaesthesia*. 2020;75(6):724-32. doi:10.1111/anae.15049.
- 33. Cook TM, McGuire B, Mushambi M, Misra U, Carey C, Lucas N et al. Airway management guidance for the endemic phase of COVID-19. *Anaesthesia*. 2020. doi:10.1111/anae.15253.**
- 34. Herman JA, Urits I, Kaye AD, Urman RD, Viswanath O. COVID-19: Recommendations for regional anesthesia. *J Clin Anesth*. 2020;65:109885. doi:10.1016/j.jclinane.2020.109885.**
35. Schwartz R, Urits I, Kaye AD, Urman RD, Viswanath O. Recommendations for perioperative coronavirus (COVID-19) testing. *Perioper Care Olper Room Manag*. 2020;20:100122. doi:10.1016/j.pccorm.2020.100122.
36. Herman JA, Urits I, Kaye AD, Urman RD, Viswanath O. COVID-19: Anesthesia management recommendations. *J Clin Anesth*. 2020:109840. doi:10.1016/j.jclinane.2020.109840.
37. Herman JA, Urits I, Kaye AD, Urman RD, Viswanath O. COVID-19: Obstetric anesthesia care considerations. *J Clin Anesth*. 2020;65:109860. doi:10.1016/j.jclinane.2020.109860.
38. Kim HJ, Ko JS, Kim TY. Recommendations for anesthesia in patients suspected of COVID-19 Coronavirus infection. *Korean J Anesth*. 2020;73(2):89-91. doi:10.4097/kja.20110.

39. Leone M, Einav S, Chiumello D, Constantin JM, De Robertis E, De Abreu MG et al. Noninvasive respiratory support in the hypoxaemic peri-operative/periprocedural patient: a joint ESA/ESICM guideline. *Intensive Care Med.* 2020;46(4):697-713. doi:10.1007/s00134-020-05948-0.
40. Oliveira JESL, Cabrera D, Barrionuevo P, Johnson RL, Erwin PJ, Murad MH et al. Effectiveness of apneic oxygenation during intubation: A systematic review and meta-analysis. *Ann Emerg Med.* 2017;70(4):483-94.e11. doi:10.1016/j.annemergmed.2017.05.001.
41. Lewis SR, Butler AR, Parker J, Cook TM, Schofield-Robinson OJ, Smith AF. Videolaryngoscopy versus direct laryngoscopy for adult patients requiring tracheal intubation: a Cochrane systematic review. *Br J Anaesth.* 2017;119(3):369-83.
42. De Jong A, Molinari N, Conseil M, Coisel Y, Pouzeratte Y, Belafia F et al. Video laryngoscopy versus direct laryngoscopy for orotracheal intubation in the intensive care unit: a systematic review and meta-analysis. *Intensive Care Med.* 2014;40(5):629-39. doi:10.1007/s00134-014-3236-5.
43. Chen X, Liu Y, Gong Y, Guo X, Zuo M, Li J et al. Perioperative management of patients infected with the novel coronavirus: Recommendation from the Joint Task Force of the Chinese Society of Anesthesiology and the Chinese Association of Anesthesiologists. *Anesthesiology.* 2020;132(6):1307-16. doi:10.1097/aln.0000000000003301.
44. Zdravkovic M, Berger-Estilita J, Sorbello M, Hagberg CA. An international survey about rapid sequence intubation of 10,003 anaesthetists and 16 airway experts. *Anaesthesia.* 2020;75(3):313-22. doi:10.1111/anae.14867.
- 45. Brown J, Gregson FKA, Shrimpton A, Cook TM, Bzdek BR, Reid JP et al. A quantitative evaluation of aerosol generation during tracheal intubation and extubation. *Anaesthesia.* 2020. doi:10.1111/anae.15292.**
46. Meier I, Vogt AP, Meineri M, Kaiser HA, Luedi MM, Braun M. Point-of-care ultrasound in the preoperative setting. *Best Pract Res Clin Anaesthesiol.* 2020;34(2):315-24. doi:10.1016/j.bpa.2020.04.010.
47. Frerk C, Mitchell VS, McNarry AF, Mendonca C, Bhagrath R, Patel A et al. Difficult Airway Society 2015 guidelines for management of unanticipated difficult intubation in adults. *Br J Anaesth.* 2015;115(6):827-48. doi:10.1093/bja/aev371.
48. Cook TM, Woodall N, Harper J, Benger J. Major complications of airway management in the UK: results of the Fourth National Audit Project of the Royal College of Anaesthetists and the Difficult Airway Society. Part 2: intensive care and emergency departments. *Br J Anaesth.* 2011;106(5):632-42. doi:10.1093/bja/aer059.
49. Chrimes N. The Vortex: a universal 'high-acuity implementation tool' for emergency airway management. *Br J Anaesth.* 2016;117 Suppl 1:i20-i7. doi:10.1093/bja/aew175.
50. Otten D, Liao MM, Wolken R, Douglas IS, Mishra R, Kao A et al. Comparison of bag-valve-mask hand-sealing techniques in a simulated model. *Ann Emerg Med.* 2014;63(1):6-12 e3. doi:10.1016/j.annemergmed.2013.07.014.
51. Lim KS, Nielsen JR. Objective description of mask ventilation. *Br J Anaesth.* 2016;117(6):828-9. doi:10.1093/bja/aew368.
52. Patwa A, Shah A, Garg R, Divatia JV, Kundra P, Doctor JR et al. All India difficult airway association (AIDAA) consensus guidelines for airway management in the operating room during the COVID-19 pandemic. *Indian J Anaesth.* 2020;64(Suppl 2):S107-s15. doi:10.4103/ija.IJA_498_20.
53. D'Silva DF, McCulloch TJ, Lim JS, Smith SS, Carayannis D. Extubation of patients with COVID-19. *Br J Anaesth.* 2020;125(1):e192-e5. doi:10.1016/j.bja.2020.03.016.
54. Patino Montoya M, Chitilian HV. Extubation barrier drape to minimise droplet spread. *Br J Anaesth.* 2020;125(1):e195-e6. doi:10.1016/j.bja.2020.03.028.
55. Tung A, Fergusson NA, Ng N, Hu V, Dormuth C, Griesdale DEG. Medications to reduce emergence coughing after general anaesthesia with tracheal intubation: a systematic review and network meta-analysis. *Br J Anaesth.* 2020. doi:10.1016/j.bja.2019.12.041.
56. Esquinas AM, Egbert Pravinkumar S, Scala R, Gay P, Soroksky A, Girault C et al. Noninvasive mechanical ventilation in high-risk pulmonary infections: a clinical review. *Eur Respir Rev.* 2014;23(134):427-38. doi:10.1183/09059180.00009413.

57. Schünemann HJ, Khabsa J, Solo K, Khamis AM, Brignardello-Petersen R, El-Harakeh A et al. Ventilation techniques and risk for transmission of coronavirus disease, including COVID-19: A living systematic review of multiple streams of evidence. *Ann Intern Med.* 2020;173(3):204-16. doi:10.7326/m20-2306.
58. Patel M, Gangemi A, Marron R, Chowdhury J, Yousef I, Zheng M et al. Retrospective analysis of high flow nasal therapy in COVID-19-related moderate-to-severe hypoxaemic respiratory failure. *BMJ Open Respir Res.* 2020;7(1). doi:10.1136/bmjresp-2020-000650.
59. Koga Y, Kaneda K, Fujii N, Tanaka R, Miyauchi T, Fujita M et al. Comparison of high-flow nasal cannula oxygen therapy and non-invasive ventilation as first-line therapy in respiratory failure: a multicenter retrospective study. *Acute Med Surg.* 2020;7(1):e461. doi:10.1002/ams2.461.
60. Leung CCH, Joynt GM, Gomersall CD, Wong WT, Lee A, Ling L et al. Comparison of high-flow nasal cannula versus oxygen face mask for environmental bacterial contamination in critically ill pneumonia patients: a randomized controlled crossover trial. *J Hosp Infect.* 2019;101(1):84-7. doi:10.1016/j.jhin.2018.10.007.
61. Lyons C, Callaghan M. The use of high-flow nasal oxygen in COVID-19. *Anaesthesia.* 2020;75(7):843-7. doi:10.1111/anae.15073.
62. Pfeifer M, Ewig S, Voshaar T, Randerath WJ, Bauer T, Geiseler J et al. Position paper for the state-of-the-art application of respiratory support in patients with COVID-19. *Respiration; international review of thoracic diseases.* 2020;99(6):521-42. doi:10.1159/000509104.
63. Cinesi Gómez C, Peñuelas Rodríguez Ó, Luján Torné M, Egea Santaolalla C, Masa Jiménez JF, García Fernández J et al. Clinical consensus recommendations regarding non-invasive respiratory support in the adult patient with acute respiratory failure secondary to SARS-CoV-2 infection. *Archivos de bronconeumologia.* 2020;56 Suppl 2:11-8. doi:10.1016/j.arbres.2020.03.005.
64. Porter RJ, Douglas K, Knight RG. Monitoring of cognitive effects during a course of electroconvulsive therapy: Recommendations for clinical practice. *J ECT.* 2008;24(1).
65. Agarwal A, Basmaji J, Muttalib F, Granton D, Chaudhuri D, Chetan D et al. High-flow nasal cannula for acute hypoxemic respiratory failure in patients with COVID-19: systematic reviews of effectiveness and its risks of aerosolization, dispersion, and infection transmission. *Can J Anesth.* 2020;67(9):1217-48. doi:10.1007/s12630-020-01740-2.
66. Raimondi N, Vial MR, Calleja J, Quintero A, Cortes Alban A, Celis E et al. Evidence-based guides in tracheostomy use in critical patients. *Med Intensiva.* 2017;41(2):94-115. doi:10.1016/j.medin.2016.12.001.
67. Scales DC, Thiruchelvam D, Kiss A, Redelmeier DA. The effect of tracheostomy timing during critical illness on long-term survival. *Crit Care Med.* 2008;36(9):2547-57. doi:10.1097/CCM.0b013e31818444a5.
68. Sommer DD, Engels PT, Weitzel EK, Khalili S, Corsten M, Tewfik MA et al. Recommendations from the CSO-HNS taskforce on performance of tracheotomy during the COVID-19 pandemic. *J Otolaryngol Head Neck Surg.* 2020;49(1):23. doi:10.1186/s40463-020-00414-9.
69. Aviles-Jurado FX, Prieto-Alhambra D, Gonzalez-Sanchez N, de Osso J, Arancibia C, Rojas-Lechuga MJ et al. Timing, complications, and safety of tracheotomy in critically ill patients with COVID-19. *JAMA Otolaryngol Head Neck Surg.* 2020. doi:10.1001/jamaoto.2020.3641.
70. Pichi B, Mazzola F, Bonsembiante A, Petrucci G, Zocchi J, Moretto S et al. CORONA-steps for tracheotomy in COVID-19 patients: A staff-safe method for airway management. *Oral Oncol.* 2020;105:104682. doi:10.1016/j.oraloncology.2020.104682.
71. Ahmad I, El-Boghdadly K, Bhagrath R, Hodzovic I, McNarry AF, Mir F et al. Difficult Airway Society guidelines for awake tracheal intubation (ATI) in adults. *Anaesthesia.* 2020;75(4):509-28. doi:10.1111/anae.14904.
72. Alhomary M, Ramadan E, Curran E, Walsh SR. Videolaryngoscopy vs. fiberoptic bronchoscopy for awake tracheal intubation: a systematic review and meta-analysis. *Anaesthesia.* 2018;73(9):1151-61. doi:10.1111/anae.14299.

- 73. Mascha EJ, Schober P, Schefold JC, Stueber F, Luedi MM. Staffing with disease-based epidemiologic indices may reduce shortage of intensive care unit staff during the COVID-19 pandemic. *Anesth Analg.* 2020;131(1):24-30. doi:10.1213/ane.0000000000004849.**
74. Cook TM. Personal protective equipment during the coronavirus disease (COVID) 2019 pandemic - a narrative review. *Anaesthesia.* 2020;75(7):920-7. doi:10.1111/anae.15071.
75. Tian Y, Tu X, Zhou X, Yu J, Luo S, Ma L et al. Wearing a N95 mask increases rescuer's fatigue and decreases chest compression quality in simulated cardiopulmonary resuscitation. *Am J Emerg Med.* 2020. doi:10.1016/j.ajem.2020.05.065.
76. De Ioris MA, Scarselli A, Ciofi Degli Atti ML, Rava L, Smarrazzo A, Concato C et al. Dynamic viral severe acute respiratory syndrome coronavirus 2 RNA shedding in children: Preliminary data and clinical consideration from a Italian Regional Center. *J Pediatric Infect Dis Soc.* 2020;9(3):366-9. doi:10.1093/jpids/piaa065.
77. Parri N, Lenge M, Buonsenso D. Children with Covid-19 in pediatric emergency departments in Italy. *N Engl J Med.* 2020;383(2):187-90. doi:10.1056/NEJMc2007617.
78. Gotzinger F, Santiago-Garcia B, Noguera-Julian A, Lanaspá M, Lancellata L, Calo Carducci FI et al. COVID-19 in children and adolescents in Europe: a multinational, multicentre cohort study. *Lancet Child Adolesc Health.* 2020;4(9):653-61. doi:10.1016/S2352-4642(20)30177-2.
79. Ramgolam A, Hall GL, Zhang G, Hegarty M, von Ungern-Sternberg BS. Inhalational versus intravenous induction of anesthesia in children with a high risk of perioperative respiratory adverse events: A randomized controlled trial. *Anesthesiology.* 2018;128(6):1065-74. doi:10.1097/ALN.0000000000002152.
80. Riva T, Pedersen TH, Seiler S, Kasper N, Theiler L, Greif R et al. Transnasal humidified rapid insufflation ventilatory exchange for oxygenation of children during apnoea: a prospective randomised controlled trial. *Br J Anaesth.* 2018;120(3):592-9. doi:10.1016/j.bja.2017.12.017.
81. Riva T, Theiler L, Jaquet Y, Giger R, Nisa L. Early experience with high-flow nasal oxygen therapy (HFNOT) in pediatric endoscopic airway surgery. *Int J Pediatr Otorhinolaryngol.* 2018;108:151-4. doi:10.1016/j.ijporl.2018.02.035.
82. Lee YC, Kim JM, Ko HB, Lee SR. Use of laryngeal mask airway and its removal in a deeply anaesthetized state reduces emergence agitation after sevoflurane anaesthesia in children. *J Int Med Res.* 2011;39(6):2385-92. doi:10.1177/147323001103900639.
83. Engelhardt T, Fiadjoe JE, Weiss M, Baker PA, Bew S, Piedad Echeverry M et al. A framework for the management of the pediatric airway. *Pediatr Anesth.* 2019;29:985-92. doi:10.1111/pan.13716.
84. Fiadjoe JE, Nishisaki A, Jagannathan N, Hunyady AI, Greenberg RS, Reynolds PI et al. Airway management complications in children with difficult tracheal intubation from the Pediatric Difficult Intubation (PeDI) registry: a prospective cohort analysis. *Lancet Respir Med.* 2016;4(1):37-48. doi:10.1016/s2213-2600(15)00508-1.
85. Morray JP, F. GJ, Caplan RA, Posner KL, Gild WM, Cheney FW. A comparison of pediatric and adult anesthesia closed malpractice claims. *Anesthesiology.* 1993;78(3):461-7.
86. Ulmer F, Lennertz J, Greif R, Butikofer L, Theiler L, Riva T. Emergency front of neck access in children: a new learning approach in a rabbit model. *Br J Anaesth.* 2020;125(1):e61-e8. doi:10.1016/j.bja.2019.11.002.
87. Umek N, Hodzovic I, Damjanovska M, Cvetko E, Zel J, Seliskar A et al. Rescue oxygenation success by cannula or scalpel-bougie emergency front-of-neck access in an anaesthetised porcine model. *PLoS one.* 2020;15(5):e0232510. doi:10.1371/journal.pone.0232510.
88. McNiven ND, Pracy JP, McGrath BA, Robson AK. The role of scalpel-bougie cricothyroidotomy in managing emergency front of neck airway access. A review and technical update for ENT surgeons. *Clin Otolaryngol.* 2018;43(3):791-4. doi:10.1111/coa.13120.
89. Simpson JP, Wong DN, Verco L, Carter R, Dzidowski M, Chan PY. Measurement of airborne particle exposure during simulated tracheal intubation using various proposed aerosol containment devices during the COVID-19 pandemic. *Anaesthesia.* 2020. doi:10.1111/anae.15188.

90. Gould CL, Alexander PDG, Allen CN, McGrath BA, Shelton CL. Protecting staff and patients during airway management in the COVID-19 pandemic: are intubation boxes safe? *Br J Anaesth.* 2020;125(3):e292-e3. doi:10.1016/j.bja.2020.05.001.
91. Sorbello M, Rosenblatt W, Hofmeyr R, Greif R, Urdaneta F. Aerosol boxes and barrier enclosures for airway management in COVID-19 patients: a scoping review and narrative synthesis. *Br J Anaesth.* 2020. doi:10.1016/j.bja.2020.08.038.
92. Kangas-Dick AW, Swearingen B, Wan E, Chawla K, Wiesel O. Safe extubation during the COVID-19 pandemic. *Respiratory Med.* 2020;170:106038. doi:10.1016/j.rmed.2020.106038.
93. Kearsley R. Intubation boxes for managing the airway in patients with COVID-19. *Anaesthesia.* 2020;75(7):969. doi:10.1111/anae.15081.

Journal Pre-proof

DECLARATION OF COMPETING INTEREST

Alexander Fuchs: None declared under financial, general, and institutional competing interests

Daniele Lanzi: None declared under financial, general, and institutional competing interests

Christian M Beilstein: None declared under financial, general, and institutional competing interests

Thomas Riva: None declared under financial, general, and institutional competing interests

Richard D. Urman: Declared research funding from Merck, Medtronic, Acacia, AcelRx and fees from Takeda and Heron.

Markus M Luedi: None declared under financial, general, and institutional competing interests

Matthias Braun: None declared under financial, general, and institutional competing interests