

Establishing the Injury Severity of Subaxial Cervical Spine Trauma: Validating the Hierarchical Nature of the AO Spine Subaxial Cervical Spine Injury Classification System

Gregory D. Schroeder, MD¹, Jose A. Canseco, MD, PhD¹, Parthik D. Patel, MD¹, Srikanth N. Divi, MD¹, Brian A. Karamian, MD¹, Frank Kandziora, MD, PhD², Emiliano N. Vialle, MD³, F. Cumhur Oner, MD, PhD⁴, Klaus J. Schnake, MD⁵, Marcel F. Dvorak, MD⁶, Jens R. Chapman, MD⁷, Lorin M. Benneker, MD⁸, Shanmuganathan Rajasekaran, MS(Ortho), PhD⁹, Christopher K. Kepler, MD, MBA¹, Alexander R. Vaccaro, MD, PhD, MBA¹, AO Spine Cervical Classification Validation Group

¹Rothman Orthopaedic Institute at Thomas Jefferson University, Philadelphia, PA, USA

²Center for Spinal Surgery and Neurotraumatology, Berufsgenossenschaftliche Unfallklinik, Frankfurt am Main, Germany

³Cajuru Hospital, Catholic University of Parana, Curitiba, Brazil

⁴University Medical Center, Utrecht, Netherlands

⁵Malteser Waldkrankenhaus St. Marien, Erlangen, Germany

⁶Vancouver General Hospital, Vancouver, British Columbia, Canada

⁷Harborview Medical Center, Seattle, WA, USA

⁸Sonnenhofspital Bern, Bern, Switzerland

⁹Department of Orthopaedic Surgery, Ganga Hospital, Coimbatore, Tamilnadu, India

Corresponding author:

Jose A. Canseco, MD, PhD

Rothman Orthopaedic Institute at Thomas Jefferson University Hospital

925 Chestnut St, 5th Floor, Philadelphia, PA 19107

jose.canseco@rothmanortho.com

The manuscript submitted does not contain information about medical device(s)/drug(s).

This study was organized and funded by AO Spine through the AO Spine Knowledge Forum Trauma, a focused group of international Trauma experts.

Relevant financial activities outside the submitted work: consultancy, grants, royalties, stocks, payment for lecture.

This is an open access article distributed under the Creative Commons Attribution License 4.0, (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mini Abstract:

This study validates the hierarchical nature of the AO Spine Subaxial Cervical Spine Injury Classification by surveying surgeons' perceived subtype injury severity scores. The study suggests this system is generalizable by geography and experience, and is a step forward towards developing a universally accepted subaxial cervical spine trauma treatment algorithm.

Structured Abstract

Study Design: Global cross-sectional survey

Objective: To validate the AO Spine Subaxial Cervical Spine Injury Classification by examining the perceived injury severity by surgeon across AO geographical regions and practice experience.

Summary of Background Data: Previous subaxial cervical spine injury classifications have been limited by subpar interobserver reliability and clinical applicability. In an attempt to create a universally validated scheme with prognostic value, AO Spine established a subaxial cervical spine injury classification involving four elements: (1) injury morphology, (2) facet injury involvement, (3) neurologic status, and (4) case-specific modifiers.

Methods: A survey was sent to 272 AO Spine members across all geographic regions and with a variety of practice experience. Respondents graded the severity of each variable of the classification system on a scale from zero (low severity) to 100 (high severity). Primary outcome was to assess differences in perceived injury severity for each injury type over geographic regions and level of practice experience.

Results: A total of 189 responses were received. Overall, the classification system exhibited a hierarchical progression in subtype injury severity scores. Only three subtypes showed a significant difference in injury severity score among geographic regions: F3 (floating lateral mass fracture, $p:0.04$), N3 (incomplete spinal cord injury, $p:0.03$), and M2 (critical disk herniation, $p:0.04$). When stratified by surgeon experience, pairwise comparison showed only 2 morphological subtypes, B1 (bony posterior tension band injury, $p:0.02$) and F2 (unstable facet fracture, $p:0.03$), and one neurologic subtype (N3, $p:0.02$) exhibited a significant difference in injury severity score.

Conclusions: The AO Spine Subaxial Cervical Spine Injury Classification System has shown to be reliable and suitable for proper patient management. The study shows this classification is substantially generalizable by geographic region and surgeon experience; and provides a consistent method of communication among physicians while covering the majority of subaxial cervical spine traumatic injuries.

Key words: AO Spine Subaxial Cervical Spine Injury, Classification System, injury severity score, cervical spine, trauma, validation

Level of Evidence: 4

Cervical spine trauma can be catastrophic, with timely identification and optimal treatment key to achieving successful outcomes. Efficient inter-provider communication is essential in the management of cervical spine injuries. Classification systems are tools utilized by physicians to simplify communication, and are best when their use is simple and reproducible, while at the same time able to transmit comprehensive information from diagnosis to prognosis and clinical management.^{1,2} The subaxial cervical spine is composed of all elements from C3 to C7, and various classification schemes have been developed over the years to describe traumatic injuries affecting this region of the spine.^{3,4} Among the first systems established, Allen and Ferguson proposed a scheme based on radiographic findings and inferred mechanisms of injury, postulating 6 main categories each with a series of anatomic severity stages: flexion-compression, flexion-distraction, extension-compression, extension-distraction, vertical compression, and lateral flexion.³⁻⁵ Harris et al. expanded upon this scheme adding rotational components to the mechanisms considered.^{4,6-8} Although this system and its modification were comprehensive, they lack reliability and clinical applicability.^{4,7-9}

More recently, in an attempt to circumvent the shortcomings of the previous systems, the AO Spine Trauma Study Group developed the Subaxial Injury Classification System (SLIC).⁸ In addition to considering the morphology of the injury, SLIC incorporated the neurological status of the patient, as well as the integrity of the disco-ligamentous complex, producing an algorithm that can be used to recommend conservative versus operative management.^{2,8,10} Studies have shown the SLIC system to be valid and reliable.⁸ Various reports have suggested it has good interobserver and intraobserver reliability when compared to older systems, with total score intraclass correlation coefficients ranging from 0.71 to 0.79 and 0.83 to 0.97, respectively.^{9,11,12} Nevertheless, persistent provider disagreements on the morphological definitions postulated by SLIC are also suggested in the literature.¹²

To date, no single scheme has been fully accepted by experts as the ideal subaxial cervical spine classification system. In an effort to achieve widespread agreement, the AO Spine Subaxial Cervical Spine Injury Classification System was developed (**Figure 1**).⁷ This system categorizes injuries based on four major criteria: (1) injury morphology, (2) facet injury involvement, (3) neurological status, and (4) case-specific modifiers.⁷ First, morphology is divided into 3 major types: Type A (compression injuries), Type B (tension band injuries), and Type C (translational injuries in any axis). Types A and B, in turn, have subcategories. Type A0 for minor injury or no bony involvement; A1 for single endplate compression fractures with no posterior vertebral body involvement; A2 for pincer fractures of both endplates without posterior wall involvement; A3 for burst fractures of one endplate; and A4 for sagittal split fractures of both endplates. Type B1 describes a bony posterior tension band injury; B2 is a complete posterior tension band injury that includes soft tissue components; and B3 is an anterior tension band injury. Type C injuries do not have subcategories, but if there are associated injuries of the A or B types, it should be noted as a subtype after describing a Type C injury.⁷

Additionally, this more comprehensive classification system incorporates facet injuries within the Type F category.⁷ The F1 subtype for non-displaced facet fractures, F2 for

facet fractures with instability, F3 for floating lateral mass injuries, and F4 for pathologic subluxation or dislocated facet injuries. Moreover, the neurological status of the patient is described by the N descriptor, with N0 representing a neurologically intact patients, N1 for transient neurologic deficits that have resolved, N2 for radiculopathies, N3 for incomplete cord injuries, and N4 for complete cord injuries; note the NX descriptor is used for patients unable to be examined due to secondary causes.⁷ Finally, the scheme incorporates case-specific modifiers to more thoroughly describe injuries. The M1 modifier describes partial disruption of the posterior soft tissue complex, while the M2 modifier depicts a critical disk herniation, and the M3 modifier signals the presence of an associated metabolic bone disorder. M4 describes the involvement of a vertebral artery.

This AO Spine subaxial cervical spine injury classification system has shown acceptable reliability, with various studies reporting kappa coefficients for overall interobserver reliability ranging from 0.57 to 0.64, and intraobserver reproducibility ranging from 0.54 to 0.95.^{1,7,13-15} Nevertheless, there still exist differences among surgeons and countries on the management of subaxial cervical spine injuries, and the development of consistent and universal treatment strategies is desirable to achieve uniform outcomes.¹⁶ The purpose of this study was to describe the severity of injury perceived by a surgeon as it relates to the scheme postulated by the AO Spine subaxial cervical spine injury classification, with the ultimate goal of developing a universally validated classification scheme.

Methods

Data Collection

As previously established¹⁷, a survey (**Supplemental Digital Content 1**, <http://links.lww.com/BRS/B694>) was sent to the members of the AO Spine Cervical Classification Validation Group. The group is composed of spine surgeons located in six different geographic regions (North America, South America, Europe, Africa, Asia, and the Middle East). For each variable (including types and subtypes) of the AO Spine Subaxial Cervical Spine Injury Classification System, respondents were asked to provide a numerical severity grade, including the morphology of the injury, the neurological status, and the case-specific modifiers. A grade of zero was assigned to a minimally severe injury, while a grade of 100 belonged to injuries with the highest severity possible. Only questionnaires with at least one valid answer, in addition to the demographic information, were included in the final analysis. Note, years of practice experience was collected as < 5 years, 5 – 10 years, 11 – 20 years, 20+ years.

Statistical Analysis

Based on prior studies¹⁷, surgeon responses were used to produce a perceived injury severity score (ISS) for each fracture type and subtype of the classification system, including neurological status and modifiers. ISS were generated for the total samples, as well as for subgroups according to geographical region and surgeon experience. A descriptive statistical analysis was performed for categorical and continuous data. For categorical data, frequencies were calculated based on the number of non-missing replies. Continuous data were analyzed using the following descriptive statistics: mean, standard deviation, median, minimum, and

maximum. Median ISS for the same fracture type, neurological status, and modifier were tested for equality across all geographic regions using a Kruskal-Wallis test. Years of surgeon experience was re-grouped (\bullet 10 years, $>$ 10 years), and fracture type, neurological status, and case-specific modifiers were tested for equality by a Wilcoxon rank sum test. Post-hoc analysis by geographic region of surgeons was done only for fracture types, neurological status, and modifiers which had an observed borderline significance ($p \bullet 0.10$) upon Kruskal-Wallis testing. Regarding practice experience of surgeons, the post-hoc analysis was done for all fracture types, neurological status, and case-specific modifiers via pairwise comparison of injury severity scores utilizing a Tukey-Kramer adjustment. The significance level was defined at $\bullet = 0.05$. All analysis was performed using the statistical software SAS version 9.4 (SAS Institute Inc., Cary, NC, USA).

Results

A total of 195 out of 272 members of the AO Spine Cervical Classification Validation Group responded to the survey. Six surveys did not meet inclusion criteria and were excluded from analysis, for a total of 189 surveys considered in the final analysis. Respondent characteristics are summarized in Table 1. The overall injury severity score of each variable (including type/subtype) of the classification is presented in Table 2. The results show a hierarchical nature of the classification system, with only a B1 injury (bony posterior tension band, 60 [45.0, 70.0]) being perceived slightly less severe than an A4 fracture (two endplate sagittal split, 60 [50,80]) based on interquartile range. All other injury types/subtypes were found to have increasing ISS based on median and interquartile range as the classification progressed.

Subgroup analysis for classification morphology demonstrated that there were no statistically significant differences in ISS based on geographic region across fracture types/subtypes, with the exception of an F3 (floating lateral mass) fracture ($p: 0.04$) (Table 3). Additionally, no statistically significant difference based on region was found across neurological status grades, with the exception of the N3 (incomplete spinal cord injury) subtype ($p: 0.03$) (Table 3). Finally, in terms of case-specific modifier, only the M2 (critical disk herniation) subtype exhibited a statistically significant difference across geographical regions ($p: 0.04$) (Table 3). Further subgroup analysis with post-hoc pairwise comparison utilizing Tukey-Kramer adjustments showed that within the F3 subtype, there was a statistically significant difference in perceived severity between surgeons from Europe and Latin/South America ($p: 0.01$). Similarly, there was a significant difference in perceived severity between Middle Eastern and North American surgeons for N3 ($p: 0.03$), and European and Latin/South American surgeons for M2 ($p: 0.03$). No other regions exhibited significant differences via pairwise comparison.

Subgroup analysis of injury severity scores based on re-grouped years of surgeon experience (\bullet 10 years and $>$ 10 years) revealed that 5 morphological subtypes, and the N3 ($p: 0.02$) neurological status subtype exhibited statistically significant differences (Table 4). For morphology, the A0 (minor injury), A1 (single endplate fracture), A2 (pincer fracture), B1 (bony posterior tension band injury), and F2 (unstable facet fractures) fracture subtypes showed significant differences with respect to surgeon experience ($p: 0.04, 0.02, 0.02, 0.03,$

0.04, respectively). Subgroup analysis, including all experience groups (< 5 years, 5-10 years, 11-20 years, and 20+ years), with post-hoc comparison utilizing Tukey-Kramer adjustments showed a statistically significant difference within the B1 subtype in perceived severity by experience between surgeons with < 5 years of practice and those with 20+ years of practice (p: 0.02). Furthermore, for the F2 and N3 subtypes, there was a significant difference in perceived severity between surgeons with < 5 years of experience and those with 20+ years (p: 0.03), and between those with 5-10 years of experience and those with 11-20 years (p: 0.02), respectively. Post-hoc analysis did not exhibit statistically significant differences for A0, A1 or A2 subtypes.

Discussion

The ultimate goal of a classification system for spine trauma is to be validated and uniformly utilized around the world. To date, no universally accepted spine trauma classification scheme exists, and the literature is awash with studies showing the variability in severity assessment and management of traumatic spine injuries.¹⁷⁻²⁵ The current study intended to describe the variations in injury severity perception by spine surgeons of the AO Spine Subaxial Cervical Spine Injury Classification System based on geographic region, as well as by a surgeon's experience in practice. Overall, the results show that the severity score of each variable within the classification system increases as the subtype increases, suggesting validity in the hierarchical progression of individual subtypes from A to C, and F1 to F4. Interestingly, there was only one instance in the entire classification scheme in which the score's hierarchical advancement was not consistent, the progression from an A4 (two endplate sagittal split) fracture to a B1 (bony posterior tension band injury, bony chance) fracture, with overall scores of 60.0 [50.0, 80.0] and 60.0 [45.0, 70.0], respectively. Although the results did not show a statistically significant difference in perceived injury severity across all geographic regions for either A4 or B1 fractures independently, it was not entirely surprising to find the transition from A4 to B1 to be inconsistent based on injury severity score. Cervical spine bony chance fractures are extremely rare, and surgeons may not be familiar with the management of said injuries.²⁶ This fact is supported by our finding that a surgeon's years of experience did show a statistically significant difference in terms of perceived injury severity of B1 fractures, with surgeons having less than 5 years of practice experience giving a B1 fracture an average score of 67.5 [50.0, 75.0], and injury severity scores progressively decreasing as a surgeon's years of experience advanced, with surgeons having 20 or more years of experience averaging a B1 injury severity score of 50.0 [30.0, 60.0]. In fact, subgroup analysis via pairwise comparison revealed a statistically significant difference in perceived injury severity for B1 subtypes only between the aforementioned groups < 5 years and 20+ years (p: 0.02). While commonly observed in the thoracolumbar spine transition between T10-L2, bony chance fractures involving the cervical spine are generally most often seen in the setting of ankylosing spondylitis²⁶⁻³¹, and the lack of experience with this type of injury leads to the observed inconsistent scoring. Furthermore, even in the more common thoracolumbar burst fracture, Schroeder et al reported that AO Spine thoracolumbar B1 fractures are considered less severe than A4 fractures¹⁷; highlighting the observation that within both, the cervical spine and thoracolumbar spine, there is no clear perceived severity difference between A4 and B1 fracture subtypes.

In terms of subgroup variations, with the exception of an F3 fracture ($p: 0.04$), the current results did not show any significant difference in morphology-based injury severity score based on a surgeon's geographical area. Given the complex nature of the F3 subtypes, it was unsurprising to find regional variations in perceived injury severity. Floating lateral mass fractures (F3) are relatively rare, unstable injuries, difficult to diagnose and generally resulting from high-energy mechanisms.³²⁻³⁴ Various studies in spine trauma discuss the persistently controversial management of cervical facet fractures, with F3 injuries generally requiring surgical intervention for instability.³²⁻³⁴ In a recent case series, Manoso and colleagues identified 60 consecutive cases of floating lateral mass cervical facet fractures between C3-C7, and found that all patients treated non-operatively developed instability, whereas all patients treated with two level fusions did not show evidence of radiographic failure and maintained alignment.³² Comparatively, Vedantam et al. retrospectively reviewed 35 patients with subaxial cervical facet fractures, six of whom had F3 injuries.³³ Even though 2/6 patients failed conservative management, the authors advocate that nondisplaced injuries warrant a trial of non-operative interventions. Regarding other AO Spine classification subtypes with notable differences across geographical regions, only N3 injuries ($p: 0.03$) and an M2 modifier ($p: 0.04$) showed a statistically significant difference across geographic areas. Again, this was foreseeable, given the continually contentious diagnostic and management recommendations of patients with incomplete spinal cord injuries and critical disc herniations in the setting of facet dislocations.³⁵ Overall, with limited exceptions, our observations suggest that the AO Spine Subaxial Cervical Spine Injury Classification System is geographically generalizable, similar to our group's findings regarding the AO Spine Thoracolumbar Spine Injury Classification scheme.¹⁷

When considering a surgeon's years of practice experience, after subgroup analysis adjustments, only morphology-related injury subtypes B1 and F2 appeared to exhibit a significant difference in injury severity scores. As previously discussed, B1 injuries are relatively rare and case experience is likely to play a factor in perceived injury severity. For facet fractures with instability (F2), similar inferences can be made given the controversial nature of facet fracture management particularly when unstable.³²⁻³⁴ Additionally, for other variables in the classification, only the N3 ($p: 0.02$) neurological grade showed significant variation across surgeon experience, with variability noted in mid-career surgeons, likely secondary to the aforementioned controversies in the management of incomplete spinal cord injuries.

As seen in equivalent analysis of the AO Spine Thoracolumbar Spine Injury Classification, this study's findings suggest that regional and experiential variations observed in cervical trauma management are possibly influenced by factors outside of the interpretation of an injury's perceived severity, such as cultural differences, healthcare infrastructure and costs.¹⁷ This is the reason behind basing the classification system on morphological features and not treatment algorithms.⁷ In terms of subspecialty training, while there is a higher proportion of orthopaedic spine surgeons compared to neurosurgeons in our study, previous reports highlight high agreement on primary management decisions among spine specialists.^{36,37} Although further studies with proportional subspecialty respondents will be helpful, perceived severity scores are unlikely to change.

The current study is not without limitations. First, the survey study design provides a small sample of surgeons with uneven numbers across geographical regions and subspecialty training. Moreover, the questionnaire administration followed the preestablished order of the classification scheme, leaving open the possibility of bias when surgeons assigned perceived injury severity scores to each ascending type/subtype in the scheme. Additionally, this report does not establish guidelines for the surgical vs. non-surgical management of fracture types/subtypes within the classification, instead it establishes their perceived severity of injury.¹⁷ Finally, all respondents of the survey had affiliations with academic institutions, which is unlikely to represent all spine surgeons practicing in a specific geographical region where community hospitals may be more numerous.

Based on our findings, and the system's considerable interobserver and intraobserver reliability, the AO Spine Subaxial Cervical Spine Injury Classification System is a sound foundation for the development of a universally accepted treatment algorithm for subaxial cervical spine trauma. With few exceptions within controversial injury subtypes, this classification system is considerably generalizable by geographic region and surgeon experience.

Acknowledgements

This study was organized and funded by AO Spine through the AO Spine Knowledge Forum Trauma, a focused group of international Trauma experts. The AO Spine Injury Classification Systems were developed and funded by AO Spine through the AO Spine Knowledge Forum Trauma, a focused group of international spine trauma experts. AO Spine is a clinical division of the AO Foundation, which is an independent medically-oriented not-for-profit organization. Study support was provided directly through the AO Spine Research Department and AO Innovation Translation Center. Figure 1 provided by © AO Foundation, Switzerland with permission. The authors would like to thank Olesja Hazenbiller (AO Spine) for her editorial and administrative assistance, and Christian Knoll and Cordula Blohm (AO Innovation Translation Center) for their support with statistical analysis.

AO Spine Cervical Classification Validation Group members:

Ahmed Shawky Abdelgawaad, Waheed Abdul, Asmatullah Abdulsalam, Mbarak Abeid, Nissim Ackshota, Olga Acosta, Yunus Akman, Osama Aldahamshah, Abduljabbar Alhammoud, Hugo Aleixo, Hamish Alexander, Mahmoud Alkharsawi, Wael Alsammak, Hassame Amadou, Mohamad Amin, Jose Arbatin, Ahmad Atan, Alkinoos Athanasiou, Paloma Bas, Pedro Bazan, Thami Benzakour, Sofien Benzarti, Claudio Bernucci, Aju Bosco, Joseph Butler, Alejandro Castillo, Derek Cawley, Wong Chek, John Chen, Christina Cheng, Jason Cheung, Chun Chong, Stipe Corluka, Jose Corredor, Bruno Costa, Cloe Curri, Ahmed Dawoud, Juan Delgado-Fernandez, Serdar Demiroz, Ankit Desai, Maximo Diez-Ulloa, Noe Dimas, Sara Diniz, Bruno Direito-Santos, Johnny Duerinck, Tarek El-Hewala, Mahmoud El-Shamly, Mohammed El-Sharkawi, Guillermo Espinosa, Martin Estefan, Taolin Fang, Mauro Fernandes, Norbert Fernandez, Marcus Ferreira, Alfredo Figueiredo, Vito Fiorenza, Jibin Francis, Seibert Franz, Brett Freedman, Lingjie Fu, Segundo Fuego, Nitesh Gahlot, Mario

Ganau, Maria Garcia-Pallero, Bhavuk Garg, Sandeep Gidvani, Bjoern Giera, Amauri Godinho Jr., Morshed Goni, Maria Gonzalez, Richard Gonzalez, Dilip Gopalakrishnan, Andrey Grin, Samuel Grozman, Marcelo Gruenberg, Alon Grundshtein, Joana Guasque, Oscar Guerra, Alfredo Guiroy, Shafiq Hackla, Colin Harris, James Harrop, Waqar Hassan, Amin Henine, Zachary Hickman, Cristina Igualada, Andrew James, Chumpon Jetjumnong, Ariel Kaen, Balgopal Karmacharya, Cumur Kilincer, Zdenek Klezl, John Koerner, Christian Konrads, Ferdinand Krappel, Moyo Kruyt, Fernando Krywinski, Raghuraj Kundangar, Federico Landriel, Richard Lindtner, Daniela Linhares, Rafael Llombart-Blanco, William Lopez, Raphael Lotan, Juan Lourido, Luis Luna, Tijjani Magashi, Catalin Majer, Valentine Mandizvidza, Rui Manilha, Francisco Mannara, Konstantinos Margetis, Fabricio Medina, Jeronimo Milano, Naohisa Miyakoshi, Horatiu Moisa, Nicola Montemurro, Juan Montoya, Joao Morais, Sebastian Morande, Salim Msuya, Mohamed Mubarak, Robert Mulbah, Yuvaraja Murugan, Mansouri Nacer, Nuno Neves, Nicola Nicassio, Thomas Niemeier, Mejabi Olorunsogo, David Orosco, Kubilay Ozdener, Rodolfo Paez, Ripul Panchal, Konstantinos Paterakis, Emilija Pemovska, Paulo Pereira, Darko Perovic, Jose Perozo, Andrey Pershin, Phedy Phedy, David Picazo, Fernando Pitti, Uwe Platz, Mauro Pluderi, Gunasaelan Ponnusamy, Eugen Popescu, Selvaraj Ramakrishnan, Alessandro Ramieri, Brandon Rebholz, Guillermo Ricciardi, Daniel Ricciardi, Yohan Robinson, Luis Rodriguez, Ricardo Rodrigues-Pinto, Itati Romero, Ronald Rosas, Salvatore Russo, Joost Rutges, Federico Sartor, Babak Shariati, Jeevan Sharma, Mahmoud Shoaib, Sean Smith, Yasunori Sorimachi, Shilanant Sribastav, Craig Steiner, Jayakumar Subbiah, Panchu Subramanian, Tarun Suri, Chadi Tannoury, Devi Tokala, Adetunji Toluse, Victor Ungurean, Joachim Vahl, Marcelo Valacco, Cristian Valdez, Alejo Vernengo-Lezica, Andrea Veroni, Rian Vieira, Arun Viswanadha, Scott Wagner, David Wamae, Alexander Weening, Simon Weidert, Wen-Tien Wu, Meng-Huang Wu, Haifeng Yuan, Sung-Joo Yuh, Ratko Yurac, Baron Zarate-Kalfopulos, Alesksei Ziabrov, Akbar Zubairi.

Key Points:

- The AO Spine Subaxial Cervical Spine Injury Classification System exhibited a hierarchical progression in subtype injury severity scores.
- Only F3, N3 and M3 classification subtypes showed a significant difference in injury severity score among AO geographic regions.
- When stratified by surgeon experience, only B1, F2 and N3 subtypes exhibited a significant difference in injury severity score.
- With few exceptions, the perceived injury severity scores of the classification system subtypes are independent of geographic region and surgeon experience.

References

1. Schnake KJ, Schroeder GD, Vaccaro AR, et al. AOSpine Classification Systems (Subaxial, Thoracolumbar). *J Orthop Trauma* 2017;31:S14–23.
2. Feuchtbaum E, Buchowski J, Zebala L. Subaxial cervical spine trauma. *Curr Rev Musculoskelet Medicine* 2016;9:496–504.
3. Kwon BK, Vaccaro AR, Grauer JN, et al. Subaxial cervical spine trauma. *J Am Acad Orthop Surg* 2006;14:78–89.
4. Divi SN, Schroeder GD, Oner FC, et al. AOSpine—Spine Trauma Classification System: The Value of Modifiers: A Narrative Review With Commentary on Evolving Descriptive Principles. *Global Spine J* 2019;9:77S–88S.
5. Allen BL, Ferguson RL, Lehmann TR, et al. A Mechanistic Classification of Closed, Indirect Fractures and Dislocations of the Lower Cervical Spine. *Spine* 1982;7:1–27.
6. Harris JH, Edeiken-Monroe B, Kopaniky DR. A practical classification of acute cervical spine injuries. *Orthop Clin North Am* 1986;17:15–30.
7. Vaccaro AR, Koerner JD, Radcliff KE, et al. AOSpine subaxial cervical spine injury classification system. *European Spine J Official Publ European Spine Soc European Spinal Deformity Soc European Sect Cerv Spine Res Soc* 2016;25:2173–84.
8. Whang PG, Patel AA, Vaccaro AR. The development and evaluation of the subaxial injury classification scoring system for cervical spine trauma. *Clin Orthop Relat R* 2011;469:723–31.
9. Stone AT, Bransford RJ, Lee MJ, et al. Reliability of classification systems for subaxial cervical injuries. *Evidence-based Spine-care J* 2010;1:19–26.
10. Joaquim AF, Lawrence B, Daubs M, et al. Evaluation of the subaxial injury classification system. *J Craniovertebral Junction Spine* 2011;2:67–72.
11. Lee WJ, Yoon SH, Kim YJ, et al. Interobserver and Intraobserver Reliability of Sub-Axial Injury Classification and Severity Scale between Radiologist, Resident and Spine Surgeon. *J Korean Neurosurg S* 2012;52:200.
12. Middendorp JJ van, Audigé L, Bartels RH, et al. The Subaxial Cervical Spine Injury Classification System: an external agreement validation study. *Spine J* 2013;13:1055–63.
13. Urrutia J, Zamora T, Campos M, et al. A comparative agreement evaluation of two subaxial cervical spine injury classification systems: the AOSpine and the Allen and Ferguson schemes. *Eur Spine J* 2016;25:2185–92.
14. Silva OT da, Sabba MF, Lira HIG, et al. Evaluation of the reliability and validity of the newer AOSpine subaxial cervical injury classification (C-3 to C-7). *J Neurosurg Spine* 2016;25:303–8.
15. Urrutia J, Zamora T, Yurac R, et al. An Independent Inter- and Intraobserver Agreement Evaluation of the AOSpine Subaxial Cervical Spine Injury Classification System. *Spine* 2017;42:298–303.

16. Pishnamaz M, Curfs I, Uhing D, et al. Two-Nation Comparison of Classification and Treatment of Subaxial Cervical Spine Fractures: An Internet-Based Multicenter Study Among Spine Surgeons. *World Neurosurg* 2019;123:e125–32.
17. Schroeder GD, Vaccaro AR, Kepler CK, et al. Establishing the Injury Severity of Thoracolumbar Trauma. *Spine* 2015;40:E498–503.
18. Schroeder GD, Kepler CK, Koerner JD, et al. Is there a regional difference in morphology interpretation of A3 and A4 fractures among different cultures? *J Neurosurg Spine* 2016;24:332–9.
19. Rajasekaran S, Kanna RM, Schroeder GD, et al. Does the Spine Surgeon's Experience Affect Fracture Classification, Assessment of Stability, and Treatment Plan in Thoracolumbar Injuries? *Global Spine J* 2017;7:309–16.
20. Sadiqi S, Oner FC, Dvorak MF, et al. The Influence of Spine Surgeons' Experience on the Classification and Intraobserver Reliability of the Novel AOSpine Thoracolumbar Spine Injury Classification System-An International Study. *Spine* 2015;40:E1250–6.
21. Alcala-Cerra G, Diaz-Becerra C, Fernandes-Joaquim A, et al. Orthosis for thoracolumbar burst fractures without neurologic deficit: A systematic review of prospective randomized controlled trials. *J Craniovertebral Junction Spine* 2014;5:25.
22. Bailey CS, Dvorak MF, Thomas KC, et al. Comparison of thoracolumbosacral orthosis and no orthosis for the treatment of thoracolumbar burst fractures: interim analysis of a multicenter randomized clinical equivalence trial. *J Neurosurg Spine* 2009;11:295–303.
23. Bakhsheshian J, Dahdaleh NS, Fakurnejad S, et al. Evidence-based management of traumatic thoracolumbar burst fractures: a systematic review of nonoperative management. *Neurosurg Focus* 2014;37:E1.
24. Reinhold M, Knop C, Beisse R, et al. Operative treatment of 733 patients with acute thoracolumbar spinal injuries: comprehensive results from the second, prospective, Internet-based multicenter study of the Spine Study Group of the German Association of Trauma Surgery. *Eur Spine J* 2010;19:1657–76.
25. Schnake KJ, Stavridis SI, Kandziora F. Five-year clinical and radiological results of combined anteroposterior stabilization of thoracolumbar fractures. *J Neurosurg Spine* 2014;20:497–504.
26. Eghbal K, Abdollahpour HR, Ghaffarpasand F. Traumatic Chance Fracture of Cervical Spine: A Rare Fracture Type and Surgical Management. *Asian J Neurosurg* 2018;13:906–9.
27. Meena RK, Dhandapani S. Chance fracture in ankylosing spondylitis—not merely by chance! Danger of injudicious use of skull traction. *Spine J* 2015;15:e47–8.
28. Wong ASH, Yu DHY. Cervical spine fracture in a patient with ankylosing spondylitis causing a C2-T9 spinal epidural hematoma- Treatment resulted in a rapid and complete recovery from tetraplegia: Case report and literature review. *Asian J Neurosurg* 2015;10:53.
29. Gilard V, Curey S, Derrey S, et al. Cervical spine fractures in patients with ankylosing spondylitis: Importance of early management. *Neurochirurgie* 2014;60:239–43.

30. Budorick TE, Anderson PA, Rivara FP, et al. Flexion-distraction fracture of the cervical spine. A case report. *J Bone Jt Surg Am Volume* 1991;73:1097–100.
31. Xiang L-B, Yu H-L, Liu J, et al. One-stage surgery by a combined anterior-posterior approach to treat ankylosing spondylitis complicated by a multiple-level cervical vertebral chance fracture. *Mod Rheumatology Jpn Rheumatism Assoc* 2015;25:282–5.
32. Manoso MW, Moore TA, Agel J, et al. Floating Lateral Mass Fractures of the Cervical Spine. *Spine* 2016;41:1421–7.
33. Vedantam A, Fridley JS, Navarro JC, et al. Management of Acute Unilateral Nondisplaced Subaxial Cervical Facet Fractures. *Oper Neurosurg* 2017;14:104–11.
34. Pehler S, Jones R, Staggers JR, et al. Clinical Outcomes of Cervical Facet Fractures Treated Nonoperatively With Hard Collar or Halo Immobilization. *Global Spine J* 2018;9:48–54.
35. Divi SN, Schroeder GD, Mangan JJ, et al. Management of Acute Traumatic Central Cord Syndrome: A Narrative Review. *Global Spine J* 2019;9:89S-97S.
36. Grauer JN, Vaccaro AR, Beiner JM, et al. Similarities and Differences in the Treatment of Spine Trauma Between Surgical Specialties and Location of Practice. *Spine* 2004;29:685–96.
37. Canseco JA, Schroeder GD, Patel PD, et al. Regional and experiential differences in surgeon preference for the treatment of cervical facet injuries: a case study survey with the AO Spine Cervical Classification Validation Group. *Eur Spine J* 2020;1–7.

Figure Legends

Figure 1. AO Spine Subaxial Cervical Spine Injury Classification.^{1,7} Reprinted with permission from AO Spine International. © AO Foundation, Switzerland.

AO Spine Subaxial Classification System

Type A Compression Injuries

A0 Minor, nonstructural fractures
No bony injury or minor injury such as an isolated lamina fracture or spinous process fracture.

A1 Wedge-compression
Compression fracture involving a single endplate without involvement of the posterior wall of the vertebral body.

A2 Split
Coronal split or pincher fracture involving both endplates without involvement of the posterior wall of the vertebral body.

A3 Incomplete burst
Burst fracture involving a single endplate with involvement of the posterior vertebral wall.

A4 Complete burst
Burst fracture or sagittal split involving both endplates.

Type B Tension Band Injuries

B1 Posterior tension band injury (bony)
Physical separation through fractured bony structures only.

B2 Posterior tension band injury (bony capsuloligamentous, ligamentous)
Complete disruption of the posterior capsuloligamentous or bony capsuloligamentous structures together with a vertebral body, disk, and/or facet injury.

B3 Anterior tension band injury
Physical disruption or separation of the anterior structures (vertebrae) with tethering of the posterior elements.

Type C Translation Injuries

C Translational injury in any axis-displacement or translation of one vertebral body relative to another in any direction.

Type F Facet Injuries

F1 Nondisplaced facet fracture
With fragment <1cm in height, <40% of lateral mass.

F2 Facet fracture with potential for instability
With fragment >1cm, > than 40% lateral mass, or displaced.

F3 Floating lateral mass

F4 Pathologic subluxation or perched/dislocated facet

BL Bilateral Injuries

BL Bilateral Injury

Algorithm for morphologic classification

Neurology

Type	Description
N0	No neurologic deficit
N1	Transient neurologic deficit
N2	Permanent neurologic deficit
N3	Radicular symptoms
N4	Incomplete spinal cord injury or any degree of cauda equina injury
N5	Complete spinal cord injury
N6	Conical transected
N7	Continued spinal cord compression

Classification Nomenclature

C6-C7: C (C7: A1) → Primary injury → C6-C7: B2* → Secondary/tertiary injury → C6-C7: B2**

* If there are multiple injuries to the same level - for example, small fracture (F1) and dislocation (F4) - only the highest level level injury is indicated (F4).

** If only focal injuries are identified - no A, B, or C injury - they are listed first after the burst injury.

Modifiers

Type	Description
M1	Posterior Capsuloligamentous Cordset injury without cordset disruption
M2	Critical disk formation
M3	Differing vertebral body masses (e.g. DISH, AS, OPLL, OLF)
M4	Vertebral artery abnormality

© 2012 AO Spine International. This is a licensed white AO Spine International AO Spine International AO Spine International AO Spine International AO Spine International. To the extent of the law, this document is provided as a service to the user. The AO Spine International logo is a registered trademark of AO Spine International.



Further information:
www.aospine.org/classification

Table 1: Surgeon Demographic Characteristics

Characteristics	Total Responders (n = 189)
Subspecialty*	
Orthopaedic Spine	131 (69.3)
Neurosurgery	58 (30.7)
Region*	
North America	18 (9.5)
Latin/South America	40 (21.2)
Europe	70 (37.0)
Africa	12 (6.3)
Asia	34 (18.0)
Middle East	15 (7.9)
No. of Years in Practice*	
< 5 years	50 (26.5)
5 – 10 years	61 (32.3)
11 – 20 years	50 (26.5)
> 20 years	28 (14.8)
Work Setting*	
Academic	78 (41.3)
Hospital Employed	88 (46.6)
Private Practice	23 (12.2)
No. of Spine Trauma Patients Treated per year**	50 (20; 100)
Time to Obtain an MRI at Home Institution*	
< 2 hours	52 (27.5)
2 – 12 hours	62 (32.8)
12 – 24 hours	28 (14.8)
> 24 hours	42 (22.2)
Cannot Obtain	5 (2.6)

* Proportions presented as: Number of Responders (%)

** Number presented as: Median (Interquartile Range)

Table 2: Global Analysis of Injury Severity Score Based on Fracture Type, Neurologic Status, and Case-Specific Modifier

Type	No. Of Responders	Median (IQR)
A0	178	5.0 (0.0; 10.0)
A1	179	20.0 (10.0; 25.0)
A2	179	30.0 (20.0; 50.0)
A3	179	50.0 (30.0; 60.0)
A4	179	60.0 (50.0; 80.0)
B1	179	60.0 (45.0; 70.0)
B2	179	80.0 (70.0; 85.0)
B3	179	80.0 (70.0; 90.0)
C	178	100.0 (100.0; 100.0)
F1	179	20.0 (10.0; 30.0)
F2	179	40.0 (30.0; 50.0)
F3	179	50.0 (40.0; 70.0)
F4	179	100.0 (85.0; 100.0)
N0	178	0.0 (0.0; 0.0)
N1	178	20.0 (10.0; 30.0)
N2	178	40.0 (30.0; 50.0)
N3	178	80.0 (70.0; 100.0)
N4	178	100.0 (85.0; 100.0)
NX	178	80.0 (50.0; 100.0)
M1	178	40.0 (30.0; 60.0)
M2	178	70.0 (50.0; 80.0)
M3	178	70.0 (60.0; 80.0)
M4	178	60.0 (50.0; 80.0)

IQR: Interquartile Range

Table 3: Regional Analysis of Injury Severity Score Based on Fracture Type, Neurologic Status, and Case-Specific Modifier

Type	_ North America _		Latin/South America		Europe		Africa		_____ Asia _____		_ Middle East _		P-value*
	N	Median (IQR)	N	Median (IQR)	N	Median (IQR)	N	Median (IQR)	N	Median (IQR)	N	Median (IQR)	
A0	18	5.0 (1.0; 10.0)	39	5.0 (0.0; 10.0)	67	5.0 (0.0; 10.0)	10	3.0 (0.0; 10.0)	30	5.0 (1.0; 10.0)	14	10.0 (0.0; 10.0)	0.81
A1	18	12.5 (6.0; 20.0)	39	15.0 (10.0; 20.0)	68	20.0 (10.0; 22.5)	10	20.0 (5.0; 25.0)	30	20.0 (10.0; 30.0)	14	20.0 (10.0; 20.0)	0.27
A2	18	30.0 (20.0; 50.0)	39	30.0 (20.0; 45.0)	68	30.0 (20.0; 40.0)	10	32.5 (30.0; 50.0)	30	37.5 (25.0; 60.0)	14	30.0 (25.0; 50.0)	0.45
A3	18	50.0 (25.0; 50.0)	39	45.0 (30.0; 50.0)	68	40.0 (30.0; 50.0)	10	50.0 (30.0; 60.0)	30	50.0 (40.0; 70.0)	14	45.0 (30.0; 75.0)	0.53
A4	18	67.5 (50.0; 75.0)	39	60.0 (50.0; 70.0)	68	60.0 (45.0; 70.0)	10	70.0 (50.0; 90.0)	30	72.5 (50.0; 90.0)	14	62.5 (50.0; 90.0)	0.29
B1	18	50.0 (40.0; 75.0)	39	55.0 (45.0; 70.0)	68	60.0 (45.0; 70.0)	10	55.0 (40.0; 75.0)	30	60.0 (50.0; 80.0)	14	60.0 (50.0; 70.0)	0.97
B2	18	80.0 (70.0; 90.0)	39	80.0 (70.0; 80.0)	68	80.0 (67.5; 80.0)	10	72.5 (70.0; 80.0)	30	80.0 (60.0; 90.0)	14	70.0 (60.0; 85.0)	0.53
B3	18	80.0 (70.0; 90.0)	39	80.0 (75.0; 90.0)	68	80.0 (70.0; 90.0)	10	87.5 (80.0; 95.0)	30	80.0 (70.0; 95.0)	14	80.0 (60.0; 90.0)	0.43
C	17	100.0 (100.0; 100.0)	39	100.0 (100.0; 100.0)	68	100.0 (100.0; 100.0)	10	100.0 (100.0; 100.0)	30	100.0 (100.0; 100.0)	14	100.0 (100.0; 100.0)	0.61
F1	18	12.5 (10.0; 25.0)	39	20.0 (10.0; 30.0)	68	20.0 (10.0; 25.0)	10	20.0 (10.0; 30.0)	30	20.0 (10.0; 30.0)	14	20.0 (15.0; 20.0)	0.64
F2	18	41.0 (30.0; 50.0)	39	40.0 (30.0; 50.0)	68	30.0 (25.0; 40.0)	10	50.0 (30.0; 60.0)	30	40.0 (30.0; 50.0)	14	40.0 (30.0; 50.0)	0.06
F3	18	52.5 (40.0; 70.0)	39	70.0 (40.0; 75.0)	68	40.0 (40.0; 60.0)	10	65.0 (50.0; 75.0)	30	52.5 (40.0; 60.0)	14	50.0 (40.0; 60.0)	0.04
F4	18	100.0 (86.0; 100.0)	39	100.0 (90.0; 100.0)	68	100.0 (85.0; 100.0)	10	100.0 (80.0; 100.0)	30	97.5 (80.0; 100.0)	14	95.0 (80.0; 100.0)	0.89
N0	18	0.0 (0.0; 1.0)	39	0.0 (0.0; 0.0)	67	0.0 (0.0; 0.0)	10	0.5 (0.0; 10.0)	30	0.0 (0.0; 0.0)	14	0.0 (0.0; 0.0)	0.29
N1	18	20.0 (10.0; 30.0)	39	20.0 (15.0; 40.0)	67	20.0 (10.0; 30.0)	10	20.0 (10.0; 30.0)	30	20.0 (10.0; 25.0)	14	20.0 (10.0; 20.0)	0.63
N2	18	30.0 (20.0; 50.0)	39	40.0 (30.0; 60.0)	67	40.0 (25.0; 50.0)	10	50.0 (40.0; 55.0)	30	35.0 (25.0; 50.0)	14	40.0 (30.0; 40.0)	0.09
N3	18	99.5 (80.0; 100.0)	39	90.0 (80.0; 100.0)	67	80.0 (70.0; 100.0)	10	77.5 (70.0; 100.0)	30	80.0 (70.0; 90.0)	14	75.0 (60.0; 80.0)	0.03
N4	18	100.0 (99.0; 100.0)	39	100.0 (90.0; 100.0)	67	100.0 (90.0; 100.0)	10	92.5 (75.0; 100.0)	30	100.0 (80.0; 100.0)	14	100.0 (80.0; 100.0)	0.57
NX	18	87.5 (70.0; 100.0)	39	80.0 (50.0; 100.0)	67	70.0 (50.0; 100.0)	10	100.0 (95.0; 100.0)	30	80.0 (50.0; 100.0)	14	87.5 (50.0; 100.0)	0.21
M1	18	30.0 (25.0; 70.0)	39	50.0 (40.0; 60.0)	67	40.0 (20.0; 50.0)	10	50.0 (25.0; 75.0)	30	40.0 (30.0; 60.0)	14	40.0 (20.0; 60.0)	0.10
M2	18	77.5 (65.0; 80.0)	39	80.0 (70.0; 95.0)	67	70.0 (50.0; 80.0)	10	60.0 (50.0; 75.0)	30	70.0 (50.0; 80.0)	14	72.5 (50.0; 80.0)	0.04
M3	18	75.0 (60.0; 90.0)	39	70.0 (50.0; 80.0)	67	70.0 (50.0; 80.0)	10	80.0 (75.0; 80.0)	30	80.0 (60.0; 90.0)	14	72.5 (60.0; 80.0)	0.18
M4	18	60.0 (20.0; 80.0)	39	70.0 (50.0; 85.0)	67	60.0 (50.0; 75.0)	10	85.0 (75.0; 100.0)	30	65.0 (50.0; 80.0)	14	70.0 (50.0; 75.0)	0.07

Note: Only questionnaires with at least one or more answers, in addition to demographic characteristics, were included

* Comparison using Kruskal-Wallis test with significance established at $p < 0.05$

Table 4: Analysis of Injury Severity Score Based on Fracture Type, Neurologic Status, and Case-Specific Modifier by Experience

Type	___ < 5 years ___		___ 5 – 10 years ___		___ 11 – 20 years ___		___ > 20 years ___		• 10 vs. > 10 years
	N	Median (IQR)	N	Median (IQR)	N	Median (IQR)	N	Median (IQR)	p-value
A0	44	5.0 (1.0; 10.0)	59	5.0 (1.0; 10.0)	48	1.5 (0.0; 10.0)	27	5.0 (1.0; 10.0)	0.04
A1	44	20.0 (10.0; 27.5)	59	20.0 (10.0; 30.0)	49	10.0 (10.0; 20.0)	27	20.0 (10.0; 20.0)	0.02
A2	44	30.0 (30.0; 50.0)	59	40.0 (25.0; 50.0)	49	30.0 (20.0; 35.0)	27	30.0 (20.0; 50.0)	0.02
A3	44	40.0 (30.0; 50.0)	59	50.0 (35.0; 60.0)	49	50.0 (30.0; 60.0)	27	40.0 (30.0; 55.0)	0.80
A4	44	60.0 (50.0; 70.0)	59	65.0 (50.0; 80.0)	49	60.0 (50.0; 80.0)	27	60.0 (45.0; 80.0)	0.63
B1	44	67.5 (50.0; 75.0)	59	60.0 (50.0; 70.0)	49	55.0 (50.0; 60.0)	27	50.0 (30.0; 60.0)	0.03
B2	44	80.0 (70.0; 82.5)	59	80.0 (70.0; 90.0)	49	80.0 (70.0; 90.0)	27	70.0 (60.0; 80.0)	0.79
B3	44	80.0 (72.5; 90.0)	59	80.0 (70.0; 90.0)	49	80.0 (70.0; 95.0)	27	75.0 (65.0; 85.0)	0.99
C	44	100.0 (100.0; 100.0)	58	100.0 (100.0; 100.0)	49	100.0 (100.0; 100.0)	27	100.0 (100.0; 100.0)	0.53
F1	44	20.0 (12.5; 30.0)	59	20.0 (10.0; 30.0)	49	20.0 (10.0; 30.0)	27	20.0 (10.0; 20.0)	0.77
F2	44	40.0 (30.0; 52.5)	59	40.0 (30.0; 50.0)	49	30.0 (30.0; 50.0)	27	30.0 (20.0; 40.0)	0.04
F3	44	52.5 (40.0; 70.0)	59	60.0 (40.0; 70.0)	49	50.0 (40.0; 70.0)	27	50.0 (40.0; 60.0)	0.18
F4	44	100.0 (90.0; 100.0)	59	100.0 (80.0; 100.0)	49	100.0 (90.0; 100.0)	27	90.0 (80.0; 100.0)	0.84
N0	44	0.0 (0.0; 0.0)	59	0.0 (0.0; 0.0)	49	0.0 (0.0; 0.0)	26	0.0 (0.0; 0.0)	0.89
N1	44	20.0 (12.5; 25.0)	59	20.0 (10.0; 25.0)	49	20.0 (10.0; 40.0)	26	20.0 (10.0; 30.0)	0.29
N2	44	40.0 (30.0; 50.0)	59	40.0 (25.0; 50.0)	49	40.0 (20.0; 60.0)	26	40.0 (20.0; 50.0)	0.58
N3	44	80.0 (72.5; 100.0)	59	80.0 (70.0; 90.0)	49	90.0 (80.0; 100.0)	26	80.0 (70.0; 100.0)	0.02
N4	44	100.0 (80.0; 100.0)	59	100.0 (90.0; 100.0)	49	100.0 (90.0; 100.0)	26	100.0 (90.0; 100.0)	0.15
NX	44	90.0 (50.0; 100.0)	59	80.0 (50.0; 100.0)	49	80.0 (50.0; 100.0)	26	90.0 (50.0; 100.0)	0.34
M1	44	47.5 (25.0; 60.0)	59	40.0 (25.0; 60.0)	49	50.0 (30.0; 60.0)	26	40.0 (30.0; 50.0)	0.61
M2	44	70.0 (50.0; 90.0)	59	70.0 (50.0; 80.0)	49	75.0 (60.0; 80.0)	26	70.0 (50.0; 80.0)	0.47
M3	44	75.0 (55.0; 80.0)	59	70.0 (50.0; 80.0)	49	70.0 (60.0; 90.0)	26	72.5 (60.0; 80.0)	0.51
M4	44	70.0 (50.0; 90.0)	59	60.0 (40.0; 75.0)	49	60.0 (50.0; 80.0)	26	70.0 (50.0; 80.0)	0.94

Note: Only questionnaires with at least one or more answers, in addition to demographic characteristics, were included

* Comparison using Wilcoxon rank sum test with significance established at $p < 0.05$