

CORRESPONDENCE

Open Access



# Comment on “Comparison of analgesic efficacy between rectus sheath blockade, intrathecal morphine with bupivacaine, and intravenous patient-controlled analgesia in patients undergoing robot-assisted laparoscopic prostatectomy: a prospective, observational clinical study”

Lukas M. Löffel<sup>\*</sup> , Dominique Engel and Patrick Y. Wuethrich

Dear Editor.

We read with great interest the article “Comparison of analgesic efficacy between rectus sheath blockade, intrathecal morphine with bupivacaine, and intravenous patient-controlled analgesia in patients undergoing robot-assisted laparoscopic prostatectomy: a prospective, observational clinical study” by Shim et al. [1] Yet we think that there are relevant methodological issues and therefore we raise concerns about the potential for generalization of the results for other centers.

The observational nature of the trial with patients being able to choose their analgesic regimen themselves poses a significant risk of bias. The patients opting for intrathecal analgesia may be different in their general perception of pain and satisfaction compared to the other groups. According to the flowchart, only 103 patients had to be evaluated for inclusion, of which 13 dropped out because of fulfilling the exclusion criteria. In our experience, it seems unlikely that the remaining 90 participants could be distributed exactly equal to three groups of 30 the trial aimed for, even though they

were reported to be able to choose themselves unrestrictedly. The authors state that there were no adverse events such as respiratory depression, which is a well known complication of intrathecal morphine [2]. This occurs with a delay of up to 24 h postoperatively. How were patients monitored for this complication? Furthermore, it is not clear how the sample size analysis was performed. Shim et al. state that it was based on drug consumption of 20 patients on postoperative day 1. There is uncertainty how these patients were identified. Last but not least, the reviewers of this paper also raised concerns that the study is not comprehensive and that the conclusions cannot be sufficiently supported [3].

With our ongoing three-arm randomized controlled trial “Impact of Perioperative Analgesia in Prostatectomy Patients on Early Quality of Recovery” (“SPITALIDO”, registered on [clinicaltrials.gov](https://clinicaltrials.gov) on August 7, 2018 with the identifier [NCT03618693](https://clinicaltrials.gov/ct2/show/study?term=NCT03618693)) we hope to omit these methodological issues and to provide further data for the studied patient population [4].

## Authors' contributions

LML wrote the letter. LML, DE, PYW edited and revised the letter. All authors read and approved the final manuscript.

## Funding

Not applicable.

This comment refers to the article available at <https://doi.org/10.1186/s12871-020-01208-2>.

<sup>\*</sup> Correspondence: [lukas.loeffel@insel.ch](mailto:lukas.loeffel@insel.ch)

Department of Anaesthesiology and Pain Medicine, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Availability of data and materials**

Not applicable.

**Declarations****Ethics approval and consent to participate**

Not applicable.

**Consent for publication**

Not applicable.

**Competing interests**

The authors declare that they have no competing interests.

Received: 22 January 2021 Accepted: 10 March 2021

Published online: 17 March 2021

**References**

1. Shim JW, Cho YJ, Kim M, Hong SH, Moon HW, Hong SH, Chae MS, et al. Comparison of analgesic efficacy between rectus sheath blockade, intrathecal morphine with bupivacaine, and intravenous patient-controlled analgesia in patients undergoing robot-assisted laparoscopic prostatectomy: a prospective, observational clinical study. *BMC Anesthesiol.* 2020;20(1):291. <https://doi.org/10.1186/s12871-020-01208-2>.
2. Meylan N, Elia N, Lysakowski C, Tramèr MR. Benefit and risk of intrathecal morphine without local anaesthetic in patients undergoing major surgery: meta-analysis of randomized trials. *Br J Anaesth.* 2009;102(2):156–67. <https://doi.org/10.1093/bja/aen368>.
3. BMC Anesthesiology 2020. Available from: <https://bmcanesthesiol.biomedcentral.com/articles/10.1186/s12871-020-01208-2/peer-review>, accessed Nov 26, 2020.
4. SPITALIDO Trial. Available from: <https://clinicaltrials.gov/ct2/show/NCT03618693?term=spitalido&draw=2&rank=1>, accessed Nov 26, 2020.

**Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

