



Effect of Brønsted acids on the thiophenol-mediated radical addition–translocation–cyclization process for the preparation of pyrrolidine derivatives

Valentin Soulard, Fabrice Dénès & Philippe Renaud

To cite this article: Valentin Soulard, Fabrice Dénès & Philippe Renaud (2016) Effect of Brønsted acids on the thiophenol-mediated radical addition–translocation–cyclization process for the preparation of pyrrolidine derivatives, Free Radical Research, 50:sup1, S2-S5, DOI: 10.1080/10715762.2016.1223294

To link to this article: <https://doi.org/10.1080/10715762.2016.1223294>



© 2016 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 24 Oct 2016.



Submit your article to this journal [↗](#)



Article views: 1015



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 3 View citing articles [↗](#)

Effect of Brønsted acids on the thiophenol-mediated radical addition–translocation–cyclization process for the preparation of pyrrolidine derivatives

Valentin Soulard^a, Fabrice Dénès^b  and Philippe Renaud^a 

^aDepartment of Chemistry and Biochemistry, University of Bern, Bern, Switzerland; ^bCEISAM UMR 6230 – UFR des Sciences et des Techniques, Université de Nantes, Nantes, Cedex 3, France

ABSTRACT

A thiophenol-mediated method for the conversion of propargylamines to pyrrolidines under acidic conditions is described. This cascade reaction involves addition of a thiyl radical to the terminal alkyne followed by a 1,5-hydrogen transfer (radical translocation) and a rapid cyclization affording the pyrrolidine ring. Our studies reveal that complete protonation of the tertiary amine with 10 equivalents of trifluoroacetic acid avoids undesired hydrogen atom abstractions by the thiyl radicals.

ARTICLE HISTORY

Received 5 July 2016
Revised 5 August 2016
Accepted 6 August 2016

KEYWORDS

Radical reaction; hydrogen atom transfer; C–H activation; thiyl radicals; propargylamines; pyrrolidines

Radical reactions are now part of the arsenal of reactions that synthetic organic chemists use for the preparation of complex molecules [1,2]. Their mildness and functional group tolerance make them particularly attractive for the synthesis of highly functionalized compounds and their reactivity has permitted some unusual transformations. Among all radical reactions, the activation of aliphatic C–H bonds via hydrogen atom transfer (HAT) is a truly unique synthetic tool [3–7]. The preparation of five-membered ring via radical translocation–cyclization (RTC) processes involving transient alkenyl radicals proved to be a particularly efficient process [8]. Alkenyl radicals can be generated directly from alkenyl halides [9,10] or by radical addition to a terminal alkyne [11–19]. The thiophenol-mediated radical addition–translocation–cyclization (RATC) process [20–23] is preparatively very useful since terminal alkynes are easily accessible [24]. It has been used to prepare simple and fused cyclopentanes as well as spirocyclic compounds such as (–)-erythrodiene [22,23,25]. Starting from homopropargylic amines and amides, it was also possible to prepare the heterocyclic systems related to the indolizidine and pyrrolizidine alkaloid skeleton (Scheme 1) [26].


However, all attempts to extend the thiophenol-mediated RATC reaction to the easily prepared propargylic amines failed and a complex mixture of products was obtained (Scheme 2). This result was interpreted as a consequence of undesired hydrogen atom abstraction from the propargylic position leading to stabilized 1-aminosubstituted propargylic radicals. This degradation pathway is well established since Bertrand et al. have described the isomerization of allylic amines under closely related reaction conditions [27,28].

Interestingly, Bietti et al. have recently demonstrated that the rate of hydrogen atom abstraction from the α -position of tertiary alkylamines by the cumyloxy radical can be lowered by protonation of the amine (Scheme 3) [5,29]. The authors showed that the rate constant for hydrogen atom abstraction from the α -C–H bonds decreased drastically upon protonation of the nitrogen. This observation was rationalized by a decrease in hyperconjugative overlap between the α -C–H σ^* orbital and the nitrogen lone-pair. Therefore, we decided to examine the effect of acid additives on the thiophenol mediated RATC process involving propargylic amines.


CONTACT Philippe Renaud  philippe.renaud@dcb.unibe.ch  Department of Chemistry and Biochemistry, University of Bern, Freiestrasse 3, CH-3012 Bern, Switzerland; Fabrice Dénès  fabrice.denes@univ-nantes.fr  CEISAM UMR 6230 – UFR des Sciences et des Techniques, Université de Nantes, 2 rue de la Houssinière, BP 92208, 44322 Nantes, Cedex 3, France

© 2016 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group
This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.


N,N-diisobutylpropargylamine (**1**) was chosen as a model substrate for the preliminary study. This compound was easily prepared by the alkylation of diisobutylamine with propargyl bromide in the presence of potassium carbonate. In order to be compatible with the presence of acid additives, acetonitrile was used as a solvent instead of *tert*-butanol employed for the thiophenol-mediated RATC processes [23]. For the optimization process, reaction yields were determined by gas chromatography and the results are summarized in Table 1. Without additive, the starting material was fully consumed and a complex mixture of products was obtained, with the expected cyclopentane product **2** obtained in only 4% yield. Adding four equivalents of acetic acid did not improve the reaction (Table 1, entry 2).



Scheme 1



Scheme 2




Scheme 3

This is not fully surprising since in acetonitrile, acetic acid ($pK_a(\text{CH}_3\text{CN}) = 22.3$) is less acidic than the trialkylammonium ion ($pK_a(\text{CH}_3\text{CN}) = 18.5$) [30]. However, running the reaction in pure acetic acid as a solvent led to an increased yield of 25% for **2** (Table 1, entry 3). Stronger Brønsted acids were tested next. Trifluoroacetic acid ($pK_a(\text{CH}_3\text{CN}) = 12.7$) [30] influenced positively the reaction when 1.1 equivalents were used (Table 1, entry 4, 18% yield). The yield increased further when five equivalents were used (Table 1, entry 5, 54% yield) and the best yield was achieved with 10 equivalents (Table 1, entry 6, 78% yield). Larger excesses (Table 1, entries 7 and 8) did not increase further the yield of the reaction. The use of trifluoromethanesulfonic acid ($\text{CF}_3\text{SO}_3\text{H}$, $pK_a(\text{CH}_3\text{CN}) = 0.7$) [30] was examined next. A low yield of 16% was obtained with 0.9 equivalent (Table 1, entry 9). With 1.0 and 1.5 equivalents, the reaction proceeded more efficiently (Table 1, entries 10 and 11, 68% and 70% yield, respectively). Larger excess of trifluoromethanesulfonic acid did not help. Indeed, in the presence of 10 equivalent of trifluoromethanesulfonic acid, the product was formed in 49% yield (Table 1, entry 12). Based on all these experiments, the use of trifluoroacetic acid (10 equivalents) was selected to run the final solvent optimization. Acetic acid and 1,2-dichloroethane were tried first and gave low yields (Table 1, entries 13 and 14). Benzene (Table 1, entry 15) gave a better result (65% yield) but still inferior to the one obtained in acetonitrile (Table 1, entry 6).

Table 1. Thiophenol promoted RATC reaction with model substrate **1**.

Entry	solvent	acid	equivalents	Yield (GC)
1	CH_3CN	–	–	4%
2	CH_3CN	$\text{CH}_3\text{CO}_2\text{H}$	4	4%
3	$\text{CH}_3\text{CO}_2\text{H}$	–	solvent	25%
4	CH_3CN	$\text{CF}_3\text{CO}_2\text{H}$	1.1	18%
5	CH_3CN	$\text{CF}_3\text{CO}_2\text{H}$	5	54%
6	CH_3CN	$\text{CF}_3\text{CO}_2\text{H}$	10	78%
7	CH_3CN	$\text{CF}_3\text{CO}_2\text{H}$	20	74%
8	CH_3CN	$\text{CF}_3\text{CO}_2\text{H}$	50	73%
9	CH_3CN	$\text{CF}_3\text{SO}_3\text{H}$	0.9	16%
10	CH_3CN	$\text{CF}_3\text{SO}_3\text{H}$	1	68%
11	CH_3CN	$\text{CF}_3\text{SO}_3\text{H}$	1.5	70%
12	CH_3CN	$\text{CF}_3\text{SO}_3\text{H}$	10	49%
13	$\text{CH}_3\text{CO}_2\text{H}$	$\text{CF}_3\text{CO}_2\text{H}$	10	25%
14	$\text{ClCH}_2\text{CH}_2\text{Cl}$	$\text{CF}_3\text{CO}_2\text{H}$	10	14%
15	benzene	$\text{CF}_3\text{CO}_2\text{H}$	10	65%



Scheme 4

The fact that good yields were obtained either with a large excess of $\text{CF}_3\text{CO}_2\text{H}$ (10 equivalents) or with at least one equivalent of $\text{CF}_3\text{SO}_3\text{H}$ indicates that full protonation of the amine is necessary for an efficient reaction. As demonstrated by the reaction run with only 0.9 equivalent of $\text{CF}_3\text{SO}_3\text{H}$ (Table 1, entry 9), a small amount of the free amine (10 mol% in this case) is sufficient to cause a massive decrease in the yield.

The best reaction conditions identified for substrate **1** based on GC yields (Table 1, entry 6), that is, the use of an excess of thiophenol (2 equivalents), AIBN (2 equivalents) and trifluoroacetic acid (10 equivalents) in refluxing acetonitrile as solvent, were used for preparative reactions using the four substrates **1**, **3-5** (Scheme 4). In all four cases, the desired cyclized products were isolated in good yields despite the fact the cyclic amines **2**, **6-8** are difficult to purify by flash chromatography due to their basic character.

In conclusion, the results presented here demonstrate that the concept of acid-mediated inhibition of undesired hydrogen atom transfers between cumyloxyl radicals and tertiary amines developed by Bietti and Salamone can be extended to thiyl radicals and it represents a key strategy to control reactions of tertiary amines prone to facile hydrogen atom abstraction. When a complex reaction cascade such as the RATC process is involved, full protonation of the amino group by a 10-fold excess of trifluoroacetic acid is required to overcome the strong hydrogen atom donor ability of propargylic tertiary amines.

Disclosure statement

The authors report no declarations of interest. The authors alone are responsible for the content and writing of the paper.

Funding

The Swiss National Science Foundation [Project 200020_152782], the State Secretariat for Education, Research and Innovation [SBFI Nr. C14.0096] and the Agence National de La Recherche (Project ANR Blanc 13 BS07-0010-01 "UZFuL-Chem") are gratefully acknowledged for financial support. The COST Action CM1201 has generously supported a short-term scientific mission of Valentin Soulard. We are also very thankful to Massimo Bietti and Michela Salamone for stimulating discussion during scientific meetings of the COST Action CM1201.

Agence Nationale de la Recherche, 10.13039/501100001665 [ANR Blanc 13 BS07-0010-01 "UZFuL-Chem"], European Cooperation in Science and Technology, 10.13039/501100000921 [CM1201], Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung, 10.13039/501100001711 [200020_152782], Staatssekretariat für Bildung, Forschung und Innovation, 10.13039/501100007352 [C14.0096].

ORCID

Fabrice Dénès  <http://orcid.org/0000-0002-9791-3177>
Philippe Renaud  <http://orcid.org/0000-0002-9069-7109>

References

1. Chatgililoglu C, Studer A. Encyclopedia of radicals in chemistry, biology, and materials. Chichester, West Sussex; Hoboken, NJ: John Wiley & Sons; 2012.
2. Renaud P, Sibi MP. Radicals in organic synthesis. 1st ed. Weinheim; New York, NY: Wiley-VCH; 2001.
3. Feray L, Kuznetsov N, Renaud P. Hydrogen atom abstraction. Radicals in organic synthesis. Weinheim: Wiley-VCH Verlag GmbH; 2001: 246–278.
4. Robertson J, Pillai J, Lush RK. Radical translocation reactions in synthesis. Chem Soc Rev 2001;30:94–103.
5. Salamone M, Bietti M. Tuning reactivity and selectivity in hydrogen atom transfer from aliphatic C-H bonds to

- alkoxyl radicals: role of structural and medium effects. *Acc Chem Res* 2015;48:2895–2903.
- Nechab M, Mondal S, Bertrand MP. 1,*n*-Hydrogen-atom transfer (HAT) reactions in which $n \neq 5$: an updated inventory. *Chemistry* 2014;20:16034–16059.
 - Cekovic Z. Reactions of carbon radicals generated by 1,5-trans-position of reactive centers. *J Serb Chem Soc* 2005;70:287–318.
 - Dénès F, Beaufils F, Renaud P. Preparation of five-membered rings via the translocation-cyclization of vinyl radicals. *Synlett* 2008;2008:2389–2399.
 - Curran DP, Shen W. Radical translocation reactions of vinyl radicals: substituent effects on 1,5-hydrogen-transfer reactions. *J Am Chem Soc* 1993;115:6051–6059.
 - Curran DP, Kim D, Liu HT, Shen W. Translocation of radical sites by intramolecular 1,5-hydrogen atom transfer. *J Am Chem Soc* 1988;110:5900–5902.
 - Heiba E-Al, Dessau RM. Free-radical isomerization. I. Novel rearrangement of vinyl radicals. *J Am Chem Soc* 1967;89:3772–3777.
 - Beaufils F, Dénès F, Renaud P. Dimethyl phosphite mediated hydrogen atom abstraction: a tin-free procedure for the preparation of cyclopentane derivatives. *Angew Chem Int Ed Engl* 2005;44:5273–5275.
 - Lamarque C, Beaufils F, Dénès F, Schenk K, Renaud P. Preparation of 5-membered rings via radical addition-translocation-cyclization (RATC) processes mediated by diethyl thiophosphites. *Adv Synth Catal* 2011;353:1353–1358.
 - Bogen S, Fensterbank L, Malacria M. Study of a radical cyclizations cascade leading to bicyclo[3.1.1]heptanes. *J Org Chem* 1999;64:819–825.
 - Bertrand MP, Crich D, Nougier R, Samy R, Stien D. Diastereoselective cyclizations of 1,3-dioxan-2-yl radicals: chiral acyl radical equivalents. *J Org Chem* 1996;61:3588–3589.
 - Clive DLJ, Cantin M. 5-Endo-trigonal radical cyclisation—a general procedure for making five-membered rings. *J Chem Soc Chem Commun* 1995:319–320.
 - Martinez-Grau A, Curran DP. Stereoselective synthesis of alkenes by using unimolecular chain transfer reactions of silicon hydrides. *J Org Chem* 1995;60:8332–8333.
 - Bosch E, Bachi MD. Free-radical annelation in the synthesis of bicyclic beta-lactams. 7. A one-pot, four-step, sequential reaction. *J Org Chem* 1993;58:5581–5582.
 - Alcaide B, Rodríguez-Campos IM, Rodríguez-López J, Rodríguez-Vicente A. Stereoselective synthesis of fused bicyclic β -lactams through radical cyclization of enyne-2-azetidiones. *J Org Chem* 1999;64:5377–5387.
 - Burke SD, Jung KW. 2,3-Disubstituted tetrahydrofuran synthesis via radical and anionic cyclization. *Tetrahedron Lett* 1994;35:5837–5840.
 - Renaud P, Beaufils F, Feray L, Schenk K. Diastereoselective radical mediated hydrogen atom abstraction. *Angew Chem Int Ed Engl* 2003;42:4230–4233.
 - Beaufils F, Dénès F, Renaud P. Thiophenol-mediated hydrogen atom abstraction: an efficient tin-free procedure for the preparation of cyclopentane derivatives. *Org Lett* 2004;6:2563–2566.
 - Beaufils F, Dénès F, Becattini B, Renaud P, Schenk K. Thiophenol-mediated 1,5-hydrogen atom abstraction: easy access to mono- and bicyclic compounds. *Adv Synth Catal* 2005;347:1587–1594.
 - Dénès F, Pichowicz M, Povie G, Renaud P. For an extensive review and the use of thiyl radical in organic synthesis. Thiyl radicals in organic synthesis. *Chem Rev* 2014;114:2587–2693.
 - Lachia M, Dénès F, Beaufils F, Renaud P. Highly diastereoselective formation of spirocyclic compounds via 1,5-hydrogen transfer: a total synthesis of (-)-erythrodiene. *Org Lett* 2005;7:4103–4106.
 - Dénès F, Beaufils F, Renaud P. Thiophenol-mediated 1,5-hydrogen transfer for the preparation of pyrrolizidines, indolizidines, and related compounds. *Org Lett* 2007;9:4375–4378.
 - Bertrand MP, Escoubet S, Gastaldi S, Timokhin VI. Tandem radical and non-radical reactions mediated with thiols—a new method of cleavage of allylic amines. *Chem Commun* 2002:216–217.
 - Escoubet S, Gastaldi S, Timokhin VI, Bertrand MP, Siri D. Thiyl radical-mediated cleavage of allylic C–N bonds: scope, limitations, and theoretical support to the mechanism. *J Am Chem Soc* 2004;126:12343–12352.
 - Salamone M, Giammaroli I, Bietti M. Tuning hydrogen atom abstraction from the aliphatic C–H bonds of basic substrates by protonation. Control over selectivity by C–H deactivation. *Chem Sci* 2013;4:3255.
 - Espinosa S, Bosch E, Rosés M. Acid–base constants of neutral bases in acetonitrile–water mixtures. *Anal Chim Acta* 2002;454:157–166.