Corrigendum to

Short Communication

A second *KRT71* allele in curly coated dogs

Anina Bauer[1,2], Sheida Hadji Rasouliha[1,2], Magdalena T. Brunner[2,3], Vidhya Jagannathan[1,2], Inga Bucher[2,4], Jeanette Bannoehr[2,4,5], Katarina Varjonen[6,7], Ross Bond[6], Kerstin Bergvall[8], Monika M. Welle[2,3], Petra Roosje[2,4], Tosso Leeb[1,2]

[1]Institute of Genetics, Vetsuisse Faculty, University of Bern, 3001 Bern, Switzerland
[2]Dermfocus, Vetsuisse Faculty, University of Bern, 3001 Bern, Switzerland
[3]Institute of Animal Pathology, Vetsuisse Faculty, University of Bern, 3001 Bern, Switzerland
[4]Division of Clinical Dermatology, Department of Clinical Veterinary Medicine, Vetsuisse Faculty, University of Bern, 3001 Bern, Switzerland
[5]Dermatology Department, Animal Health Trust, Lanwades Park, Kentford, Newmarket, Suffolk CB8 7UU, United Kingdom
[6]Department Veterinary Clinical Sciences, Royal Veterinary College, Hatfield AL9 7TA, United Kingdom
[7]Anicura Albano Animal Hospital, 18236 Danderyd, Sweden
[8]Department of Clinical Sciences, Swedish University of Agricultural Sciences, Box 7054, Uppsala, 750 07, Sweden

Address for correspondence

Tosso Leeb
Institute of Genetics
Vetsuisse Faculty
University of Bern
Bremgartenstrasse 109a
3001 Bern
Switzerland

Phone: +41-31-6312326
E-mail: Tosso.Leeb@vetsuisse.unibe.ch

Corrigendum

Bauer, A., Hadji Rasouliha, S., Brunner, M.T., Jagannathan, V., Bucher, I., Bannoehr, J., Varjonen, K., Bond, R., Bergvall, K., Welle, M.M., Roosje, P., Leeb, T. (2019). A second KRT 71 allele in curly coated dogs. *Animal genetics, 50*(1), 97-100. https://doi.org/10.1111/age.12743

In the above article published in *Animal Genetics* (https://doi.org/10.1111/age.12743), the authors identified a new *KRT71* allele in curly coated dogs and reported *KRT71* diplotypes for 1286 dogs in Table 1.

In the meantime, the authors discovered that due to redundantly genotyped duplicate samples, the true number of dogs is only 1281 - the number of Lagotto Romagnolo dogs with the c^1/c^1 diplotype was incorrectly reported as 559 instead of 555, and the number with the c^1/c^2 diplotype was incorrectly reported as 172 instead of 171. Furthermore, the diplotypes of 44 curly coated Barbets were incorrectly given as *wt/wt* instead of c^1/c^1. These errors do not affect the main conclusions of the publication. A corrected version of Table 1 is given below.

Table 1. *KRT71* diplotypes in 1281 dogs from 15 different breeds with curly or wavy hair. We refer to the mutant alleles at the previously described c.451C>T missense variant as c^1 and the c.1266_1273delinsACA variant as c^2. In five of the tested breeds, both alleles were segregating.

Breed	***KRT71* diplotype**							**N**
	wt/wt	c^1/wt	c^1/c^1	c^2/wt	c^1/c^2	c^2/c^2	c^1/c^1c^2	
Airedale Terrier			5					5
Barbet		2	44					46
Bergamasco Shepherd dog	8	4						12
Bolonka Zwetna	8	8	8					24
Cão de Serra de Aires	15							15
Chesapeake Bay Retriever			3		9	1[1]		13
Curly Coated Retriever						125		125
Lagotto Romagnolo		14	555	3	171	12[1]		755
Mudi			11		18	3	2	34
Perro de Agua Español			73		20	2[1]		95
Poodle			89		6	1[2]		96

Portugese Water dog		3		3
Puli			1	1
Schapendoes	44	4		48
Soft Coated Wheaten Terrier	7	2		9
				1281

[1]The cohort contained 3 dogs, which had been diagnosed with follicular dysplasia. They all had the diplotype c^2/c^2.
[2]This poodle had pronounced alopecia on the body and an almost completely hairless tail. Unfortunately no histopathological examination of a skin biopsy was performed to confirm the suspected follicular dysplasia.