

The effect of a succession of ocean ventilation changes on radiocarbon

T. F. Stocker, D. G. Wright

published in *Radiocarbon* 40, 351 – 366, (1998)

Abstract

Using the model of *Stocker and Wright* [1996], we investigate the effect of a succession of ocean ventilation changes on the atmospheric concentration of radiocarbon, $\Delta^{14}\text{C}^{\text{atm}}$, the surface reservoir ages, the top-to-bottom age differences, and the calendar-radiocarbon age relationships in different regions of the ocean. The model includes a representation of the cycling of radiocarbon through the atmosphere, the ocean and the land biosphere. Ocean ventilation changes are triggered by increasing rates of freshwater discharge into the North Atlantic which are determined according to a simple feedback mechanism between the melting rates and the climatic state of the North Atlantic region. The results demonstrate that ventilation changes can cause $\Delta^{14}\text{C}^{\text{atm}}$ fluctuations of 25‰, surface reservoir age fluctuations of 100 yr (200 yr) in the Pacific (Atlantic) and top-to-bottom age variations of 500 yr (1000 yr) in the Pacific (Atlantic). We also show that ^{14}C age estimates based on marine organisms which live in the near-surface region of the ocean and take up the signal of surface- ^{14}C can result in apparent age reversals if the assumption of a constant reservoir age is made.

1 Introduction

High-resolution radiocarbon dating has been recognized as an important tool in the investigation of climatic change. Its primary use is in the estimation of chronologies for past events, but radiocarbon dates also contain information

on changes in the production rate of atmospheric radiocarbon, the globally integrated ventilation rate of the ocean, atmosphere-ocean exchange rates and land biomass. Radiocarbon chronologies [Becker *et al.*, 1991; Björck *et al.*, 1996; Hughen *et al.*, 1998] are particularly suitable for the reconstruction of such changes as they allow us to construct time series of $\Delta^{14}\text{C}^{\text{atm}}$ in the relevant reservoir. Climate reconstructions based on dynamical models which include radiocarbon as a tracer can use this information to provide a quantitative test of the validity of these reconstructions and the assumptions on which they are based (e.g. reservoir ages). In addition, such models can provide estimates of anomalous variations in radiocarbon age that are associated with climatic variations.

The purpose of this paper is to estimate the effect of a succession of ocean ventilation changes on $\Delta^{14}\text{C}^{\text{atm}}$, and to show how quantities that are derived from high-resolution radiocarbon dating can be affected by such events. The quantities considered are surface reservoir ages, almost always assumed constant in paleoceanographic reconstructions, top-to-bottom age differences and chronologies. It should be noted that our discussion pertains primarily to marine or lacustrine paleoclimatic archives which are subject to a reservoir age effect. This problem does not arise in tree rings since, in this case, accurate reconstruction of $\Delta^{14}\text{C}^{\text{atm}}$ can be determined by taking into account changes of $\delta^{13}\text{C}$.

High-resolution radiocarbon records contain numerous phases of faster and slower advance of the radiocarbon “clock” indicating increasing and decreasing concentrations of the atmospheric concentration of radiocarbon. Part of these fluctuations can be associated with cyclic changes in the production rate of radiocarbon due to changes in the Earth’s magnetic field or cosmic radiation [Damon *et al.*, 1978; Stuiver and Braziunas, 1993]. However, some of these variations are particularly rapid or occur in association with contemporaneous climatic changes recorded in these same archives, suggesting that additional processes influence the radiocarbon variations. Recently documented examples include the changes of tree ring width in the German chronology [Björck *et al.*, 1996] and a phase of slower increase of the radiocarbon “clock” at the end of Younger Dryas (YD), and changes of the color scale and a contemporaneous strong speed-up of the radiocarbon “clock” during the beginning of Younger Dryas as found in the varved marine sediments of Cariaco Basin [Hughen *et al.*, 1998]. Such results suggest a climatic origin for some part of these $\Delta^{14}\text{C}^{\text{atm}}$ fluctuations.

Stocker and Wright [1996] and *Mikolajewicz* [1998] have investigated the effect of a collapse of the Atlantic thermohaline circulation on $\Delta^{14}\text{C}^{\text{atm}}$ and found fluctuations of up to 30‰ and, if extended changes in the sea ice cover occur, up to 60‰. When the circulation is strongly reduced or shut down, $\Delta^{14}\text{C}^{\text{atm}}$ increases because the radiocarbon that is continually produced in the atmosphere is less efficiently taken up by the ocean, and hence a greater fraction of it remains in the atmospheric and biospheric reservoirs. When the circulation resumes, the deep ocean, that is by now depleted in radiocarbon, is ventilated again and efficiently takes up the excess radiocarbon from the atmosphere. This leads to a period of reduced radiocarbon age variation in both the atmosphere and the upper ocean (a so called age plateau). Fluctuations of $\Delta^{14}\text{C}^{\text{atm}}$ as reconstructed by *Hughen et al.* [1998] during the beginning of YD and by *Björck et al.* [1996] at the end of YD are consistent with rapid turn off and on of the thermohaline circulation. The situation, however, appears to be more complicated during the intervening years. *Hughen et al.* [1998] found that $\Delta^{14}\text{C}^{\text{atm}}$ started to decrease already soon after the initiation of YD. This suggests that a new ventilation source must have become active during YD; possibilities are intermediate water formation at mid-latitudes in the North Atlantic or the Southern Ocean. Detailed simulations of the YD event that reflect such changes have not yet been achieved by dynamical models. Apart from model shortcomings, we urgently need to know where, when and how much meltwater was discharged into the ocean during deglaciation [*Clark et al.*, 1996].

2 Model Description and Experiments

We use a zonally averaged ocean-atmosphere climate model including a simple representation of the land biosphere. The ocean component includes a representation of the dynamics [*Wright et al.*, 1995] of the thermohaline circulation in three basins (Pacific, Atlantic, Indian) interconnected by a circumpolar Southern Ocean. The atmosphere is represented by a linear energy balance formulation [*Stocker et al.*, 1992] and the biosphere is described by a simple four-box model *Siegenthaler and Oeschger* [1987]. Radiocarbon is included as a tracer which is produced in the atmosphere and exchanged between all three reservoirs. The production rate is determined from the steady state conditions at the end of a run in which atmospheric radiocarbon was

held fixed at $\Delta^{14}\text{C}^{\text{atm}}=0\text{‰}$. Subsequently, the rate of production was held constant so that the changes in radiocarbon considered here are entirely due to changes of the ventilation rate of the ocean. *Stocker and Wright* [1996] describe the model in detail, and the parameter values used here are identical to those used in this earlier study.

In our previous experiments we treated the freshwater flux perturbation as an externally specified forcing: feedback mechanisms, e.g. between the Northern Hemisphere temperatures and melting rates were ignored. Here we consider a crude feedback mechanism defined as follows:

$$F(t) = \begin{cases} R \cdot (t - t_0), & \text{if } F_H(32^\circ\text{N}) \geq 0.1\text{PW}, \\ 0, & \text{otherwise,} \end{cases}$$

where $F(t)$ is the perturbation freshwater flux which we take to be discharged at 50°N in the Atlantic basin. When the meridional heat flux in the Atlantic is small or absent (collapsed circulation) there is no melting, i.e. $F(t) = 0$. When the Atlantic heat flux at 32°N , $F_H(32^\circ\text{N})$, first exceeds 0.1PW ($1\text{PW}=10^{15}\text{W}$) at $t = t_0$, melting begins and it increases at the constant rate R (in m^3s^{-2}) as long as the heat flux remains above 0.1PW . This represents the simplest possible feedback mechanism between a climatic variable simulated by the model and the freshwater perturbation. Although we cannot give a quantitative justification for this choice, we note that a binge-purge mechanism was proposed for partly disintegrating ice sheets [*MacAyeal*, 1993b; *MacAyeal*, 1993a] with the effect that ice discharge rates increase progressively with time. For the present experiments we select $R = 2.4 \cdot 10^{-6}\text{m}^3\text{s}^{-2} = 0.075\text{Sv/kyr}$ ($1\text{Sv} = 10^6\text{m}^3\text{s}^{-1}$). Note that the continually increasing rate of input of fresh water will eventually cause the Atlantic overturning circulation to collapse. At this time, the supply of heat to the high northern latitudes will be strongly reduced, and melting will be stopped.

It is not immediately clear what the long term effect of this feedback mechanism on the deep circulation will be; there are several possibilities that might be consistent with the model dynamics. First, a quasi-steady state could develop in which the fresh water perturbation keeps the overturning circulation fluctuating around small values with meridional heat fluxes on the order of 0.1PW . A second possibility is that the circulation might break down completely and a new steady state establish with little or no Atlantic overturning. In this scenario, the heat flux would never again exceed the

critical value of 0.1 PW and there would be no further melting. The third possibility is that the circulation might collapse but recover. As the circulation recovers and the heat flux exceeds 0.1 PW, there might be little effect of the meltwater input until it grows to a critical value at which the circulation collapses again, and the cycle repeats.

The third possibility discussed above depends critically on the recovery phase, which might, for example, result from the significant cooling of high northern latitudes that would accompany the collapse of the overturning circulation, or it might be induced by increased high-latitude surface salinity after the meltwater input is terminated. In this regard, it should be noted that recovery is encouraged by the fact that the meltwater input is terminated as soon as the circulation collapses. This is a significant difference from previous experiments in which the meltwater input was specified independently of the model's climatic state.

Fig. 1 illustrates a typical model response, which is clearly of the third type. Meltwater input is initiated at $t=1000$ kyr and increases linearly at the specified rate of 0.075 Sv/kyr. After about 2 kyr the Atlantic circulation decreases abruptly (Fig. 1a), then recovers temporarily and completes its collapse by the time the meltwater input reaches about 0.26 Sv. At this point, the meltwater input is reduced to zero and soon after the circulation recovers rapidly to maximum strength. Subsequently, the cycle is repeated almost identically. Thus, the model responds with a succession of gradual coolings and abrupt warmings that are strongest in the northern hemisphere. The time scale of the oscillation is determined by R , but the qualitative nature of the variability is not sensitive to the value of R .

The model atmospheric temperature at 72°N gradually decreases as the meltwater input increases until the circulation collapses, at which point the temperature decreases abruptly, increases again as the circulation recovers and then decreases to near zero as the circulation completes its collapse (Fig. 1b). Once the Atlantic thermohaline circulation shut down and the meridional heat flux drops below 0.1 PW, the meltwater input stops. A few hundred years later, the circulation resumes again, resulting in an abrupt warming. It is remarkable that time scales of cooling and warming much like those seen in Fig. 1b are found to be a common feature of the Dansgaard/Oeschger cycles found in marine sediments [Bond *et al.*, 1993] and Greenland ice cores [GRIP Members, 1993]. In particular, our model results suggest that warming events follow times of maximum discharge with a lag,

a scenario which is not unlike what is indicated by the paleoclimatic record of Heinrich events.

The abrupt changes in ocean circulation are reflected in associated changes in the content of atmospheric radiocarbon (Fig. 1c); the signal is low-pass filtered with respect to high-latitude temperature (Fig. 1b) due to the damping effects of exchanges of radiocarbon between the surface reservoir and both the atmosphere and the deep ocean. $\Delta^{14}\text{C}^{\text{atm}}$ increases during the time of gradual cooling in the North Atlantic and continues to increase after the collapse until the circulation recovers. The maximum increase of $\Delta^{14}\text{C}^{\text{atm}}$ is about 25‰. The abrupt warming is due to the reinitiation of the overturning circulation, which also initiates a rapid decrease of $\Delta^{14}\text{C}^{\text{atm}}$ and leads to a period of reduced speed of the radiocarbon “clock” (a plateau).

3 Results

We now discuss quantities related to radiocarbon that can be measured in marine paleoclimatic archives. We select three “coring sites” in the model: 39°N in both Pacific and Atlantic and 60°N in the latter. The maximum depth across each of these sections is 4000 m, but we assume that the cores are taken at a local water depth of 2250 m, and we consider time series of surface reservoir ages and top-to-bottom age differences that would be recorded at these sites.

Fig. 2 shows time series of the surface reservoir age at the three “coring sites”. It is evident that these ages do not remain constant but vary by 100 to 200 years depending on the location. Reduced deep ventilation results in higher radiocarbon content of the atmosphere. This signal is rapidly transmitted to the surface waters which lowers the apparent age and consequently speeds up the radiocarbon “clock” during the adjustment period (Fig. 2a, b). Reinitiation of the ventilation increases the surface reservoir ages again. The increase is very rapid in the Atlantic where the strongest circulation changes occur. At high latitudes in the Atlantic, this increase occurs already during the cold phase because of the large changes in the vertical mixing ((Fig. 2)c). This is consistent with the findings of *Bard et al.* [1994] who report an increase of the surface reservoir age during the cooling phase of YD. However, the increase is much smaller than in earlier experiments [*Stocker and Wright*, 1996, Fig. 13] because here the maximum increase of $\Delta^{14}\text{C}^{\text{atm}}$ is smaller (cf.,

standard experiment C2 of *Stocker and Wright, 1996*).

Although the Pacific experiences only small circulation changes, Fig. 2a shows that the radiocarbon signal is still substantial there. This is primarily due to the fact that the changes in atmospheric radiocarbon associated with the reduced ventilation are efficiently transmitted to the ocean surface reservoir all around the globe. This again suggests that YD-type signals, which originate in the North Atlantic, are also to be expected in remote areas, since they can be efficiently transmitted through the atmosphere [*Stocker and Wright, 1996; Mikolajewicz et al., 1997*].

The temporal evolution of top-to-bottom age differences is presented in Fig. 3. In the Pacific the age difference increases by about 400 years during the cooling. About 100 years can be explained by a decrease of the surface reservoir age; the rest is due to the cessation of the inflow of younger Atlantic water into the deep Pacific. The two Atlantic sites exhibit more dramatic variations: age differences increase by about 1000 years, most of which is due to the ageing of the deep Atlantic associated with the reduced exchange with the surface reservoir when the Atlantic overturning circulation is sluggish (Fig. 3b, c). It must be noted, however, that this increase during the cooling period, and the subsequent rapid decrease following the warming happen within a relatively short time interval. Therefore, high-resolution radiocarbon dating (resolution of a century or better) will be required in order to resolve this signal. Changes in regions away from the North Atlantic occur on somewhat slower time scales but are significantly reduced in magnitude (Fig. 3a).

Finally, we address the possibility of apparent age reversals. These can occur when the surface reservoir ages increase rapidly at the time of switch-on of the circulation. This period is associated with increasing levels of convection which mixes up deep waters that are depleted in radiocarbon. In order to quantify the maximum possible effects, we base our considerations on the phase of rapid switch-on of the circulation in experiment I5 of *Stocker and Wright [1996]* (their Fig. 12d). This experiment shows the largest transient changes in $\Delta^{14}\text{C}^{\text{atm}}$ (up to +58‰), because of enhanced sea ice cover, and hence reduced air-sea exchange, during the time of collapsed circulation.

Fig. 4 shows three chronologies of radiocarbon age versus real time including a climate indicator (North Atlantic sea surface temperature at 60°N). The atmospheric chronology is most often discussed because of the absolute time scales provided by tree rings and varved lake and marine sediments.

The model's atmospheric chronology is shown in Fig. 4a. A plateau of about 290 years is visible which straddles the time of rapid climatic warming. If we construct a chronology based on radiocarbon dates of a marine near-surface core in the Pacific, and assume a constant reservoir age of 670 years, we obtain a result similar to the atmospheric chronology. This is as expected from the previous discussion of the atmospheric influence on the radiocarbon content of the surface reservoir in the Pacific. However, assuming a constant reservoir age of 430 years for a marine near-surface core in the Atlantic results in an apparent age reversal. This age reversal is a direct consequence of the exchange with the radiocarbon-depleted deep reservoir when convection and the overturning circulation are re-initiated in the North Atlantic. However, the signal is muted by exchange with the atmosphere, so that the maximum time offset does not exceed about 325 years, and the time interval of the age reversal lasts for less than 150 years.

4 Conclusions

Using a dynamical ocean-atmosphere model, including a simple land biosphere and a representation of the cycling of radiocarbon, we have investigated the effect of a succession of gradual Atlantic circulation collapses (coolings) and abrupt recoveries (warmings) on the radiocarbon concentration of the atmosphere and related quantities in the ocean. Such climatic fluctuations are characteristic of the last glacial: their signatures in ice and sediment cores are commonly referred to as Dansgaard/Oeschger and Heinrich events.

In the experiments considered here, we found that $\Delta^{14}\text{C}^{\text{atm}}$ changes by about 25‰. If larger parts of the sea surface become ice covered this value may increase to about 60‰ [Stocker and Wright, 1996]. The corresponding changes seen in the ocean surface reservoir ages exhibit substantial spatial variation. Reservoir ages increase by about 200 yr in the North Atlantic immediately following the abrupt warming; the response in the Pacific is slightly delayed and less pronounced. Top-to-bottom age differences document the changes in the distribution of radiocarbon most clearly. During the cooling they may increase by about 1 kyr (400 yr) in the Atlantic (Pacific), only to decrease again at the time of the abrupt warming. Changes are particularly fast in the Atlantic so that their detection in marine records would require

high temporal resolution. Predicted changes in the Pacific are somewhat less abrupt and reduced in amplitude. Finally, we showed that if age estimates are determined under the assumption that surface reservoir ages are constant, then apparent age reversals may occur in the marine records coincident with age plateaux in the atmospheric records. These age reversals are removed if the correct evolution of the surface reservoir age is taken into account.

Acknowledgements: This study was supported by the Swiss National Science Foundation and the Natural Sciences and Engineering Research Council of Canada through the Climate System History and Dynamics Project. We are grateful to O. Marchal and an anonymous reviewer for useful comments.

References

- Bard, E., M. Arnold, J. Mangerud, M. Paterne, L. Labeyrie, J. Duprat, M.-A. Mélières, E. Sønstegaard, and J.-C. Duplessy, The North Atlantic atmosphere–sea surface ^{14}C gradient during the Younger Dryas climatic event, *Earth Planet. Sci. Lett.*, *126*, 275–287, 1994.
- Becker, B., B. Kromer, and P. Trimborn, A stable-isotope tree-ring timescale of the Late Glacial/Holocene boundary, *Nature*, *353*, 647–649, 1991.
- Björck, S., B. Kromer, S. Johnsen, O. Bennike, D. Hammarlund, G. Lemdahl, G. Possnert, T. L. Rasmussen, B. Wohlfahrt, C. U. Hammer, and M. Spurk, Synchronised terrestrial-atmospheric deglacial records around the North Atlantic, *Science*, *274*, 1155–1160, 1996.
- Bond, G., W. S. Broecker, S. J. Johnsen, J. McManus, L. Labeyrie, J. Jouzel, and G. Bonani, Correlations between climate records from North Atlantic sediments and Greenland ice, *Nature*, *365*, 143–147, 1993.
- Clark, P. U., R. B. Alley, L. D. Keigwin, J. M. Licciardi, S. J. Johnsen, and H. Wang, Origin of the first global meltwater pulse, *Paleoceanogr.*, *11*, 563–577, 1996.
- Damon, P. E., J. C. Lerman, and A. Long, Temporal fluctuations of atmospheric ^{14}C : Causal factors and implications, *Annu. Rev. Earth Planet. Sci.*, *6*, 457–494, 1978.
- GRIP Members, Climate instability during the last interglacial period recorded in the GRIP ice core, *Nature*, *364*, 203–207, 1993.

Hughen, K. A., J. T. Overpeck, S. C. Lehman, M. Kashgarian, J. Southon, L. C. Peterson, R. Alley, and D. M. Sigman, Deglacial changes in ocean circulation from an extended radiocarbon calibration, *Nature*, 391, 65–68, 1998.

MacAyeal, D. R., Binge/purge oscillations of the Laurentide ice sheet as a cause of the North Atlantic’s Heinrich event, *Paleoceanogr.*, 8, 775–784, 1993a.

MacAyeal, D. R., A low-order model of the Heinrich event cycle, *Paleoceanogr.*, 8, 767–773, 1993b.

Mikolajewicz, U., A meltwater-induced collapse of the ‘conveyor belt’ thermohaline circulation and its influence on the distribution of $\Delta^{14}\text{C}$ and $\delta^{18}\text{O}$ in the oceans, *J. Geophys. Res.*, (in press), 1998.

Mikolajewicz, U., T. J. Crowley, A. Schiller, and R. Voss, Modelling teleconnections between the North Atlantic and North Pacific during the Younger Dryas, *Nature*, 387, 384–387, 1997.

Siegenthaler, U., and H. Oeschger, Biospheric CO_2 emissions during the past 200 years reconstructed by convolution of ice core data, *Tellus*, 39B, 140–154, 1987.

Stocker, T. F., and D. G. Wright, Rapid changes in ocean circulation and atmospheric radiocarbon, *Paleoceanogr.*, 11, 773–796, 1996.

Stocker, T. F., D. G. Wright, and L. A. Mysak, A zonally averaged, coupled ocean-atmosphere model for paleoclimate studies, *J. Clim.*, 5, 773–797, 1992.

Stuiver, M., and T. F. Braziunas, Modeling atmospheric ^{14}C influences and ^{14}C ages of marine samples to 10,000 BC, *Radiocarbon*, 35, 137–189, 1993.

Wright, D. G., C. B. Vreugdenhil, and T. M. Hughes, Vorticity dynamics and zonally averaged ocean circulation models, *J. Phys. Oceanogr.*, 25, 2141–2154, 1995.




Figure 1: (a) Evolution of the maximum overturning in the North Atlantic as a result of the meltwater flux perturbations (dashed line). When the meltwater flux increases at a rate of $0.075 \text{ Sv}/1000\text{yr}$ the weakening of the thermohaline circulation occurs on the same time scale and is gradual, but punctuated by some shorter cold events. The maximum meltwater discharge is 0.26 Sv . A few hundred years after the meltwater discharge is stopped, the thermohaline circulation switches on abruptly. (b) Evolution of surface air temperature at 72°N as a result of changing meridional heat flux associated with the thermohaline circulation. (c) Evolution of $\Delta^{14}\text{C}^{\text{atm}}$, the atmospheric concentration of radiocarbon. During the coolings $\Delta^{14}\text{C}^{\text{atm}}$ increases by about 25‰ and then decreases quickly during the abrupt warmings. The latter decrease leads to age plateaux.




Figure 2: Evolution of the surface reservoir age in the Pacific at 39°N (a) and the Atlantic at 39°N (b), and 60°N (c). The age decreases during the cooling because of the reduced ventilation of the deep ocean. At the time of abrupt warming, deep waters that are depleted in radiocarbon are rapidly mixed to the surface increasing the surface reservoir age in the Pacific and mid-latitude Atlantic. There is a slight increase of the surface reservoir age in the Atlantic at 60°N during the cooling period. Remote regions may also be influenced by local convection changes, but the changes in the North Atlantic are rapidly communicated to all regions through the atmosphere. Note the scales for the reservoir ages at different locations.




Figure 3: Evolution of the radiocarbon age difference between 2250 m depth and the surface in the Pacific at 39°N (a) and the Atlantic at 39°N (b) and at 60°N (c). Both decreasing surface reservoir ages and increasing bottom ages contribute to an overall increase of the age difference during the cooling. Changes in the Atlantic are sharp so that high temporal resolution would be required to resolve such signals in marine records. Note the scales for the age difference at different locations.




Figure 4: Radiocarbon chronology (thick solid line) during an event of abrupt warming. The atmospheric chronology (a) is based on the evolution of $\Delta^{14}\text{C}^{\text{atm}}$. Chronologies can also be constructed from marine records from the surface Pacific (b) and the surface Atlantic (c). The usual assumption of constant reservoir ages (670 yr and 430 yr for the Pacific and Atlantic, respectively) leads to apparent age reversals at locations where the ventilation changes significantly (Atlantic, c). A change in the values of the constant reservoir ages simply leads to a vertical shift of the curves. The straight thin solid line indicates the undisturbed chronology (no changes of $\Delta^{14}\text{C}^{\text{atm}}$); the thin solid curve is a the sea surface temperature in the Atlantic during the termination of the cold event.