

RESEARCH ARTICLE

Understanding the psychological impact of the COVID-19 pandemic and containment measures: An empirical model of stress

Bartholomäus Wissmath ^{1,2,3*}, Fred W. Mast ^{1,3}, Fabian Kraus ^{1,2}, David Weibel ^{1,2,3}

1 Institute of Psychology, University of Bern, Bern, Switzerland, 2 w hoch 2 GmbH, Bern, Switzerland, 3 Swiss Distance Learning University, Brig, Switzerland

* wissmath@psy.unibe.ch



Abstract

Epidemics such as COVID-19 and corresponding containment measures are assumed to cause psychological stress. In a survey during the lockdown in Switzerland (n = 1565), we found substantially increased levels of stress in the population. In particular, individuals who did not agree with the containment measures, as well as those who saw nothing positive in the crisis, experienced high levels of stress. In contrast, individuals who are part of a risk group or who are working in healthcare or in essential shops experienced similar stress levels as the general public. The psychological mechanisms that determine stress, caused by the COVID-19 pandemic and containment measures, are not yet clear. Thus, we conducted a path analysis to gain a deeper understanding of the psychological mechanisms that lead to stress. Experiencing fear of the disease is a key driver for being worried. Our model further shows that worries about the individual, social, and economic consequences of the crisis, strongly boost stress. The infection rate in the canton (i.e., state) of residence also contributes to stress. Positive thinking and perceived social, organizational, and governmental support mitigate worries and stress. Our findings indicate that containment measures increase worries and stress, especially for those who feel that these measures either are not sufficient or go too far. Thus, highlighting positive aspects of the crisis and convincing people of the effectiveness and necessity of mitigation measures can, not only promote compliance, but also reduce stress. Our model suggests that people who feel protected by the authorities have fewer worries, which can, in turn, limit the negative impact of the crisis on mental health.

Introduction

COVID-19 evokes stress in patients, healthcare professionals, and relatives [1]. The impact may not be limited to those directly affected; health worries and uncertainty are assumed to generate fear, anxiety, and severe stress in the general population [2]. Hence, mental health practitioners anticipate a sharp rise in the need for mental health services [3]. Some individuals may be more vulnerable to psychosocial consequences; risk factors include certain personality

OPEN ACCESS

Citation: Wissmath B, Mast FW, Kraus F, Weibel D (2021) Understanding the psychological impact of the COVID-19 pandemic and containment measures: An empirical model of stress. *PLoS ONE* 16(7): e0254883. <https://doi.org/10.1371/journal.pone.0254883>

Editor: Rosemary Frey, University of Auckland, NEW ZEALAND

Received: June 3, 2020

Accepted: July 6, 2021

Published: July 29, 2021

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0254883>

Copyright: © 2021 Wissmath et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Understanding the psychological impact of the COVID-19 pandemic and containment measures: an empirical model of

stress [Dataset]. Available from: <https://osf.io/grvwa/>. Deposited May 13, 2020.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

traits (e.g., neuroticism, alexithymia), mental or physical pre-existing conditions, as well as previous traumatic experiences (e.g., childhood maltreatment), or the individual's current situation, such as working in healthcare, social isolation, or poverty [4–10].

Switzerland was among the countries first and most affected by COVID-19 in Europe. To curb the spread of the virus, the Swiss government enacted a set of containment measures: encouraging hygiene, giving shelter-in-place orders, as well as closing borders, child-care facilities, schools, restaurants, bars, leisure facilities, and non-essential shops. Events and gatherings of more than five individuals were banned.

These measures directly affect the general population and may come with side effects. Reduced social interactions are risk factors for mental disorders such as major depression [11], and the containment measures may trigger an economic downturn, thus adding further stress [12]. Early studies on COVID-19 observed increased stress levels [13]. However, little is known about the psychological mechanisms that determine individual stress in this crisis. It is unclear which individuals are particularly affected by stress, which factors have an increasing or decreasing influence on stress, and how these factors interact. In particular, the interplay between fear of the virus, infection rate, risk factors, individual and collective resources, perception of the containment strategy, and psychological stress has not yet been fully understood. The aim of the present study is to address these questions by using path analysis. We assume that fear of COVID-19 and the local infection rate trigger stress [14, 15]. Individual, economic, and societal worries are also expected to increase stress levels. Agreement with containment measures, perceived support, [14, 16], and being optimistic about the crisis [17], are likely to decrease stress. During the lockdown, we surveyed 1565 individuals in Switzerland online.

Materials and methods

Participants

From March 27 to April 26, 2020, a total of 1565 individuals completed the online survey. Participants were recruited via academic institutions, Facebook, and LinkedIn. Posts on social media platforms were published on March 31, 2020 (German) and April 1 (French and Italian). Posts on Facebook were boosted by paid promotion, thereby reaching 1515 (German), 420 (French) and 1006 (Italian) post engagements. Before starting the survey, participants were provided with information about the purpose of the study, data protection policy, and the institution and authors responsible for the study. Individuals from German-speaking ($n = 1206$), French-speaking ($n = 247$), and Italian-speaking ($n = 112$) regions were surveyed. A total of 215 individuals had not completed or had cancelled the survey. The data of three participants were removed, since they had requested the deletion of their data after completion of the survey. An examination of the data for repeated or non-human responses did not reveal any indication of such responses. According to their demographic information, all participants resided in a Swiss canton (as opposed to living abroad). Further information about the sample is provided in Table 1. The study was reviewed and approved by the Ethics Committee of the Faculty of Human Sciences, University of Bern, Switzerland.

Measurements

Stress. We used the German [18], French [19], and Italian [20] version of the Perceived Stress Scale (PPS-10) (10-items) [21, 22]. The widely used PSS is based on the stress-related components of unpredictable, uncontrollable, and overloading life events (e.g., “In the last month, how often have you been angry because of things that were outside your control?”), (0 = never, 1 = almost never, 2 = sometimes, 3 = fairly often, 4 = very often). A PSS score was calculated for each participant by summing up across all scale items. Reliability, calculated for

Table 1. Group comparisons of experienced stress.

Variable	Sample statistics (N = 1565)				ANOVA statistics		
	n	Percentage (%)	Mean PSS Score	SD	F	p	Effect size
Sex⁺					12.6	< .001*	$\eta^2 = 0.016$
(A) Female	1043	66.6	16.1	6.24			
(B) Male	516	33.0	14.4	6.84			
(C) Other / not specified	6	0.4	17.5	10.4			
Age⁺					8.84	< .001*	$\eta^2 = 0.028$
(A) <25	126	8.1	17.4	6.87			
(B) 25–34	371	23.7	16.1	6.10			
(C) 35–44	380	24.3	16.2	6.29			
(D) 45–54	295	18.8	14.9	6.78			
(E) 55–64	261	16.7	15.1	6.67			
(F) > 64	132	8.4	12.9	5.95			
Education					2.66	0.014	$\eta^2 = 0.01$
(A) no finished education	5	0.3	13.2	4.02			
(B) mandatory school time (9 years)	40	2.6	15.8	6.77			
(C) apprenticeship, vocational school, commercial school	311	19.9	15.3	6.97			
(D) High School	99	6.3	17.6	6.9			
(E) Higher technical or vocational school	298	19.0	14.8	6.28			
(F) University	778	49.7	15.6	6.28			
(G) Other	34	34	16.7	6.94			
Type of household					2.19	0.042	$\eta^2 = 0.008$
(A) Single person household	327	20.9	15.5	7.14			
(B) Non-family household with multiple persons	127	8.1	15.8	6.05			
(C) Couple without kids in the household	440	28.1	14.8	6.29			
(D) Couple with kids in the household	469	30.0	15.8	6.14			
(E) Single parent with kids in the household	143	9.1	16.7	6.86			
(F) Multiple-family household	58	3.7	15.4	6.93			
(G) Retirement or nursing home	1	0.1	25.0	N/A (n = 1)			
Having Kids					1.58	0.209	d = 0.077
Yes	472	30.2	15.9	6.33			
No	1093	69.8	15.4	6.58			
Employment at the beginning of crisis					0.59	0.442	d = -0.046
Yes	1252	80.0	15.5	6.31			
No	313	20.0	15.8	7.24			
Working in healthcare¹⁾					0.42	0.516	d = 0.048
Yes	235	15.0	15.7	5.82			
No	1017	65.0	15.4	6.42			
Working in an essential shop¹⁾					0.83	0.361	d = 0.111
Yes	91	5.8	16.1	5.57			
No	1161	74.2	15.4	6.36			
Part of a risk group²⁾					4.77	0.009	$\eta^2 = 0.006$
(A) Yes	265	16.9	15.4	7.27			
(B) No	1180	75.4	15.4	6.30			
(C) Don't know	120	7.7	17.3	6.52			
Agreement with government's containment strategy⁺					48.8	< .001*	$\eta^2 = 0.036$
(A) Agreement	1036	66.2	14.8	6.10			
(B) No agreement	425	27.2	16.7	6.46			

(Continued)

Table 1. (Continued)

Variable	Sample statistics (N = 1565)				ANOVA statistics		
	n	Percentage (%)	Mean PSS Score	SD	F	p	Effect size
(C) No agreement at all	104	6.6	18.8	8.55			
Following shelter-in-place orders stringently					3.19	0.074	d = 0.108
Yes	1041	66.5	15.8	6.53			
No	524	33.5	15.1	6.44			
News consumption					3.23	0.012	$\eta^2 = 0.008$
(A) Never	15	1.0	17.1	8.20			
(B) 1–2 times per week	126	8.1	14.6	6.53			
(C) 3–4 times per week	157	10.0	15.5	5.81			
(D) Once per day	634	40.5	15.1	6.21			
(E) Multiple times per day	633	40.4	16.2	6.86			
Estimated duration of the crisis					0.90	0.496	$\eta^2 = 0.003$
(A) 2 weeks	30	1.9	14.8	7.40			
(B) 1 month	163	10.4	14.8	6.32			
(C) 2 months	555	35.5	15.6	6.47			
(D) 3–5 months	484	30.9	15.8	6.20			
(E) 6 months	161	10.3	15.1	6.45			
(F) 1 year	74	4.7	15.8	7.23			
(G) longer than a year	98	6.3	16.2	7.60			
Seeing positive aspects of the crisis					87.6	< .001*	d = 0.767
Yes	1415	90.4	15.1	6.15			
No	150	9.6	20.2	7.85			

PSS: Perceived Stress Scale; SD: Standard Deviation; p = significance of ANOVA main effects;

+ = significant post hoc Bonferroni tests ($p < .001$): Sex: (A) > (B), Age: (A) > (F), Agreement with containment strategy: (A) < (B) < (C);

* = significant after Bonferroni correction ($\alpha = 0.05/14 = 0.00357$);

¹⁾ unemployed participants were excluded from this analysis.

²⁾ at the time defined by the Swiss authorities as being either > 65 years old or being adult and having high blood pressure, diabetes, cardiovascular disease, respiratory disease or cancer.

<https://doi.org/10.1371/journal.pone.0254883.t001>

all participants and all versions of the scale (German, French, and Italian), was high (Cronbach's $\alpha = .86$).

Amount of worries. We assessed worries about the following aspects (0 = not at all, 1 = little, 2 = medium, 3 = strongly, 4 = very strongly): physical health, mental health, health of family and friends, personal safety, social life, private life and personal needs, financial situation, job security, economic situation, healthcare, and basic supply. A principal component analysis on these items suggests a one-factorial solution. We computed the mean value across all items.

Fear of COVID-19. Fear was measured with a single item: "How afraid are you of the coronavirus (COVID-19)?" (0 = not at all, 1 = little, 2 = medium, 3 = strongly, 4 = very strongly).

Support. Participants assessed perceived support from the following: family, friends and social network, neighbors, employer, authorities, day-care centers, schools, church or religious community, primary care physician, and hospitals (0 = not at all, 1 = little, 2 = partially, 3 = strongly, 4 = very strongly). For the subsequent analysis, a perceived support score was calculated for each participant by summing up all items.

Infection rate. The cases per 100,000 inhabitants in the participants' resident canton (state) were used for the path model [23].

Agreement with containment measures. We asked the participants whether they thought that the measures were (a) not at all strict enough (b) not strict enough (c) just right (d) too strict (e) much too strict. For the analyses, the categories (a) and (e) were combined as “no agreement at all,” and the categories (b) and (d) were combined as “no agreement.” These combined categories were contrasted with (c) “agreement.”

Following shelter-in-place orders stringently. We asked the participants if they followed the shelter-in-place orders stringently (1 = No; 2 = Yes).

Seeing positive aspects. We asked the respondents whether there were any positive aspects for them in the current situation (1 = No; 2 = Yes).

Participants' individual situation and demographics. We assessed sex, age, education, type of household, having kids, employment, working in healthcare or essential shops with customer interaction, part of a risk group (at the time, defined as >64 years old or adult with high blood pressure, diabetes, cardiovascular disease, respiratory disease, or cancer), media consumption, and participants' estimation of the length of the crisis.

Statistical analysis

In a first step, a one-sample z-test was conducted to compare the stress level of the study sample to a representative community sample under normal conditions [22]. Before the test was carried out, we checked whether the stress values were normally distributed: the stress scores were approximately normally distributed, as the skewness equaled 0.43 (a normal distribution can be assumed for skewness values between -0.5 and 0.5 [24]). Thus, the data allowed for z-test as well as ANOVAs (step 2 below).

In a second step, an ANOVA was carried out to compare different subgroups of the study sample. Because of multiple comparisons, Bonferroni's adjustments were made to prevent Type I error inflation ($\alpha = 0.05/14 = 0.00357$). IBM SPSS Statistics for Windows, Version 25.0 was used to run this analysis [25].

In a third step, we tested how stress is related to various variables that are assumed to affect well-being in the context of the pandemic. A path analysis was conducted with stress as the target variable. The path analysis was carried out with IBM SPSS Amos, Version 25.0 [26], using the maximum likelihood method. To test for normality prior to the analysis, a descriptive approach was used. In the context of SEM or path analysis, kurtosis values of > 7 indicate a substantial deviation from normality [27, 28]. In terms of skewness values of > 3 indicate extreme levels of skewness [29]. For both skewness and kurtosis, our values were below these thresholds and thus, the use of the maximum likelihood method was appropriate. To test the significance of the path coefficients, t-tests were calculated. The following variables were included in the path model: stress, fear of COVID-19, worries, support, and infection rate. These variables were selected because previous findings suggest that they affect stress. In addition, "agreement with containment measures" as well as "positive aspects" were included, because the second step of analysis (see above) showed that these variables exert the strongest influence on stress.

We used $p < 0.05$ as a priori level of significance. However, we reported lower p-values when appropriate (e.g., $p < 0.001$). Effect sizes were reported as small, medium, or strong according to the heuristics of Eid, Gollwitzer and Schmitt [30].

Results

First, we compared the observed stress levels in our sample ($M = 15.58$; $SD = 6.65$) to a representative community sample under normal conditions ($M = 12.57$; $SD = 6.24$) [22]. A one-

sample *z*-test showed higher stress levels during the lockdown period ($z = 19.08, p < .001$). The effect ($d = .48$) was medium to strong.

In a second step, we computed multiple comparisons (ANOVA) of different subgroups (see [Table 1](#)). Women expressed higher stress levels than men ($p < .001$), as has been found in non-epidemic situations [21]. Young individuals reported the highest stress levels, whereas the levels were lowest for individuals older than 65 years ($p < .001$; small to medium effect); the other age groups did not differ significantly. The majority of respondents (66%) agreed with the measures of the Swiss authorities. These individuals scored lower on stress than individuals who felt that the measures were either not sufficient or too extreme ($p < .001$). Individuals who strongly disagreed experienced even more stress (medium effect). Individuals who reported that the lockdown situation also has positive aspects (89.4%) had substantially lower stress levels compared to those who did not see any positive aspects at all ($p < .001$; strong effect).

No other assessed variable influenced stress (cf. [Table 1](#)). Interestingly, individuals identifying as members of a risk group ($p = 0.009$; non-significant after Bonferroni-correction; $\alpha = 0.05/14 = 0.00357$), as well as individuals working in healthcare ($p = 0.516$) or in essential shops with customer contact ($p = 0.361$), did not experience more stress than the rest of the population. Also, whether or not participants followed the shelter-in-place orders stringently was not related to stress ($p = .074$). The expected duration of the pandemic ($p = 0.496$) and amount of news consumption ($p = 0.012$, non-significant after Bonferroni-correction; see above) had no influence on stress.

In a third step, we conducted a path analysis with stress as the target variable. [Fig 1](#) depicts the resulting model, showing a good fit.

In total, the measured variables explain 30% of variance in the endogenous variable, stress. Worries contribute the most to stress, and worries mediate between fear of COVID-19 and stress. Seeing positive aspects of the crisis, as well as agreement with the government's containment strategy, are correlated, and both mitigate worries and boost perceived support. In addition, positive thinking directly minimizes stress. Perceived support from family, friends, organizations, and authorities mitigates worries and reduces stress.

Discussion

Our results show that the COVID-19 pandemic and the mitigation strategies increase the population's stress level. We present a model that builds upon existing theories of stress and explains 30% of the variance in stress.

In our model, fear of the virus is the most important booster of worries and the amount of worries is the most important driver of stress. Agreeing with the authorities' containment strategy and seeing positive aspects of the crisis are important factors mitigating stress. Correspondingly, feeling that the containment measures are not sufficient or too extreme is associated with more stress. Support provided by family, friends, organizations, and authorities are also important protective factors against stress.

In our study, risk factors for high stress levels are a high amount of worries, perceived lack of support and, to a lesser extent, high infection rates. Eleven percent of the respondents did not see anything positive about the crisis. This group experienced substantially more stress than individuals who saw at least one positive aspect. Although seeing something positive about the crisis may be associated with the individual situation, personality traits, and respondent's coping behavior [17], our findings indicate that recognizing positive aspects alleviates stress. Some individuals, however, did not see anything positive about the crisis. This raises the question as to why their assessment of the situation is so negative. For example, childhood




Fig 1. A path model showing the influence of various determinants on stress. Model fit statistics: $\chi^2(10) = 42.85$, $p < .01$, GFI = .99, CFI = .97, NFI = .96, TLI = .94, RMSEA = .05. All path coefficients turned out to be highly significant (all paths $p < .001$, except path between Support and Worries: $p < .005$). Error variances appear in small circles.

<https://doi.org/10.1371/journal.pone.0254883.g001>

maltreatment is associated with hopelessness [31] and psychological distress [5]. A recent study in connection with COVID-19 indicates that pre-pandemic maltreatment makes people particularly vulnerable to negative psychosocial consequences of the pandemic [32]. In addition, other factors and traits—such as pessimistic thinking [e.g., 33] or generalized anxiety [e.g., 34]—could account for the lack of perspective. Surprisingly, members of the risk groups, old respondents, and individuals working in healthcare or essential shops with customer contact, did not score higher on stress. One reason could be that extensive protection measures had been mandated. In addition, healthcare providers in Switzerland have, so far, not been overwhelmed by cases of COVID-19. In contrast to agreement with the government's containment strategies, there was no association between following the shelter-in-place orders stringently and perceived stress. About a third of the respondents had indicated that they did not follow the orders stringently. Previously, some COVID-19-containment measures (e.g., social distancing) were found to reduce stress, while others (e.g., more days at home) led to isolation and increased stress [35]. These findings could account for the lack of an association between these two variables in our study.

Our model could be useful in understanding and addressing the psychological impact of possible new waves of COVID-19 cases and other epidemics. Mitigation measures boost worries and stress, particularly for those individuals who feel that these measures are not sufficient or go too far. Highlighting positive aspects about the crisis and convincing people of the effectiveness and the necessity of containment measures may not only boost compliance, but also decrease stress, since individuals feel protected by the authorities and experience less worries. This, in turn, will hopefully limit the impact on mental health as a consequence of the crisis.

Providing support is another important way to mitigate worries, enable coping, and reduce stress.

Taken together, to mitigate stress, authorities should explain containment measures well, highlight positive aspects of the crisis, address worries, and facilitate support. Since agreement with the containment measures may decrease stress, it is crucial that the measures are well-explained and their importance emphasized, and that measures are backed up by scientific evidence. Since the protection of others is an important motivating factor to restrict one's own everyday life [36], the authorities could thus emphasize that compliance with containment measures helps to protect those in need of protection. Furthermore, our analyses show that support through communities, relatives, and employers seems to be key in preventing stress; thus, these support networks should be encouraged to provide help. We identified young individuals as a group experiencing high stress levels. A recent study suggests that social media are particularly well-suited to reach out to adolescents [37]. Since we show that worries are a key driver for stress, our study also provides arguments for economic support, such as stimulus checks or short-time work. In addition, our study has identified groups that are particularly affected by the crisis. This information could help to distribute resources and target efforts.

There are some limitations to the study. First, the cross-sectional design of the study limited the ability to make inferences about the directions of causality. Second, the sample is not representative in terms of gender; as in most studies using the PSS, women are somewhat overrepresented. Third, we could not assess all individual characteristics that could have an influence on stress levels. For example, neuroticism or alexithymia could be relevant traits that make individuals particularly vulnerable to crises such as a pandemic [38]. It would be interesting to investigate the role of these characteristics in future studies.

Supporting information

S1 Appendix. Detailed description of materials and methods. This appendix contains all materials and translations used in the survey, which were developed by the authors. (PDF)

Author Contributions

Conceptualization: Bartholomäus Wissmath, Fred W. Mast, Fabian Kraus.

Data curation: David Weibel.

Formal analysis: Fabian Kraus, David Weibel.

Investigation: Bartholomäus Wissmath, Fabian Kraus.

Methodology: Bartholomäus Wissmath, Fred W. Mast, David Weibel.

Project administration: Bartholomäus Wissmath, Fabian Kraus, David Weibel.

Supervision: David Weibel.

Writing – original draft: Bartholomäus Wissmath, Fred W. Mast, David Weibel.

References

1. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *The Lancet*. 2020; 395(10227):912–20.
2. Pfefferbaum B, North CS. Mental Health and the Covid-19 Pandemic. *N Engl J Med*. 2020; 383:510–512. <https://doi.org/10.1056/NEJMp2008017> PMID: 32283003

3. Fiorillo A, Gorwood P. The consequences of the COVID-19 pandemic on mental health and implications for clinical practice. *Eur. Psychiatry*. 2020; 63(1):e32,1–2. <https://doi.org/10.1192/j.eurpsy.2020.35> PMID: 32234102
4. Liu S, Lithopoulos A, Zhang CQ, Garcia-Barrera MA, Rhodes RE. Personality and perceived stress during COVID-19 pandemic: Testing the mediating role of perceived threat and efficacy. *Pers Individ Dif*. 2021; 168:110351. Epub 2020 Aug 21. <https://doi.org/10.1016/j.paid.2020.110351> PMID: 32863508
5. De Berardis D, Fornaro M, Orsolini L, Valchera A, Carano A, Vellante F, et al. Alexithymia and suicide risk in psychiatric disorders: a mini-review. *Front Psychiatry*. 2017; 8:148. <https://doi.org/10.3389/fpsy.2017.00148> PMID: 28855878
6. Pompili M, Innamorati M, Lamis DA, Erbuto D, Venturini P, Ricci F, et al. The associations among childhood maltreatment, "male depression" and suicide risk in psychiatric patients. *Psychiatry Res*. 2014; 220(1–2):571–8. <https://doi.org/10.1016/j.psychres.2014.07.056> PMID: 25169890
7. Limcaoco RSG, Mateos EM, Fernandez JM, Roncero C. Anxiety, worry and perceived stress in the world due to the COVID-19 pandemic, March 2020. Preliminary results. medRxiv [Preprint]. 2020 [cited 2020 Jun 2]. Available from: <https://www.medrxiv.org/content/10.1101/2020.04.03.20043992v1> <https://doi.org/10.1101/2020.04.03.20043992>
8. Pellecchia U, Crestani R, Decroo T, Van den Bergh R, Al-Kourdi Y. Social consequences of Ebola containment measures in Liberia. *PLoS One*. 2015; 10(12):e0143036. <https://doi.org/10.1371/journal.pone.0143036> PMID: 26650630
9. Varshney M, Parel JT, Raizada N, Sarin SK. Initial psychological impact of COVID-19 and its correlates in Indian community: An online (FEEL-COVID) survey. *PLoS One*. 2020; 15(5):e0233874. <https://doi.org/10.1371/journal.pone.0233874> PMID: 32470088
10. Santiago CD, Wadsworth ME, Stump J. Socioeconomic status, neighborhood disadvantage, and poverty-related stress: Prospective effects on psychological syndromes among diverse low-income families. *J Econ Psychol*. 2011; 32(2):218–230.
11. Bruce ML, Hoff RA. Social and physical health risk factors for first-onset major depressive disorder in a community sample. *Soc. Psychiatry. Psychiatr. Epidemiol*. 1994; 29(4):165–71. <https://doi.org/10.1007/BF00802013> PMID: 7939965
12. Mehta K, Kramer H, Durazo-Arvizu R, Cao G, Tong L, Rao M. Depression in the US population during the time periods surrounding the great recession. *J. Clin. Psychiatry*. 2015; 76(4):e499–504. <https://doi.org/10.4088/JCP.14m09637> PMID: 25919842
13. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *Int J Environ Res Public Health*. 2020; 17(5):1729. <https://doi.org/10.3390/ijerph17051729> PMID: 32155789
14. Kelloway EK, Mullen J, Francis L. The stress (of an) epidemic. *Stress and Health*. 2012; 28(2):91–7. <https://doi.org/10.1002/smi.1406> PMID: 22281817
15. Barbisch D, Koenig KL, Shih F-Y. Is there a case for quarantine? Perspectives from SARS to Ebola. *Disaster Med. Public Health Prep*. 2015; 9(5):547–53. <https://doi.org/10.1017/dmp.2015.38> PMID: 25797363
16. Wong TY, Koh GC, Cheong SK, Fong YT, Sundram M, Koh K. Concerns, perceived impact and preparedness in an avian influenza pandemic—a comparative study between healthcare workers in primary and tertiary care. *Ann Acad Med Singapore*. 2008; 37(2):7.
17. Naseem Z, Khalid R. Positive thinking in coping with stress and health outcomes: Literature review. *Journal of Research & Reflections in Education (JRRE)*. 2010; 4(1):42–61. <https://doi.org/10.1155/2021/7814159> PMID: 33815741
18. Schneider EE, Schönfelder S, Domke-Wolf M, Wessa M. Measuring stress in clinical and nonclinical subjects using a German adaptation of the Perceived Stress Scale. *Int J Clin Health Psychol*. 2020; 20(2):173–181. <https://doi.org/10.1016/j.ijchp.2020.03.004> PMID: 32550857
19. Ben Loubir D, Serhier Z, Otmani N, Housbane S, Mouddene NA, Agoub M, et al. Le stress perçu: validation de la traduction d'une échelle de mesure de stress en dialecte marocain [Perceived stress: validation of the translation of a stress measuring scale in Moroccan dialect]. *Pan Afr Med J*. 2015; 21:280. <https://doi.org/10.11604/pamj.2015.21.280.6430> PMID: 26587130
20. Fossati A. Traduzione Italiana della scala per lo stress percepito [Italian translation of the Perceived Stress Scale]. 2010 [cited 2020 October 15]. Available from: http://www.pensierocritico.eu/files/Italian_PSS_10_with_info.pdf.
21. Cohen S, Williamson G. Perceived stress in a probability sample of the U.S. In: Spacapan S, Oskamp S, editors. *The social psychology of health: claremont symposium on applied social psychology*. Newbury Park, CA: Sage Publications; 1988. pp. 31–67.

22. Klein EM, Brähler E, Dreier M, Reinecke L, Müller KW, Schmutzer G, et al. The German version of the Perceived Stress Scale—psychometric characteristics in a representative German community sample. *BMC Psychiatry*. 2016; 16(1):159.
23. Probst D. COVID-19 Information for Switzerland. 2020 [cited 2020 June 2]. Available from: <https://www.corona-data.ch/>.
24. Lienert GA, Raatz U, Testaufbau und Testanalyse [Test construction and test analysis]. Weinheim: Psychologie Verlags Union; 1998.
25. IBM Corp. IBM SPSS Statistics for Windows, Version 25.0 [Computer Program] Armonk, NY: IBM Corp; 2017.
26. Arbuckle JL. Amos, Version 25.0 [Computer Program]. Chicago: IBM SPSS; 2020.
27. West SG, Finch JF, Curran PJ. Structural equation models with nonnormal variables: Problems and remedies. In: Hoyle RH, editor. *Structural equation modeling: Concepts, issues, and applications*. Thousand Oaks: Sage Publications; 1995. pp. 56–75.
28. Byrne BM. *Structural equation modeling with EQS: Basic concepts, applications, and programming*. New York: Routledge; 2013.
29. Kline RB. Convergence of structural equation modeling and multilevel modeling. In: Williams M, Vogt WP, editors. *The SAGE handbook of innovation in social research methods*. Thousand Oaks: Sage Publications; 2011. pp. 562–589.
30. Eid M, Gollwitzer M, Schmitt M. *Statistik und Forschungsmethoden. [Statistics and research methods]* 5th ed. Weinheim: Beltz; 2017.
31. Wanklyn SG, Day DM, Hart TA, Girard TA. Cumulative childhood maltreatment and depression among incarcerated youth: impulsivity and hopelessness as potential intervening variables. *Child Maltreat*. 2012; 17(4):306–17. <https://doi.org/10.1177/1077559512466956> PMID: 23180865
32. Guo J, Fu M, Liu D, Zhang B, Wang X, van IJzendoorn MH. Is the psychological impact of exposure to COVID-19 stronger in adolescents with pre-pandemic maltreatment experiences? A survey of rural Chinese adolescents. *Child Abuse Negl*. 2020; 20:104667. <https://doi.org/10.1016/j.chiabu.2020.104667> PMID: 32859393
33. Norem JK. Defensive pessimism, optimism, and pessimism. In: Chang EC, editor, *Optimism & pessimism: Implications for theory, research, and practice*. Washington DC: American Psychological Association; 2001. p. 77–100.
34. Miranda R, Mennin DS. Depression, generalized anxiety disorder, and certainty in pessimistic predictions about the future. *Cognit Ther Res*. 2007; 31(1):71–82.
35. Zhao SZ, Wong JYH, Wu Y, Choi EPH, Wang MP, Lam TH. Social distancing compliance under Covid-19 pandemic and mental health impacts: A population-based study. *INT J ENV RES PUB HE*. 2020; 17(18):6692. <https://doi.org/10.3390/ijerph17186692> PMID: 32937929
36. Betsch C. How behavioural science data helps mitigate the COVID-19 crisis. *Nat Hum Behav*. 2020; 4:438. <https://doi.org/10.1038/s41562-020-0866-1> PMID: 32221514
37. Andrews JL, Foulkes L, Blakemore SJ. Peer influence in adolescence: Public-Health implications for COVID-19. *Trends Cogn Sci*. 2020; 24(8):585–587. <https://doi.org/10.1016/j.tics.2020.05.001> PMID: 32444200
38. Tang W, Hu T, Yang L, Xu J. The role of alexithymia in the mental health problems of home-quarantined university students during the COVID-19 pandemic in China. *Pers Individ Dif*. 2020; 165:110131. <https://doi.org/10.1016/j.paid.2020.110131> PMID: 32518435