

Association between self-reported motivation to quit smoking with effectiveness of smoking cessation intervention among patients hospitalized for acute coronary syndromes in Switzerland

Inge Worni-Schudel¹, Vasilis Tzalis¹, Julian Jakob^{1,2}, Kali Tal¹, Lauriane Gilgien-Dénéreaz³, Baris Gencer⁴, Christian M Matter⁵, Thomas Felix Lüscher^{6,7}, Stephan Windecker⁸, François Mach⁴, Jean-Paul Humair⁸, Nicolas Rodondi^{1,10}, David Nanchen³, Reto Auer^{1,3}

Affiliations:

- ¹ Institute of Primary Health Care (BIHAM), University of Bern, Switzerland
- ² Department of Paediatrics, University Hospital Bern (Inselspital, Bern), Switzerland
- ³ Center for Primary Care and Public Health (Unisanté), University of Lausanne, Lausanne, Switzerland
- ⁴ Cardiology Division, Geneva University Hospitals, Geneva, Switzerland
- ⁵ Department of Cardiology, University Hospital Zurich, Zurich, Switzerland
- ⁶ Center for Molecular Cardiology, University of Zurich, Switzerland
- ⁷ Royal Brompton and Harefield Hospitals and Imperial College, London, United Kingdom
- ⁸ Cardiology, University Hospital Bern (Inselspital, Bern), Switzerland
- ⁹ Faculty of Medicine, Division of Primary Care Medicine, Geneva University Hospitals, Geneva, Switzerland
- ¹⁰ Department of General Internal Medicine, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland

eFig 1: Flow-chart of inclusions and exclusions

