



Interpatient heterogeneity in hepatic microvascular blood flow during vascular inflow occlusion (Pringle manoeuvre)

Eric Felli¹, Edoardo Maria Muttillio², Emanuele Felli^{3,4}

¹Hepatology, Department of Biomedical Research, Inselspital, University of Bern, Bern, Switzerland; ²Department of General Surgery, Sant'Andrea Hospital, Sapienza University of Rome, Rome, Italy; ³Digestive and Endocrine Surgery, Nouvel Hopital Civil, University of Strasbourg, Strasbourg, France; ⁴IHU Strasbourg, Institute of Image-Guided Surgery, Strasbourg, France

Correspondence to: Eric Felli, PhD. Hepatology, Department of Biomedical Research, Inselspital, University of Bern, Bern 3010, Switzerland. Email: eric.felli@dbmr.unibe.ch.

Comment on: Shen L, Uz Z, Verheij J, *et al.* Interpatient heterogeneity in hepatic microvascular blood flow during vascular inflow occlusion (Pringle manoeuvre). *Hepatobiliary Surg Nutr* 2020;9:271-83.

Submitted Feb 28, 2021. Accepted for publication Mar 17, 2021.

doi: 10.21037/hbsn-21-91

View this article at: <http://dx.doi.org/10.21037/hbsn-21-91>

Hepatic inflow occlusion (Pringle manoeuvre), is a widely used technique, consisting in temporary intermittent or continuous clamping of the hepatic pedicle. This technique was described in 1908 by Pringle to minimize blood loss during emergency surgery for liver trauma. Therefore, in hepatobiliary surgery, the Pringle manoeuvre assumes a primary role especially in complex liver resections where an intraoperative blood loss could be significant. Afterwards, selective vascular occlusions, notably the glissonian approach described by Takasaki (1), were proposed as an alternative to the Pringle manoeuvre to decrease oxidative stress and postoperative complications in major and minor liver resections. However, this technique also presents some drawbacks as it can increase post-hepatectomy liver failure (PHLF) secondary to prolonged ischemia, especially on fibrotic or cirrhotic livers.

After the inflow clamping, the re-establishment of blood circulation worsens the ischemic damage in a process called ischemia reperfusion injury (IRI) (2). Briefly, the initial hypoxia together with the lack of biomechanical stimulus is followed by an accumulation of reactive oxygen species (ROS) and damage associated molecular patterns (DAMPS) as well as with the increasing in shear stress after the restoration of the oxygen supply, leading the infiltration of neutrophils. In liver transplantation, IRI is a dreadful vascular complication responsible for up to 10% of early transplant failures (3). Additionally, IRI mechanism alters the normal function of hepatic regeneration especially in

patients undergoing extensive resections for which the risk of PHLF may be increased. Moreover, reperfusion damage was experimentally shown to be particularly detrimental in aged liver where the role of statins was successful in reducing IRI damage (4).

The article proposed by Shen *et al.* (5), shows a heterogeneous response in hepatic microvascular flow to Pringle manoeuvre in two homogeneous patient groups called partial responders and full responders. The authors highlight that residual flow in the hepatic microcirculation was still observed despite the total vascular inflow occlusion. Furthermore, backflow from the hepatic venous system and the presence of portal-systemic and arterial collaterals are proposed as possible explanation of the heterogeneous response to the manoeuvre. In particular it has to be noticed that the left and right triangular ligaments have constantly a minimal arterial supply, that may increase in certain conditions such as cirrhosis. These aspects, which remained unclear to date, could be the subject of future studies especially if focused on the role of the endothelial phenotype impairment in PHLF. While this could explain the heterogeneity of response to the Pringle manoeuvre, at the same time it may explain how the outcome of the regenerative process varies depending on different surgical techniques and the different associated procedures such as portal vein embolization (PVE), liver vein deprivation (LVD) and Associating Liver Partition and Portal vein ligation for Staged hepatectomy (ALPPS). Additionally, ALPPS

presents the theoretical advantage of the suppression of the collateral circulation, if the remnant liver is fully mobilized and the partition is complete.

In advanced liver disease numerous intra- and extra-hepatic changes in the circulation contribute in the heterogeneity of response to IRI. These vascular remodelling, especially in cirrhosis and fibrosis, arise from an altered molecular interaction that disrupts the normal balance in mechanical force between cells and extracellular matrix. Among the many actors that are involved (such as hepatocytes, Kupffer cells, hepatic stellate cells, macrophages, platelets and neutrophils), liver endothelial cells (LSECs) are important targets due to their protective role controlling vascular homeostasis (2). Consequently, an intraoperative and non-invasive monitoring of hepatic microcirculation during the Pringle manoeuvre (ischemic time) is a promising technique. Moreover, different phenotypes of LSECs and the respective subpopulations among patients may contribute to the heterogeneity of the response to the ischemic insult discussed in the article (5). In addition, the lactate analysis was found to increase in both groups of patients during the whole surgical procedure. As coherently stated by the authors, the analysis of intrahepatic (capillary) lactate would furnish a better correlation between the local oxidative stress and the lobular microvascular dysfunction. This analysis would help in validating incident dark field imaging (IDFI) providing a quantitative relation between optical and biological properties. Temporary sinusoidal collapse was correctly found at the end of VIO with IDFI and with the microvascular flow index as well. Thus, a proper mathematical correlation between the capillary lactate and the quantitative assessment via IDFI would be highly suggested in order to find similarities in different liver diseases. Finally, the authors conclude that the cytoCam IDFI may be a suitable tool for the assessment of hepatic microcirculation during vascular inflow occlusion (VIO). Overall, we found that this could be an interesting technique that would deserve additional studies and validations. Nevertheless, excluding the ultrasound which represent a clinical standard, other non-invasive and innovative techniques showed positive results in the intraoperative assessment of hepatic microcirculation such as hyperspectral imaging (HSI) and confocal endomicroscopy performed over the Glisson's capsule (6,7). IDFI presents a limitation due to the relatively small field of view when compared with the hyperspectral imaging which is able to furnish an immediate oxygenation map of the liver surface that was shown to be representative of the overall ischemic

lobe. In conclusion, a more detailed and comparative analysis in a preclinical setting would be therefore highly suggested. The authors coherently concluded that this proof of the concept would need further studies.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office of *Hepatobiliary Surgery and Nutrition*. The article did not undergo external peer review.

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/hbsn-21-91>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Takasaki K. Glissonian pedicle transection method for hepatic resection: a new concept of liver segmentation. *J Hepatobiliary Pancreat Surg* 1998;5:286-91.
2. Gracia-Sancho J, Casillas-Ramirez A, Peralta C. Molecular pathways in protecting the liver from ischaemia/reperfusion injury: a 2015 update. *Clin Sci (Lond)* 2015;129:345-62.
3. Rampes S, Ma D. Hepatic ischemia-reperfusion injury in liver transplant setting: mechanisms and protective strategies. *J Biomed Res* 2019;33:221-34.
4. Hide D, Warren A, Fernandez-Iglesias A, et al. Ischemia/

- Reperfusion Injury in the Aged Liver: The Importance of the Sinusoidal Endothelium in Developing Therapeutic Strategies for the Elderly. *J Gerontol A Biol Sci Med Sci* 2020;75:268-77.
5. Shen L, Uz Z, Verheij J, et al. Interpatient heterogeneity in hepatic microvascular blood flow during vascular inflow occlusion (Pringle manoeuvre). *Hepatobiliary Surg Nutr* 2020;9:271-83.
 6. Felli E, Al-Taher M, Collins T, et al. Hyperspectral evaluation of hepatic oxygenation in a model of total vs. arterial liver ischaemia. *Scientific Reports* 2020;10:15441.
 7. Urade T, Felli E, Barberio M, et al. Hyperspectral enhanced reality (HYPER) for anatomical liver resection. *Surg Endosc* 2021;35:1844-50.

Cite this article as: Felli E, Muttillio EM, Felli E. Interpatient heterogeneity in hepatic microvascular blood flow during vascular inflow occlusion (Pringle manoeuvre). *HepatoBiliary Surg Nutr* 2021;10(3):413-415. doi: 10.21037/hbsn-21-91