

Present and future of CONNECT:
a new and compelling project of modern medicine

Giovambattista Capasso^{1,2} and Christoph Wanner³; the CONNECT Action (Cognitive Decline in Nephro-Neurology European Cooperative Target)

¹Department of Translational Medical Sciences, University of Campania Luigi Vanvietlli, Naples, Italy

²Biogem Institute of Molecular Biology and Genetics, Ariano Irpino, Italy

³Department of Medicine, Division of Nephrology, University of Würzburg, Germany

Correspondence to: Giovambattista Capasso; E-mail: gb.capasso@unicampania.it

Epidemiological data collected in different countries indicate a strong increase in the number of patients with chronic kidney diseases (CKD). This phenomenon is evident at all ages, but is particularly significant among elderly people. CKD is a complex and potentially fatal illness affecting all organs and altering many fundamental physiological parameters, such as plasma volume, electrolytes and acid–base balance, hormones and protein metabolism. Due to associated comorbidities, the treatment of choice requires a multidisciplinary approach involving the appropriate use of drugs and nutrient-specific interventions (1).

Although dialysis and transplantation have widely contributed to extend the life of CKD patients, the quality of life is still sub-optimal. This is due to the numerous comorbidities related to the several compromised organs, including the nervous system.

In 30-60% of advanced CKD cases, patients have been diagnosed with cognitive injuries. In these patients, both the central and peripheral nervous systems are frequently affected by several symptoms, such as cognitive decline, peripheral neuropathies and epileptic seizures. The most frequently observed deficiency is Mild Cognitive Impairment (MCI), which is characterized by neurological injuries and cognitive dysfunctions. MCI is already present in the early phase of kidney failure and is significantly more common in CKD patients than in the age-matched general population (2).

To date, interdisciplinary collaboration aimed at interpreting the nature and origin of MCI-CKD has yet been very limited. Indeed, many of the fundamental questions about this pathology have remained unanswered, or even worse, they have been neglected and

never addressed. The dramatic consequence is the lack of any pathogenetic therapy until now.

We thus express deep concern over the current state, and strongly intend to propose effective actions. Concerning the brain field nowadays there are many novel tools that have considerably enhanced our knowledges, including new imaging technologies applicable both for animal models (2-photon microscopy, optogenetics, functional neuroanatomy) and for humans (functional MRI, brain tractography, brain resting state). In addition, innovative cognitive tests are available and Omics approaches are also widely used in neuroscience.

Precisely to achieve this goal, the European CONNECT (Cognitive decline in Nephro-Neurology European Cooperative Target Action) project was established and financed through the COST Action program. As it is well described in the Memorandum of Understanding (MoU), CONNECT comprised of five working groups has the clear aim to fill this gap. It engages clinical nephrologists and neurologists, scientists with pre-clinical expertise in the field of nephrology and neuroscience, and scientists with outstanding knowledge in clinical trials, epidemiology, and data sciences (Fig. 1). To date, CONNECT gathers seventy-two researchers with wide range of scientific and clinical interests from twenty-seven European countries plus the United States. The number is even expected to grow, as it holds enormous potential to involve other scientific fields such as geriatrics, infectious diseases, and many basic sciences.

One of the main objectives of CONNECT, specifically in this first year of activity, is to target clinicians and scientists through publications, reports, conferences and satellite events. To this end, we believe that a good starting point would be to launch a special issue of NDT dedicated to our current understanding of the different topics in the complex kidney-brain relationship. We therefore brainstormed among all members of CONNECT to suggest themes to be discussed and to actively participate in drafting the manuscripts. Below is a brief presentation of the six articles published.

Imenez Silva et al (3) dealt with the very important issue of a possible effect of metabolic acidosis, a characteristic clinical sign of CKD, on cerebral cognitive capacity and motor function. The review, written with the collaboration of excellent experts in the field, highlights the evidence of the correlation between acidosis and cognitive abilities and describes some of the mechanisms that entail an action of acidosis on brain functions.

Pepin et al (4) have written a clinical review on the large topic of neurocognitive disorders in patients with CKD (REF). They described the tests currently used to diagnose the disorders and verify the level of severity. The authors deserve credit for having tried to summarize a large amount of data that has been collected in recent years, although it is understandable that many aspects of this intricate relationship are still far from being completely clarified.

Liebeuf et al (5) reviewed the factors that could influence cognitive abilities during CKD, focusing on uremic toxins (REF). Their timely and exhaustive review has not only the merit of having identified some specific neuro-damaging toxins, but also of having hypothesized their mechanisms of action, allowing a therapeutic perspective for patients.

Bikbov et al (6) had the undoubted merit of addressing one of the newest and most debated topics in the kidney-brain relationship, namely the possibility that an increase in albuminuria could be a direct factor responsible of altering brain cognition (REF). After recalling all the epidemiological studies concerning this relationship, the authors, including leading experts in the field, point out the still unclear aspects and suggest possible actions to overcome this knowledge gap.

Viggiano et al (7) tackled a completely new topic: the potential effect of tubulopathies on brain function with particular regard to cognitive capacity (REF). Although in literature there are only a few data, it is clear that the tubulopathies causing electrolytes disturbances, and in particular those able of altering the water metabolism, affect various brain functions. Certainly in the near future this new chapter of medicine will provide us with much more knowledges, also because nowadays we have the required technologies to explore in detail the function of these two organs.

Zoccali et al (8) discussed the role of neuropeptide Y (NPY) in the regulation of brain function (REF). NPY is currently considered as a key regulator in the gut-brain cross-talk. Its plasma concentration increases with the progress of kidney failure and could play a crucial role in the impairment of cognitive abilities of patients with CKD and in dialysis patients. This hypothesis, if confirmed, could pave the way for new pharmacological treatments.

FUNDING

This article is published as part of a supplement financially supported by the COST Action CA19127-Cognitive Decline in Nephro-Neurology: European Cooperative Target (CONNECT).

CONFLICT OF INTEREST STATEMENT

None declared.

ORIGINAL UNEDITED MANUSCRIPT

APPENDIX

CONNECT collaborators are

Giovambattista Capasso

Alexandre Andrade

Maie Bachmann

Inga Bumblyte

Adrian Constantin Covic

Pilar Delgado

Nicole Endlich

Andreas Engvig

Denis Fouque

Casper Franssen

Sebastian Frische

Liliana Garneata

Loreto Gesualdo

Konstantinos Giannakou

Dimitrios Goumenos

Ayşe Tuğba Kartal

Laila-Yasmin Mani

Hans-Peter Marti

Christopher Mayer

Rikke Nielsen

Vesna Pešić

Merita Rroji (Molla)

Giorgos Sakkas

Goce Spasovski

Kate Stevens

Evgueniy Vazelov

Davide Viggiano

Lefteris Zacharia

ORIGINAL

UNEDITED MANUSCRIPT

Ana Carina Ferreira

Jolanta Malyszko

Ewout Hoorn

Andreja Figurek

Robert Unwin

Carsten Wagner

Christoph Wanner

Annette Bruchfeld

Marion Pepin

Andrzej Wiecek

Dorothea Nitsch

Ivo Fridolin

Gaye Hafez

Maria José Soler Romeo

Michelangela Barbieri

Bojan Batinić

Laura Carrasco

ORIGINAL UNEDITED MANUSCRIPT

Sol Carriazo
Ron Gansevoort
Gianvito Martino
Francesco Mattace Raso
Ionut Nistor
Alberto Ortiz
Giuseppe Paolisso
Daiva Rastenyte
Gabriel Stefan
Gioacchino Tedeschi
Ziad Massy
Boris Bikbov
Karl Hans Endlich
Olivier Godefroy
Jean-Marc Chillon
Anastassia Kossioni
Justina Kurganaite
Norberto Perico
Giuseppe Remuzzi
Tomasz Grodzicki
Francesco Trepiccione
Carmine Zoccali
Mustafa Arici
Peter Blankestijn
Kai-Uwe Eckardt
Danilo Fliser
Eugenio Gutiérrez Jiménez
Maximilian König
Ivan Rychlik
Michela Deleidi
George Reusz


UNEDITED MANUSCRIPT

ORIGINAL

REFERENCES

1. Viggiano D, Wagner CA, Martino G, et al. *Mechanisms of cognitive dysfunction in CKD*. **Nat Rev Nephrol**. 2020 Aug;16(8):452-469. doi: 10.1038/s41581-020-0266-9. Epub 2020 Mar 31.
2. Viggiano D, Wagner CA, Blankestijn PJ, et al. *Mild cognitive impairment and kidney disease: clinical aspects*. **Nephrol Dial Transplant**. 2020 Jan 1;35(1):10-17. doi: 10.1093/ndt/gfz051.
3. Imenez Silva PH, Unwin R, Hoorn EJ, et al. *Acidosis, cognitive dysfunction and motor impairments in patients with kidney disease*. on behalf of CONNECT Action (Cognitive decline in Nephro-Neurology European Cooperative Target) **Nephrol Dial Transplant** 2021; : this issue.
4. Pépin M, Ferreira AC, Arici M, et al. *Cognitive disorders in patients with chronic kidney disease: specificities of clinical assessment* on behalf of the Cognitive decline in Nephro-Neurology European Cooperative Target Action; **Nephrol Dial Transplant** 2021; : this issue.
5. Liabeuf S, Pepin M, Franssen CFM, et al. *Chronic kidney disease and neurological disorders: are uremic toxins the missing piece of the puzzle?* on behalf of the Cognitive decline in Nephro-Neurology European Cooperative Target Action. **Nephrol Dial Transplant** 2021; : this issue.
6. Bikbov B, Soler MJ, Pešić V, et al. *Albuminuria as a risk factor for mild cognitive impairment and dementia – what is the evidence?* on behalf of the Cognitive decline in Nephro-Neurology European Cooperative Target Action
7. Viggiano D, Bruchfeld A, Carriazo S, et al. *Brain dysfunction in tubular and tubulointerstitial kidney diseases*. For the CONNECT ACTION participants **Nephrol Dial Transplant** 2021; : this issue.
8. Zoccali C, Ortiz A, Bumblyte IA, et al. *Neuropeptide Y as a risk factor for cardiorenal disease and cognitive dysfunction in CKD: translational opportunities and challenges*. **Nephrol Dial Transplant** 2021; : this issue.

Figure 1



RIPT

ORIGINAL UNEDITED