

Prevalence of latent structural heart disease in Nepali schoolchildren

Journal:	<i>Cardiology in the Young</i>
Manuscript ID	CTY-21-Jul-586
Manuscript Type:	Brief Report
Date Submitted by the Author:	25-Jul-2021
Complete List of Authors:	Pilgrim, Thomas; Inselspital Universitätsspital Bern, Cardiology Shrestha, Nikesh ; Department of Cardiology, Neuro Cardio and Multispeciality Hospital, Biratnagar, Nepal Uranw , Surendra ; BP Koirala Institute of Health Sciences Karki, Prahlad ; BP Koirala Institute of Health Sciences Bastola , Santosh ; BP Koirala Institute of Health Sciences Mahto , Rajan ; BP Koirala Institute of Health Sciences Sherpa , Kunjang ; BP Koirala Institute of Health Sciences Dhungana , Sahadeb ; Nobel Medical College Teaching Hospital P Ltd Gurung , Keshar ; Department of Cardiology, Neuro Cardio and Multispeciality Hospital, Biratnagar, Nepal Pandey , Naveen ; BP Koirala Institute of Health Sciences Agrawal , Krishna ; BP Koirala Institute of Health Sciences Bartkowiak, Joanna; Inselspital Universitätsspital Bern Shah , Prashant ; BP Koirala Institute of Health Sciences Rothenbühler , Martina; Inselspital Universitätsspital Bern
Keywords:	Congenital heart disease, Rheumatic heart disease, Prevalence
Manuscript Category:	General Cardiology
Subspeciality Collections:	
Diagnostic and Procedural Collections:	
Abstract:	<p>Background The present study aimed to quantify the burden of structural heart disease in Nepali children.</p> <p>Methods We performed a school-based cross-sectional echocardiographic screening study with cluster random sampling among children 5 to 16 years of age.</p> <p>Results Between December 2012 and January 2019, 6,573 children (mean age 10.6 ± 2.9 years) from 41 randomly selected schools underwent echocardiographic screening. Structural heart disease was detected in 14.0 per 1000 children (95% CI 11.3 to 17.1) and was congenital in 3.3 per 1000 (95% CI 2.1 to 5.1) and rheumatic in 10.6 per 1000 (95% CI 8.3 to 13.4). Rates of rheumatic heart disease were higher among children attending public as compared to private schools (OR 2.8, 95% CI 1.6 to 5.2, $p=0.0001$).</p> <p>Conclusion Rheumatic heart disease accounted for three out of four cases of structural heart disease and was more common among children attending public as compared to private schools.</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



SCHOLARONE™
Manuscripts

Prevalence of latent structural heart disease in Nepali schoolchildren

ClinicalTrials.gov Identifier: NCT01550068

Nikesh Raj Shrestha, MD¹; Surendra Uranw, MPH²; Prahlad Karki, MD²; Santosh Bastola, MSc²; Rajan Mahato, MD²; Kunjang Sherpa, MD²; Sahadeb Dhungana, MD³; Keshar Gurung, MD¹; Naveen Pandey, MD²; Krishna Agrawal, MD²; Joanna Bartkowiak MD⁴; Prashant Shah, MD²; Martina Rothenbühler, PhD⁴; Thomas Pilgrim, MD, MSc⁴

¹Department of Cardiology, Neuro Cardio and Multispeciality Hospital, Biratnagar, Nepal;

²Department of Internal Medicine and Cardiology, BP Koirala Institute of Health Sciences, Dharan, Nepal; ³Noble Medical College Teaching Hospital and Research Center, Biratnagar, Nepal;

⁴Department of Cardiology, Inselspital, Bern University Hospital, University of Bern, Switzerland

Word count: 894

Corresponding author:

Thomas Pilgrim, MD, MSc
Department of Cardiology
Swiss Cardiovascular Center
Bern University Hospital
CH-3010 Bern
Phone: 0041 31 632 21 11
Fax: 0041 31 632 47 70
Mail: thomas.pilgrim@insel.ch

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Background The present study aimed to quantify the burden of structural heart disease in Nepali children.

Methods We performed a school-based cross-sectional echocardiographic screening study with cluster random sampling among children 5 to 16 years of age.

Results Between December 2012 and January 2019, 6,573 children (mean age 10.6 ± 2.9 years) from 41 randomly selected schools underwent echocardiographic screening. Structural heart disease was detected in 14.0 per 1000 children (95% CI 11.3 to 17.1) and was congenital in 3.3 per 1000 (95% CI 2.1 to 5.1) and rheumatic in 10.6 per 1000 (95% CI 8.3 to 13.4). Rates of rheumatic heart disease were higher among children attending public as compared to private schools (OR 2.8, 95% CI 1.6 to 5.2, $p=0.0001$).

Conclusion Rheumatic heart disease accounted for three out of four cases of structural heart disease and was more common among children attending public as compared to private schools.

Keywords: Congenital heart disease; rheumatic heart disease; prevalence

48 INTRODUCTION

49 Structural heart disease in children comprises a wide spectrum of congenital and acquired pathologies.
50 Cyanotic congenital heart defects manifest early in life, while acyanotic congenital defects often
51 remain latent throughout childhood and adolescence. Acquired heart disease during childhood is
52 predominantly rheumatic in origin. The prevalence of rheumatic heart disease is distributed unequally
53 across and within regions of socioeconomic development.¹ We aimed to quantify the burden of
54 acquired relative to congenital structural heart disease among children and adolescents in Nepal.

55 METHODS

56 *Study design and data collection*

57 Children and adolescents between the ages of 5 to 16 years attending randomly selected public and
58 private schools in urban and rural areas were prospectively enrolled into a cross-sectional
59 echocardiographic screening program in the Sunsari district in Nepal. The study was originally designed
60 to detect evidence of latent rheumatic heart disease. Details of the sampling frame, sampling strategy,
61 eligibility criteria and informed consent have been reported previously.^{2,3} In brief, cluster random
62 sampling stratified by the location (rural or urban) and administration (public or private) of schools
63 applied to reflect the socioeconomic demographic distribution in the district. All children attending
64 selected schools underwent screening echocardiography by use of a battery-operated portable
65 ultrasound machine (Samsung Medison MySonoU6). Children suspected to have structural heart
66 disease underwent independent confirmation at B.P. Koirala Institute of Health Sciences in Dharan,
67 Nepal. The study was approved by the institutional review board of the B.P. Koirala Institute of Health
68 Sciences and the Nepal Health Research Council. Informed consent was obtained from each patient
69 and the study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki as
70 reflected in a priori approval by the institution's human research committee. The protocol for data
71 acquisition, confirmation, electronic data capturing and central monitoring has been outlined
72 previously.^{2,4}

73 *Definitions*

1
2
3 74 The primary outcome of the present analysis was the prevalence of congenital or acquired structural
4
5 75 heart disease. Acquired heart disease consisted of definite or borderline rheumatic heart disease
6
7 76 defined according to the World Heart Federation (WHF) criteria for individuals 20 years or younger.⁵
8
9
10 77 Congenital heart disease included incidentally detected structural heart disease present since birth
11
12 78 with the exception of bicuspid aortic anatomy, which was not systematically assessed.
13
14

15 79 *Statistical Analysis*
16

17 80 Baseline characteristics and clinical findings are presented as numbers and frequencies (%) for
18
19 81 categorical variables and as means (\pm standard deviation) for continuous variables. Categorical
20
21 82 variables were compared by means of the Fisher's exact or Chi-squared test, continuous ones by two-
22
23 83 sample Student's t-tests. All statistical analyses were performed with Stata software (version 14.2).
24
25
26

27 84 **RESULTS**
28

29
30 85 Between December 2012 and January 2019, 6,573 children from 41 randomly selected schools
31
32 86 underwent echocardiographic screening. The mean age of the children was 10.6 ± 2.9 years and 48.5%
33
34 87 were girls; 50.6% attended public and 49.4% of children attended private schools. Latent structural
35
36 88 heart disease was detected in 92 children (14.0 per 1000, 95% CI 11.3 to 17.1) and was congenital in
37
38 89 22 (3.3 per 1000, 95% CI 2.1 to 5.1) and acquired in 70 (10.6 per 1000, 95% CI 8.3 to 13.4). Among the
39
40 90 children with congenital heart disease, 18 children had an ostium secundum atrial septal defect, four
41
42 91 children were found to have a ventricular septal defect, and one child had Ebstein's anomaly. Among
43
44 92 children with acquired valvular heart disease, 38 children had definite and 32 children borderline
45
46 93 rheumatic heart disease according to the WHF criteria.
47
48
49 94 Prevalence of structural heart disease was more common among children attending public (18.6 per
50
51 95 1000, 95% CI 14.3 to 23.8) as compared to private (9.2 per 1000, 95% CI 6.2 to 13.1) schools (OR 2.0
52
53 96 (95% CI 1.3 to 3.3, $p=0.001$) ([Figure 1](#)). The difference was driven by higher rates of rheumatic heart
54
55 97 disease in children attending public (15.6 per 1000, 95% CI 11.7 to 20.4) as compared to private (5.5
56
57 98 per 1000, 95% CI 3.8 to 8.8) schools (OR 2.8, 95% CI 1.6 to 5.2, $p=0.0001$). The prevalence of congenital
58
59
60

99 heart disease was comparable in children attending public (3.0 per 1000, 95% CI 1.4 to 5.5) or private
100 (3.7 per 1000, 95% CI 1.9 to 6.5) schools (OR 0.8, 95% CI 0.3 to 2.1, $p=0.6256$).

101 **DISCUSSION**

102 Our findings indicate a substantial burden of latent structural heart disease among children in Nepal.
103 In three out of four cases of structural heart disease, or in 1 out of 100 children, there was an acquired
104 valvular defect consistent with rheumatic heart disease. The burden of acquired but not congenital
105 heart disease was higher in children attending public schools as compared to those attending private
106 schools, indicating a gradient of socioeconomic disparity within a region with an endemic pattern of
107 rheumatic heart disease.

108 Our findings need to be interpreted in light of several limitations. As a consequence of dismal survival
109 of newborns with cyanotic heart defects in a region with limited healthcare resources, we found no
110 children with cyanotic structural heart defects in our cohort of schoolchildren. Furthermore, numbers
111 are small to reflect robust prevalence estimates of rare structural defects.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

ACKNOWLEDGMENTS

none

CONTRIBUTIONS

TP conceived the study. TP, NRS and PK had responsibility for the design of the study. NRS, SU, KS, RM, SD, KA, KG and NP were responsible for acquisition of the data. MR did the analysis and interpreted the results in collaboration with TP. TP wrote the first draft of the report. All authors critically revised the report for important intellectual content and approved the final version.

DISCLOSURES

TP reports research grants to the institution from Boston Scientific, Biotronik, and Edwards Lifesciences, personal fees from Biotronik, Boston Scientific, and HighLife SAS outside of this study.

FUNDING

Funded by a dedicated research grant from the UBS Optimus Foundation, Switzerland.

REFERENCES

1. Rothenbühler M, O'Sullivan CJ, Stortecky S, Stefanini GG, Spitzer E, Estill J, Shrestha NR, Keiser O, Jüni P, Pilgrim T. Active surveillance for rheumatic heart disease in endemic regions: a systematic review and meta-analysis of prevalence among children and adolescents. *Lancet Glob Health* 2014;**2**:e717–e726.
2. Pilgrim T, Kalesan B, Karki P, Basnet A, Meier B, Urban P, Shrestha NR. Protocol for a population-based study of rheumatic heart disease prevalence and cardiovascular outcomes among schoolchildren in Nepal. *BMJ Open* 2012;**2**.
3. Karki P, Uranw S, Bastola S, Mahato R, Shrestha NR, Sherpa K, Dhungana S, Odutayo A, Gurung K, Pandey N, Agrawal K, Shah P, Rothenbühler M, Jüni P, Pilgrim T. Effectiveness of Systematic Echocardiographic Screening for Rheumatic Heart Disease in Nepalese Schoolchildren: A Cluster Randomized Clinical Trial. *JAMA Cardiol* 2021; online ahead of print.
4. Shrestha NR, Karki P, Mahto R, Gurung K, Pandey N, Agrawal K, Rothenbühler M, Urban P, Jüni P, Pilgrim T. Prevalence of Subclinical Rheumatic Heart Disease in Eastern Nepal: A School-Based Cross-sectional Study. *JAMA Cardiol* 2016;**1**(1):89-96.
5. Reményi B, Wilson N, Steer A, Ferreira B, Kado J, Kumar K, Lawrenson J, Maguire G, Marijon E, Mirabel M, Mocumbi AO, Mota C, Paar J, Saxena A, Scheel J, Stirling J, Viali S, Balekundri VI, Wheaton G, Zühlke L, Carapetis J. World Heart Federation criteria for echocardiographic diagnosis of rheumatic heart disease-an evidence-based guideline. *Nat Rev Cardiol* 2012;**9**:297–309.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE LEGENDS

Figure 1. Prevalence of congenital (blue) and acquired (red) structural heart disease in Nepali schoolchildren from public and private schools.

For Peer Review

FIGURES

Figure 1

