



# STandard Reporting of CAries Detection and Diagnostic Studies (STAR CARDDS)

Klaus W. Neuhaus<sup>1,2</sup> · Florin Eggmann<sup>1</sup> · Jan Kühnisch<sup>3</sup>  · Svetlana Kapor<sup>3</sup> · Mila Janjic Rankovic<sup>4</sup> · Ina Schüler<sup>5</sup> · Felix Krause<sup>6</sup> · Adrian Lussi<sup>7,8</sup> · Stavroula Michou<sup>9</sup> · Kim Ekstrand<sup>9</sup> · Marie-Charlotte Huysmans<sup>10</sup>

Received: 7 September 2020 / Accepted: 31 August 2021  
© The Author(s) 2021

## Abstract

**Aim** The aim of this paper is to present recommendations from an international workshop which evaluated the methodology and reporting of caries diagnostic studies. As a unique feature, this type of studies is focused on caries lesion detection and assessment, and many of them are carried out in vitro, because of the possibility of histological validation of the whole caries spectrum. This feature is not well covered in the existing reporting STARD guideline within the EQUATOR Network.

**Participants and methods** An international working group of 13 cariology researchers was formed. The STARD checklist was reviewed and modified for caries detection and diagnosis purposes, in a three-step process of evaluation, consensual modification, and delivery during three 2-day workshops over 18 months. Special attention was paid to reporting requirements of caries studies that solely focus on reliability.

**Results** The STARD checklist was modified in 14/30 items, with an emphasis on issues of sample selection (tooth selection in in vitro studies), blinding, and detailed reporting of results.

**Conclusion** Following STARCARDDS (STandard Reporting of CAries Detection and Diagnostic Studies) is expected to result in complete reporting of study design and methodology in future caries diagnosis and detection experiments both in vivo and in vitro, thus allowing for better comparability of studies and higher quality of systematic reviews.

**Clinical relevance** Standardization of caries diagnostic studies leads to a better comparability among future studies, both in vivo and in vitro.

**Keywords** Caries detection · Caries diagnostics · Caries assessment · Methodology · Laboratory studies · Reference standard

✉ Klaus W. Neuhaus  
klaus.neuhaus@unibas.ch

<sup>1</sup> Department of Periodontology, Endodontology and Cariology, University Center for Dental Medicine Basel UZB, University of Basel, Mattenstrasse 40, 4058 Basel, CH, Switzerland

<sup>2</sup> Department of Dermatology, Inselspital - Bern University Hospital, Bern, Switzerland

<sup>3</sup> Department of Conservative Dentistry and Periodontology, University Hospital, Ludwig-Maximilians-Universität München, Munich, Germany

<sup>4</sup> Department of Orthodontics and Dentofacial Orthopedics, University Hospital, Ludwig-Maximilians-Universität München, Munich, Germany

<sup>5</sup> Department of Orthodontics, Section of Preventive and Paediatric Dentistry, University Hospital, Jena, Germany

<sup>6</sup> Department Clinic for Operative Dentistry, Periodontology and Preventive Dentistry, University Hospital RWTH Aachen, Aachen, Germany

<sup>7</sup> Department of Operative Dentistry and Periodontology, Faculty of Dentistry, University Medical Centre, Freiburg, Germany

<sup>8</sup> School of Dental Medicine, University of Bern, Bern, Switzerland

<sup>9</sup> Department of Odontology, University Copenhagen, Copenhagen, Denmark

<sup>10</sup> Department of Dentistry, Radboud University Medical Center, Nijmegen, The Netherlands

## Introduction

There is a wide range of methods currently available for caries detection and diagnosis [1]. In addition, novel diagnostic approaches are being developed while existing methods undergo refinement as the diagnostic thresholds and cutoffs may change over time to meet the requirements of the prevailing caries management principles that they underpin [2, 3].

Caries detection and diagnostic studies aim to assess the accuracy of a diagnostic method, or combinations thereof, with commonly used statistical measures such as sensitivity and specificity, or the reliability of a diagnostic workup or both. Systematic reviews indicate, however, that the quality of reporting in caries detection and diagnostic studies is frequently low [3–9]. Incomplete and inconsistent reporting can preclude a comprehensive evaluation of study methods and the potential of bias. As a consequence, both critical appraisal and replication of a study may be challenging, if not impossible [10–12]. Moreover, systematic reviews and meta-analyses may be severely hampered by insufficient reporting of data in primary studies [12].

To improve the completeness and transparency of reporting in diagnostic accuracy studies, STAndards for Reporting of Diagnostic Accuracy Studies (STARD), a checklist of essential items that ought to be reported in diagnostic studies, was compiled and published in numerous journals in 2003 [13]. This checklist was later revised and the updated version, including 30 essential items, was published in 2015 [12]. The STARD checklist is a valuable methodological tool to enhance, safeguard, and assess the quality of reporting in diagnostic studies. As such, it provides assistance to authors, editors, reviewers, and the readership of diagnostic studies. The CONSORT (Consolidated Standards of Reporting Trial) statement, like STARD part of the EQUATOR Network, provides guidelines to improve the reporting of randomized, controlled trials [14]. Using the CONSORT checklist is strongly recommended to ensure complete, clear, and transparent reporting of randomized, controlled trials. However, randomized, controlled trials investigating caries detection and diagnostic methods are currently scarce. There is a need for a reporting guideline — applicable regardless of the study design — that assists authors in writing reports of caries detection and diagnostic studies.

STARD is applicable to all types of diagnostic tests [12]. The STARD checklist does not, however, fully address some features that are specific to caries diagnostic studies. For instance, there is a majority of studies concentrating on the accuracy of caries lesion detection and (staging) assessment usually in an *in vitro* setting, but also *in vivo* in a selected sample of teeth [3, 5, 7, 8].

The composition of the sample of teeth with regard to the presence and depth of caries lesions can vary greatly [3, 5, 9]. So-called spectrum effects that derive from the specific case mix in a sample have a profound impact on the diagnostic performance of a test [15]. Thus, the applicability of the results may be severely restricted whenever the sample is unrepresentative of the target population [15]. A thorough assessment of any spectrum effects therefore relies on the accurate and complete description of the tooth sample. This requires a particular level of detail in the study report, which is not readily identified by the STARD criteria.

The STARD group has welcomed the development of additional instructions for informative reporting for specific applications [12]. The objective of this study was therefore to identify aspects of reporting that are crucial in caries detection/diagnostic studies and, based on that, to develop a STARD extension for this specific domain. This STARD extension, termed STARCARDDS (short for STAndard Reporting of CARies Detection and Diagnostic Studies), should promote and facilitate the completeness and transparency of reporting of future studies in the field of caries detection and diagnosis, both *in vivo* and *in vitro*.

## Methods

An international, 3-member steering committee, including JK, KWN, and IS, was responsible for coordinating the development of the STARD extension STARCARDDS. This team secured funding, identified, and invited potential participants for the development process and organized meetings, both in person and virtual. Under the leadership of the steering committee, an international working group, comprising 13 researchers in cariology, was formed. The participating members were from Europe and were internationally recognized experts in the field of caries detection and diagnostic studies. They needed to consent in a three-step process and needed to be physically available at the group meetings. All authors of the present report were members of the STARCARDDS working group. The working group held three 2-day consensus workshops, taking place in Bern, Switzerland (October 16–17, 2017), Berlin, Germany (September 8–9, 2018), and Frauenwörth, Germany (March 25–26, 2019).

To establish a consensus, underpinned by cariologic evidence, on reporting standards for caries detection and diagnosis studies, the STARCARDDS group adopted a step-wise development approach [16]. This development process comprised three broad phases: (1) evaluation, (2) drafting with discussion and feedback, and (3) delivery.

**Phase 1, evaluation** A systematic review was undertaken to assess the risk of bias in caries detection and diagnosis studies [17, 18]. The quality of reporting of included studies was comprehensively evaluated. Based on the findings of the systematic review, the applicability of the existing reporting guideline, STARD, was assessed and STARD items that may benefit from additional or modified reporting recommendations were identified.

Data of the systematic reviews are reported in detail elsewhere [17, 18]. In brief, within established methodological frameworks, two separate systematic reviews were undertaken: one on occlusal surface caries detection and diagnosis and one on proximal surface caries detection and diagnosis. Studies pertaining to primary teeth or teeth with restorations, secondary caries, or artificially induced caries lesions were excluded.

The systematic review of the literature on *occlusal surface* caries detection and diagnosis included in vitro and in vivo diagnostic studies that tested the diagnostic accuracy and/or reliability/reproducibility of different diagnostic methods for primary caries detection and assessment in human permanent posterior teeth. The following index tests were included in the search: visual examination, conventional and digital bitewing radiography, laser fluorescence measurements, fiber-optic transillumination, and quantitative light-induced fluorescence. Use of a reference standard was a requisite feature of studies to be eligible for inclusion. The systematic review of the literature on occlusal surface caries detection and diagnosis initially identified 140 studies out of a total of 1090 screened records. A total of 103 publications needed to be excluded owing to a high risk of bias or insufficient data reporting. Finally, 29 in vitro and 8 in vivo studies were selected according to a stepwise eligibility assessment.

The systematic review of the literature on *proximal surface* caries detection and diagnosis included in vivo and in vitro caries diagnostic studies that tested the diagnostic performance of the following caries diagnostic methods: visual examination with and without tactile examination, conventional and digital bitewing radiography, laser fluorescence measurement, and fiber-optic transillumination. Only studies assessing primary caries on the proximal surfaces of permanent posterior teeth were considered for inclusion. The actual status of the tooth surface had to be confirmed by a reference standard. In order to be included, at least one of the following outcomes had to be assessed: diagnostic test accuracy or reliability/reproducibility.

In total, out of 851 screened records, 129 studies met the inclusion criteria in the first selection step of the systematic review of the literature on proximal surface caries detection. When additionally considering those studies with a low/moderate risk of bias, the number of includable studies decreased to 43, of which 7 studies had to be excluded owing to low quality of data reporting. Finally, 31

laboratory studies and 5 clinical studies were included in the meta-analysis.

**Phase 2, drafting with discussion and feedback** Supplementary recommendations, specific for caries detection and diagnosis studies, were drafted and discussed. A consensual draft version of STARCARDDS checklist was decided upon on March 26, 2019. After the last face-to-face meeting, the working group produced a final draft, collecting feedback and holding further discussions throughout the drafting process.

**Phase 3, delivery** Publication in an international peer reviewed journal.

## Results

In result of this consensus process, the STARCARDDS checklist was developed which can be taken from Table 1.

## Discussion

It was found that the STARD checklist needed refinement for caries detection and caries diagnosis studies in order to specifically target the research needs in cariology. Justifications are given below.

*Item 1:* On top of common accuracy studies, for which sensitivity, specificity, positive or negative predictive value, or area under the ROC curve (AUC) are appropriate measures, in cariology, there also exist a number of studies addressing solely reliability. This type of studies aims at assessing objective criteria for reliability performance for new diagnostic methods [19] or, e.g., at assessing the influence of training and calibration [20]. Furthermore, because primary and permanent teeth may have different diagnostic thresholds, and because buccal smooth surface caries is easier to detect than interdental caries or occlusal caries, it is advisable to include information about the dentition (deciduous, permanent) and the assessed tooth surfaces of the study material. The title should describe which type of diagnostic study was conducted.

*Item 5:* In addition to the STARD checklist, the type of study conducted should be specified (validation and/or reliability study, see item 1).

*Items 6–9:* For in vitro caries diagnostic studies, full reporting is needed about the exclusion and inclusion criteria for tooth selection, storage of teeth, and any processing before execution of the study. This is especially important because it may represent a major source of

**Table 1** Recommendations and checklist for the STAndard Reporting of CAries Detection and Diagnostic Studies (STARCARDDS)

Section and Topic	Item No.	STARD Recommendations	STARCARDDS specifications
<b>Title</b>	1	Identification as a study of diagnostic accuracy using at least one measure of accuracy (such as sensitivity, specificity, predictive values, or AUC)	Indicate the type of study: in vitro or in vivo; validation and/or reliability; surfaces type; primary or permanent teeth.
<b>Abstract</b>	2	Structured summary of study design, methods, results, and conclusions	
<b>Introduction</b>		Introduce & justify	
Background/rationale	3	Scientific and clinical background, including the intended use and clinical role of the index test	
Objectives	4	Study objectives and hypotheses	
<b>Methods</b>		Describe	
Study design	5	Whether data collection was planned before the index test and reference standard were performed (prospective study) or after (retrospective study)	Study design & type. Validation and/ or reliability study.
Participants	6	Eligibility criteria	<i>In vitro studies:</i> Exclusion and inclusion criteria for tooth selection, storage of teeth, cleaning and processing prior beginning of the study. Account all teeth which were included, processed or lost. Report the number of included teeth/ surfaces/ occlusal sites separately for each tooth type.
	7	On what basis potentially eligible participants were identified (such as symptoms, results from previous tests, inclusion in registry)	
	8	Where and when potentially eligible participants were identified (setting, location and dates)	
	9	Whether participants formed a consecutive, random or convenience series	<i>In vivo studies:</i> A) Describe the study setting, relevant dates, time intervals including periods of recruitment and follow-ups. B) Exclusion and inclusion criteria for patients. Give the sources of patients, eligibility criteria, screening and selection procedures. Report numbers of individuals at each stage of study, e.g. numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analyzed. Give reasons for non-participation at each stage. C) Describe exclusion and inclusion criteria for teeth.
Test methods	10a	Index test, in sufficient detail to allow replication	Indicate all used test methods. Provide details about the (clinical) setting, workflow, illumination, cleaning, drying, instruments, devices incl. software versions, exposure data, diagnostic criteria etc. for each method. Report the level of clinical and/or diagnostic research experience of trainers and trainees. e.g. years of relevant clinical experience. Visual acuity of the examiners should be reported. Details of theoretical and practical training, calibration, training setting and results (e.g. Kappa) should be reported.
	10b	Reference standard, in sufficient detail to allow replication	Indicate the reference method. Provide details about the (clinical) setting, workflow of the validation technique including instruments & machines, caries staining and imaging, image handling, criteria etc.

Table 1 (continued)

	11	Rationale for choosing the reference standard (if alternatives exist)	
Study material/ population	12a	Definition of and rationale for test positivity cut-offs or result categories of the index test, distinguishing pre-specified from exploratory	
	12b	Definition of and rationale for test positivity cut-offs or result categories of the reference standard, distinguishing pre-specified from exploratory	
	13a	Whether clinical information and reference standard results were available to the performers/readers of the index test	Whether or not the readers of the index test and reference standard were blind (masked) to the results of the other test and describe any other (clinical) information available to the readers. Report the time intervals between the readings. Which clinical/ experimental procedure was performed by which researcher
	13b	Whether clinical information and index test results were available to the assessors of the reference standard	
	14	Methods for estimating or comparing measures of diagnostic accuracy	A) Describe all statistical methods for validity and/ or reliability testing. B) Describe diagnostic & histological thresholds used to compute diagnostic performance. Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why
	15	How indeterminate index test or reference standard results were handled	
	16	How missing data on the index test and reference standard were handled	
	17	Any analyses of variability in diagnostic accuracy, distinguishing pre-specified from exploratory	
	18	Intended sample size and how it was determined	
<b>Results</b>		Report	
Population and/or teeth	19	Flow of participants, using a diagram	
Descriptive	20	Baseline demographic and clinical characteristics of participants	Characterize the study population (age, female/male ratio, dental health status) and/ or teeth sample.
Outcome data	21a	Distribution of severity of disease in those with the target condition	
	21b	Distribution of alternative diagnoses in those without the target condition	
	22	Time interval and any clinical interventions between index test and reference standard	
	23	Cross tabulation of the index test results (or their distribution) by the results of the reference standard	<i>Validation testing:</i> Full cross tabulation of the results of the index tests and the reference standard; for continuous results the distribution of the test results by the results of the reference standard. <i>Reliability testing:</i> Report numbers of the included patients, teeth and/ or examiners.

Table 1 (continued)

	24	Estimates of diagnostic accuracy and their precision (such as 95% confidence intervals)	<i>Validation testing:</i> Estimates of diagnostic accuracy and measures of statistical uncertainty (e.g. 95% confidence intervals). <i>Reliability testing:</i> Report reliability data according to appropriate statistical measures.
	25	Any adverse events from performing the index test or the reference standard	
<b>Discussion</b>		Discuss	
Key results	26	Study limitations, including sources of potential bias, statistical uncertainty, and generalizability	Consider strengths & limitations of the study. Discuss both direction and magnitude of any potential bias.
Interpretation	27	Implications for practice, including the intended use and clinical role of the index test	Discuss the (clinical) relevance of the study results.
<b>Other information</b>			
Conflicts of interest	28	Registration number and name of registry	If applicable (clinical studies)
	29	Where the full study protocol can be accessed	If applicable (clinical studies)
	30	Sources of funding and other support; role of funders	

bias. For instance, a caries diagnostic study may be carried out on teeth with an open apex, or teeth which have clearly been impacted and never been in function, teeth which would have a very low likelihood of having caries lesions. If this is obvious to the examiners, for instance because roots are left uncovered, then this information will likely lead to bias. Similar to other studies where pre-testing failures need to be reported [21, 22], also in caries detection studies, all teeth that were included, processed, or lost need to be fully accounted for. As different tooth types (e.g., premolars vs. molars) or tooth surfaces (e.g., smooth surface vs. occlusal surface) are not equally demanding for caries detection, the number of assessed teeth/surfaces should be given separately for each tooth type.

For in vivo studies, the STARD criteria are valid, but also require some further refinement. The study setting, relevant dates, time intervals, including periods of recruitment, and follow-ups need to be described. Furthermore, STARCAR DDS elaborates on how to report the numbers of patients at each stage of the conducted study. Especially in patients with a mixed dentition, it seems to be important to exactly describe the reasons for inclusion and exclusion criteria of teeth. Special attention should be paid to how missing teeth are handled (missing due to caries vs. missing due to physiological exfoliation).

*Item 10a:* In order to be able to compare the results of caries detection trials, the study settings should be standardized to the best of knowledge. It seems to be essential to particularly provide sufficient details about the clinical setting, the workflow, illumination conditions, eventual cleaning and/or drying of the tooth surfaces, and/or the supportive use of any instruments. If medical devices are used, their software version or exposure data should be provided. Because each diagnostic method in cariology has its own diagnostic criteria, they should be given in full detail. Furthermore, as clinical experience [20] and visual acuity are crucial factors in caries diagnostic studies [23, 24], it is indispensable to also report on the levels of clinical experience (years after graduation; specialist training) and on the visual acuity of the researchers.

*Item 10b:* Histologic validation of caries varies between applied methods (e.g., [25, 26]). It seems crucial that, whenever some sort of histology serves as a reference standard, the workflow of the validation technique must be described in sufficient detail, including the use of instruments/machines, any staining of caries, the imaging process, the handling of the imaging, and the applied assessment criteria. Even though an absolute determination of the extension of the caries process is not possible (partly because of an ongoing discussion on the nature and level of tissue change that is relevant/critical in deter-

mining lesion progression stage), taking care of reporting all relevant details of the respective applied reference method helps to avoid study heterogeneity.

*Item 11:* In in vivo caries trials, there is a problem how to validate healthy tooth surfaces or those tooth surfaces that do not need to be opened up. The latter case allows for immediate inspection of the depth of a lesion; the drilling of a lesion thus equals taking a biopsy in medicine. Though restorative caries treatment still represents a substantial part of everyday dental care provision, it is neither ethical nor helpful to operatively open up early stages of the caries process or even healthy surfaces. Therefore, many clinical diagnostic studies rely on imperfect reference standards such as radiography [27]. However, it seems to be an accepted method to directly inspect tooth surfaces after temporary tooth separation in order to validate early stages of caries or healthy surfaces [28]. While direct inspection of interdental caries after tooth separation has perfect specificity, the depth of more advanced carious lesions cannot be assessed by this procedure. It is therefore legitimate to discuss, for clinical caries studies, the use of a composite or hybrid reference standard [10] which on the one hand fully allows for assessment of healthy tooth surfaces and initial caries, and on the other hand for precise estimates of lesion depth. There have been concerns that composite reference standards may represent bias themselves because of their dependency of disease prevalence and their possibly underestimating or overestimating diagnostic accuracy [29]. However, using latent class models could render more precise estimates when using composite reference standards in in vivo caries diagnostic trials [30].

*Items 13a and 13b:* The aspect of blinding is important also in caries diagnostic studies but often reported in insufficient detail. The steps to ensure blinding should be reported in a full and comprehensible way. Special attention should be paid to whether or not the readers of the index test and reference standard were blind (masked) to the results of the other test. Because in caries laboratory studies teeth are often assessed more than once, starting at sample selection and preparation, the time interval between the respective assessments needs to be reported [31]. Furthermore, the clinical and/or experimental procedures should be clearly attributable to the respective researchers who performed these steps.

*Item 14:* Because in many caries diagnostic studies not only accuracy is assessed, but also reliability, all statistical methods for testing both validity and/or reliability have to be reported. With respect to the diagnostic and histological thresholds, a proper description is necessary. Because some diagnostic tests yield quantitative data, their statistical analysis must be reported as well. If grouping/clustering of data seems necessary (e.g., enamel

caries vs. dentine caries), doing so has to be described and justified. Testing reliability between more than 2 examiners may require more sophisticated statistical methods (e.g., Fleiss' Kappa [32], bootstrapping [33]) and these might be applied as well.

*Item 20:* In in vitro experiments, the teeth used for the study have to be specified. Due to their characteristic appearance, it makes a diagnostic difference, if they have served in the oral cavity, or if they are newly erupted or impacted (see also items 6–9).

*Item 23:* Next to full cross-tabulations, for reliability testing, the numbers of the included patients, teeth, and/or examiners need to be reported. In the case of multiple examiners, it can sometimes be observed that the diagnostic data are collapsed, or that “a consensus decision” was derived at. For the sake of transparency and confirmability, all data should be reported separately, or at least made public as an online supplemental content.

*Item 24:* Next to testing of validation, where estimates of diagnostic accuracy and measures of statistical uncertainty (e.g., 95% confidence intervals) must be provided, reliability data also needs to be reported as well according to the appropriate statistical measures applied. This becomes especially relevant in trials with more than two observers.

*Item 25:* Usually, no adverse effects can be noted in caries diagnostic trials. However, some methods use ionizing radiation or the application of dyes and thus could evoke adverse reactions. This item therefore needs to be addressed, too.

## Reporting in future diagnostic studies

There is evidence that insufficient reporting contributes to a higher risk of bias and to an increased heterogeneity between diagnostic accuracy studies [10]. Standardized reporting requirements are an important means to reduce heterogeneity among diagnostic accuracy studies. Most scientific journals try to standardize the way of reporting in their author guidelines in order to reduce heterogeneity of the reported data. STARCARDDS aims into the same direction but is specific with respect to caries diagnostic and detection studies, including reliability studies. STARCARDDS can, moreover, be used together with established guidelines such as the CONSORT statement [14] when reporting randomized clinical trials investigating caries detection and diagnostic methods. It is our hope that following the STARCARDDS checklist leads to a more complete reporting of study methodology and results, and will thus result in better comparability of future diagnostic studies in cariology.

**Funding** This project was funded by a grant for creating a scientific network obtained from the German Research Foundation (Deutsche Forschungsgemeinschaft, SCHU-3217/1–1).

## Declarations

**Ethical approval** This article does not contain any studies with human participants or animals performed by any of the authors.

**Informed consent** For this type of study, formal consent is not required.

**Conflict of interest** The authors declare no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Neuhaus KW, Lussi A (2018) Carious lesion diagnosis: methods, problems, thresholds. *Monographs Oral Sci* 27:24–31
2. Innes NPT, Chu CH, Fontana M, Lo ECM, Thomson WM, Uribe S, Heiland M, Jepsen S, Schwendicke F (2019) A century of change towards prevention and minimal intervention in cariology. *J Dent Res* 98:611–617
3. Gimenez T, Piovesan C, Braga MM, Raggio DP, Deery C, Ricketts DN, Ekstrand KR, Mendes FM (2015) Visual inspection for caries detection: a systematic review and meta-analysis. *J Dent Res* 94:895–904
4. Gimenez T, Braga MM, Raggio DP, Deery C, Ricketts DN, Mendes FM (2013) Fluorescence-based methods for detecting caries lesions systematic review, meta-analysis and sources of heterogeneity. *PLoS One* 8:e60421
5. Gimenez T, Piovesan C, Braga MM, Raggio DP, Deery C, Ekstrand RDN, KR, Mendes FM, (2015) Clinical relevance of studies on the accuracy of visual inspection for detecting caries lesions: a systematic review. *Caries Res* 49:91–98
6. Schwendicke F, Tzschoppe M, Paris S (2015) Radiographic caries detection: a systematic review and meta-analysis. *J Dent* 43:924–933
7. Brouwer F, Askar H, Paris S, Schwendicke F (2016) Detecting secondary caries lesions: a systematic review and meta-analysis. *J Dent Res* 95:143–151
8. Ekstrand KR, Gimenez T, Ferreira FR, Mendes FM, Braga MM (2018) The International Caries Detection and Assessment System - ICDAS: a systematic review. *Caries Res* 52:406–419
9. Schwendicke F, Elhennawy K, El Shahawy O, Maher R, Gimenez T, Mendes FM, Willis BH (2018) Visual and radiographic caries detection: a tailored meta-analysis for two different settings. Egypt and Germany *BMC Oral Health* 18:105
10. Rutjes AWS, Reitsma JB, Di Nisio M, Smidt N, van Rijn JC, Bossuyt PM (2006) Evidence of bias and variation in diagnostic accuracy studies. *Can Med Assoc J* 174:469–476
11. Glasziou P, Altman DG, Bossuyt PM, Boutron I, Clarke M, Julious S, Michie S, Moher D, Wager E (2014) Reducing waste from incomplete or unusable reports of biomedical research. *The Lancet* 383(9913):267–276
12. Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig LM et al (2015) STARD 2015: an updated list of essential items for reporting diagnostic accuracy studies. *Radiol* 277:826–832
13. Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig LM et al (2003) Towards complete and accurate reporting of studies of diagnostic accuracy: the STARD Initiative. *Radiol* 226:24–28
14. Schulz KF, Altman DG, Moher D (2010) CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *Trials* 11:32. <https://doi.org/10.1186/1745-6215-11-32>
15. Willis Brian H (2008) Spectrum bias - why clinicians need to be cautious when applying diagnostic test studies. *Fam Pract* 25:390–396
16. Noel-Storr AH, McCleery JM, Richard E, Ritchie CW, Flicker L, Cullum SJ, Davis D, Quinn TJ, Hyde C, Rutjes AW, Smailagic N, Marcus S, Black S, Blennow K, Brayne C, Fiorivanti M, Johnson JK, Köpke S, Schneider LS, Simmons A, Mattsson N, Zetterberg H, Bossuyt PM, Wilcock G, McShane R (2014) Reporting standards for studies of diagnostic test accuracy in dementia: the STARDdem Initiative. *Neurol* 83:364–373
17. Janjic Rankovic M, Kapor S, Khazaei Y, Crispin A, Schüler I, Krause F, Ekstrand K, Michou S, Eggmann F, Lussi A, Huysmans MC, Neuhaus KW, Kühnisch J (submitted) Systematic review and meta-analysis of caries diagnostic studies on proximal surfaces. *Clin Oral Investig* (submitted)
18. Kapor S, Janjic Rankovic M, Khazaei Y, Schüler I, Krause F, Lussi A, Neuhaus KW, Eggmann F, Michou S, Ekstrand K, Huysmans MC, Crispin A, Kühnisch J (submitted) Systematic review and meta-analysis of caries diagnostic methods on occlusal surfaces. *Clin Oral Investig* (2021) online ahead of print
19. Litzenburger F, Heck K, Pitchika V, Neuhaus KW, Jost FN, Hickel R, Jablonski-Momeni A, Welk A, Lederer A, Kühnisch J (2018) Inter- and intraexaminer reliability of bitewing radiography and near-infrared light transillumination for proximal caries detection and assessment. *Dentomaxillofac Radiol* 47:20170292
20. Diniz MB, Rodrigues JA, Neuhaus KW, Cordeiro RC, Lussi A (2010) Influence of examiner's clinical experience on the reproducibility and accuracy of radiographic examination in detecting occlusal caries. *Clin Oral Invest* 14:515–523
21. De Munck J, Shirai K, Yoshida Y, Inoue S, Van Landuyt K, Lambrechts P, Suzuki K, Shintani H, van Meerbeek B (2006) Effect of water storage on the bonding effectiveness of 6 adhesives to class I cavity dentin. *Oper Dent* 31:456–465
22. Swissmedic. Swiss interpretation of the GLP principles 2019. [www.anmeldestelle.admin.ch](http://www.anmeldestelle.admin.ch) (access 22 May 2020)
23. Eichenberger M, Perrin P, Neuhaus KW, Bringolf U, Lussi A (2013) Visual acuity of dentists under simulated clinical conditions. *Clin Oral Invest* 17:725–729
24. Neuhaus KW, Jost F, Perrin P, Lussi A (2015) Impact of different magnification levels on visual caries detection with ICDAS. *J Dent* 43:1559–1564
25. Jacobsen JH, Hansen B, Wenzel A, Hintze H (2004) Relationship between histological and radiographic caries lesion depth measured in images from four digital radiography systems. *Caries Res* 38:34–38
26. Rodrigues JA, Neuhaus KW, Diniz MB, Hug I, Stich H, Karlsson L, Lussi A (2012) Comparison among gold standard techniques used for the validation of methods for occlusal caries detection. *Microsc Res Tech* 75:605–608
27. Machiulskiene V, Nyvad B, Baelum V (2004) Comparison of diagnostic yields of clinical and radiographic caries examinations in children of different age. *Eur J Paediatr Dent* 5:157–162



28. Novaes TF, Matos R, Braga MM, Imparato JC, Raggio DP, Mendes FM (2009) Performance of a pen-type laser fluorescence device and conventional methods in detecting approximal caries lesions in primary teeth—in vivo study. *Caries Res* 43:36–42
29. Collins J, Huynh M (2014) Estimation of diagnostic test accuracy without full verification: a review of latent class methods. *Stat Med* 33(24):4141–4169
30. Dendukuri N, Schiller I, de Groot J, Libman M, Moons K, Reitsma J, van Smeden M (2018) Concerns about composite reference standards in diagnostic research. *BMJ*. 18(360):j5779
31. Jablonski-Momeni A, Ricketts DN, Weber K, Ziomek O, Heinkel-Gutenbrunner M, Schipper HM, Stoll R, Pieper K (2010) Effect of different time intervals between examinations on the reproducibility of ICDAS-II for occlusal caries. *Caries Res* 44:267–271
32. Fleiss JL (1971) Measuring nominal scale agreement among many raters. *Psychol Bulletin* 76:378–382
33. Efron B (1979) Bootstrap methods: another look at the jackknife. *Ann Statist* 7:1–26

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.