

RESEARCH ARTICLE


Assessment of feeding, ruminating and locomotion behaviors in dairy cows around calving – a retrospective clinical study to early detect spontaneous disease appearance

Mahmoud Fadul¹✉, Luigi D'Andrea²✉, Maher Alsaad¹, Giuliano Borriello², Antonio Di Lori², Dimitri Stucki¹, Paolo Ciaramella^{2*}✉, Adrian Steiner^{1*}, Jacopo Guccione²

1 Vetsuisse-Faculty, Department of Clinical Veterinary Medicine, Clinic for Ruminants, University of Bern, Bern, Switzerland, **2** Department of Veterinary Medicine and Animal Productions, University of Napoli 'Federico II,' Napoli, Italy

✉ These authors contributed equally to this work.

* paociara@unina.it (PC); adrian.steiner@vetsuisse.unibe.ch (AS)



Abstract

The study aims to verify the usefulness of new intervals-based algorithms for clinical interpretation of animal behavior in dairy cows around calving period. Thirteen activities associated with feeding-ruminating-locomotion-behaviors of 42 adult Holstein-Friesian cows were continuously monitored for the week (wk) -2, wk -1 and wk +1 relative to calving (overall 30'340 min/animal). Soon after, animals were retrospectively assigned to group-S (at least one spontaneous diseases; n = 24) and group-H (healthy; n = 18). The average activities performed by the groups, recorded by RumiWatch® halter and pedometer, were compared at the different weekly intervals. The average activities on the day of clinical diagnosis (dd0), as well as one (dd-1) and two days before (dd-2) were also assessed. Differences of dd0 vs. dd-1 ($\Delta D1$), dd0 vs. wk -1 ($\Delta D2$), and wk +1 vs. wk -1 ($\Delta weeks$) were calculated. Variables showing significant differences between the groups were used for a univariate logistic regression, a receiver operating characteristic analysis, and a multivariate logistic regression model. At wk +1 and dd0, *eating-* and *ruminating-time*, *eating-* and *ruminant-chews* and *ruminating boluses* were significantly lower in group-S as compared to group-H, while *other activity time* was higher. For $\Delta D2$ and $\Delta weeks$, the differences of *eating-* and *ruminating-time*, as well as of *eating-* and *ruminant-chews* were significantly lower in group-S as compared to group-H. Concerning the locomotion behaviors, the *lying time* was significantly higher in group-S vs. group-H at wk +1 and dd-2. The *number of strides* was significantly lower in group-S compared to group-H at wk +1. The model including *eating-chews*, *ruminant-chews* and *other activity time* reached the highest accuracy in detecting sick cows in wk +1 (area under the curve: 81%; sensitivity: 73.7%; specificity: 82.4%). Some of the new algorithms for the clinical interpretation of cow behaviour as described in this study may contribute to monitoring animals' health around calving.

OPEN ACCESS

Citation: Fadul M, D'Andrea L, Alsaad M, Borriello G, Di Lori A, Stucki D, et al. (2022) Assessment of feeding, ruminating and locomotion behaviors in dairy cows around calving – a retrospective clinical study to early detect spontaneous disease appearance. PLoS ONE 17(3): e0264834. <https://doi.org/10.1371/journal.pone.0264834>

Editor: Juan J. Loo, University of Illinois, UNITED STATES

Received: June 28, 2021

Accepted: February 17, 2022

Published: March 4, 2022

Copyright: © 2022 Fadul et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its files.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Introduction

The peripartum period represents a crucial phase for dairy cows' life cycle because of the significant increase of energy requirements and the severe metabolic adjustments mainly due to the exponential fetal growth, calving and the onset of lactation [1–3]. Throughout this period of physiological and behavioral changes, the natural drop in dry matter intake, as parturition approaches, makes the adaptation phase even more difficult, culminating in a negative energy balance (NEB) status; this is recognized as a risk factor for the development of puerperal diseases [2,4,5]. A successful management of this period, therefore, represents a key factor of profitable farms [6]. As the latter continue to increase in size, and supervision of individual cows gets more difficult, the use of precision dairy farming (PDF) technologies may provide essential support to manage the herd. Their use may increase the overall farm's efficiency, reduce the time spent for animal management, improve animals' health and well-being, minimize adverse environmental impact, and sustain high quality of products of animal origin [7–9]. In the past, veterinary involvement in herd health management has been mainly the consequence of farmers' experience and judgment to identify animals' behavioral changes [7,10]. Although this skill is still invaluable, it can be easily influenced by human perception and animal's clinical status (e.g., clinical symptoms not obvious to the human eye or clinical signs only shown at a late stage of disease) [11]. Thus, technologies for monitoring cows' activities and behavior may have a great impact to support the observations of skilled herdspeople and to allow for early diagnosis improving the success rate of therapeutic measures.

Although the electronic technologies have been primarily developed in attempts to improve estrus detection efficiency, pointing out changes in physical and/or mounting activities (pedometers), the list of devices dedicated to PDF and used for cow status monitoring and management continues to increase, day-to-day [9,12]. Thanks to the rapid development of new technologies and supporting applications, several digital systems have been proposed and validated by different companies and research teams with exciting results [13–15]. Nevertheless, despite widespread availability, the number of information provided by these devices is often limited, and the combined use of multiple technologies becomes less economically feasible for dairy farms requiring a higher level of multiple analytical capabilities at the same time [10].

In this context, RumiWatch® halter (RWh) and the RumiWatch® pedometer (RWp) equipped with three-dimensional (3D)-accelerometers fulfil these criteria, having unique features that—to the best of our knowledge—are not offered by any other accelerometer currently available on the market [16,17]. Indeed, as reported in our previous studies, the combined use of the two devices provides meaningful and accurate information regarding walking, feeding and rumination behaviors in dairy ruminants [18–20]. Sick and healthy cows seem to express different levels of activity [21], including parameters such as number of lying bouts [22], time spent lying down [23], length of strides [24], walking speed [24], chewing activity [25] feeding and rumination time [26,27]. Despite the amount of data available continues to increase, clinical consideration that may arise from them and consequently the systems for the early disease detection still show wide margins for improvement [9]. Promising studies have been focused on changes in feeding behavior as indicators for cows at risk of postpartum disease [28,29]; nevertheless, data analysis investigating the predictive values of pre-partum behaviors on diseases after calving are still incomplete [30,31].

For all these reasons, the current study was initiated. Our hypothesis was that the parameters recorded by the RumiWatch® in dairy cows naturally experiencing the around calving period may represent meaningful tools for reliable animals' behavior assessment in such a critical period, if associated with the use of new algorithms for data interpretation.

Based on the previous considerations, the goals of the present investigation were: (i) to analyze retrospectively the behaviors of free-stall housed adult dairy cows, before and after calving in states of health and disease, (ii) to compare the RumiWach® output with the findings based on clinical observations, and (iii) to assess the usefulness of newly developed algorithms based on the devices' data to detect disease.

Material and methods

General and ethical animal care

The current investigation was carried out between spring and the beginning of summer 2018 (three months) in a dairy farm located in Caserta district (southern Italy). All clinical and diagnostic procedures performed in this study received institutional approval by the Ethical Animal Care and Use Committee of the University of Naples Federico II [n°2016/0052972] and were performed abiding by the *common good clinical practices* [32] by expert clinicians. Moreover, the farm manager's consent for scheduled measures and methods used was received before the beginning of the study.

Farm and management

The study farm was characterized by a free-stall housing system and by an average milk yield/cow of 11127 ± 1558 (kg \pm SD, 305-days). Forced by the farm routine, animals enrolled were moved through 4 different pens belonging to the same building, during the entire investigation: (i) close-up pen (over a period of three weeks before calving), (ii) calving pen (24h before calving up to the end of the calving event), (iii) post-partum pen (immediately after calving up to 3 days post-partum), (iv) fresh pen (from 4 to 30 days post-partum). All pens were designed to host up to 150 adult dairy cows and to provide at least 10 m² per animal of space (including bedding, feeding and loafing areas). During the entire study period, the stocking density was $\leq 80\%$ of the maximal capacity, the feeding space was ~ 0.9 m/cow, the linear space for water provision > 10 cm/cow and the width of passageways if present ≥ 3 m. Cows were ad-libitum fed, twice a day (d), with a total mixed ration specific for the respective reproductive/productive phases (close-up and fresh cows, supplementary file—S1 Table). After calving, all animals were milked 3 times/d, in a side-by-side milking parlour (total of 80 places).

At the investigation time, the farm was already routinely implementing a claw-health monitoring program including: (i) claw trimming 2 times/year; (ii) digital data recording of foot disorders/diseases observed (i.e., location, type and severity); (iii) regular (every three months) locomotion scoring (LS) of the herd according to Sprecher et al. [33]; (iv) in any case of obvious lameness (score ≥ 3), examination and treatment in the trimming chute was initiated within 48h, for severely lame cows (score = 4 or 5) within 24h. Moreover, stockpersons regularly observed close-up/lactating cows (3 times/d) and cows in the calving pen (every hour, night included) for abnormal behaviors and other signs of disease identification.

Study design, animal selection, and findings based on clinical observations

The study was designed as observational, longitudinal, and retrospective. The assessment of the clinical usefulness of the output originating from thirteen—continuously recorded—RumiWach® activities (ITIN + HOCH GmbH, Fütterungstechnik, Liestal, Switzerland) was performed employing 42 healthy, multiparous, Holstein-Friesian, dairy cows [parity = 2.39 ± 1.22 ; mean \pm standard deviation (SD)]. The latter were selected by convenience sampling, therefore, the chosen cows represent all of the available subjects (17.5% of the calving cows) that respected the following eligibility criteria at the selection-time: (i) belonging

to the cohort of dry-off cows present in the farm during recruitment (restricted period to minimize seasonal influences on cows' behaviors) and ranged between 22 to 26 d before the planned calving date (about 80 cows/month); (ii) being classified as healthy and without health problems since the previous transition period time according to the historical data (i.e., free from systemic diseases or from such affecting individual organs); (iii) having a LS ≤ 2.5 (mean value of 3 independent observers, attributed after observation of a minimum of 10 consecutive strides according to Flower and Weary, [34]); (iv) being without signs of a claw horn lesion or an infectious disease process of the foot, excluded by a functional claw trimming and a complete clinical examination of the locomotor system. Moreover, the good health status was confirmed by (v) a complete veterinary clinical examination (including temperature, respiratory rate, pulse, etc.) [35] including body condition scoring (BCS), and (vi) a blood sampling (coccygeal venipuncture) for a hemato-biochemical profile analysis [36].

Actions and timing regarding the findings based on clinical observations are reported in detail in Fig 1. Briefly, the health status was assessed at different time-intervals: (i) daily, by regular health monitoring (observation of cows' behaviour) routinely performed by the farm-staff (recording abnormalities), and by the investigators during the daily check of the devices; (ii) every other day, by complete clinical examination with particular focus on the overall general status, respiratory, gastro-intestinal, locomotor and genital systems (cows locked at the feeding rack <20min/d, after morning milking); (iii) weekly, by means of BCS [35,37], cleanliness and locomotion scores [34,38], and hemato-biochemical blood analyses (sampling by coccygeal venipuncture) performed directly in farm after collection (BHB and Glucose: FreeStyle Optium, Abbott, Chicago, Illinois, US; iCa²⁺ and blood gas analysis: i-STAT, Abbott, Chicago, Illinois, US, EG7+ cartridges) and within 1 hour (h) at the University Veterinary Teaching Hospital of the Department of Veterinary Medicine and Animal Productions of Napoli (complete blood cell count—HeCo C—Hematology, Radim Seac, Italy). Exact clinical criteria for diagnosis of the various health disorders observed are given in supplementary file—S3 Table.

Any anomalous cows' behavior or finding identified by the investigators (during the daily devices-control or the regular clinical examinations), as well as by the farm-staff was further investigated within 1h through in-depth veterinary clinical procedures. Moreover, for the enrolled animals, calving assistance was carried out by the stock-persons and the veterinary surgeons; the presence of delivery problems, potentially modifying animals' behavior (i.e. unproductive straining for at least 1 h or dystocia) were communicated to the investigators. Finally, data regarding the regular monitoring programs involving the chosen animals (e.g. reproduction, udder health, lameness, nutrition, etc.) and abnormal events were immediately digitally recorded by the farm-staff and made available to the investigators.

RumiWatch[®] devices and data handling

Cows included were mounted with both the RWh and RWp d from the 21th d before calving up to the 7th after calving (overall 28d) and received a continuous recorders' monitoring of 40'320 minutes (min). The first 7 d were considered as adaptation phase (10.080 min, from - 21d to - 15d, data not used), while the following 21d (30'240 min, from - 14d to + 7d) were considered as data acquisition phase (data were used for the analysis). Activities performed with the devices were conducted similarly for all cows, and details regarding their timing are reported in detail in Fig 1. Proper functioning of both devices was daily checked by real time monitoring of the cows' activities transmitted by Wi-Fi connection from the devices attached to the cows to the RumiWatchManager2 software downloaded to a barn-side laptop computer. Both, the RWh and RWp were equipped with an integrated micro SD Memory Card (Swissbit




Fig 1. Visual representation and timing of the different actions performed on each cow enrolled. AE = animal enrolment; EI = equipment installation; OMP = overall monitoring phase; AdP = adaptation phase; AcP = acquisition phase; DD = data download; CE = clinical examination; HMFs = health monitoring by farm-staff three time/d HBI = hemato-biochemical investigation; BCS = body condition scoring; CS = cleanliness scoring; LS = locomotion scoring; ER = equipment removal.

<https://doi.org/10.1371/journal.pone.0264834.g001>

AG, Bronschhofen, Switzerland), where raw data were continuously stored. Weekly, data recorded were transferred from both instruments to the computer by means of a USB cable (supplied by the manufacturer) and stored using RumiWatch[®] Manager 2 dedicated software (ITIN + HOCH, Liestal, Switzerland). Raw data were converted from '.RAW' filename extension (automatically generated by the software) to '.CSV' filename extension by means of the RumiWatch[®] Converter V0.7.3.6 (dedicated software, ITIN + HOCH, Liestal, Switzerland). *Eating time, ruminating time, ruminating boluses, eating chews, ruminate chews, other chews* (i.e., number of chews not attributable to any ruminating, feeding take, or drinking activity), *other activity time* (i.e., time not engaged in eating, ruminating or drinking activities), *lying time, standing time, walking time, lie down, stand up, number of strides* were the parameters identified by the algorithm and finally reported by the software, as published in previous studies [15,18]), and described in detail in the supplementary file—S2 Table.

Statistical analyses

Data were expressed as absolute numbers, percentages, or mean \pm SD. RumiWatch[®] output were converted into 1'440-min summaries (24h), and days around calving (4'320 min, d -1, d 0 and d +1) were excluded for the analyses. At the end of the clinical monitoring phase, cows were retrospectively allocated either to group-S (animals affected by ≥ 1 health disorder) or group-H (no signs of disease observed during the entire study period). The mean values of week-2 (10'080 min, wk. -2; d -14 to d -8), week -1 (8'640 min, wk. -1; d -7 to d -2) and week +1 (8'640 min; wk. +1; d +2 to d +7) relative to calving were calculated. Additionally, the day a disease was first clinically diagnosed (dd0), the day before disease diagnosis (dd-1) and two days before disease diagnosis (dd-2) were defined in group-S. The differences between dd0 and dd-1 ($\Delta D1$), dd0 and week -1 ($\Delta D2$), and between week +1 and week -1 (Δ weeks) were also calculated.

Shapiro-Wilk test was performed to check for normality distribution of the variables for each time point and calculated differences separately, and the natural logarithm was calculated for not normally distributed data. To compare between group-H and group-S within wk. -2, -1, +1, dd0, dd-1, dd-2 and for $\Delta D1$, $\Delta D2$ and Δ weeks, separately, the equal-variance T-test and Aspin-Welch unequal-variance test for normally distributed variables with equal and unequal variance, respectively were performed. Moreover, for the inter-group comparison, activities performed by cows belonging to group-S at dd0, dd-1, dd-2 were compared with mean daily activities performed by group-H during the corresponding days (for every RW parameter separately). The alpha level of significance was defined as $\alpha \leq 0.05$ for all tests; the false discovery rate was considered using the Benjamini-Hochberg procedure, to account for the testing of multiple hypotheses.

Only variables that showed significant differences in the T-test or Aspin-Welch unequal-variance test between group-H and group-S were used for further analysis to determine their usefulness in disease prediction (i.e., detecting cows with a health disorder prior to the traditional clinical diagnosis) or disease detection (i.e., at the very same day as the disease was first clinically diagnosed). Subsequently, univariate logistic regression models were employed to reduce the amount of further potential predictors. To determine the sensitivity (Se) and specificity (Sp) of the model at a given cut-off, a receiver operating characteristic (ROC) analysis was performed. Then, significant variables were combined into multivariate logistic regression models. Only variables moderately or not correlated with each other were combined in the same model (Spearman correlation coefficient > -0.7 and < 0.7). Variables were eliminated from the model by stepwise backward selection. Additionally, the ROC analyses before and after removing a variable were compared to determine how much the variable added to the sensitivity and specificity. Statistical analyses were performed with NCSS® (NCSS12 Statistical Software 2018, LLC. Kaysville, Utah, USA, [ncss.com/software/ncss](https://www.ncss.com/software/ncss)).

Results

Animals and findings based on clinical observations

Cows of both groups did not differ in age, parity, milk yield in the preceding lactation and LS around calving; a slight numerical increase in the LS from wk -2 to wk +1, for both groups, was instead observed (Table 1).

In total, clinical procedures included: $n = 3'528$ behavioral observations; $n = 559$ clinical examinations, $n = 215$ hemato-biochemical investigations, $n = 215$ body condition-, locomotion-, and cleanliness scorings, as well as $n = 43$ blood gas analyses. A total of $n = 47$ diseases were diagnosed and included: puerperal metritis ($n = 15$), subclinical hypocalcemia ($n = 7$), retained fetal membrane ($n = 6$), digital dermatitis ($n = 4$), cecal dilation ($n = 3$), tracheobronchitis ($n = 3$), ketosis ($n = 2$), sole bruise ($n = 2$), subacute ruminal acidosis ($n = 3$), interdigital hyperplasia ($n = 1$), sole ulcer ($n = 1$). In all cows, the RW devices were well tolerated, and no skin lesions were observed. All health disorders were diagnosed at wk +1, while none at wk -2 and wk -1. Therefore, at the end of the acquisition phase, $n = 24$ cows were assigned to group-S and $n = 18$ to group-H. Ten of 24 cows belonging to group-S (41.7%) were categorized as affected by one health disorder and 14 of 24 by more than one health disorders (58.3%). Overall, a least one diagnosis was made at day 1 after calving in 11 of 24 animals (45.8%).

RumiWacht® data analysis

The overall amount of data used to create new intervals-based algorithms for clinical interpretation of cows behavior originated from 14'206'920 min of continuous recording [(13

Table 1. Age, parity, milk yield and lameness score of healthy cows) and those cows diseased during week +1 relative to calving.

Variables	Group-H ^a (n = 18)				Group-S ^b (n = 24)				P-value
	Mean	SD	Median	IQR ^c	Mean	SD	Median	IQR	
Age	3.70	0.98	3.39	1.12	3.70	1.11	3.44	1.01	0.99
Parity	2.66	1.32	3.00	1.25	2.25	1.15	2.00	1.75	0.28
Milk yield ^d	11.07	1.61	11.55	1.65	10.58	1.74	10.61	1.80	0.55
LS ^e week -2	2.14	0.23	2.00	0.50	2.17	0.29	2.00	0.50	0.75
LS ^e week -1	2.17	0.24	2.00	0.50	2.20	0.25	2.00	0.50	0.59
LS ^e week +1	2.33	0.34	2.50	0.50	2.29	0.36	2.25	0.50	0.71

^aGroup-H = healthy cows during the entire study period^bGroup-S = cows diagnosed with at least one health disorder in the first week after calving^cIQR = interquartile range^dMilk yield = milk yield (kg) in the preceding lactation (*1000)^eLS = locomotion score.<https://doi.org/10.1371/journal.pone.0264834.t001>

activities*42 cows*30'340 min)-(13 activities*42 cows*4'320 min around calving)]. In particular, 8'118'240 and 6'088'680 min were obtained by RW_H and RW_P, respectively.

Regarding the behavioral intragroup differences between weeks (wks -2, -1 and +1), the overall mean values for feeding, ruminating and locomotion behaviors of both groups S and H, are given in detail in Table 2A and 2B, respectively (descriptive data).

Briefly, feeding, ruminating and locomotion behaviors did not differ between weeks -2 and -1 for both groups. In group-H, feeding and ruminating behaviors did not differ between wk -1 and wk +1. The locomotion behaviors *standing time*, *walking time*, and *strides* were

Table 2a. Mean values of variables of RumiWatch® halters and pedometers of group-H cows at week -2, -1 and +1.

Variables	Week -2 ^a		Week -1 ^b		P-value	Week -1		Week +1 ^c		P-value
	Mean	SD	Mean	SD		Mean	SD	Mean	SD	
Halter										
Other activity time (min/24 hrs)	525.5	125.2	612.3	206.5	0.15	612.3	206.5	680.5	164.5	0.28
Ruminate time (min/24 hrs)	540.6	84.8	487.8	134.7	0.18	487.8	134.7	439.3	90.9	0.22
Eat time (min/24 hrs)	365.9	78.9	330.7	102.8	0.27	330.7	102.8	311.9	106.6	0.59
Other chews (1000/24 hrs)	1.5	0.7	1.9	0.7	0.07	1.9	0.7	2.0	1.0	0.81
Ruminate chews (1000/24 hrs)	35.0	7.9	30.8	9.3	0.17	30.8	9.3	27.4	6.7	0.21
Eat chews (1000/24 hrs)	24.1	8.1	20.9	9.4	0.29	20.9	9.4	19.8	8.2	0.72
Bolus (bolus/24 hrs)	565.7	82.4	506.4	142.4	0.15	506.4	142.4	426.5	111.9	0.07
Pedometer										
Lying time (min/24 hrs)	742.5	124.0	741.0	117.0	0.97	741.0	117.0	582.1	100.2	0.0001
Standing time (min/24 hrs)	659.7	115.8	662.8	112.5	0.93	662.8	112.5	783.4	89.8	0.001
Walking time (min/24 hrs)	38.2	11.8	36.6	9.7	0.67	36.6	9.7	74.9	22.0	< 0.0001
Stand up (unit/24 hrs)	9.4	2.0	9.9	2.4	0.49	9.9	2.4	9.9	2.5	0.99
Lie down (unit/24 hrs)	9.4	1.9	10.0	2.5	0.41	10.0	2.5	9.7	2.8	0.71
Strides (unit/24 hrs)	870.3	273.2	836.1	228.8	0.68	836.1	228.8	2191.4	584.0	< 0.0001

Group-H: Healthy cows during the entire study period

^aWeek-2: From d -14 to d -8 relative to calving date^bWeek -1: From d -7 to d -2 relative to calving date^cWeek +1: From d +2 to d +7 relative to calving date.<https://doi.org/10.1371/journal.pone.0264834.t002>

Table 2b. Mean values of variables of RumiWatch® halters and pedometers group-S cows (sick cows) at week -2, -1 and +1.

Variables	Week -2 ^a		Week -1 ^b		P-value	Week -1		Week +1 ^c		P-value
	Mean	SD	Mean	SD		Mean	SD	Mean	SD	
Halter										
Other activity time (min/24 hrs)	531.6	128.4	537.3	113.4	0.88	537.3	113.4	814.5	145.9	<0.0001
Ruminate time (min/24 hrs)	534.2	98.0	524.6	72.8	0.73	524.6	72.8	383.2	105.3	0.00002
Eat time (min/24 hrs)	368.9	69.9	370.6	71.8	0.94	370.6	71.8	234.2	81.3	< 0.0001
Other chews (1000/24 hrs)	1.5	1.0	1.6	0.59	0.91	1.5	0.6	1.9	1.2	0.27
Ruminate chews (1000/24 hrs)	34.1	8.3	33.4	6.8	0.77	33.4	6.8	23.0	7.2	0.00004
Eat chews (1000/24 hrs)	23.9	7.6	23.9	7.7	0.99	23.9	7.7	13.6	6.3	0.00005
Bolus (bolus/24 hrs)	544.7	114.9	531.0	66.1	0.64	531.0	66.1	367.5	110.1	< 0.0001
Pedometer										
Lying time (min/24 hrs)	789.5	83.8	774.3	75.3	0.52	774.3	75.3	676.0	90.5	0.0002
Standing time (min/24 hrs)	612.6	79.5	627.5	70.2	0.5	627.5	70.2	700.5	82.1	0.002
Walking time (min/24 hrs)	38.4	8.5	38.6	12.6	0.95	38.6	12.6	63.8	17.4	< 0.0001
Stand up (unit/24 hrs)	9.6	2.2	10.6	3.3	0.28	10.6	3.3	11.4	2.7	0.32
Lie down (unit/24 hrs)	9.7	2.2	10.6	3.3	0.28	10.6	3.3	11.4	2.8	0.35
Strides (unit/24 hrs)	889.8	206.0	887.2	294.8	0.97	887.2	294.8	1803.3	498.1	< 0.0001

Group-S: Cows diagnosed with at least one health disorder in the first week after calving

^aWeek -1: From d -14 to d -8 relative to calving date

^bWeek -1: From d -7 to d -2 relative to calving date

^cWeek +1: From d +2 to d +7 relative to calving date.

<https://doi.org/10.1371/journal.pone.0264834.t003>

significantly higher and *lying time* significantly lower in wk +1 compared to wk -1. In group-S, *ruminating time*, *eating time*, *eating chews*, *ruminate chews*, and *ruminating boluses* were significantly lower and *other activity time* significantly higher in wk +1 compared to wk -1. The locomotion behaviors standing time, walking time, and strides were significantly higher and lying time significantly lower in wk +1 compared to wk -1.

Comparative data between groups within wks regarding feeding, ruminating and locomotion behaviors are given in Table 3.

No significant differences were detected between the two groups within wks -2 and -1. Instead, during wk +1, cows of group-S spent significantly less time *eating* and performed fewer *eating chews* compared to group-H. Moreover, sick animals spent significantly more time doing *other activity* compared to the healthy ones. Concerning the locomotion behavior, cows of group-S walked a significantly lower *number of strides* and spent significantly less time *standing* as compared to group-H. Consequently, sick cows spent significantly more time *lying* than healthy ones.

RumiWatch® vs. clinical observation based activity

Regarding behavior at dd0, dd-1 and dd-2, data are reported in detail in Table 4.

At dd-2, cows of group-S spent more time *lying* and less time *standing* as compared to cows of group-H. At dd-1, no significant differences were detected for any of the behaviors considered. At dd0, cows of group-S spent significantly less time *eating* and *ruminating* and showed significantly lower numbers of *eating chews*, *ruminate chews* and *ruminating boluses* as compared to group-H; *other activity time* was instead significantly increased. Differences between groups concerning locomotion behavior were not found at dd0.

Data regarding $\Delta D1$, $\Delta D2$ and Δ weeks are given in Table 5.

Table 3. Mean values of variables of RumiWatch® halters and pedometers of group-H and group-S at weeks -2, -1 and +1 relative to calving.

Weeks relative to the calving date															
Variables	Week -2 ^a					Week -1 ^b					Week +1 ^c				
	Group-H ^c		Group-S ^d		P-value	Group-H		Group-S		P-value	Group-H		Group-S		P-value
	Mean	SD	Mean	SD		Mean	SD	Mean	SD		Mean	SD	Mean	SD	
Halter															
Other activity time (min/24 hrs)	525.5	125.2	531.6	128.4	0.88	612.3	206.5	537.3	113.4	0.16	680.5	164.5	814.5	145.9	0.01
Ruminate time (min/24 hrs)	540.6	84.8	534.2	98.0	0.83	487.8	134.7	524.6	72.8	0.29	439.3	90.9	383.2	105.3	0.09
Eat time (min/24 hrs)	365.9	78.9	368.9	69.9	0.90	330.7	102.8	370.6	71.8	0.17	312.0	106.6	234.3	81.3	0.01
Other chews (1000/24 hrs)	1.5	0.7	1.5	1.0	0.88	1.9	0.7	1.6	0.6	0.08	2.0	1.0	1.9	1.2	0.81
Ruminate chews (1000/24 hrs)	35.0	78.8	34.1	8.3	0.76	30.8	9.3	33.4	6.8	0.33	27.4	6.7	23.0	7.2	0.06
Eat chews (1000/24 hrs)	24.1	8.1	23.9	7.6	0.93	20.9	9.4	23.9	7.7	0.28	19.8	8.2	13.7	6.3	0.01
Bolus (bolus/24 hrs)	565.7	82.4	544.7	114.9	0.54	506.4	142.4	531.0	66.1	0.49	426.5	111.9	367.5	110.1	0.11
Pedometer															
Lying time (min/24 hrs)	742.5	124.0	789.5	83.8	0.16	741.0	117.0	774.3	75.31	0.27	582.1	100.2	676.0	90.5	0.003
Standing time (min/24 hrs)	659.7	115.8	612.5	79.5	0.13	662.8	112.5	627.5	70.23	0.22	783.4	89.8	700.5	82.1	0.003
Walking time (min/24 hrs)	38.2	11.8	38.4	8.5	0.94	36.6	9.7	38.6	12.60	0.59	74.9	22.0	63.8	17.4	0.07
Stand up (unit/24 hrs)	9.4	2.0	9.6	2.2	0.72	9.9	2.4	10.6	3.28	0.49	9.9	2.5	11.4	2.7	0.07
Lie down (unit/24 hrs)	9.4	1.9	9.7	2.2	0.68	10.0	2.5	10.6	3.31	0.55	9.7	2.8	11.4	2.8	0.05
Strides (unit/24 hrs)	870.3	273.2	7.1	31.8	0.80	836.1	228.8	887.2	294.9	0.54	2191.4	584.0	1803.3	498.8	0.02

^aWeek -2: From d -14 to d -8 relative to calving date^bWeek -1: From d -7 to d -2 relative to calving date^cWeek +1: From d +2 to d +7 relative to calving date^dGroup-H: Healthy cows during the entire study period^eGroup-S: Cows diagnosed with at least one health disorder in the first week after calving.<https://doi.org/10.1371/journal.pone.0264834.t004>

Significant differences between groups for $\Delta D1$ were not found. For $\Delta D2$ and Δ weeks, cows of group-S showed significantly higher differences for *eating time*, *eating chews*, *ruminate chews* and *other activity time* compared to cows of group-H. For *ruminate time*, a significant difference between the two groups was only found for $\Delta D2$. For Δ weeks, cows of group-S had a significantly smaller difference in the *number of strides*, *walking time* and *lying time*.

Results of univariable logistic regression models at wk +1, dd-2, dd0 and for $\Delta D2$ and Δ weeks are shown in Table 6.

Although most of the univariable logistic regression models showed a significant association of the feeding, ruminating and locomotion variables with the presence of health disorders in wk +1, they had a sensitivity or specificity of less than 80%. *Eating chews* and *standing time* during wk +1 were the variables with the highest sensitivity and specificity for differentiation between groups. Indeed, the models revealed an area under the ROC-curve (AUC) of 0.72 and 0.75, a sensitivity of 70.6% and 83.3% and specificity of 85% and 69.6%, for the two variables considered. The univariable models for *eating chews*, showed an AUC of 0.73, 0.79 and 0.79 for dd0, $\Delta D2$ and Δ weeks, respectively. For the same parameter, a sensitivity of 70.6%, 76.5%, and 82.4% and a specificity of 82.4%, 68.4%, and 63.2%, respectively, was found.

The results of the multivariable logistic regression models are shown in Table 7.

For $\Delta D2$, the model considering the combination of the variables *eating chews* and *ruminate chews* revealed the highest accuracy in detecting cows with health disorders with an AUC of 0.78, a sensitivity of 63.2% and specificity of 88.2%. Adding the variable *other activity time* from the noseband sensor, the AUC and sensitivity increased up to 0.81 and 73.7%, respectively, while specificity decreased to 82.4%. For Δ weeks, the model with the highest accuracy to

Table 4. Mean values of variables of RumiWatch® halters and pedometers of group-H and group-S cows at dd0, dd-1, and dd-2.

Days relative to the first day of disease diagnosis															
Variables	dd-2 ^a					dd-1 ^b					dd0 ^c				
	Group-H ^d		Group-S ^e		P-value	Group-H ^d		Group-S ^e		P-value	H group		Group-S ^e		P-value
	Mean	SD	Mean	SD		Mean	SD	Mean	SD		Mean	SD	Mean	SD	
Halter															
Other activity time (min/24 hrs)	697.2	192.5	711.0	136.3	0.88	684.6	186.7	747.9	163.6	0.39	666.7	160.8	840.7	179.8	0.005
Ruminate time (min/24 hrs)	423.3	117.0	429.5	60.6	0.91	454.4	113.6	421.3	58.7	0.41	438.8	104.8	345.0	147.4	0.04
Eat time (min/24 hrs)	309.1	125.5	294.1	148.7	0.82	291.1	131.2	265.1	140.2	0.64	328.2	126.1	250.3	71.8	0.03
Other chews (1000/24 hrs)	2.2	1.3	1.7	1.1	0.38	2.0	1.2	2.0	1.6	0.9	1.8	1.0	2.3	1.4	0.29
Ruminate chews (1000/24 hrs)	26.6	8.7	25.9	3.7	0.87	28.4	8.3	26.1	4.8	0.45	27.4	7.2	20.7	9.4	0.02
Eat chews (1000/24 hrs)	18.9	9.2	19.2	9.5	0.96	17.8	9.4	16.2	8.8	0.67	21.1	9.3	14.1	5.3	0.01
Bolus (bolus/24 hrs)	417.3	135.9	384.8	40.1	0.6	442.7	135.9	405.6	73.0	0.44	425.5	107.4	335.4	146.3	0.04
Pedometer															
Lying time (min/24 hrs)	559.6	115.6	681.5	69.0	0.01	603.4	167.1	637.6	136.2	0.56	604.6	112.8	642.5	161.1	0.41
Standing time (min/24 hrs)	798.3	108.4	691.9	55.7	0.01	763.5	152.2	730.2	119.1	0.53	763.8	92.8	721.5	134.6	0.27
Walking time (min/24 hrs)	82.4	24.3	66.9	24.2	0.15	73.4	24.8	72.4	26.7	0.91	72.1	33.6	76.3	35.5	0.71
Stand up (unit/24 hrs)	10.3	3.3	11.1	2.0	0.51	10.5	4.2	10.8	2.0	0.84	9.8	2.8	11.2	3.5	0.16
Lie down (unit/24 hrs)	10.1	3.1	11.2	1.7	0.33	10.1	3.9	10.6	1.7	0.67	10.0	2.9	11.1	3.6	0.30
Strides (unit/24 hrs)	2374.6	732.6	1972.6	682.4	0.2	2161.4	736.0	2081.8	726.3	0.77	2116.8	959.6	2149.1	975.3	0.91

^add-2: Two days before disease diagnosis^bdd-1: One day before disease diagnosis^cdd0: The day, a disease was first clinically diagnosed^dGroup-H: Healthy cows during the entire study period^eGroup-S: Cows diagnosed with at least one health disorder in the first week after calving.<https://doi.org/10.1371/journal.pone.0264834.t005>

detect cows with one or more health disorder(s) considered variables from both the noseband sensor and the pedometer; the combination of *other activity time* and *strides* allowed to reach an AUC of 0.80, a sensitivity of 83.3% and specificity of 64.7% for the differentiation between groups S and H.

Discussion

Estimation under field conditions of the clinical usefulness of the output originating from the novel dairy farming technology can represent a stimulating challenge in cows' medicine, where the knowledge regarding behavioral changes over time is still incomplete, even more, when associated with spontaneous diseases appearance. Therefore, the goal of the present investigation was to evaluate whether the RumiWatch® output around calving may represent supporting bases for new meaningful clinical considerations if analyzed with new time intervals and compared to findings based on clinical observation.

During the entire study period, one of the main positive aspects was represented by the good tolerance of the cows to RWh and RWp. Indeed, none of the cows showed signs of lacerations or lesions due to their long-term positioning of the instruments (mainly employed for scientific purposes), even longer in place than previously described in the literature [19,39]. Although an apparently low number of animals has been enrolled, a relevant amount of data has been produced during the study both by the findings based on clinical observations and examinations and by the continuous digital monitoring based on thirteen variables/cow of two tools simultaneously used (Supplementary file—S2 Table). Despite the fact that the major part

Table 5. Mean values of variables of RumiWatch® halters and pedometers of group-H and group-S cows at ΔD1, ΔD2, and Δweeks.

Differences between time period relative to the first day of disease diagnosis															
Variables	Δweeks ^a					ΔD2 ^b					ΔD1 ^b				
	Group-H ^c		Group-S ^d		P-value	Group-H		Group-S		P-value	Group-H		Group-S		P-value
	Mean	SD	Mean	SD		Mean	SD	Mean	SD		Mean	SD	Mean	SD	
Halter															
Other activity time (min/24 hrs)	111.4	158.5	279.6	145.4	0.002	97.5	152.3	294.2	206.7	0.003	-41.7	87.8	39.0	225.5	0.23
Ruminate time (min/24 hrs)	-76.7	113.7	-148.7	108.9	0.06	-74.5	130.3	-176.5	151.5	0.04	7.9	65.7	-35.4	147.9	0.34
Eat time (min/24 hrs)	-33.7	91.2	-131.1	82.2	0.001	-20.2	102.2	-114.0	95.0	0.01	36.2	78.9	-2.5	99.0	0.29
Other chews (1000/24 hrs)	0.08	0.96	0.42	1.2	0.34	-0.1	0.91	0.73	1.5	0.06	-1.5	0.5	0.03	1.0	0.56
Ruminate chews (1000/24 hrs)	-5.2	7.6	-11.0	8.0	0.03	-5.0	8.5	-12.8	10.4	0.02	0.4	4.1	-2.6	8.8	0.27
Eat chews (1000/24 hrs)	-2.1	7.7	-10.2	6.5	0.001	-1.0	8.3	-9.9	9.4	0.01	2.9	5.6	-0.4	7.5	0.23
Bolus (bolus/24 hrs)	-109.3	130.9	-168.5	112.7	0.15	-109.4	132.3	-195.6	150.0	0.08	1.2	66.1	-27.2	150.6	0.53
Pedometer															
Lying time (min/24 hrs)	-158.9	114.3	-98.3	86.0	0.05	-135.6	139.1	-126.9	151.3	0.85	16.1	132.3	33.6	190.1	0.78
Standing time (min/24 hrs)	120.7	104.4	73.0	76.1	0.09	99.8	128.7	90.0	127.3	0.81	-12.0	131.2	-33.8	164.8	0.71
Walking time (min/24 hrs)	38.3	22.8	25.3	20.1	0.05	35.8	35.6	36.9	36.8	0.92	-4.0	25.9	0.24	33.7	0.71
Stand up (unit/24 hrs)	-0.01	2.0	0.9	3.4	0.33	-0.2	2.9	0.7	4.7	0.50	-0.6	3.2	1.1	3.8	0.23
Lie down (unit/24 hrs)	-0.3	2.5	0.8	3.5	0.23	0.02	2.8	0.6	4.7	0.66	0.02	3.1	1.3	3.2	0.33
Strides (unit/24 hrs)	1355.2	608.4	916.0	552.0	0.02	1289.8	1001.0	1242.1	1003.6	0.88	-104.7	737.7	36.8	980.3	0.67

^aΔWeeks: Differences between week +1 and week -1^bΔD2: Differences between the day of dd0 and week -1^cΔD1: Differences between the day of dd0 and dd-1^dGroup-H: Healthy cows during the entire study period^eGroup-S: Cows diagnosed with at least one health disorder in the first week after calving.<https://doi.org/10.1371/journal.pone.0264834.t006>

of previous studies concerned sensors-based validations [15,40,41], observations of physiological behaviors [20,42,43] or comparison between PDF technologies outputs and the available animals' health parameters derived from the farms' databases [29,44,45], only a few studies focused on a strict comparison between the devices' output and individual clinical-diagnostic activities [46–48], such as performed in the current investigation (Fig 1 and Supplementary file–S3 Table).

Based on clinical observations, group-S was found to be heterogeneous concerning the diseases, because several common and concomitant diseases of the peripartum period were observed. Although, a retrospective categorization, according to a single pathology was not possible, the analysis of the RumiWatch® output revealed some interesting differences between the two groups. Our findings appeared in accordance with what was described by several authors, assessing significant differences regarding ruminating and feeding behaviors, not obviously dependent on the type and severity of the diseases [19,49–51].

The comparison between intra-group weekly activities, made possible to draft interesting considerations. Although the current study did not show any intra-group differences in feeding, ruminating and locomotion behaviors for the whole population studied in wk -2 and wk -1, changes occurred from wk-1 to wk+1 (Table 2A and 2B). It was already demonstrated that *lying time* and *lying bouts*' duration significantly decrease, while *standing time* increases, in healthy cows from wk -1 to wk +1 [49,52]. Our results, comparing the locomotion activities between the same periods, additionally indicated that *walking time* and *number of strides* significantly increased in both groups (Table 2A and 2B). The resumption of milk production (associated with the rise of nutritional requirements and time spent searching for food), the

Table 6. Results of the univariable logistic regression models and receiver operating characteristics (ROC) analysis of cows being diseased using different Rumi-Watch® halter and pedometer variables and the cut-off values with highest sensitivity and specificity.

Variable	AUC ^b (95% CI)	Cut-off	Sensitivity %	Specificity %
Week +1				
Other activity time	0.71 (0.50–0.85)	705.6	85.0	47.1
Eat time	0.74 (0.52–0.87)	262.2	64.7	80.0
Eat chews	0.72 (0.49–0.86)	17729.8	70.6	85.0
Lying time	0.76 (0.57–0.87)	652.1	65.2	83.3
Standing time	0.75 (0.56–0.87)	726.0	83.3	69.6
Lie down	0.68 (0.47–0.81)	9.2	87.0	50.0
Strides	0.66 (0.45–0.80)	2260.5	38.9	91.3
dd-2				
Lying time	0.84 (0.55–0.95)	662.6	66.7	92.3
Standing time	0.82 (0.54–0.94)	719.7	84.6	66.7
dd0				
Other activity time	0.75 (0.53–0.87)	657.7	94.1	52.9
Ruminate time	0.67 (0.44–0.82)	391.2	76.5	58.8
Eat time	0.69 (0.45–0.84)	287.2	64.7	82.4
Ruminate chews	0.71 (0.49–0.85)	23303.0	82.4	52.9
Eat chews	0.73 (0.48–0.87)	17174.0	70.6	82.4
Bolus	0.68 (0.45–0.83)	432.0	58.8	76.5
ΔD2				
Other activity time	0.77 (0.57–0.89)	159.4	79.0	89.5
Ruminate time	0.68 (0.46–0.82)	-23.0	47.1	58.8
Eat time	0.79 (0.59–0.90)	-54.2	64.7	79.0
Ruminate chews	0.70 (0.48–0.83)	-8292.6	70.6	63.2
Eat chews	0.79 (0.59–0.90)	-7320.4	76.5	68.4
Δweeks				
Other activity time	0.77 (0.57–0.89)	253.1	58.0	82.4
Eat time	0.79 (0.59–0.90)	-54.2	64.7	79.0
Ruminate chews	0.70 (0.48–0.83)	-9315.1	76.5	57.9
Eat chews	0.79 (0.59–0.90)	-8062.4	82.4	63.2
Lying time	0.64 (0.43–0.79)	-142.5	73.9	50.0
Walking time	0.63 (0.43–0.78)	18.3	88.9	43.5
Strides	0.66 (0.45–0.80)	731.8	88.9	47.8

^aAUC: Area under the receiver operating characteristics curve^b CI = confidence Interval<https://doi.org/10.1371/journal.pone.0264834.t007>

necessity to walk to the milking parlor 3 times/d, as well as the regrouping of cows in other barns might explain the significant changes for these parameters observed within both groups. The authors realized how such an organized herd management system might paradoxically mitigate the early diagnostic ability of the PDF technologies. Indeed, sharing the same environment, having the same time budget to perform certain behaviors and duties, or suffering the same competition from other cows represented some hindrances that influenced cows' behaviors regardless of their state of health. For rumination and feeding activities, some intra-group differences were instead detected over time (from wk -1 to wk +1). A significant decrease in feeding (i.e., *eating time*, *eating chews*, *boluses*) and ruminating activities (i.e., *ruminating time*, *ruminate chews*, *other activity time*) was indeed recorded in group-S ([Table 2A and 2B](#)), but no

Table 7. Combination of the different Rumiwatch® noseband sensor and pedometer variables as predictors of cows being sick in multivariable logistic regression and receiver characteristics analysis on different cut-off values with corresponding sensitivity and specificity.

Variable	AUC ^a (95% CI ^b)	Cut-off	Sensitivity %	Specificity %
Week +1				
Eat Time	0.75 (0.53–0.88)	262.2	64.7	84.2
+ Lying time	0.74 (0.53–0.86)	652.1	57.9	88.2
	0.77 (0.56–0.89)	0.5	79.0	64.7
Eat Time	0.75 (0.53–0.88)	262.2	64.7	84.2
+ Lying time	0.74 (0.53–0.86)	652.1	57.9	88.2
+ Strides	0.62 (0.39–0.77)	2225.5	41.2	89.5
	0.78 (0.57–0.89)	0.5	73.7	70.6
dd0				
Eat chews	0.67 (0.44–0.82)	391.2	76.5	58.8
+ Ruminant time	0.73 (0.48–0.87)	17174.0	70.6	82.4
	0.76 (0.54–0.88)	0.4	82.4	64.7
ΔD2				
Eat chews	0.79 (0.59–0.90)	-8062.4	82.4	63.2
+ Ruminant chews	0.70 (0.48–0.83)	-9315.1	76.5	57.9
	0.78 (0.56–0.90)	0.6	63.2	88.2
Eat chews	0.79 (0.59–0.90)	-8062.4	82.4	63.2
+ Ruminant chews	0.70 (0.48–0.83)	-9315.1	76.5	57.9
+ Other activity time	0.77 (0.57–0.89)	197.0	68.4	70.6
	0.81 (0.60–0.92)	0.5	73.7	82.4
Δweeks				
Other activity time	0.79 (0.59–0.90)	159.4	83.3	58.8
+ Strides	0.62 (0.39–0.78)	731.8	88.2	38.9
	0.80 (0.59–0.91)	0.4	83.3	64.7

^aAUC: Area under the receiver operating characteristics curve

^b CI = confidence Interval.

<https://doi.org/10.1371/journal.pone.0264834.t008>

difference of eating and ruminating variables was found for group-H. Our results seem to be in line with what was described by other authors: sick cows are likely to change their feeding behavior several days before the clinical diagnosis [3,7,29], even if affected by a mild form of disease [49–51]. Based on the previous statements, changes of feeding and ruminating activities between wk. -1 and wk. +1 seem to be candidates to play a role in disease prediction.

Concerning the inter-group comparisons of the weekly activities considered, significant differences were only found for the parameter *other chews* at wk -1 (Table 3). The latter indicates chews attributable to the activity of tongue and mouth in the frame of allogrooming, self-grooming, or licking of surfaces. Miedema and collaborations [53], reporting a consistent increase of ground-licking activity before calving, proved for the first time the usefulness of this behavior as calving predictor. To the authors' knowledge, a predictive role for disease appearance was not hypothesized so far; therefore, further studies confirming the usefulness of this interesting parameter should be performed in the future. As expected, some feeding (i.e., *eating time*, *eating chews* and *other activity times*) and locomotion behaviors (i.e., *standing time*, *lying time*, *lie down*, and *number of strides*) appeared significantly different between the two groups at week +1, because of the various diseases occurring in group-S (Table 3). The assumption is also supported by the inter-group differences observed for Δweeks regarding some feeding (i.e., *eating time and chews*, *other activity time*), ruminating (i.e., *ruminant chews*)

and walking behaviors (i.e., *number of strides*, *lying time*) (Table 5). As well known [54], cows affected by distress or disease may show several abnormal behaviors and changes of general appearance (e.g., separation from the group, sluggish reaction or indifference to normal stimuli, postural changes, etc.), altering the daily interaction with the environment and consequently influencing the parameters considered in this study. This finding recently received further confirmation, as several authors demonstrated how clinical (e.g., lameness, retained fetal membranes, metritis, etc.) and subclinical diseases (e.g., subclinical ketosis and hypocalcemia, acidosis, etc.) may negatively influence several feeding [7,55], ruminating [29,51,56] and walking behaviors [19,57,58].

About the differences relative to the day of the disease diagnosis based on clinical examination, at dd0 cows of group-S showed different feeding and ruminating behaviors as compared to group-H (i.e., lower eating and ruminating time, eating chews, ruminate chews and ruminating boluses, as well as higher other activity time) (Table 4). Therefore, according to our findings, the RW seems to be the more reliable device to identify anomalous behaviors originating from one or more diseases, as compared to RWp. To the authors' knowledge, also other activity time was never considered as potential disease-predictor so far, except for an association with a non-specific discomfort status pre-calving by Fadul and collaborators [39]. Therefore, our outcomes confirm the usefulness of the parameter as effective expression of discomfort-stress in sick animals and its potential to support the in-field clinical examination. At dd-2, two RW variables already indicated an upcoming disease (both from RWp), while at dd0, six variables were significantly different (5 from RW and 1 from RWp) (Table 4). Nevertheless, the comparison of dd-1 and $\Delta D1$ between group-H and group-S did not reveal any significant difference (Tables 4 and 5). The reason for this observation may be the consequence of a substantial overlap between the three days relative to calving (-1, 0, and +1, excluded from the study) and those of the clinical diagnoses, reducing the overall number of diagnosis useful for the analysis. However, the decision was also endorsed by the farm routine, which planned to move cows from close-up to calving pen 24h before delivery. Indeed, Schirmann and collaborators [21] proved negative effects on rumination and feeding behaviors in cows regrouped immediately before calving. Therefore, based on the previous statement, the poor diagnostic performance of the parameters dd-1 and $\Delta D1$ may be mainly due to the study design rather than RW's deficiencies. Extending the period of comparison to the entire week before the day of diagnosis ($\Delta D2$), RW seems instead to be able to predict a pathological status with five of the seven parameters belonging to RW (Table 5). The results of multivariable logistic regression definitively confirm that sick cows differ in a set of behavioral variables from the healthy ones. The analysis revealed satisfying results for $\Delta D2$ (AUC = 0.81, Se = 73.7% and Sp = 82.4%, including *eat chews*, *ruminate chews* and *other activity time*) and $\Delta weeks$ (AUC = 0.80, Se = 83.3% and Sp = 64.7%, including *other activity time* and *number of strides*), supporting the usefulness of the combined use of RW and RWp for early diagnosis of clinical diseases in the first week after calving. As recently stated by Knight [12], although the ideal PDF technology still does not exist, those instruments available may actively support herdspeople to identify cows' behavioral changes, improving their well-being and the overall farm efficiency if immediate adequate measures are taken. Indeed, for example, in order to distinguish between sound and lame cows using RWp and RW, Beer and collaborators [19] described a very high level of accuracy to discriminate even slightly from non-lame cows by different variables (AUC = 0.96, Se = 100.0% and Sp = 66.7%, including *walking speed calculation*, *standing bouts* and *eating time*). The explanation of the difference to the current study might be that group-S cows were much more heterogeneous concerning the diseases involved as compared to the lameness study. Therefore, the presence of several pathologies (sometimes even concurrent) may have influenced differently the data originating from the RW sensors. According to

the findings of the current investigation, the $\Delta D2$ algorithm seems to adequately anticipate disease diagnoses based on clinical veterinary examination, and it may represent a useful tool supportive to traditional clinical considerations. However, further studies should be performed to confirm this observation.

If one hand, our data analyses open exciting new considerations regarding the clinical use of these algorithms under field conditions, on the other the study has its limits. The first is the unavoidable necessity to exclude $d - 1$, 0 , and $+1$ relative to calving. As known since a long time [59], the myometrial contractions associated with the fetal movements can significantly increase during the 24 hours before calving. The onset of these changes frequently produces signs of discomfort-mild colic, restlessness (with elevated heart and respiratory rates), as well as a fall of the body temperature, potentially influencing animals' behavior. These findings have been recently confirmed by studies employing PDF technologies and observing how this period can significantly affect rumination and feeding behaviors in dairy cows [21,39]. The second limit was instead represented by the relatively low number of disease-diagnoses and the concomitant presence of multiple diseases affecting the same animals. The current study was based on the observation of the natural behaviors and occurrence of spontaneous diseases of free stall housed adult cows around calving; therefore type, severity and timing of the disease's appearance were not predictable. Whilst it is true that the study met the necessities exposed by recent manuscripts that suggest both the assessment of the performance of PDF technologies in real situations and the development of new studies focused on the automatic diagnosis of health issues [60,61], it is nevertheless necessary to consider the present study as one of the first examples based on a retrospective clinical trial. The outputs observed should be confirmed in prospective studies, under varying different feeding and husbandry conditions with a larger number of disease events to further assess the reliability of the new algorithms for the clinical interpretation of cow behaviour under field conditions.

Conclusions

The current retrospective study offers clinical considerations concerning the usefulness of novel algorithms of digitally recorded data of feeding, ruminating and locomotion behaviors in dairy cows naturally experiencing the around calving period. The study revealed that the combined use of RWh and RWp may represent a supportive instrument for clinical interpretation of cows' behavior, showing alterations of several feeding, rumination and locomotion behaviors at the very same day as the disease was first clinically diagnosed. Moreover, the multivariable logistic regression model of this study revealed that the parameters *eating chews*, *ruminate chews* and *other activity time* achieved the highest accuracy in detecting cows with a health disorder prior to the traditional clinical diagnosis based on veterinary examinations.

The analyses revealed that some of the new algorithms for the clinical interpretation of cow behaviour used may represent a starting point for prospective studies focused on monitoring animals' health and well-being. In this regard, further studies should be performed to assess the performance of the described algorithms for the clinical interpretation of cow behaviour under field condition.

Supporting information

S1 Table. Ration formulated for close-up and fresh dairy cows by the farm.
(DOCX)

S2 Table. Definition of the activities recorded by RumiWatch® noseband sensors and 3D-accelerometers.

(DOCX)

S3 Table. Criteria for clinical diagnosis of the diseases detected in group-S during the entire study period.

(DOCX)

S1 Raw data.

(XLSX)

Acknowledgments

The Authors are very grateful to ITIN+HOCH GmbH for providing the RumiWatch® devices, to the farm and to all the farm staff involved. The Authors would like especially to thank Dr. Paolo Grendene, Dr. Nicola Malgieri and Dr. Renato Di Fonzo for their precious support.

Author Contributions

Conceptualization: Paolo Ciaramella, Adrian Steiner, Jacopo Guccione.

Data curation: Mahmoud Fadul, Luigi D'Andrea, Jacopo Guccione.

Formal analysis: Mahmoud Fadul, Luigi D'Andrea, Maher Alsaaod, Giuliano Borriello, Dimitri Stucki, Paolo Ciaramella, Jacopo Guccione.

Investigation: Luigi D'Andrea, Giuliano Borriello, Adrian Steiner, Jacopo Guccione.

Methodology: Mahmoud Fadul, Luigi D'Andrea, Maher Alsaaod, Paolo Ciaramella, Adrian Steiner, Jacopo Guccione.

Project administration: Jacopo Guccione.

Software: Mahmoud Fadul, Luigi D'Andrea, Maher Alsaaod, Dimitri Stucki, Jacopo Guccione.

Supervision: Paolo Ciaramella, Adrian Steiner, Jacopo Guccione.

Validation: Maher Alsaaod, Paolo Ciaramella, Jacopo Guccione.

Writing – original draft: Mahmoud Fadul, Luigi D'Andrea, Jacopo Guccione.

Writing – review & editing: Paolo Ciaramella, Adrian Steiner, Jacopo Guccione.

References

1. Drackley J. K. Biology of Dairy Cows During the Transition Period: the Final Frontier? J. Dairy Sci. 1999. 82:2259–2273. [https://doi.org/10.3168/jds.s0022-0302\(99\)75474-3](https://doi.org/10.3168/jds.s0022-0302(99)75474-3) PMID: 10575597
2. Roche J.R., Macdonald K.A., Schutz K.E., Matthews L.R., Verkerk G.A., Meier S., et al. Calving body condition score affects indicators of health in grazing dairy cows. J. Dairy Sci. 2013. 96:5811–5825. <https://doi.org/10.3168/jds.2013-6600> PMID: 23871378
3. Sepúlveda-Varas P., Weary D.M., and von Keyserlingk M.A.G. Lying behavior and postpartum health status in grazing dairy cows. J. Dairy Sci. 2014. 97:6334–6343. <https://doi.org/10.3168/jds.2014-8357> PMID: 25151885
4. Dubuc J., and Denis-Robichaud J. A dairy herd-level study of postpartum diseases and their association with reproductive performance and culling. J. Dairy Sci. 2017. 100:3068–3078. <https://doi.org/10.3168/jds.2016-12144> PMID: 28161186

5. Agrawal A., Khan M.J., Graugnard D.E., Vailati-Riboni M., Rodriguez-Zas S.L., Osorio J.S., et al. Prepartal Energy Intake Alters Blood Polymorphonuclear Leukocyte Transcriptome During the Peripartal Period in Holstein Cows. *Bioinform and Biol Insights*. 2017. 11:1–17. <https://doi.org/10.1177/1177932217704667> PMID: 28579762
6. Overton T.R., and Waldron M.R. Nutritional Management of Transition Dairy Cows: Strategies to Optimize Metabolic Health. *J. Dairy Sci.* 2004. 87:105–119.
7. González L., Tolkamp B.J., Coffey M.P., Ferret A., and Kyriazakis I. Changes in feeding behavior as possible indicators for the automatic monitoring of health disorders in dairy cows. *J Dairy Sci.* 2008. 91:1017–1028. <https://doi.org/10.3168/jds.2007-0530> PMID: 18292258
8. Gargiulo J. I., Eastwood C. R., Garcia S. C., and Lyons N. A. Dairy farmers with- larger herd sizes adopt more precision dairy technologies. *J. Dairy Sci.* 2018. 101:5466–5473. <https://doi.org/10.3168/jds.2017-13324> PMID: 29525319
9. Eckelkamp E.A. Invited Review: current state of wearable precision dairy technologies in disease detection**this review is based on a presentation at the ruminant nutrition symposium: management and nutrition of dairy cattle in the new era of automation at the 2018 ADSA annual meeting in Knoxville, Tennessee. *J. Appl. Anim. Sci.* 2019. 35, 209–220.
10. Caja G., Castro-Costa A., and Knight C.H. Engineering to support wellbeing of dairy animals. *J Dairy Res.* 2016. 83:136–147. <https://doi.org/10.1017/S0022029916000261> PMID: 27210489
11. Valros A., and Hänninen L. Animal Ethical Views and Perception of Animal Pain in Veterinary Students. *Anim.* 2018. 8:220–228. <https://doi.org/10.3390/ani8120220> PMID: 30477084
12. Knight CH. Review: Sensor techniques in ruminants: more than fitness trackers. *Animal.* 2020. 14: s187–s195. <https://doi.org/10.1017/S1751731119003276> PMID: 32024562
13. Braun U., Trösch L., Nydegger F., and Hässig M. Evaluation of eating and rumination behavior in cows using a noseband pressure sensor. *BMC Vet Res.* 2013. 9:164–171. <https://doi.org/10.1186/1746-6148-9-164> PMID: 23941142
14. Borchers M.R., Chang Y.M., Tsai I.C., Wadsworth B.A., and Bewley J.M. A validation of technologies monitoring dairy cow feeding, ruminating, and lying behaviors. *J Dairy Sci.* 2016. 99(9):7458–7466. <https://doi.org/10.3168/jds.2015-10843> PMID: 27423949
15. Zehner N., Umstätter C., Niederhauser J.J., and Schick M. System specification and validation of a noseband pressure sensor for measurement of ruminating and eating behavior in stable-fed cows. *Computers and Electronics in Agriculture*, 2017. 136, pp.31–41.
16. D'Andrea L., Guccione J., Alsaad M., Deiss R., Di Loria A., Steiner A., et al. Validation of a pedometer algorithm as a tool for evaluation of locomotor behavior in dairy Mediterranean buffalo. *J Dairy Res.* 2017. 84:391–394. <https://doi.org/10.1017/S0022029917000668> PMID: 29154738
17. Guccione J., D'Andrea L., Alsaad M., Borriello G., Steiner A., and Ciaramella P. Validation of a noseband pressure sensor algorithm as a tool for evaluation of feeding behavior in dairy Mediterranean buffalo (*Bubalus Bubalis*). *J Dairy Res.* 2019. 86:40–42. <https://doi.org/10.1017/S0022029919000074> PMID: 30729911
18. Alsaad M., Niederhauser J.J., Beer G., Zehner N., Schuepbach-Regula G., and Steiner A. Development and validation of a novel pedometer algorithm to quantify extended characteristics of the locomotor behavior of dairy cows. *J Dairy Res.* 2015. 98:6236–6242. <https://doi.org/10.3168/jds.2015-9657> PMID: 26142842
19. Beer G., Alsaad M., Starke A., Schuepbach-Regula G., Müller H., Kohler P., et al. Use of Extended Characteristics of Locomotion and Feeding Behavior for Automated Identification of Lamé Dairy Cows. *PLoS ONE*. 2016. 11(5):e0155796. <https://doi.org/10.1371/journal.pone.0155796> PMID: 27187073
20. Pereira G.M., Sharpe K.T., and Heins B.J. Evaluation of the RumiWatch system as a benchmark to monitor feeding and locomotion behaviors of grazing dairy cows. *J Dairy Sci.* 2021. 104:3736–3750. <https://doi.org/10.3168/jds.2020-18952> PMID: 33455761
21. Schirmann K., Weary D. M., Heuwieser W., Chapinal N., Cerri R. L. A., and von Keyserlingk M. A. G. Short communication: Rumination and feeding behaviors differ between healthy and sick dairy cows during the transition period. *J. Dairy Sci.* 2016. 99:1–8. <https://doi.org/10.3168/jds.2015-9445> PMID: 26387020
22. Ito K., von Keyserlingk M.A., Leblanc S.J., and Weary D.M. Lying behavior as an indicator of lameness in dairy cows. *J Dairy Sci.* 2010. 93:3553–3560. <https://doi.org/10.3168/jds.2009-2951> PMID: 20655423
23. Vasseur E., Rushen J., Haley D.B., and de Passillé A.M. Sampling cows to assess lying time for on-farm animal welfare assessment. *J Dairy Sci.* 2012. 95:4968–4977. <https://doi.org/10.3168/jds.2011-5176> PMID: 22916901

24. Flower F.C., Sanderson D.J., and Weary D.M. Hoof Pathologies Influence Kinematic Measures of Dairy Cow Gait. *J. Dairy Sci.* 2005. 88:3166–3173. [https://doi.org/10.3168/jds.S0022-0302\(05\)73000-9](https://doi.org/10.3168/jds.S0022-0302(05)73000-9) PMID: 16107407
25. Brandstetter V., Neubauer V., Humer E., Kröger I., and Zebeli Q. Chewing and Drinking Activity during Transition Period and Lactation in Dairy Cows Fed Partial Mixed Rations. *Anim.* 2019. 9:1088. <https://doi.org/10.3390/ani9121088> PMID: 31817555
26. Thorup V.M., Nielsen B.L., Robert P.E., Reverdin S.G., Konka J., Michie C., et al. Lameness Affects Cow Feeding But Not Rumination Behavior as Characterized from Sensor Data. *Front. Vet. Sci.* 2016. 3:37. <https://doi.org/10.3389/fvets.2016.00037> PMID: 27243025
27. Aditya S., Humer E., Pourazad P., Khiaosa-Ard R., Huber J., and Zebeli Q. Intramammary infusion of *Escherichia coli* lipopolysaccharide negatively affects feed intake, chewing, and clinical variables, but some effects are stronger in cows experiencing subacute rumen acidosis. *J. Dairy Sci.* 2017. 100:1363–1377. <https://doi.org/10.3168/jds.2016-11796> PMID: 27939552
28. Jawor P. E., Huzzey J. M., LeBlanc S. J., and von Keyserlingk M. A. G. Associations of subclinical hypocalcemia at calving with milk yield, and feeding, drinking, and standing behaviors around parturition in Holstein cows. *J. Dairy Sci.* 2012. 95:1240–1248. <https://doi.org/10.3168/jds.2011-4586> PMID: 22365207
29. Steensels M., Antler A., Bahr C., Berckmans D., Maltz E., and Halachmi I. A decision-tree model to detect post-calving diseases based on rumination, activity, milk yield, BW and voluntary visits to the milking robot. *Animal* 2016. 10:1493–1500. <https://doi.org/10.1017/S1751731116000744> PMID: 27221983
30. Vergara C. F., Döpfer D., Cook N. B., Nordlund K. V., McArt J. A. A., Nydam D. V., et al. Risk factors for postpartum problems in dairy cows: Explanatory and predictive modeling. *J. Dairy Sci.* 2014. 97:4127–4140. <https://doi.org/10.3168/jds.2012-6440> PMID: 24792805
31. Luchterhand K. M., Silva P.R., Chebel R.C., and Endres M.I. Association between Prepartum Feeding Behavior and Periparturient Health Disorders in Dairy Cows. *Front. Vet. Sci.* 2016. 3:65 <https://doi.org/10.3389/fvets.2016.00065> PMID: 27597948
32. EMA/INS/GCP/46309/2012. Classification and analysis of the GCP inspection findings of GCP inspections conducted at the request of the CHMP.
33. Sprecher D.J., Hostetler D.E., Kaneene J.B. A lameness scoring system that uses posture and gait to predict dairy cattle reproductive performance. *Theriogenology.* 1997. 47(6):1179–87. [https://doi.org/10.1016/s0093-691x\(97\)00098-8](https://doi.org/10.1016/s0093-691x(97)00098-8) PMID: 16728067
34. Flower F.C., and Weary D.M. Effect of hoof pathologies on subjective assessments of dairy cow gait. *J. Dairy Sci.* 2006. 89:139–146. [https://doi.org/10.3168/jds.S0022-0302\(06\)72077-X](https://doi.org/10.3168/jds.S0022-0302(06)72077-X) PMID: 16357276
35. Jackson P., and Cockcroft P. The General Clinical Examination of Cattle. In *Clinical Examination of Farm Animals*. 1st ed. John Wiley & Sons, New York, NY. 2002. Pages: 9–11.
36. Mulligan F.J., O'Grady L., Rice D.A., and Doherty M. A herd health approach to dairy cow nutrition and production diseases of the transition cow. *Anim Reprod Sci.* 2006. 96:331–53. <https://doi.org/10.1016/j.anireprosci.2006.08.011> PMID: 16979858
37. Ferguson J.D., Galligan D.T., and Thomsen N. Principal descriptors of body condition score in Holstein cows. *J Dairy Sci.* 1994. 77:2695–703. [https://doi.org/10.3168/jds.S0022-0302\(94\)77212-X](https://doi.org/10.3168/jds.S0022-0302(94)77212-X) PMID: 7814740
38. Guccione J., Carcasole C., Alsaaod M., D'Andrea L., Di Loria A., De Rosa A., et al. Assessment of foot health and animal welfare: clinical findings in 229 dairy Mediterranean Buffaloes (*Bubalus bubalis*) affected by foot disorders. *BMC Vet Res.* 2016. 12:107. <https://doi.org/10.1186/s12917-016-0726-4> PMID: 27297174
39. Fadul M., Bogdahn C., Alsaaod M., Hüsler J., Starke A., Steiner A., et al. Prediction of calving time in dairy cattle. *Anim Reprod Sci.* 2017. 187:37–46. <https://doi.org/10.1016/j.anireprosci.2017.10.003> PMID: 29029873
40. Strutzke S., Fiske D., Hoffmann G., Ammon C., Heuwieser W., and Amon T. Technical note: Development of a noninvasive respiration rate sensor for cattle. *J Dairy Sci.* 2019. 102:690–695. <https://doi.org/10.3168/jds.2018-14999> PMID: 30415860
41. Pereira G.M., Heins B.J., O'Brien B., McDonagh A., Lidauer L., and Kicking F. Validation of an ear tag-based accelerometer system for detecting grazing behavior of dairy cows. *J Dairy Sci.* 2020. 103:3529–3544. <https://doi.org/10.3168/jds.2019-17269> PMID: 32089298
42. Arai S., Okada H., Sawada H., Takahashi Y., Kimura K., and Itoh T. Evaluation of ruminal motility in cattle by a bolus-type wireless sensor. *J Vet Med Sci.* 2019. 81:1835–1841. <https://doi.org/10.1292/jvms.19-0487> PMID: 31685723

43. Miller G.A., Mitchell M., Barker Z.E., Giebel K., Codling E.A., Amory J.R., et al. Using animal-mounted sensor technology and machine learning to predict time-to-calving in beef and dairy cows. *Animal*. 2020. 14:1304–1312. <https://doi.org/10.1017/S1751731119003380> PMID: 31928536
44. Steeneveld W., Vernooij J.C., and Hogeveen H. Effect of sensor systems for cow management on milk production, somatic cell count, and reproduction. *J Dairy Sci*. 2015. 98:3896–905. <https://doi.org/10.3168/jds.2014-9101> PMID: 25841965
45. Post C., Rietz C., Büscher W., and Müller U. Using Sensor Data to Detect Lameness and Mastitis Treatment Events in Dairy Cows: A Comparison of Classification Models. *Sensors (Basel)*. 2020. 20:3863. <https://doi.org/10.3390/s20143863> PMID: 32664417
46. Demba S., Hoffmann G., Ammon C., and Rose-Meierhöfer S. Sensor-Based Detection of the Severity of Hyperkeratosis in the Teats of Dairy Cows. *Sensors (Basel)*. 2018. 18:3925 <https://doi.org/10.3390/s18113925> PMID: 30441773
47. Alsaad M., Fadul M., Deiss R., Bucher E., Rehage J., Guccione J., et al. Use of validated objective methods of locomotion characteristics and weight distribution for evaluating the efficacy of ketoprofen for alleviating pain in cows with limb pathologies. *PLoS One*. 2019. 14:e0218546. Erratum in: *PLoS One*. 2019 14:e0225565. <https://doi.org/10.1371/journal.pone.0218546> PMID: 31211805
48. Sturm V., Efrosinin D., Öhlschuster M., Gusterer E., Drillich M., and Iwersen M. Combination of Sensor Data and Health Monitoring for Early Detection of Subclinical Ketosis in Dairy Cows. *Sensors (Basel)*. 2020. 20:1484. <https://doi.org/10.3390/s20051484> PMID: 32182701
49. Huzzey J.M., Veira D.M., Weary D.M., and von Keyserlingk M.A. Prepartum behavior and dry matter intake identify dairy cows at risk for metritis. *J Dairy Sci*. 2007. 90:3220–3233. <https://doi.org/10.3168/jds.2006-807> PMID: 17582105
50. Soriani N., Trevisi E., and Calamari L. Relationships between rumination time, metabolic conditions, and health status in dairy cows during the transition period. *J. Anim. Sci*. 2012. 90:4544–4554. <https://doi.org/10.2527/jas.2012-5064> PMID: 23255819
51. Calamari L., Soriani N., Panella G., Petrera F., Minuti A., and Trevisi E. Rumination time around calving: an early signal to detect cows at greater risk of disease. *J Dairy Sci*. 2014. 97:3635–47. <https://doi.org/10.3168/jds.2013-7709> PMID: 24731630
52. Kaufman E.I., LeBlanc S.J., McBride B.W., Duffield T.F., and DeVries T.J. Short communication: Association of lying behavior and subclinical ketosis in transition dairy cows. *J Dairy Sci*. 2016a. 99:7473–7480.
53. Miedema H.M., Cockram M.S., Dwyer C.M., and Macrae A.I. Changes in the behaviour of dairy cows during the 24 h before normal calving compared with behaviour during late pregnancy. *Applied Animal Behaviour Science*. 2011. 131:8–14.
54. Radostits O.M., Gay C.C., Hinchcliff K.W., and Constable P.D. Clinical examination and making a diagnosis. in *Veterinary Medicine, a textbook of the diseases of cattle, horses, sheep, pigs and goats*. Tenth Edition, ed. Saunders Elsevier, Philadelphia, USA. 2007. Pages 3–38.
55. Goldhawk C., Chapinal N., Veira D.M., Weary D.M., and von Keyserlingk M.A. Prepartum feeding behavior is an early indicator of subclinical ketosis. *J Dairy Sci*. 2009. 92:4971–4977. <https://doi.org/10.3168/jds.2009-2242> PMID: 19762814
56. Stangaferro M. L., Wijma R., Quinteros C., Medrano M., Masello M., and Giordano J. Use of a rumination and activity monitoring for the identification of dairy cows with health disorders: Part II. Mastitis. *J. Dairy Sci*. 2016. 99:7411–7421. <https://doi.org/10.3168/jds.2016-10908> PMID: 27372584
57. Kaufman E.I., LeBlanc S.J., McBride B.W., Duffield T.F., and DeVries T.J. Association of rumination time with subclinical ketosis in transition dairy cows. *J. Dairy Sci*. 2016b. 99:5604–5618.
58. Itle A.J., Huzzey J.M., Weary D.M., and von Keyserlingk M.A.G. Clinical ketosis and standing behavior in transition cows. *J. Dairy Sci*. 2015. 98:128–134. <https://doi.org/10.3168/jds.2014-7932> PMID: 25465623
59. Gillette D.D., and Holm L. Prepartum and postpartum uterine and abdominal contractions in cows. *Am J Physiol*. 1963. 204:115–121.
60. Stygar A.H., Gómez Y., Berteselli G.V., Dalla Costa E., Canali E., Niemi J.K., et al. Systematic Review on Commercially Available and Validated Sensor Technologies for Welfare Assessment of Dairy Cattle. *Front Vet Sci*. 2021. 8:634338. <https://doi.org/10.3389/fvets.2021.634338> PMID: 33869317
61. Stachowicz J., Umstätter C. Do we automatically detect health- or general welfare-related issues? A framework. *Proc Biol Sci*. 2021. 288:20210190. <https://doi.org/10.1098/rspb.2021.0190> PMID: 33975474